

# Homework 7

Elliott Pryor

29 March 2021

**Problem 1** 7.6.3 Problem 2

If  $|f_n(x) - f_n(y)| \leq M|x - y|^\alpha$  for some fixed  $M$  and  $\alpha > 0$  and all  $x, y$  in a compact interval. Show that  $\{f_n\}$  is uniformly equicontinuous

---

PROOF.

By definition of uniform equicontinuity we need to show:  $\forall 1/m \exists 1/n \text{ st } |x - y| < 1/n \implies |f_k(x) - f_k(y)| < 1/m \forall k$ . We have from the definition of  $f_n$  that  $|f_n(x) - f_n(y)| \leq M|x - y|^\alpha$ . So  $|f_k(x) - f_k(y)| \leq M|x - y|^\alpha \leq M(1/n)^\alpha$  which we need to be  $< 1/m$ . So we choose  $1/n < \sqrt[\alpha]{\frac{1}{Mm}}$  since  $M, \alpha$  are defined and  $\alpha > 0$  we can compute this for any  $1/m$  as required.  $\square$

**Problem 2** 7.6.3 Problem 5

Give an example of a sequence that is uniformly equicontinuous but not uniformly bounded

---

$f_n = x + n$ . Is uniformly equicontinuous since can choose  $n = m$ . But is not uniformly bounded since  $f_m \rightarrow \infty$  as  $n \rightarrow \infty$ .

**Problem 3** 7.6.3 Problem 6

Prove that the family of all polynomials of degree  $\leq N$  with coefficients in  $[-1, 1]$  is uniformly bounded and uniformly equicontinuous on any compact interval.

**Hint:** Let the compact interval be  $[a, b]$ , and each polynomial in the family has the form

$$p(x) = c_0 + c_1x + c_2x^2 + \cdots + c_Nx^N \quad x \in [a, b], \quad c_i \in [-1, 1], \quad 0 \leq i \leq N$$

To prove uniform equicontinuity, show that the derivatives of all polynomials in the family are uniformly bounded.

---

PROOF.

Consider the derivatives of the family of functions:  $p'(x) = c_1 + 2c_2x + 3c_3x^2 + \cdots + Nc_Nx^{N-1}$ . Then let  $g$  be the polynomial with all coefficients = 1:  $g(x) = 1 + x + x^2 + \cdots + x^N$ , and  $g'(x) = 1 + 2x + 3x^2 + \cdots + Nx^{N-1}$ .

Then clearly outside of  $(-1, 1)$   $p \leq g$  and  $p' \leq g'$  for any  $p$  in the family. And inside  $(-1, 1)$ ,  $p'(x) \leq \sum_{i=0}^n i$  since for any term  $i \cdot c_i x^{i-1} \leq i \cdot x^{i-1} \leq i$  inside  $x \in (-1, 1)$ . We call this summation  $e = \sum_{i=0}^n i$ . And let  $f = \sup_{x \in [a, b]} g'(x)$  which we know exists since  $g'$  is continuous.

Then let  $M = \max(e, f)$ . So then clearly  $p'$  is uniformly bounded by  $M$ .

Then copied from the proof of Corollary 7.6.1: by MVT  $\forall x, y \in [a, b] \quad \forall k \quad |f_k(x) - f_k(y)| = |f'_k(z)(x - y)| \leq |f'_k(z)||x - y| \leq M|x - y| < 1/m$  by choosing  $1/n > \frac{1}{mM}$ . Thus all  $p$  are equicontinuous  $\square$

**Problem 4** 7.6.3 Problem 9

Give an example of a uniformly bounded and uniformly equicontinuous sequence of functions on the whole line that does not have any uniformly convergent subsequences.

**Hint:** Consider the following sequence of functions on  $\mathbb{R}$

$$f_n(x) = \begin{cases} 0 & x \leq n-1 \\ x - (n-1) & n-1 < x \leq n \\ 1 & x > n \end{cases}$$

---