

Homework 1

Elliott Pryor

31 Jan 2021

Problem 1 6.2.4 Problem 6

Prove that if f is Riemann integrable on $[a, b]$ and $g(x) = f(x)$ for every x except for a finite number, then g is Riemann integrable.

Hint: Mimic the proof of Theorem 6.2.3

PROOF.

Let a_1, a_2, \dots, a_N denote the points where $g(a_i) \neq f(a_i)$. Given any $1/n$ surround each a_k by an interval I_k such that $|I_k| < 1/n$. Then, g is continuous and Riemann integrable on $[a, b]$ with $\cup_{k=1}^N I_k$ removed.

Then, we estimate the contribution to the oscillation from the I_k intervals. First, f must be bounded since it is Riemann-integrable. Then since g differs from f at a finite number of locations, it must also be bounded. Then let $M = \sup_{x \in [a, b]} g(x)$ and $m = \inf_{x \in [a, b]} g(x)$. Then the contribution of any of the intervals: I_k is at most $1/n \cdot (M - m)$. There are N such intervals, so the total contribution to oscillation is $N/n \cdot (M - m)$.

Then for the remaining intervals: $[a, b] \setminus \cup_{k=1}^N I_k$, $g = f$. Since f is Riemann-integrable, the oscillation on these intervals can be made sufficiently small by choosing the partition size sufficiently small. In other words, the total oscillation on a partition P' of $[a, b] \setminus \cup_{k=1}^N I_k$, $g = f$ can be, $\forall 1/n \exists 1/m$ st $Osc(g, P') < 1/n$ for $|P'| < 1/m$ (part b of Theorem 6.2.1)

Then the total oscillation $Osc(g, P) < 1/n + N/n \cdot (M - m)$. Since N, M, m are constant, there exists a sequence of partitions such that $Osc(g, P_j) \rightarrow 0$ as $j \rightarrow \infty$ by selecting n large enough.

□

Problem 2 6.2.4 Problem 9, part b

If f is Riemann integrable on $[a, b]$, prove $F(x) = \int_a^x f(t)dt$ satisfies a Lipschitz condition

PROOF. We need to show that $|F(x) - F(x_0)| \leq M|x - x_0|$ for some M .

Without loss of generality assume $x > x_0$. Then by the definition of F . We have

$$|F(x) - F(x_0)| = \int_a^x f(t)dt - \int_a^{x_0} f(t)dt = \int_{x_0}^x f(t)dt$$

We say $M = \sup_{x \in [a, b]} f(x)$. Then

$$|F(x) - F(x_0)| = \int_{x_0}^x f(t)dt \leq M|x - x_0|$$

as required. □

Problem 3 6.2.4 Problem 10

If f is Riemann integrable on $[a, b]$ and continuous at x_0 , prove that $F(x) = \int_a^x f(t)dt$ is differentiable at x_0 and $F'(x_0) = f(x_0)$. Show that if f has a jump discontinuity at x_0 , then F is not differentiable at x_0 .

Hint: Refer to the proof of Theorem 6.1.2, note that $f(x)$ being continuous at x_0 can be written as $\forall 1/m, \exists 1/n$ st. $\forall x \in [a, b], |x - x_0| < 1/n$, we have $|f(x) - f(x_0)| < 1/m$ or $f(x_0) - 1/m < f(x) < f(x_0) + 1/m$
