

Homework # 5

Elliott Pryor

9/18/2020

Problem 1

Statement

3.121 - (clearly show steps to achieve a written probability statement, then write the code you would use to find the probability in R/find probability using R)

Let Y denote a random variable that has a Poisson distribution with mean $\lambda = 2$. Find

1. $P(Y = 4)$
 2. $P(Y \geq 4)$
 3. $P(Y < 4)$
 4. $P(Y \geq 4|Y \geq 2)$
-

Solution

$$f_Y(y) = \frac{\lambda^y e^{-\lambda}}{y!}$$

```
poisson_pmf <-function(y) {# generate a sequence of 100 independent Bernoulli trials
  p <- ((2^y) * exp(-2))/(factorial(y))
  return(p)}
```

1. $f_Y(4) = \frac{2^4 e^{-2}}{4!} = \frac{16e^{-2}}{24} = 2/3e^{-2}$

```
2/3 * exp(-2)
```

```
## [1] 0.09022352
```

2. This is the probability of $P(Y \geq 4) = 1 - P(Y < 4) = 1 - (P(Y = 0) + P(Y = 1) + P(Y = 2) + P(Y = 3))$

```
1 - (poisson_pmf(0) + poisson_pmf(1) + poisson_pmf(2) + poisson_pmf(3))
```

```
## [1] 0.1428765
```

3. $P(Y < 4) = P(Y = 0) + P(Y = 1) + P(Y = 2) + P(Y = 3)$

```
(poisson_pmf(0) + poisson_pmf(1) + poisson_pmf(2) + poisson_pmf(3))
```

```
## [1] 0.8571235
```

4. This is very similar to problem 2, but since we know that $Y \geq 2$ we can ignore the $P(Y = 0)$ and $P(Y = 1)$ terms. So we get $P(Y \geq 4|Y \geq 2) = 1 - (P(Y = 2) + P(Y = 3))$

```
1 - (poisson_pmf(2) + poisson_pmf(3))
```

```
## [1] 0.5488824
```

Problem 2

Statement

3.128

Cars arrive at a toll both according to a Poisson process with mean 80 cars per hour. If the attendant makes a one-minute phone call, what is the probability that at least 1 car arrives during the call?

Solution

This can be represented as $1 - P(\text{no cars arrive at the booth}) = 1 - f_Y(0)$. We must scale the unit effort from hour to minute, so $\lambda' = \lambda/60 = 4/3$. Following the Poisson pmf distribution $f_Y(y) = \frac{\lambda^y e^{-\lambda}}{y!}$ results in:

$$1 - \frac{\lambda^0 e^{-\lambda}}{0!} = 1 - e^{-\lambda} = 1 - e^{-4/3} = 0.7364$$

Problem 3

Statement

First, run the example R code provided, then answer the following question. On the same plotting region, plot the distribution of $Y \sim \text{Binomial}(n, p)$ and $W \sim \text{Poisson}(np)$ for $p = 0.1$ and varying values of n . Create a maximum of 3 plots and explain what you notice about Y and W as $n \rightarrow \infty$

Solution

```
# set the largest y for which you want to plot P(Y = y)
max_y <- 50
y <- 0:max_y

1. Binomial distribution


v1 <- 100
v2 <- 50
v3 <- 200

plot(y, dbinom(y, v1, 0.1), type="h",
     xlim=c(0, max_y), ylim=c(0,0.2),
     xlab = "y", ylab = "p(Y = y)",
     main = "Y ~ Binom(y, p)",
     col = "darkgreen")
# add points and lines for the distribution of Y if Y ~ Geom(p) for different p
points(y, dbinom(y, v1, 0.1), pch = 20, col = "darkgreen")

lines(y, dbinom(y, v2, 0.1), type = "h", col = "magenta")
points(y, dbinom(y, v2, 0.1), pch = 20, col = "magenta")

lines(y, dbinom(y, v3, 0.1), type = "h", col = "blue")
points(y, dbinom(y, v3, 0.1), pch = 20, col = "blue")
# add a legend
legend("topright", legend = c(paste0("n= ", v1), paste0("n=", v2), paste0("n=", v3)),
     col = c("darkgreen", "magenta", "blue"),
     pch = c(20,20,20), bty = "n")
```

$Y \sim \text{Binom}(y, p)$



```
# set the largest y for which you want to plot P(Y = y)
max_y <- 50
y <- 0:max_y
```


2. Poisson distribution

```
v1 <- 100
v2 <- 50
v3 <- 200

plot(y, dpois(y, v1 * 0.1), type="h",
     xlim=c(0, max_y), ylim=c(0,0.2),
     xlab = "y", ylab = "p(Y = y)",
     main = "Y ~ Poisson(np)",
     col = "darkgreen")
# add points and lines for the distribution of Y if Y ~ Geom(p) for different p
points(y, dpois(y, v1 * 0.1), pch = 20, col = "darkgreen")

lines(y, dpois(y, v2 * 0.1), type = "h", col = "magenta")
points(y, dpois(y, v2 * 0.1), pch = 20, col = "magenta")

lines(y, dpois(y, v3 * 0.1), type = "h", col = "blue")
points(y, dpois(y, v3 * 0.1), pch = 20, col = "blue")
# add a legend
legend("topright", legend = c(paste0("n= ", v1), paste0("n=", v2), paste0("n=", v3)),
     col = c("darkgreen", "magenta", "blue"),
     pch = c(20,20,20), bty = "n")
```



Extra Credit

Statement

3.85

Find $E(Y(Y-1))$ for a geometric random variable Y by finding $\frac{d^2}{dq^2}(\sum_{y=1}^{\infty} q^y)$. Use this result to find the variance of Y .

Solution