

Praxia Update

Daily orthopaedic literature digest · Saturday, 08 August 2026 · 488 new articles across 10 journals



American Journal of Sports Medicine

Volume 54, Issue 10 · August 2026 · 28 new articles

The Rise of Remplissage.

Owens BD

[PubMed](#) · [DOI](#)

Risk of Revision and Patient-Reported Outcomes After ACL Reconstruction: Influence of Concomitant MCL Injury and Graft Choice: Analysis of 35,139 Reconstructions From the Norwegian Knee Ligament Register.

Birkenes T, Chahla J, Lygre SHL, Williams A, Inderhaug E

BACKGROUND: Concomitant medial collateral ligament (MCL) injury is common in anterior cruciate ligament (ACL) tears, but optimal management of these injury components remains controversial.

PURPOSE: To compare patient-reported outcome measures and the risk of ACL revision surgery in patients with concomitant MCL injuries (treated operatively and nonoperatively) versus isolated ACL injuries and to assess whether MCL treatment strategy or ACL graft choice influenced outcomes.

STUDY DESIGN: Cohort study; Level of evidence 3.

METHODS: Data were obtained from the Norwegian Knee Ligament Register. Patients undergoing primary ACL reconstruction (ACLR) between 2004 and 2024 were included for graft-survival analyses, and those operated by December 31, 2022, were eligible for 2-year Knee injury and Osteoarthritis Outcome Score (KOOS) outcomes. Patients with concomitant ligament injuries other than to the MCL, ACL repairs, or those with unknown ACL graft type were excluded. Patients were stratified into isolated ACL and ACL + MCL injury groups. Kaplan-Meier survival analyses (unadjusted), multivariable Cox regression, and linear/logistic regression models were used, adjusted for descriptive data, body mass index, meniscal/cartilage injury, time to surgery, smoking, pivoting sports, graft choice, and baseline KOOS.

RESULTS: A total of 35,139 ACLRs were included (2410 ACL + MCL and 32,7 [...])

[PubMed](#) · [DOI](#)

Short-term Patient-Reported Outcomes After Primary Anterior Cruciate Ligament Reconstruction With and Without Lateral Extra-articular Tenodesis: A Matched-Cohort Analysis From the Swedish Knee Ligament Registry.

Pruneski JA, Zsidai B, Öttl F, Heder Ternell K, Cristiani R, Musahl V, Senorski EH, Samuelsson K et al.

BACKGROUND: Despite reduced graft failure rates reported in randomized controlled trials comparing anterior cruciate ligament reconstruction (ACL-R) with and without lateral extra-articular tenodesis (LET), the impact of LET on postoperative patient-reported outcomes (PROs) remains poorly defined.

PURPOSE: To compare short-term PROs, as well as anterior cruciate ligament (ACL) revision rates and clinical failure rates (defined as Knee injury and Osteoarthritis Outcome Score [KOOS] Quality of Life [QoL] value <44) between patients undergoing primary ACL-R with and without concomitant LET.

STUDY DESIGN: Cohort study; Level of evidence, 3.

METHODS: Data were extracted from the Swedish Knee Ligament Registry for all patients undergoing primary ACL-R between January 1, 2005, and June 25, 2025. Patients who underwent ACL-R + LET were matched to patients who underwent ACL-R without LET using 1:4 matching based on age, sex, body mass index, graft type,

time from injury to reconstruction, and concomitant meniscal and cartilage injury status. The primary outcomes were KOOS values at the 1- and 2-year follow-ups. Secondary outcomes included revision ACL-R and clinical failure within 1 and 2 years. Categorical variables were compared using Fisher exact or chi-square tests, continuous variables were compared using Fisher nonparametric permutation tests, and within-group KOOS changes were [...]

[PubMed](#) · [DOI](#)

ACL Reconstruction With Attachment-Sparing Hamstring Autograft Results in Earlier Graft Maturation and Better Short-Term Clinical Outcome in Comparison to Free Graft: A Randomized Controlled Trial.

Sinha S, Arumugham KC, Singh DK, Kumar N, Bera AP, Uppal H, Vakharia KS

BACKGROUND: Attachment-sparing anterior cruciate ligament reconstruction (ACLR) is gaining acceptance over the use of free grafts because of reported early graft ligamentization in both animals and humans. However, the literature regarding clinical outcomes and the correlation of early graft healing is unclear.

HYPOTHESIS: Tibial attachment-sparing hamstring graft (AS) results in earlier graft maturation, which correlates with a better clinical outcome than a free hamstring graft (FG).

STUDY DESIGN: Randomized controlled trial; Level of evidence, 1.

METHODS: A total of 56 patients were randomly assigned to 2 groups at the commencement of the study. Of these patients, 25 (AS group) underwent ACLR using the attachment-sparing technique, while 25 patients (FG group) underwent ACLR using the free hamstring graft technique. Prospective sequential magnetic resonance imaging (MRI) evaluations were performed at 3, 6, and 9 months to document graft maturation, characteristics, orientation, and tunnel dilation. Clinical evaluation was done at 3, 6, 9, and 12 months to document Lysholm, International Knee Documentation Committee (IKDC), and Tegner scores and anterior tibial translation (ATT) by KT-2000 arthrometer, by independent observers.

RESULTS: The overall follow-up rate was 78% (19 cases and 20 controls of 25 in each group). MRI Figueroa scores demonstrated significant difference [...]

[PubMed](#) · [DOI](#)

MRI-Based Volumetric Analysis of the Quadriceps Muscles Correlation With Knee Extension Strength in Patients Undergoing Anterior Cruciate Ligament Reconstruction.

DeFoor MT, Knurr KA, Heiderscheit B, Riem L, Cognetti DJ, Shaw J, Blaylock M, Antosh IJ et al.

BACKGROUND: Preoperative quadriceps and hamstring muscle weakness may predispose patients to strength deficits and prolonged recovery in the setting of anterior cruciate ligament (ACL) injury. Three-dimensional (3D) magnetic resonance imaging (MRI) analysis has been developed to provide a comprehensive assessment of the thigh musculature.

PURPOSE: To (1) describe the magnitude of quadriceps and hamstring muscle weakness relative to the uninjured limb preoperatively in patients with known ACL tears, and (2) determine if preoperative quadriceps and hamstring muscle volume correlate with quadriceps knee extension strength and hamstring knee flexion strength.

STUDY DESIGN: Cross-sectional study; Level of evidence, 3.

METHODS: Patients undergoing primary ACL reconstruction (ACLR) underwent MRI scans of the bilateral lower extremities and strength testing preoperatively within 7 days of ACLR. Three-dimensional MRI-based autosegmentation was used to measure the volume of individual quadriceps and hamstring muscles. Preoperative knee extension and flexion strength were assessed isometrically at 90° and isokinetically. Linear mixed-effects models were used to assess muscle volume by limb interaction effect on strength. Pearson correlation coefficients (r) were used to assess the association between muscle volume and strength in the injured and uninjured limbs.

RESULTS: In total, 72 [...]

[PubMed](#) · [DOI](#)

Knee Synovial Fluid Biomarker Type and Concentration With Long-Term Outcomes After ACL Reconstruction With Concomitant Meniscal Injury.

Ehlers M, Estes LC, Derry K, Kaplan DJ, Strauss EJ

BACKGROUND: Anterior cruciate ligament (ACL) rupture frequently occurs alongside meniscal injury, together altering the intra-articular environment. While ACL injury triggers an inflammatory cascade within synovial fluid, meniscal injury may intensify these responses, contributing to joint degeneration. However, little is known about how combined ACL and meniscal injuries shape the synovial inflammatory milieu or whether these profiles influence long-term outcomes.

PURPOSE/HYPOTHESIS: The purpose was to characterize synovial fluid inflammatory phenotypes at the time of ACL reconstruction with concomitant meniscal injury and assess their association with patient-reported outcome measures (PROMs) and clinically meaningful improvement. It was hypothesized that (1) distinct inflammatory phenotypes would exist, (2) these phenotypes would correlate with symptom severity and recovery, and (3) individual biomarkers' effects on outcomes would vary by phenotype.

STUDY DESIGN: Cohort study; Level of evidence, 3.

METHODS: Patients undergoing arthroscopic ACL reconstruction with concomitant meniscal injury were prospectively enrolled between July 2011 and January 2024. Synovial fluid was aspirated before incision, and the concentrations of 10 biomarkers were quantified. Unsupervised k-means clustering identified high- and low-inflammation phenotypes. PROMs were collected preoperatively o [...]

[PubMed](#) · [DOI](#)

Quantitative Investigation of Tibiofemoral Cartilage Early Degeneration After ACL Reconstruction: An Integrated Biomechanical and Longitudinal MRI Analysis.

Lin J, Cheng R, Yan Y, Zeng X, Huang W, Tsai TY, Deng C, Wang S et al.

BACKGROUND: Anterior cruciate ligament reconstruction (ACLR) restores stability but is often followed by early cartilage degeneration. The contribution of altered dynamic contact pressure during gait to this degeneration remains poorly understood.

PURPOSE: To investigate the biomechanical mechanisms underlying early cartilage degeneration after ACLR, with a focus on dynamic contact pressure distribution during gait.

STUDY DESIGN: Controlled laboratory study.

METHODS: In a 3-year longitudinal magnetic resonance imaging (MRI) study of patients with ACLR (n = 30), cartilage thickness of the tibial plateau was quantitatively assessed using deep-learning-based 3-dimensional segmentation techniques. In parallel, cadaveric knees (n = 8) were tested under intact, ACL-deficient, and ACL-reconstructed conditions using a 6 degree-of-freedom robotic simulator replicating gait cycles. Tibiofemoral contact pressure and pressure center trajectories were recorded using pressure-sensitive film.

RESULTS: Quantitative MRI analyses revealed significant cartilage thinning in the posterior tibial subregion 3 years after ACLR. On the medial plateau, the central medial tibia, internal medial tibia, and posterior medial tibia subregions exhibited mean reductions of 10% (P = .027), 21% (P = .019), and 13% (P = .041), respectively. On the lateral plateau, significant decreases were observed in centra [...]

[PubMed](#) · [DOI](#)

Comparable Midterm Survival After Medial Meniscus Root Repair for High and Low Intraoperative Outerbridge Classification Grades.

Grandberg C, Boduch AN, Bilodeau RE, Cohen D, Herman ZJ, Ahrendt GM, Drain NP, Setliff JC et al.

BACKGROUND: While medial meniscus posterior root repairs (MMPRRs) have been shown to reverse harmful biomechanical effects of medial meniscus posterior root tears, it is questionable whether meaningful improvement occurs in patients with advanced chondral loss.

PURPOSE/HYPOTHESIS: The objectives of this study were to compare midterm survival of MMPRR and to determine patient-reported outcomes (PROs) after MMPRR based on intraoperative Outerbridge grades (0-2 vs 3 and 4). It was hypothesized that patients undergoing MMPRR with Outerbridge grades 3 and 4 would have a decreased rate of MMPRR survival and inferior PROs compared to patients with Outerbridge grades 0 to 2.

STUDY DESIGN: Cohort study; Level of evidence, 3.

METHODS: Consecutive patients who underwent primary isolated MMPRR performed at a single institution between July 2003 and November 2022 with a minimum follow-up of 1 year were included and categorized according to medial compartment Outerbridge grade (0-2 or 3 and 4) based on diagnostic arthroscopy. Failure was defined as reoperation of the index knee. PROs collected included International Knee Documentation Committee (IKDC), visual analog scale (VAS) for pain, and Marx activity scale scores, which were compared to the previously established minimal clinically important difference (MCID) and/or clinically meaningful change. A minimum follow-up of 2 years was use [...]

[PubMed](#) · [DOI](#)

The Effect of Smoking History on Inflammatory Biomarkers in Synovial Fluid of Patients Undergoing Arthroscopic Knee Surgery for Meniscal Injury.

Kurtz JL, Ehlers M, Montgomery SR, Kaplan DJ, Strauss EJ

BACKGROUND: Smoking has been linked to alterations in cellular inflammatory pathways and adverse outcomes in orthopaedic procedures. It is unclear how patient smoking affects the intra-articular microenvironment in the setting of symptomatic meniscal tears.

PURPOSE: To investigate the association between tobacco use and variation in cytokine concentrations in knee synovial fluid in patients undergoing arthroscopic knee surgery for symptomatic meniscal injury.

STUDY DESIGN: Cohort study; Level of evidence, 3
Methods: Patients who underwent knee arthroscopy for meniscal injury were prospectively enrolled between July 2011 and June 2019. Synovial fluid was aspirated and the concentrations of 10 biomarkers were measured by immunoassay. Smoking status, pack-years, and time since smoking cessation were collected. Log-normalized biomarker concentrations and smoking status were analyzed using analysis of variance (ANOVA) and linear regression.

RESULTS: A total of 297 patients (mean age, 45.7 ± 12.5 years; mean body mass index, 28.3 ± 5.3 kg/m²; 55.6% male) were included. Patients were divided into current smokers (n = 27), former smokers (n = 54), and nonsmokers (n = 216). No descriptive differences between groups were found. ANOVA showed significant differences (P < .01) in regulated on activation, normal T cell expressed and secreted (RANTES), monocyte chemotactic protein-1 (MCP-1), [...]

[PubMed](#) · [DOI](#)

A Prospective Clinical and Magnetic Resonance Imaging-Based Follow-up Beyond 10 Years After Arthroscopic Matrix-Induced Autologous Chondrocyte Implantation.

Ebert JR, Edwards PK, Klinken S, Wood DJ, Janes GC

BACKGROUND: Long-term studies are emerging reporting encouraging outcomes after matrix-induced autologous chondrocyte implantation (MACI). While MACI has historically been undertaken via open arthrotomy, arthroscopic techniques have been reported. However, longer term clinical outcomes and/or graft survivorship after arthroscopically performed MACI are yet to be reported.

PURPOSE: To present longer term clinical and radiological outcomes in patients after MACI performed arthroscopically on the femoral condyles.

STUDY DESIGN: Case series; Level of evidence, 4.

METHODS: A total of 45 patients were prospectively recruited and underwent arthroscopically performed MACI to the medial femoral condyle (MFC; n = 31; defect size, 1.0-7.7 cm²) or lateral femoral condyle (LFC; n = 14; defect size, 1.5-5.0 cm²). Outcomes were collected before surgery and at 1, 2, and 5 years as well as at a mean of 13.6 years (range, 11-17 years) after surgery. These included scores for a variety of patient-reported outcome measures (PROMs) including the Knee Injury and Osteoarthritis Outcome Score, Lysholm Knee Score, Tegner Activity Scale, and patient satisfaction. Active knee range of motion as well as peak isokinetic knee extensor and flexor strength were assessed, while magnetic resonance imaging (MRI) was undertaken. The Magnetic Resonance Observation of Cartilage Repair Tissue (MOCART) scoring too [...]

[PubMed](#) · [DOI](#)

A Novel Autologous Osteoperiosteal Composite for Osteochondral Regeneration.

Yao L, Zhu S, Wu Y, Ye G, Wang G, Niu Z, Ding S, Xiang Z et al.

BACKGROUND: Osteochondral defects result in persistent knee pain and functional impairment, and remain a clinical challenge to repair effectively.

PURPOSE: To evaluate the efficacy of an autologous inverted subchondral bone-periosteum composite graft for osteochondral reconstruction.

STUDY DESIGN: Controlled laboratory study.

METHODS: A total of 30 mature New Zealand White rabbits were randomly assigned to 3 groups: microfracture (MF) (n = 10), osteoperiosteal composite graft (OPC) (n = 10), and untreated control (n = 10). A 2 mm-diameter osteochondral defect was created in the control and MF groups and left untreated in the control group. In the MF group, microfracture was then performed by creating four 4 mm-deep perforations in the defect bed using Kirschner wires. A cylindrical osteochondral autograft (2 mm diameter × 4 mm height) was aseptically harvested from the intercondylar fossa using a calibrated coring drill only in the OPC group. Group-specific treatments included defect creation without reparative intervention (control), defect creation followed by subchondral perforation (MF), and inverted autograft implantation with periosteal wrapping (OPC). Animals were euthanized at 6 and 12 weeks postoperatively (n = 5 per group per time point) for gross and histological evaluation.

RESULTS: At 6 weeks, fibrocartilaginous tissue partially filled the defects in all groups [...]

[PubMed](#) · [DOI](#)

Weakening of the Active Muscle Restraint Against Valgus Loading During Repeated Baseball Pitching: Direct Evidence of Fatigue in Dynamic Valgus Stabilizers of the Elbow Joint in Collegiate Baseball Pitchers.

Yanai T, Onuma K

BACKGROUND: Although research has shown the effect of repeated pitching on throwing arm strength, no studies have examined the weakening in the overall ability of dynamic stabilizers to restrain the medial elbow against valgus loading. Fatigue in the dynamic stabilizers increases reliance on the ulnar collateral ligament (UCL) to withstand the valgus loads during pitching.

HYPOTHESIS: It was hypothesized that (1) the muscular varus strength weakens as the pitch count accumulates and (2) the muscular varus strength becomes insufficient to overcome the valgus loads in later innings.

STUDY DESIGN: Descriptive laboratory study.

METHODS: A total of 28 collegiate baseball pitchers threw 7 innings of 15 pitches in a bullpen. Seven genlocked cameras were used to record pitching performances, and the elbow valgus load was determined. Muscular varus strength was measured before the first inning and after completing the fourth and seventh innings using an isokinetic dynamometer system and an ultrasound device. A mixed-effect model was used to examine changes over the innings in the valgus load and varus strength, and t tests were used to compare the varus strengths and the largest valgus load in corresponding innings.

RESULTS: Neither the largest valgus load of the inning nor the mean valgus load of the inning significantly changed across the innings, whereas the varus strength was re [...]

[PubMed](#) · [DOI](#)

Changes in Rate and Indications for Remplissage When Performing Arthroscopic Bankart Repair for Anterior Shoulder Instability.

Brej BL, Rauck RC, Huntley KS, David T, Denard PJ, Lin A, Bishop JY

BACKGROUND: Remplissage is used as an adjunct to arthroscopic Bankart repair (ABR) to reduce recurrent instability. Despite growing evidence supporting efficacy and broader application of the procedure, the absence of a specific procedural code limits knowledge of its utilization, and current surgical indications are not well defined.

HYPOTHESIS: It was hypothesized that remplissage utilization would increase over the study period and that indications would expand beyond traditionally defined off-track Hill-Sachs lesions.

STUDY DESIGN: Cohort study; Level of evidence, 3.

METHODS: Patients undergoing arthroscopic anterior shoulder stabilization from 2012 to 2024 were retrospectively reviewed across multiple institutions. Patients were grouped by ABR alone (ABR) or with remplissage (REMP). Temporal trends were evaluated by comparing early (2012-2020) and late (2021-2024) cohorts. Preoperative magnetic resonance imaging was used to assess bipolar bone loss using established measurement techniques, including glenoid track (GT), glenoid bone loss (GBL), and distance to dislocation (DTD).

RESULTS: The cohort included 244 patients who underwent remplissage and 1100 patients who underwent ABR. The proportion of REMP increased 4.5-fold over the study period (7.7% in 2012 vs 34.7% in 2024). Later year of surgery was significantly associated with increased remplissage utilization (odd [...]

[PubMed](#) · [DOI](#)

Morphological Characteristics After Single Versus Recurrent Traumatic Posterior Shoulder Dislocation Events.

Gibbs HP, McQuade AM, Ramirez V, Feeley SM, McCarthy CF, Hoyt BW, Dickens JF

BACKGROUND: Posterior shoulder dislocations are rare but particularly challenging in high-demand populations. While anterior instability has been well studied, limited data exist on the morphological consequences of traumatic first-time and recurrent posterior dislocations.

PURPOSE: To quantify posterior glenoid bone loss (pGBL), glenoid version, reverse Hill-Sachs lesions, and acromial morphology in patients with traumatic posterior shoulder dislocations and compare these features between the single and recurrent dislocation groups.

STUDY DESIGN: Case series; Level of evidence, 4.

METHODS: A retrospective review was conducted using the Military Health System Data Repository to identify patients who sustained traumatic posterior shoulder dislocations between January 2017 and December 2024. These were defined as events requiring manual reduction with subsequent advanced imaging demonstrating a posterior labral tear or posterior bony Bankart lesion, with or without a reverse Hill-Sachs lesion. Morphological features were independently assessed by 2 raters. Measured factors included pGBL (measured by the perfect circle technique (PCT), two-thirds glenoid height methods, and one-third glenoid height method), glenoid version, reverse Hill-Sachs Lesion parameters, and acromial tilt and height. Patients were stratified into single and recurrent dislocation groups. Statistical compa [...]

[PubMed](#) · [DOI](#)

Comparison of Long-term Supraspinatus Tear Progression After Arthroscopic Isolated Subscapularis Repair With and Without Comma Tissue Preservation: A Minimum 10-Year MRI Follow-up Study.

Do WS, Chun YM, Kim SJ, Sim W, Yoon TH

BACKGROUND: Long-term outcomes after arthroscopic isolated subscapularis repair are limited, and magnetic resonance imaging (MRI)-based assessments extending to 10 years are particularly scarce. Therefore, it remains unclear whether preserving the comma tissue (in-continuity technique) provides long-term advantages over disrupting the upper subscapularis margin (margin disruption technique).

HYPOTHESIS: The in-continuity and margin disruption techniques would demonstrate comparable 10-year radiological and clinical outcomes in patients undergoing arthroscopic isolated subscapularis repair.

STUDY DESIGN: Cohort study; Level of evidence, 3.

METHODS: Among patients who underwent arthroscopic isolated subscapularis repair between 2008 and 2015, those with an intact repair on 6-month postoperative MRI and at least 2 years of clinical follow-up were identified. From this cohort, 24 patients treated with the in-continuity technique and 24 treated with the margin disruption technique were included. Ten-year MRI findings (subscapularis Goutallier grade, tendon integrity, and supraspinatus tear progression) and clinical outcomes (active range of motion [ROM], visual analog scale [VAS] pain score, Subjective Shoulder Value [SSV], American Shoulder and Elbow Surgeons [ASES] score, and University of California, Los Angeles [UCLA] score) were compared. A subgroup analysis was performed to [...]

[PubMed](#) · [DOI](#)

Intraoperative Zoledronic Acid for Arthroscopic Rotator Cuff Repair Results in Distinct Circulating Micro-RNA Profiles Indicating Improved Tendon-to-Bone Healing.

Schanda JE, Hackl M, Bischl R, Ullrich R, Boesmueller S, Mittermayr R, Redl H, Grillari J et al.

BACKGROUND: A prospective, randomized, placebo-controlled trial showed reduced retears after arthroscopic rotator cuff (RC) repair in patients without osteoporosis by a systemic single-dose of zoledronic acid. Distinct micro-ribonucleic acids (miRNAs) related to inflammation, fibrosis, and tendon-to-bone healing are associated with RC injuries.

PURPOSE: To investigate the longitudinal effects of a single-dose of zoledronic acid in patients without osteoporosis undergoing arthroscopic RC repair on circulating miRNAs in order to explore the molecular mechanism of this treatment.

STUDY DESIGN: Controlled laboratory study.

METHODS: Data were collected in the course of a single-center, prospective, randomized, placebo-controlled, triple-blinded (investigator, surgeon, patient) phase II trial. A total of 80 patients underwent arthroscopic RC repair and were intraoperatively randomized to the zoledronic acid group (n = 40) or the control group (n = 40). Circulating plasma miRNAs were assessed preoperatively, 2 days postoperatively, and 3 months postoperatively using small RNA sequencing and reverse transcription quantitative polymerase chain reaction (RT-qPCR).

RESULTS: Six miRNAs were selected for validation by RT-qPCR based on small RNA-sequencing analysis. No statistical differences in miRNA plasma levels were observed preoperatively between the 2 study groups. Two days after s [...]

[PubMed](#) · [DOI](#)

Association of Central Acetabular Osteophytes With Microinstability and Increased Combined Anteversion in Borderline Dysplasia Hips.

Zhu JB, Ding R, Huang Y, Chen YZ, Zheng GY, Uchida S, Shen C

BACKGROUND: Central acetabular osteophytes (CAOs) have been implicated in abnormal hip biomechanics. However, their relationship with microinstability in borderline developmental dysplasia of the hip (BDDH) is unclear.

PURPOSE/HYPOTHESIS: The purpose of this study is to compare radiographic parameters between BDDH hips with and without CAOs and to determine whether CAOs are associated with microinstability. It was hypothesized that CAOs would correlate with increased version abnormalities and microinstability.

STUDY DESIGN: Case-control study; Level of evidence, 3.

METHODS: We retrospectively reviewed patients with BDDH, defined as a lateral center-edge angle of 18° to 25°, who underwent hip arthroscopy from 2020 to 2024. Patients were grouped as BDDH with CAO (cBDDH) or without CAO (nBDDH). Radiographic parameters, including anterior/posterior acetabular coverage, femoral neck version, acetabular version, and combined anteversion, were compared. Intraoperative findings of microinstability and ligamentum teres (LT) pathology were analyzed. Binary logistic regression identified predictors of CAO.

RESULTS: A total of 157 patients were included (cBDDH, 76; nBDDH, 81). The cBDDH group demonstrated higher rates of microinstability and LT tears, as well as increased femoral neck, acetabular, and combined anteversion. Logistic regression showed that microinstability ($\beta = .027$; $P < [...]$)

[PubMed](#) · [DOI](#)

Minimum 5-Year Outcomes After Arthroscopic Subspine Decompression in Patients With Femoroacetabular Impingement: A Propensity-Matched Study.

Kahana-Rojkind AH, Rana K, Barda SR, Quesada-Jimenez R, Flynn ME, Sugarman EP, Domb BG

BACKGROUND: Subspine impingement (SSI) is a frequently coexisting pathology in patients undergoing hip arthroscopy for femoroacetabular impingement (FAI). While short-term outcomes after arthroscopic subspine decompression (SSD) are promising, midterm outcomes remain unclear.

PURPOSE: To evaluate minimum 5-year patient-reported outcomes (PROs), survivorship, and rates of clinically meaningful improvement after arthroscopic SSD in patients undergoing hip arthroscopy for FAI; and to compare these results to a matched cohort without SSD.

STUDY DESIGN: Cohort study; Level of evidence, 3.

METHODS: The data of patients undergoing primary hip arthroscopy for FAI between 2009 and 2019 were retrospectively reviewed. Inclusion criteria included a confirmed diagnosis of FAI and SSI and complete PROs (modified Harris Hip Score, Nonarthritic Hip Score, Hip Outcome Score-Sports-Specific Subscale, International Hip Outcome Tool-12 score, and visual analog scale score) at a minimum of 5 years. Exclusion criteria included prior ipsilateral hip surgery, workers' compensation claims, and Tönnis grade >1. Patients with SSD were matched 1:2 to a control group without SSI. Matching variables included age, sex, body mass index, follow-up duration, lateral center-edge angle, alpha angle, Tönnis grade, acetabular labrum articular disruption grade, and labral treatment. Outcomes were compared between [...]

[PubMed](#) · [DOI](#)

Revision Hip Arthroscopy: Propensity Score-Matched Comparison of Labral Reconstruction Versus Repair With Minimum 5-Year Outcomes.

Quesada-Jimenez R, Cohen MF, Rana K, Kahana-Rojkind AH, Domb BG

BACKGROUND: There is a paucity of literature comparing the midterm outcomes of arthroscopic labral reconstruction and arthroscopic labral repair in the revision setting.

PURPOSE: To evaluate the minimum 5-year patient-reported outcomes (PROs) of patients who underwent revision hip arthroscopy, comparing labral reconstruction with labral repair.

STUDY DESIGN: Cohort study; Level of evidence, 3.

METHODS: A retrospective analysis was performed for patients who underwent revision hip arthroscopy, either labral repair or labral reconstruction, to treat femoroacetabular impingement and labral tears between 2008 and 2019. Patients who underwent reconstruction were matched 1:1 to patients who underwent repair. Included patients had complete preoperative and minimum 5-year follow-up scores for at least one of the following PRO measures: modified Harris Hip Score, Non-Arthritic Hip Score, Hip Outcome Score-Sports-Specific Subscale, patient satisfaction, and visual analog scale for pain. Rates of achieving the minimal clinically important difference and patient acceptable symptom state as well as secondary surgery rates were compared between groups.

RESULTS: A total of 68 patients were included, with 34 patients who underwent labral reconstruction successfully matched to 34 patients who underwent labral repair, controlling for age, body mass index, sex, acetabular Outerbridge grade, a [...]

[PubMed](#) · [DOI](#)

Has the Protective Effect of Hormonal Contraception Been Overestimated? A Systematic Review and Meta-analysis of Oral Contraceptive Pills and ACL Injury.

Paska J, Watson SL, Larrabee GA, Ndjonko LCM, Tjong VK

BACKGROUND: Hormonal contraception (HC), including oral contraceptive pills (OCPs) and long-acting reversible contraceptives, is widely used by reproductive-aged women for various medical indications. Emerging evidence suggests a possible association between HC use and reduced anterior cruciate ligament (ACL) injury risk; however, no meta-analysis has systematically evaluated this relationship.

PURPOSE: To determine whether HC use reduces the risk of ACL injury in females through a systematic review and meta-analysis.

STUDY DESIGN: Meta-analysis; Level of evidence, 3.

METHODS: A systematic search of the PubMed, OVID (MEDLINE), and Embase databases was conducted from inception through March 2025 following PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guidelines. Studies were included if they reported ACL injury incidence among both HC users and nonusers. Data were synthesized using a random-effects meta-analysis to calculate a pooled risk ratio. Publication bias was evaluated using funnel plots and Egger regression.

RESULTS: Eight studies consisting of 18,103,890 participants met inclusion criteria. All studies evaluated OCPs, and 3 included other forms of HC. The pooled risk ratio for ACL injury among HC users versus nonusers was 0.88 (95% CI, 0.77-1.01). Heterogeneity was high ($I^2 = 97.3\%$), and Egger regression showed no significant publicatio [...]

Surgical Timing and Outcomes in Medial Meniscus Posterior Root Tear Repair: A Systematic Review.

Moews LD, Thamrongsuksiri N, Morgan JT, Vega TF, Perez Lloveras GO, Casanova F, Verma NN, Chahla J

BACKGROUND: Medial meniscus posterior root tears (MMPRTs) compromise hoop stress transmission and tibiofemoral load distribution, accelerating degenerative changes comparable with total meniscectomy. While early repair has been recommended, many patients present late because of a delayed diagnosis or an insidious onset of symptoms. The effect of delayed surgical management on outcomes remains unclear.

PURPOSE: To systematically review the literature comparing clinical outcomes, radiological progression, and complication rates of MMPRT repair performed at <3 months, 3 to 6 months, and >6 months after the injury.

STUDY DESIGN: Systematic review; Level of evidence, 4.

METHODS: A systematic search of PubMed, Embase, and Scopus was conducted in accordance with PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses) guidelines and registered with PROSPERO. A total of 11 studies (366 patients) reporting isolated transtibial pullout repair with a documented time from injury to surgery were included. Extracted data included patient-reported outcome scores (International Knee Documentation Committee [IKDC], Lysholm), osteoarthritis progression by the Kellgren-Lawrence grade, meniscal extrusion, healing status, and surgical failures or complications.

RESULTS: Across all surgical timing groups, MMPRT repair resulted in significant functional improvements. Patients t [...]

Hip Arthroscopy in Patients With Acetabular Dysplasia: A Systematic Review of Clinical Outcomes at Long-term Follow-up.

Sparks CA, Monty TL, Brinkman JC, Barton KI, Nho SJ

BACKGROUND: Hip arthroscopy has shown satisfactory outcomes for the treatment of intra-articular pathology and femoroacetabular impingement syndrome (FAIS) in the setting of acetabular dysplasia at short- and mid-term follow-up. However, the long-term outcomes remain less understood.

PURPOSE: To systematically review the literature to determine patient-reported outcomes (PROs), clinically significant outcome (CSO) achievement, reoperation rates, and predictors of failure in patients with acetabular dysplasia undergoing hip arthroscopy at long-term follow-up.

STUDY DESIGN: Systematic review; Level of evidence, 4.

METHODS: Three databases were searched around the following terms: hip arthroscopy, dysplasia, and long-term follow-up. Articles available in English, presenting original data, and reporting on a mean follow-up of ≥ 10 years after primary hip arthroscopy using either PROs, CSOs, or conversion to total hip arthroplasty (THA) and/or revision surgery were included. Quality assessment was completed using Methodological Index for Non-Randomized Studies (MINORS) assessment.

RESULTS: Five studies (555 hips) were included in the systematic review: patients had a mean age of 30 to 41 years, 13% to 88% of patients were female, and follow-up ranged from 9.6 to 12 years. MINORS assessment ranged from 11 to 21. All studies included PROs and reported statistically significant impr [...]

Psychiatric Diagnoses and Outcomes After Hip Arthroscopy: A Systematic Review and Meta-analysis.

Villegas Meza AD, Nocek M, Morales Lugo B, Mitchell BC, Philippon MJ

BACKGROUND: Psychiatric comorbidity-most commonly depression and anxiety-is frequently reported among patients undergoing hip arthroscopy and may influence postoperative trajectories, but existing analyses vary in scope and methods, limiting clinical interpretation.

PURPOSE: To determine whether recorded psychiatric diagnoses are associated with worse surgical, patient-reported, and health care utilization outcomes after hip arthroscopy.

STUDY DESIGN: Systematic review and meta-analysis; Level of evidence, 3.

METHODS: The authors conducted a PRISMA (Preferred Reporting Items for Systematic Reviews and Meta-Analyses)-compliant search (International Prospective Register of Systematic Reviews [PROSPERO] CRD420251101816) of the MEDLINE, Embase, Web of Science, and PsycINFO databases to identify studies comparing postoperative outcomes in patients undergoing hip arthroscopy with and without recorded psychiatric diagnoses. Two reviewers independently screened, extracted data, and assessed the risk of bias using the Risk of Bias in Non-randomized Studies of Interventions tool. Primary endpoints were arthroscopy failure, revision, conversion to total hip arthroplasty (THA), and readmission. Secondary endpoints included the modified Harris Hip Score (mHHS), International Hip Outcome Tool-12 (iHOT-12) score, visual analog scale (VAS) pain score, and costs. Random-effects meta-analyses [...]

[PubMed](#) · [DOI](#)

Platelet Concentration Does Not Influence Clinical Efficacy and Retear Rates of Rotator Cuff Repair With Platelet-Rich Plasma: A Systematic Review and Meta-analysis With Meta-Regression.

Lim JJ, Stepaniuk EA, Belk JW, Keeter C, Oeding JF, Dragoo JL, McCarty EC, Frank RM

BACKGROUND: Platelet-rich plasma (PRP) is emerging as a popular augmentation technique in the context of rotator cuff repair (RCR) to strengthen the repair construct and promote healing. Previous studies examining the efficacy of PRP augmentation during RCR have described heterogeneous results.

PURPOSE: To determine whether platelet concentration of a PRP injection as an adjunct of RCR impacts clinical outcomes and retear rates.

STUDY DESIGN: Systematic review and meta-analysis; Level of evidence, 2.

METHODS: A systematic review was performed by searching the PubMed, Cochrane Library, and Embase databases to identify level 1 and 2 studies that evaluated the clinical efficacy of RCR augmented with PRP. The search phrase used was "rotator cuff repair AND (PRP OR platelet rich plasma OR platelet-rich plasma)." Included randomized controlled trials were classified as utilizing high-dose or low-dose PRP based on their platelet concentration factor as a >4-fold increase over whole blood. The outcomes evaluated were the standardized mean differences of visual analog scale scores, Constant-Murley scores, American Shoulder and Elbow Surgeons scores, and University of California-Los Angeles scores, and pooled odds ratios of retear rates. Mixed-effects and random-effects meta-analyses were performed along with meta-regression to evaluate relationships in patient outcomes associated with [...]

[PubMed](#) · [DOI](#)

Episode Count Should Not Substitute for Anatomy in Arthroscopic Bankart Repair: Response.

Lin RT, Gilbert R, Lin A

[PubMed](#) · [DOI](#)

Time After ACL Reconstruction: Degenerative Clock or Vulnerable Knee Exposure? Response.

Agostinone P, Romandini I, Rotzius P, Zaffagnini S, Forssblad M, Sandon A

[PubMed](#) · [DOI](#)

Episode Count Should Not Substitute for Anatomy in Arthroscopic Bankart Repair: Letter to the Editor.

Qu X, Li D, Tang X, Chen B

[PubMed](#) · [DOI](#)

Time After ACL Reconstruction: Degenerative Clock or Vulnerable Knee Exposure? Letter to the Editor.

Tang X, Zhang J, Yu B

[PubMed](#) · [DOI](#)



The Bone & Joint Journal

Volume 108-B, Issue 8 · August 2026 · 13 new articles

Beyond BMI cut-offs : optimizing risk, not excluding patients.

Alvand A, Rudran B, Haddad FS

[PubMed](#) · [DOI](#)

Clinical efficacy to climate accountability : the case for embedding environmental outcomes in orthopaedic trials.

Farrow L, Purdie S, Evans JT, Krause T, Campbell M

There is increasing global recognition of the importance of sustainability in the design and delivery of healthcare, including trauma and orthopaedics. There has, however, to date been little focus on providing high-quality research which includes environmental outcomes to guide clinical practice. This annotation explores the importance of including sustainability outcomes using three key carbon-related domains, and highlights the urgent need for the incorporation of sustainability outcomes into future clinical trials.

[PubMed](#) · [DOI](#)

Periprosthetic joint infection : lessons from the INFORM research programme.

Moore AJ, Beswick AD, Whitehouse MR, Blom AW

Infection is one of the most devastating complications following arthroplasty. The National Institute for Health and Care Research (NIHR) funded the multinational INFection ORthopaedic Management (INFORM) research programme to study this problem comprehensively, and then funded INFORM: Evidence into Practice to translate findings into the INFORM guidelines. This annotation outlines the key findings with particular emphasis on hip and knee arthroplasty, including risk factors, impact, diagnosis, and treatment, and suggests guidelines underpinned by high-quality evidence from the programme. Finally, future directions for research are discussed.

[PubMed](#) · [DOI](#)

Glucagon-like peptide-1 receptor agonists are associated with improved 90-day outcomes after total hip and knee arthroplasty : a systematic review and meta-analysis.

Rudran B, Little CDJ, Palmer AJR, Judge A, Carli A, Price AJ, Alvand A

AIMS: Obesity and diabetes are risk factors for developing osteoarthritis and associated with worse outcomes following total hip (THA) and total knee arthroplasty (TKA). Glucagon-like peptide-1 receptor agonists (GLP-1as) improve glycaemic control and promote weight loss. The aim of this study was to assess the impact of GLP-1a treatment on 90-day surgical complications following THA and TKA.

METHODS: A systematic review was undertaken according to PRISMA guidelines. Studies were included if they reported at least one outcome of interest following primary THA or TKA. In all the studies which were included, a GLP-1a was prescribed for the management of type 2 diabetes mellitus or for weight loss in obese patients. The primary outcome was the 90-day surgical complication rate in the combined THA and TKA cohort. The 90-day surgical complications were defined as: periprosthetic joint infection (PJI), wound dehiscence, periprosthetic fracture (PPF), haematoma, nerve injury, or surgical site infection (SSI). Secondary outcomes included 90-day medical complications and readmission rates, all-cause revision, healthcare costs, and length of stay.

RESULTS: A total of 78 studies were reviewed, and ten matched cohort studies with a total of 96,356 patients (30,350 THAs and 66,606 TKAs) were included. The mean age of the patients was 61.9 years; 60.1% female (n =

57,939). Two studies had [...]

[PubMed](#) · [DOI](#)

Autologous impaction bone grafting of a cemented acetabular component in primary total hip arthroplasty in young patients: a randomized controlled trial.

de Boer DR, Pasman P, Munnik-Hagewoud R, Steinweg MJQ, Edens MA, van Driel PBAA, Ettema HB

AIMS: This randomized controlled trial (RCT) aimed to compare clinical and radiological outcomes of cemented acetabular component in primary total hip arthroplasty (THA) in patients aged under 60 years with and without autologous impaction bone grafting (IBG).

METHODS: A single-centre, triple-blinded RCT was conducted in patients aged 18 to 59 years undergoing primary THA without acetabular defects. Patients were randomized to receive either IBG or no IBG. Noninferiority of IBG was assessed for the primary endpoint, the Hip disability and Osteoarthritis Outcome Score (HOOS) sub-scale for activities of daily living (ADL) at one-year follow-up. Secondary outcomes included other patient-reported outcome measures (PROMs), perioperative outcomes, complications, and radiological outcomes. Radiographs assessed acetabular component and centre of rotation positioning, offset changes, and the presence of radiolucent lines in the bone-cement interface.

RESULTS: A total of 131 patients were randomized. No significant group differences were found for the HOOS-ADL sub-scale at one year (mean difference 1.9 points (95% CI -4.85 to 8.7); $p = 0.578$), meeting the criteria for noninferiority. Furthermore, there was no effect of group on the scores for other PROMs. Surgery duration (+ 7.5 minutes; $p < 0.001$) and blood loss (+ 100 ml; $p = 0.001$) were significantly higher in the IBG group. Complic [...]

[PubMed](#) · [DOI](#)

Increased risk of early failure of multiply revised total hip arthroplasty.

D'Apollonio TM, Hopman MLT, Sullivan T, Abrahams JM, Sheth NP, Solomon LB, Callary SA

AIMS: As the demand for revision total hip arthroplasty (rTHA) grows, understanding how successive revision procedures influence implant survivorship and patient outcomes becomes increasingly important. This study aimed to evaluate the association between revision number and implant survivorship following aseptic rTHA.

METHODS: We analyzed 845 aseptic rTHAs performed between January 2003 and December 2023. Index primary THA dates, rTHA sequence, and reoperations were sourced and verified using hospital records, national registry data, and radiological records. Cases were stratified by revision number: first (R1), second (R2), or third and beyond (R3+). Re-revision risk was evaluated using cumulative incidence functions and Fine-Gray competing-risk regression, with death treated as a competing event.

RESULTS: Of 845 rTHAs (717 patients), 838 (710 patients) with confirmed index THA were included: 598 R1, 155 R2, and 85 R3+. At two years, cumulative incidence of re-revision was 10.2% (95% CI 8.0 to 12.9) for R1, 14.2% (95% CI 9.6 to 20.8) for R2, and 17.7% (95% CI 11.0 to 27.7) for R3+. Unadjusted competing-risk regression demonstrated a higher cumulative incidence of re-revision in R2s compared with R1s. However, after multivariable adjustment, revision number was not independently associated with a statistically significant increase in re-revision risk. Across all revision gro [...]

[PubMed](#) · [DOI](#)

Preoperative predictors of patient-reported outcomes after joint-preserving forefoot surgery for rheumatoid arthritis : a case series.

Etani Y, Noguchi T, Hirao M, Okamura G, Sugimoto A, Okada S, Nakata K, Ebina K

AIMS: Forefoot deformities are common in patients with rheumatoid arthritis (RA) and often impair quality of life. Patient-reported outcome measures (PROMs), including the Self-Administered Foot Evaluation Questionnaire (SAFE-Q), are increasingly used to assess surgical outcomes; however, the preoperative conditions needed to achieve favourable postoperative PROMs remain unclear. This study aimed to identify factors associated with postoperative SAFE-Q scores (mean score across five sub-scales, maximum 100) and determined preoperative cutoff values predictive of achieving predefined postoperative SAFE-Q thresholds (≥ 70 or ≥ 90).

METHODS: In this multicentre, retrospective case series, 101 feet in 80 patients (including 21 bilateral cases) with RA who underwent forefoot surgery between July 2015 and July 2023 were analyzed (mean age at surgery 65.8 years (SD 10.7); 79 female patients). Inclusion criteria were symptomatic forefoot deformities with available SAFE-Q data pre-surgery and at $\geq$ one year post-surgery. Procedures included modified scarf osteotomy and offset shortening osteotomies for the metatarsals. SAFE-Q scores, Japanese Society for Surgery of the Foot scores, radiological parameters, and disease activity were evaluated pre- and postoperatively. Multiple regression identified preoperative factors associated with postoperative SAFE-Q scores, and significant variable [...]

[PubMed](#) · [DOI](#)

Non-medical surgical delay increases one-year mortality following hip fracture.

Busby C, Nightingale J, Deacon C, Deakin D, Valdes A, Ollivere BJ, Moran C

AIMS: This study aimed to assess the association between surgical delay due to non-medical reasons and one-year mortality in patients with hip fracture, and to explore whether a dose-response relationship exists between increasing time to surgery and mortality risk.

METHODS: A retrospective cohort analysis was performed using prospectively collected data from the Nottingham Hip Fracture Database. A total of 5,414 patients aged $\geq$ 60 years admitted with a fragility femoral fracture to a major trauma centre between April 2016 and March 2023 were reviewed. Time to theatre was measured from emergency department arrival to anaesthetic start. Delays beyond 36 hours were categorized as either medical or logistical. The primary outcome was all-cause one-year mortality with a secondary outcome being length of stay (LOS). Multivariable Cox proportional hazards models were used to assess associations, adjusting for age, sex, comorbidity, and cognition.

RESULTS: Of the cohort, 38.5% (n = 2,083) experienced surgical delay beyond 36 hours; 27.5% (n = 1,490) due to limited surgical capacity and 11.0% (n = 593) due to medical unfitness. After adjusting for age, sex, comorbidity, and cognition, delay due to logistical factors was associated with a 20% increase in one-year mortality (hazard ratio (HR) 1.20 (95% CI 1.06 to 1.36); p = 0.004). A secondary analysis excluding medically unfit patient [...]

[PubMed](#) · [DOI](#)

Use of neurolytic nerve blocks and mortality in nonoperative management of hip fracture patients : a nationwide population study in the Netherlands.

van Bremen HE, Lommerse MI, Hegeman JH, Schuijt HJ, Kallewaard JW, Wijnen HH, Seppala LJ, van der Velde N et al.

AIMS: Nonoperative management may be suitable for very frail hip fracture patients; in these instances, neurolytic nerve blocks with phenol could act as a long-term analgesic, with potentially fewer disadvantages compared with systemic analgesia. While neurolytic nerve blocks are being used more frequently in end-of-life care, there are limited data on how often they are used, and what long-term outcomes they achieve in patients with hip fractures. This study aims to evaluate the association between neurolytic nerve block use and mortality in patients with hip fractures managed nonoperatively, while also exploring the characteristics of patients who receive these blocks.

METHODS: A retrospective population-based cohort study was conducted including patients with a hip fracture aged 70 years and older, who were treated nonoperatively between 1 January 2024 and 31 December 2024. The primary outcome was mortality, comparing patients who received a neurolytic block with those who did not.

RESULTS: Of 1,415 nonoperatively managed patients (mean age 86 years (SD 6.75)), 633 (45%) received neurolytic nerve blocks. These patients were more frail, had greater levels of dependence, malnourishment, and immobility, and were diagnosed with a displaced fracture more often than those who did not receive a block. Nerve block patients also showed significantly lower mortality at seven days (2 [...])

[PubMed](#) · [DOI](#)

A matter of definition : variation in treatment success of fracture-related infections.

Reinert N, Poggioli A, Metsemakers WJ, Clauss M, Kuehl R, FRI-Basel Research Group, Morgenstern M, FRI-Basel research Group et al.

AIMS: The establishment of the fracture-related infection (FRI) consensus definition and the FRI classification system provide a foundation for research and the development of clinical recommendations. To date, however, the evaluation of treatment success is still impeded by a lack of clear and standardized outcome definitions. We

hypothesized that applying different outcome measurements to the same cohort of FRI patients results in significantly different treatment success rates.

METHODS: Adult patients undergoing curative surgery for FRI at the Center of Musculoskeletal Infections, University Hospital Basel, between 1 January 2020 and 31 December 2023 were included in this prospective cohort study. Treatment success was evaluated by review of patient charts; after one year of follow-up according to seven predefined outcome definitions (S1 to S7): no infection persistence at one year (S1); no recurrence within one year (with the same (S2) or a new (S3) pathogen); bony union (S4); no infection recurrence, or persistence and bony union (S5); no unplanned revision surgery (S6); and bony union and no infection recurrence, or persistence and no unplanned revision surgery (S7).

RESULTS: In total, 106 patients with 112 confirmed FRIs were included. At one year, treatment success varied significantly ($p < 0.001$) between the different outcome definitions: for S1 the success rate was [...]

[PubMed](#) · [DOI](#)

Reducing complications after resection of pelvic bone sarcomas : an aggregation of marginal gains.

Jeys LM, Kurisunkal V, Parry MC, Morris G, Stevenson JD, Laitinen MK

AIMS: Primary pelvic bone sarcomas are rare and their treatment remains complex, with reconstruction after resection remaining controversial due to high complication and revision rates. Both prosthetic and non-prosthetic reconstructions carry risks including infection, implant failure, wound problems, and reoperation. The aim of this study was to review the changes in the management of these patients over the 20-year period between 2003 and 2023.

METHODS: This was a retrospective cohort study including 479 patients who underwent surgery for a primary pelvic sarcoma during this time, divided into a historical (2003 to 2012) and a modern (2013 to 2023) group. Postoperative complications and the rates of return to theatre (RTT) were compared, and factors associated with improved outcomes were analyzed.

RESULTS: Early RTT within six weeks decreased significantly from 22% in the historical cohort ($n = 38$) to 14% in the modern cohort ($n = 42$; $p = 0.021$). Historically, the ilioinguinal incision was associated with a 44% early rate of RTT ($n = 8$), compared with 10% when using the question mark incision in the modern cohort ($n = 10$). In the modern cohort, the rate of early RTT was similar between those with prosthetic (17%; $n = 13$) and non-prosthetic reconstructions (13%; $n = 29$) ($p = 0.344$), although the long-term rate of RTT remained significantly higher in the prosthetic group ($p = [...]$)

[PubMed](#) · [DOI](#)

Natural history of solitary osteochondroma : a retrospective analysis of 236 patients in South Korea.

Sung MJ, Kim CH, Jung ST, Kim S

AIMS: The aim of this study was to analyze the natural history of solitary osteochondroma and determine how skeletal maturity influences tumour behaviour and treatment outcomes.

METHODS: A retrospective analysis of 236 patients with solitary osteochondroma (151 skeletally immature, 85 skeletally mature) was undertaken. Skeletal maturity was defined as bone age ≥ 18 years in males and ≥ 17 years in females. Tumour characteristics, disease progression, indications for surgery and recurrence rates were evaluated. Interobserver reliability for bone age assessment was excellent (intraclass correlation coefficient (ICC) 0.92 (95% CI 0.88 to 0.95)).

RESULTS: Spontaneous regression occurred exclusively in skeletally immature patients (17.2% ($n = 26$) vs 0%; $p < 0.001$). Skeletally immature patients predominantly presented with sessile lesions (76.2%; $n = 115$), had lower surgical needs (31.8%; $n = 48$), but higher progression rates (9.9% ($n = 15$) vs 1.2% ($n = 1$); $p = 0.013$). Skeletally mature patients showed more stable disease but required surgery more often (54.1%; $n = 46$). Among skeletally immature patients with regression, independent predictors were younger age at diagnosis (odds ratio (OR) 0.82 (95% CI 0.73 to 0.92); $p < 0.001$), proximal humeral location (OR 10.59 (95% CI 2.01 to 55.85); $p = 0.005$), and humeral shaft location (OR 38.93 (95% CI 4.81 to 315.10); $p = 0.001$). The optim [...]

[PubMed](#) · [DOI](#)

Cost-effectiveness of screening for developmental dysplasia of the hip in children born in England and Wales.

Plumpton CO, Perry DC

AIMS: The aim of this study was to evaluate the cost-effectiveness of screening programmes for identifying developmental hip dysplasia (DDH) in children born in England and Wales.

METHODS: The costs and quality-adjusted life-years (QALYs) associated with DDH screening programmes were estimated using a decision-analytic model. Six programmes were modelled from an NHS perspective: 1) Current practice - 'selective' screening: clinical examination with later ultrasound screening (USS) for those with an abnormal examination or specific risk factors (breech birth, first degree family history of DDH); 2) No hip screening: no clinical examination or risk factor assessment; 3) Clinical examination: treatment decisions based solely upon a positive Barlow or Ortolani manoeuvre; 4) Universal USS at birth: USS in the neonatal period in addition to a postnatal check; 5) Universal USS at six weeks: USS in addition to the six-week postnatal check; 6) Sex-stratified screening: clinical examination with selective USS for males and universal USS in the neonatal period for females. The incremental net monetary benefit (INMB) was estimated at a cost-effectiveness threshold of £20,000 per QALY gained. Whether the result was robust to assumptions was assessed using analyses of uncertainty. Secondary outcomes included an estimate of the value of reducing the uncertainty of the decisions.

RESULTS: Un [...]

[PubMed](#) · [DOI](#)



Clinical Orthopaedics & Related Research

Volume 484, Issue 8 · August 2026 · 38 new articles

A Conversation With ... Malcolm Willett, Cartoonist Who Looks at Us With a Wry Smile.

Leopold SS

[PubMed](#) · [DOI](#)

Editor's Spotlight/Take 5: How Does Traumatic Hand Injury Impact Patients in a Safety-net Healthcare System? A Mixed Methods Study.

Wongworawat MD

[PubMed](#) · [DOI](#)

How Does Traumatic Hand Injury Impact Patients in a Safety-net Healthcare System? A Mixed Methods Study.

Higashi RT, Lee J, Hernandez Y, Pontón M, Liao JM, Cross KC, Billig JI

BACKGROUND: Traumatic hand injuries can result in substantial harms to patients' functional abilities and financial health, disproportionately affecting individuals who are of working age, low income, and uninsured. However, little is known about how patients and their families cope with the functional limitations and economic consequences that follow, and how these injuries affect patients' mental and emotional health. Exploring clinician and staff perceptions about the impact of traumatic hand injuries on patients may provide insight into how to help address the various burdens patients face.

QUESTIONS/PURPOSES: In this paper, we aimed to explore the following questions: (1) What is the impact of a traumatic hand injury on patients' financial health? (2) How does a traumatic hand injury affect patients' physical health and disability? (3) What kind of toll does a traumatic hand injury take on patients' emotional and mental health? (4) What health system challenges function as additional stressors after a traumatic hand injury?

METHODS: This descriptive study consisted of surveys and semistructured interviews. We recruited patients from an outpatient hand surgery clinic at a safety-net hospital in a southern US city. Eligible patients were English- or Spanish-speaking, age 18 years or older, who presented to the clinic after a traumatic hand injury. Between January and April [...]

[PubMed](#) · [DOI](#)

Clinical Faceoff: Social Media for Orthopaedic Surgeons-Tool or Vice?

Fourman MS, Daniels AH, Herron PD

[PubMed](#) · [DOI](#)

Behind the Mask: Setting Expectations.

Zhang SE

[PubMed](#) · [DOI](#)

Value-based Healthcare: Moving From PROMs as Population Benchmarks to Clinical Tools for Patient Goal Setting and Shared Decision-making.

Ahlfinger Z, Bozic KJ

[PubMed](#) · [DOI](#)

Virtue Ethics in a Value-driven World: Soliciting Donations From Grateful Patients.

Humbyrd CJ

[PubMed](#) · [DOI](#)

Not the Last Word: Seers, Soothsayers, and Surgeons.

Bernstein J

[PubMed](#) · [DOI](#)

Editorial Comment: Selected Proceedings From the 2025 Musculoskeletal Infection Society Meeting.

Zalavras CG

[PubMed](#) · [DOI](#)

Can We Validate the Musculoskeletal Infection Society Criteria in Patients Suspected of Infection After Transfemoral Osseointegration Surgery?

Bozzay AB, Rivera JA, Rabin SE, Souza JM, Potter BK, Forsberg JA

BACKGROUND: Major complications that may result in the removal of osseointegrated implants in patients with limb loss include peri-implant fracture, severe bone resorption, and infection. It remains challenging to diagnose and manage peri-implant infection in patients who have undergone osseointegration surgery. The Musculoskeletal Infection Society (MSIS) developed criteria to identify periprosthetic joint infection (PJI) after total joint arthroplasty during the 2018 MSIS/International Consensus Meeting; however, there is no such validated system for patients who have received osseointegrated implants, and some of the measures on the MSIS/International Consensus Meeting criteria are not applicable to osseointegrated implants. Nevertheless, establishing an adapted version of the MSIS criteria, or a comparable standardized system, would be valuable for enabling consistent diagnosis of osseointegration-associated peri-implant infections.

QUESTIONS/PURPOSES: (1) Is there a difference in the serum parameters of the 2018 MSIS criteria between a cohort of patients without infection and a cohort with an infection? (2) Can the serum parameters of the 2018 MSIS criteria be validated in a population that has undergone osseointegration? (3) Can we develop a new, valid set of criteria to gauge the likelihood of infection from serum parameters for those who have undergone osseointegration [...]

[PubMed](#) · [DOI](#)

CORR Insights®: Can We Validate the Musculoskeletal Infection Society Criteria in Patients Suspected of Infection After Transfemoral Osseointegration Surgery?

Synthesis and Characterization of an Antimicrobial Honey-based Composite Bone Cement.

Kemp LE, Delgado-Alvarado I, Rossiter TE, Chen AF, Tower RJ, Tatara AM

BACKGROUND: Orthopaedic infections are a major cause of morbidity and mortality with increasing prevalence. Antibiotics are often added to bone cement or polymethyl methacrylate (PMMA) for local delivery to prevent or treat infection. However, PMMA is not inherently antimicrobial and has poor antibiotic elution kinetics. A biomaterial composite of PMMA and antimicrobial medical-grade honey (PMMA-H) could address some of these shortcomings of current bone cements to better inhibit bacteria.

QUESTIONS/PURPOSES: (1) How does honey affect the architecture and static mechanical properties of bone cement? (2) How does honey affect antibiotic elution? (3) What is the in vitro antimicrobial activity of PMMA-H with and without antibiotics?

METHODS: Three different formulations of bone cements were created by mixing a generic dental bone cement without honey (0 wt%), with a low amount of medical-grade manuka honey (15 wt%), or with a high amount of honey (30 wt%). Vancomycin or gentamicin were added to the three formulations at a typical cement loading dose (3 wt%) or no additional antibiotic for a total of nine formulations. We characterized the subsequent total porosity and percentage of open porosity of constructs synthesized from the nine formulations using microcomputed tomography. Both the compressive and bending yield strength and modulus were measured by mechanical testing. Con [...]

[PubMed](#) · [DOI](#)

CORR Insights®: Synthesis and Characterization of an Antimicrobial Honey-based Composite Bone Cement.

DeBaun MR

[PubMed](#) · [DOI](#)

What Is the Fate of Retained Antibiotic Spacers After First-stage Revision for Periprosthetic Joint Infection?

Sarfraz A, Khury F, McCormick K, Aziz HH, Koljaka S, Rozell JC, Schwarzkopf R, Aggarwal VK

BACKGROUND: Prolonged retention of antibiotic-loaded articulating spacers after the first stage of a two-stage revision for periprosthetic joint infection (PJI) can occur because of patient preference, surgeon preference, or medical reasons that prevent the planned second stage. Little is known about the frequency of persistent infections, mechanical complications, and functional results in patients with retained spacers.

QUESTIONS/PURPOSES: At a minimum follow-up of 2 years after spacer placement, among patients who do not undergo the second stage revision (replacement of the spacer with a definitive prosthesis of the hip or knee): (1) What was the survival of the spacer free from unplanned reoperation or removal, the cumulative incidence of symptomatic infection, and the overall (Kaplan-Meier) survivorship of the patients? (2) What is the cumulative incidence of mechanical complications (spacer fracture or dislocation)? (3) What is the ambulatory status of patients who have retained their spacers?

METHODS: Between March 2011 and July 2023, a total of 111 and 152 patients underwent first-stage revision with an articulating spacer placement as part of a planned two-stage procedure for chronic PJI after THA and TKA at our institution, respectively. Of these, 21% (23 of 111) in the THA group and 24% (37 of 152) in the TKA group did not undergo the anticipated second-stage reimp [...]

[PubMed](#) · [DOI](#)

CORR Insights®: What Is the Fate of Retained Antibiotic Spacers After First-stage Revision for Periprosthetic Joint Infection?

Hartman CW

[PubMed](#) · [DOI](#)

Can a Targeted Dissemination Strategy of the PREPARE Trial Results Change Surgical Antiseptic Practice?

Nesbit SE, Hong I, O'Hara NN, Sprague S, Slobogean GP, Levack AE

BACKGROUND: The PREPARE trial (Pragmatic Randomized trial Evaluating Pre-operative Alcohol skin solutions in Fractured Extremities), published in February 2024, demonstrated that 0.7% iodine povacrylex in 74% isopropyl alcohol reduced surgical site infections in patients with closed fractures compared with chlorhexidine gluconate in 70% isopropyl alcohol. Prior studies have shown that the translation of clinical trial findings into routine clinical practice and surgical procedures frequently takes longer than a decade, and strategies to accelerate implementation in orthopaedic surgery remain poorly defined. Engaging in a multifaceted implementation strategy with minimal barriers to entry for providers may be able to reduce this time to widespread adoption.

QUESTIONS/PURPOSES: (1) Can the low-cost single-center dissemination efforts of the PREPARE trial results change antiseptic usage? (2) If a true practice change occurs, can the change be sustained over time without additional dissemination efforts? (3) Does the adoption of new evidence vary between fellowship-trained orthopaedic trauma surgeons and orthopaedic trauma surgeons who are not fellowship-trained? (4) What is the economic impact of a practice change to iodine povacrylex on hospital purchasing costs?

METHODS: This was a retrospective, historically controlled comparative study of patients who underwent operative tre [...]

[PubMed](#) · [DOI](#)

CORR Insights®: Can a Targeted Dissemination Strategy of the PREPARE Trial Results Change Surgical Antiseptic Practice?

Ring DC

[PubMed](#) · [DOI](#)

Do Published Orthopaedic RCTs Match Their Registered Protocols? A 6-year Analysis of Leading Orthopaedic Journals.

Poursalehian M, Sahebi M, Tajvidi M, Sabaghian A, Asgari AM, Tabaie SA, Hoveidaei AH

BACKGROUND: RCTs are a key block in the evidence pyramid, but their quality relies on detailed, consistent reporting, and one best-practice standard is prospective registration. Prospective trial registration was intended to reduce publication bias, and adherence to a prespecified protocol helps limit bias from selective reporting; any protocol or end point changes should be transparently documented and justified. However, the degree to which articles published in the leading journals on orthopaedic surgery comply with this best-practice standard has, to our knowledge, not been evaluated.

QUESTIONS/PURPOSES: (1) Do RCTs published in leading, general-interest orthopaedic surgery journals comply with best practices regarding prospective clinical trial registration? (2) Do major discrepancies exist between registered protocols and published orthopaedic RCTs? (3) Are there specific study types that are more likely to demonstrate discrepancies?

METHODS: A review was performed on RCTs published in the top five general-interest orthopaedic surgery journals, based on the 2022 scientific journal rankings (from SciMago): Journal of Bone and Joint Surgery , Bone and Joint Journal , Clinical Orthopaedics and Related Research ®, Journal of the American Academy of Orthopaedic Surgeons (JAAOS) , and Acta Orthopaedica . During the study period, all journals maintained editorial policies requ [...]

[PubMed](#) · [DOI](#)

CORR Insights®: Do Published Orthopaedic RCTs Match Their Registered Protocols? A 6-year Analysis of Leading Orthopaedic Journals.

Woo EJ

[PubMed](#) · [DOI](#)

Do Personalized Recovery Trajectories Enhance the Utility of Patient-reported Outcome Measures After Orthopaedic Surgery?

Graber J, Jin X, Juarez-Colunga E, Dibblee P, Woodfield D, Minick KI, Chamberlain AM, Bade M et al.

BACKGROUND: Patient-reported outcome measures (PROMs) could be useful tools for supporting patient care after orthopaedic surgery, but they can be difficult to interpret and apply in practice. Existing methods for PROM interpretation, such as meaningful change values and population norms, have practical limitations and are unresponsive to patient-specific factors that influence recovery. Personalized recovery trajectories could help overcome these limitations, making PROM scores more clinically meaningful and actionable.

QUESTIONS/PURPOSES: (1) Does a "people-like-me" modeling approach to crafting personalized recovery trajectories provide accurate and precise trajectories of PROM recovery early after total shoulder arthroplasty (TSA)? (2) What advantages do personalized recovery trajectories provide over alternative methods for PROM interpretation (minimum detectable change, minimum clinically important difference, and normative values) when applied to individual patients?

METHODS: We used data from 28 outpatient physical therapy clinics within a single integrated health system from 2017 to 2022. We included patients who underwent unilateral anatomic or reverse TSA and had postoperative data available from the DASH questionnaire. We excluded patients who were missing key modeling variables, who had fewer than three postoperative DASH assessments, or who lacked a DASH assessm [...]

[PubMed](#) · [DOI](#)

CORR Insights®: Do Personalized Recovery Trajectories Enhance the Utility of Patient-reported Outcome Measures After Orthopaedic Surgery?

Navarro RA

[PubMed](#) · [DOI](#)

Does Liposomal Bupivacaine Provide Superior Postoperative Analgesia Compared With Ropivacaine With Dexamethasone in Suprainguinal Fascia Iliaca Compartment Block for THA? A Randomized Controlled Trial.

Wu W, Hu X, Yang C, Sun Y, Li J, Wang L

BACKGROUND: Choosing the most effective pain management is important for enhancing functional recovery after surgery. The fascia iliaca compartment block (FICB) is recommended for analgesia after THA, but whether liposomal bupivacaine is more effective for this application remains uncertain.

QUESTIONS/PURPOSES: (1) Did a single-injection suprainguinal FICB using liposomal bupivacaine provide superior pain control by a clinically important margin? (2) What was the effect of liposomal bupivacaine on perioperative opioid consumption? (3) How did liposomal bupivacaine affect the duration of the sensory block and quadriceps motor weakness in the operated limb?

METHODS: This was a prospective, parallel-group, assessor-blinded, single-center, randomized controlled clinical trial. Patients who underwent unilateral primary THA via the posterolateral approach were included in the study. Exclusion criteria included a BMI of > 32.5 kg/m² (which is the diagnostic criterion for moderate obesity in China), contraindications to peripheral nerve block, known allergy to study medications, psychiatric or cognitive disorders that could interfere with pain assessment, severe hepatic or renal dysfunction, and chronic pain necessitating long-term opioid therapy. According to these criteria, 76% (60 of 79) of screened patients met eligibility requirements and were subsequently randomized into either [...]

[PubMed](#) · [DOI](#)

CORR Insights®: Does Liposomal Bupivacaine Provide Superior Postoperative Analgesia Compared With Ropivacaine With Dexamethasone in Suprainguinal Fascia Iliaca Compartment Block for THA? A Randomized Controlled Trial.

Sadeghirad B

[PubMed](#) · [DOI](#)

Is Functional Reconstruction Feasible With Modified Hip Transposition Using a Customized 3D-printed Femoral Prosthesis After Pelvic Tumor Resection? A Preliminary Study.

Wang H, Shen J, Zuo D, Liu K, Sun W

BACKGROUND: Functional reconstruction after resection of malignant pelvic tumors involving zones I + II ± IV remains a major challenge in orthopaedic oncology. Conventional hip transposition can reduce prosthesis-related complications, but it is often associated with limb shortening and femoral head malrotation. We propose a modified hip transposition technique-femoral lengthening and retroversion hip transposition with a customized three-dimensionally (3D) printed femoral prosthesis-to address deficiencies in limb length, femoral head positioning, and fixation stability, and we evaluate its effectiveness in reducing complications and improving functional outcomes.

QUESTIONS/PURPOSES: In the context of a small, initial patient series, we asked: (1) What was the postoperative functional outcome, as assessed by Musculoskeletal Tumor Society 1993 (MSTS-93) score, after reconstruction using this technique? (2) What were the frequency and nature of complications associated with the method? (3) How well was limb length restored at a minimum follow-up of 3 years?

METHODS: Between January 2019 and December 2021, a total of 49 patients underwent resection and reconstruction for pelvic zone I + II ± IV tumors. Of these, 18% (9) received the modified hip transposition with a customized 3D-printed femoral lengthening and retroversion prosthesis. This approach was selected for patients in [...]

[PubMed](#) · [DOI](#)

CORR Insights®: Is Functional Reconstruction Feasible With Modified Hip Transposition Using a Customized 3D-printed Femoral Prosthesis After Pelvic Tumor Resection? A Preliminary Study.

Potter BK

[PubMed](#) · [DOI](#)

Does Aspiration With Culture Augmented With Percutaneous Core Biopsy Analysis Have Better Diagnostic Value Than Culture Alone for PJI in Large Joints? A Prospective Enhancement and Feasibility Study.

Borucki RJ, Leutloff D, Labs K

BACKGROUND: Existing diagnostic approaches in periprosthetic joint infections (PJIs) are primarily based on joint aspiration, with a reported sensitivity of 40% to 86%. "Dry taps," occurring in up to 40% of aspirations, reduce diagnostic accuracy. Core biopsies, which enable minimally invasive tissue sampling, are widely used across various medical specialties and represent a promising tool for application in orthopaedics.

QUESTION/PURPOSE: Can the inclusion of core biopsy analysis, as opposed to a sole joint aspiration, enhance diagnostic accuracy for PJI in large joint (hip, knee, and shoulder) arthroplasties?

METHODS: In this prospective enhancement study, 175 patients with painful large arthroplasties were recruited between September 2020 and September 2023. All patients underwent joint aspiration and core biopsy, leading to 181 primary data sets, of which 111 were excluded from final analysis for not reaching the endpoint or because of unrelated death, protocol breach, or a dry tap, which occurred in 34% (62 of 181) of examined joints. Diagnostic accuracy was assessed for sensitivity and specificity against the 2018 International Consensus Meeting (ICM) criteria, constituting the primary endpoint. Subsequent revision surgery, septic or aseptic, along with the microbiological evaluation of tissue samples and the sonicated prosthesis served as the confirmatory endpoint.

R [...]

[PubMed](#) · [DOI](#)

S1P Modulation via S1P Lyase Inhibition Enhances Bone Regeneration in a Murine Critical-size Defect Model of Atrophic Nonunion.

Reinkemeier F, Korte R, Bakri A, Wallner C, Drysch M, Schmidt SV, Fueth M, Puszcz F et al.

BACKGROUND: Atrophic nonunion is a challenging complication of fracture healing with few noninvasive therapeutic options. Given that sphingosine-1-phosphate (S1P) regulates bone remodeling and angiogenesis, systemic S1P modulation may offer a regenerative strategy; however, its efficacy in nonunion has not been established.

QUESTIONS/PURPOSES: We therefore asked whether S1P lyase inhibition (1) enhances osteogenesis and consolidation in a murine critical-size bone defect model of atrophic nonunion, (2) modulates bone remodeling (osteoblast differentiation and osteoclast activity), and (3) augments cellular proliferation and endothelial activity, and (4) whether quantitative micro-CT corroborates histologically observed improvements.

METHODS: In a validated murine critical-size bone defect model of atrophic nonunion, 12-week-old male and female C57BL/6J mice were treated by continuous systemic administration of the S1P lyase inhibitor 4-deoxypyridoxine (DOP) for 5 weeks, while a reference group remained untreated after nonunion induction. Bone regeneration was assessed by histology and immunohistochemistry to evaluate osteoblast differentiation, osteoclast activity, cellular proliferation, and angiogenesis (n = 7 per group) and by high-resolution micro-CT to quantify newly formed bone within the defect region (n = 6 per group).

RESULTS: DOP treatment resulted in increased min [...]

[PubMed](#) · [DOI](#)

CORR Insights®: S1P Modulation via S1P Lyase Inhibition Enhances Bone Regeneration in a Murine Critical-size Defect Model of Atrophic Nonunion.

Tatara AM

[PubMed](#) · [DOI](#)

Letter to the Editor: Is Displacement of the Lesser Trochanter Associated With Functional Outcome in Older Adults With Intertrochanteric Fractures?

Qin K, Xu Y, Liang W

[PubMed](#) · [DOI](#)

Reply to the Letter to the Editor: Is Displacement of the Lesser Trochanter Associated With Functional Outcome in Older Adults With Intertrochanteric Fractures?

Ratanasermsub N, Amarase C, Tantavisut S, Sirichuchnin P

[PubMed](#) · [DOI](#)

Letter to the Editor: All-inside ACL Reconstruction Offers No Advantage in Clinical Outcomes, Graft Healing, or Tunnel Widening Compared With the Complete Tibial Tunnel Technique: A Prospective Randomized Trial.

Lu J, Jiang K, Du X, Ji J

[PubMed](#) · [DOI](#)

Reply to the Letter to the Editor: All-inside ACL Reconstruction Offers No Advantage in Clinical Outcomes, Graft Healing, or Tunnel Widening Compared With the Complete Tibial Tunnel Technique: A Prospective Randomized Trial.

Okutan AE, Gürün E

[PubMed](#) · [DOI](#)

Letter to the Editor: Patients With Nonanaphylactic Penicillin Allergy Are Not at Increased Risk of Allergic Events After Receiving Prophylactic Preoperative Cefazolin Prior to Closed Fracture Repair.

Kocak S

[PubMed](#) · [DOI](#)

Letter to the Editor: Does Insertional Reattachment for Acute Sleeve Avulsion Fracture of the Achilles Tendon Provide Sustained Function and Sports Participation?

Spiezia F, Oliva F, Maffulli N

[PubMed](#) · [DOI](#)

Reply to the Letter to the Editor: Does Insertional Reattachment for Acute Sleeve Avulsion Fracture of Achilles Tendon Provide Sustained Function and Sports Participation?

Xiong S, Xie X, Guo Q

[PubMed](#) · [DOI](#)

Letter to the Editor: Equity360: Gender, Race, and Ethnicity: Sex and Fairness in Sports.

Lichtblau E

[PubMed](#) · [DOI](#)

Reply to the Letter to the Editor: Equity360: Gender, Race, and Ethnicity: Sex and Fairness in Sports.

Jain R

[PubMed](#) · [DOI](#)

Reply to the Letter to the Editor: Equity360: Gender, Race, and Ethnicity: Sex and Fairness in Sports.

O'Connor MI

[PubMed](#) · [DOI](#)



JBJS (American)

Volume 108, Issue 15 · August 2026 · 16 new articles

Introducing the JBJS Ambassadors.

Bhandari M

[PubMed](#) · [DOI](#)

Calming the Mind to Heal the Hip: Commentary on an article by Boyi Jiang, MD, et al.: " Significant Anxiolytic Effect and Enhanced Recovery Benefits of Perioperative Low-Dose Olanzapine in Patients with Anxiety Undergoing THA. A Randomized Controlled Trial ".

Wyles CC

[PubMed](#) · [DOI](#)

Large Language Models Should Be the New Gold Standard for Orthopaedic Surgical Registry Curation, but Likely Will Not Be without Clearer Direction: Commentary on an article by Linjun Yang, PhD, et al.: " Comparison of Large Language Models with Rules-Based Natural Language Processing Algorithms for Extracting Data from Operative Notes ".

Yang AJ, Ramkumar PN, Woo JJ

[PubMed](#) · [DOI](#)

Patient-Reported Outcome Thresholds for Hip and Knee Replacements: Commentary on an article by Zhaorui Wang, MD, et al.: "CMS-Proposed Substantial Clinical Benefit Thresholds

Correlate with Patient-Reported Measures After Primary Total Joint Arthroplasty. Improvement, Satisfaction, and Willingness to Repeat Surgery ".

Valsamis EM, Grammatopoulos G

[PubMed](#) · [DOI](#)

Magnetic Resonance Imaging to Diagnose Carpal Tunnel Syndrome: When Size Does Matter: Commentary on an article by Wen-Chih Liu, MD, PhD, et al.: " MRI Assessment of Median Nerve Size in Patients with Proximate Electrodiagnostic Studies ".

Slater RR

[PubMed](#) · [DOI](#)

Fertility Preservation Option Accommodations and Education in Orthopaedic Surgery Training: Commentary on an article by Grace Q. Chen, BS, et al.: " Oocyte Cryopreservation Experiences and Attitudes Among Female Orthopaedic Surgeons ".

Agel J

[PubMed](#) · [DOI](#)

The Extended Lesser Trochanteric Osteotomy (ELTO) in Complex Revision Hip Arthroplasty: An Alternative to the Extended Trochanteric Osteotomy (ETO).

Farid YR

The extended trochanteric osteotomy (ETO) has been a standard technique in complex revision hip arthroplasty. Despite the satisfactory healing rates with ETO, nonunion may occur, and failure of fixation and displacement of the trochanter may compromise abductor strength. The extended lesser trochanteric osteotomy (ELTO) is an alternative to the ETO in select patients that allows medial access to 3 sides of the femoral stem and incorporates the lesser trochanter along with a variable length of the medial femoral cortex, to which the iliopsoas tendon and adductors remain attached. The ELTO might be easier, faster, and shorter compared with the ETO while offering unobstructed exposure of 3 sides of the femoral stem. The most substantial advantage is the ability to spare the abductor attachment, eliminating the need for postoperative protected weight-bearing and abduction braces.

[PubMed](#) · [DOI](#)

The Persistent Challenges of Diagnosing Orthopaedic Implant-Related Infections.

Lum ZC, Cohen-Rosenblum A, Yao JJ, Chen AF, Landy DC, Parvizi J

Infection remains one of the most catastrophic complications following orthopaedic surgery. Despite substantial advances in molecular diagnostics, biomarker assays, and consensus definitions, accurately diagnosing orthopaedic infection continues to challenge even the most experienced clinicians. There are differences in the diagnosis and treatment of infections that are related to different anatomic regions. The difficulty arises from the inherent biological diversity of infecting organisms and surgical locations, variable host responses, and the absence of a true diagnostic "gold standard." This article summarizes the current diagnostic challenges and emerging solutions, drawing on recent high-impact evidence and consensus frameworks.

[PubMed](#) · [DOI](#)

Humility: The Essential Virtue Enhancing Patient Care, Education, Research, and Innovation.

Rizzo M, Lin I, Murray P, Sotereanos D, Stern P

The word humility is derived from the Latin word humus, meaning "ground" or "soil." Romans equated humility with being grounded. Perhaps because it contradicts our natural tendencies toward self-preservation, humility remains an often misunderstood and underappreciated virtue. It is foreign to many, especially in today's society, which is so focused on self-promotion. However, it is truly one of the most important and powerful virtues that a person can possess and practice. Humility is a just assessment of ourselves and the world around us. It is not thinking less of oneself; rather, it is actually thinking less about oneself. In looking beyond ourselves, we can obtain an honest

estimate of our limitations and what is needed to remain patient-centered. In approaching our profession with humility, we more closely honor the privileges of caring for patients, teaching others, innovating, and advancing science. The aim of this article is to show how humility nurtures our development in all aspects of medicine. A humble clinician maintains a steadfast focus on their patient, a humble educator emphasizes the learner, a humble researcher respects science through their honesty and integrity, and a humble innovator does not let pride blind them, recognizing that innovation without humility is doomed to fail. In addition to being a superpower in medicine, humility blesses us with the wis [...]

[PubMed](#) · [DOI](#)

My Lucky Break: A Rare Bone Condition, A Sudden Fracture, and the Right Team.

Sando N, Ahn J

[PubMed](#) · [DOI](#)

Heading Toward a Different Future: The Microbiome, Dysbiosis, Microbial Translocation, and Beyond.

He M, Ferrini A, Chisari E, Kolhoff F, Sehgal P, Parvizi J

➤ The relationship between the gastrointestinal tract and joint diseases has garnered increased attention over recent decades, leading to the introduction of the gut-joint axis concept. ➤ Infections at sites such as joints and the spine may originate endogenously from the gut microbiome. ➤ The idea of microbial translocation through a compromised epithelial barrier, resulting in the circulation of pathogens or their byproducts, and the idea of immune cell-mediated transport of pathogens to various sites are gaining further attention. ➤ By understanding the interaction between the immune system and gut microbiota, potential therapeutic strategies, such as the use of organoids, can be developed to restore the gut barrier integrity, to replenish gut microbiota, and to provide biodiversity. ➤ To better understand the mechanisms linking gut health and joint diseases, future basic-science research and well-designed clinical trials, exploiting advanced next-generation sequencing techniques, are needed.

[PubMed](#) · [DOI](#)

Significant Anxiolytic Effect and Enhanced Recovery Benefits of Perioperative Low-Dose Olanzapine in Patients with Anxiety Undergoing THA: A Randomized Controlled Trial.

Jiang B, Fan J, Lai Y, Cai Y, Ding Z, Luo Z, Zhou Z

BACKGROUND: Major anxiety symptoms are commonly observed in patients undergoing total hip arthroplasty (THA); these symptoms exacerbate pain and compromise hip recovery. Olanzapine demonstrates clinically meaningful efficacy in reducing anxiety; thus, we investigated its anxiolytic effect and benefits for enhancing recovery in these high-risk patients.

METHODS: We prospectively enrolled 135 patients who were scheduled for primary THA at our institution between April 2024 and March 2025 and who scored at least 40 points on the State-Trait Anxiety Inventory-State (STAI-S) before surgery. Patients randomly received oral olanzapine (2.5 mg), alprazolam (0.4 mg), or a placebo once nightly for 5 days beginning on the day of admission. The 3 groups (45 patients in the olanzapine group, 45 in the alprazolam group, and 44 in the placebo group after 1 patient was lost to follow-up) were compared postoperatively in terms of the STAI-S score, Pittsburgh Sleep Quality Index (PSQI), visual analog scale (VAS) pain score, opioid consumption, and functional recovery of the hip. Adverse events related to drugs and surgery were recorded.

RESULTS: Compared with placebo and alprazolam, olanzapine was associated with significantly lower STAI-S scores on postoperative days (PODs) 1 and 3, significantly lower resting VAS pain scores on PODs 1 to 3, and significantly lower incidence of postoperative [...]

[PubMed](#) · [DOI](#)

Comparison of Large Language Models with Rules-Based Natural Language Processing Algorithms for Extracting Data from Operative Notes.

Yang L, Mulford KL, Girod-Hoffman MM, Khela M, Khosravi A, Crossman DM, Kanabar A, Saniei S et al.

BACKGROUND: We aimed to develop automated data extraction pipelines with large language models (LLMs) to extract registry data from total hip arthroplasty (THA) operative notes and compare the performance with that of existing natural language processing (NLP) algorithms.

METHODS: We randomly sampled 1,000 primary THA cases from our institutional registry. Two human annotators manually reviewed each operative note for 3 data points: surgical approach, bearing surface, and fixation technique. All labeled THA notes were split into the development set (n = 239) and the testing set (n = 719). We developed a custom data extraction pipeline for each data point by combining an iteratively customized prompt with an LLM. The performance was compared with that of existing rules-based NLP algorithms.

RESULTS: The accuracy of LLMs was superior to that of NLP algorithms for all data points: surgical approach (96% compared with 94%), bearing surface (89% compared with 74%), and fixation technique (96% compared with 95%). Furthermore, the LLM accurately inferred the bearing surface for 80% of the notes that were ambiguous about the bearing surface.

CONCLUSIONS: We developed LLM pipelines for extracting 3 registry-relevant data points from THA operative notes, demonstrating superior performance to existing NLP algorithms.

CLINICAL RELEVANCE: LLMs have the potential to impact clinical care, [...]

[PubMed](#) · [DOI](#)

CMS-Proposed Substantial Clinical Benefit Thresholds Correlate with Patient-Reported Measures After Primary Total Joint Arthroplasty: Improvement, Satisfaction, and Willingness to Repeat Surgery.

Wang Z, Janney CA, Guo E, Carender CN, Hallstrom BR, Kheir MM

BACKGROUND: The U.S. Centers for Medicare & Medicaid Services (CMS) requires the collection of patient-reported outcome measures (PROMs) after primary total joint arthroplasty (TJA), with penalties for noncompliance affecting all Medicare reimbursement. The CMS will publish risk-standardized improvement rates based on substantial clinical benefit (SCB) thresholds of 22 points for the HOOS JR (Hip disability and Osteoarthritis Outcome Score for Joint Replacement) and 20 points for the KOOS JR (Knee injury and Osteoarthritis Outcome Score for Joint Replacement). Our aims were to determine if preoperative scores predicted postoperative PROMs, to externally validate the SCB thresholds, and to analyze them with different anchors.

METHODS: We retrospectively identified patients who underwent TJA at our institution between 2015 and 2023. The HOOS JR and KOOS JR were prospectively collected in the Michigan Arthroplasty Registry Collaborative Quality Initiative (MARCQI); all other variables were obtained from electronic medical records. Logistic regression analyses and anchor-based receiver operating characteristic curves were generated to determine threshold values and the efficacy of using preoperative and postoperative scores as predictors of patient improvement, satisfaction, and willingness to repeat surgery.

RESULTS: In total, 3,465 cases (1,498 total knee arthroplasties [TKAs] [...]

[PubMed](#) · [DOI](#)

MRI Assessment of Median Nerve Size in Patients with Proximate Electrodiagnostic Studies.

Liu WC, Wung CH, Simeone FJ, Eberlin KR, Chen NC

BACKGROUND: Carpal tunnel syndrome (CTS) diagnosis has traditionally relied on electrodiagnosis (EDX) to confirm the diagnosis and to assess severity. Ultrasound has shown potential in measuring median nerve cross-sectional area (CSA) for CTS diagnosis, and magnetic resonance imaging (MRI) can be used for wrist soft-tissue evaluation. This study explored the correlation between CTS diagnosis and median nerve CSA measured on MRI at different wrist levels.

METHODS: A retrospective review of an electronic medical record database identified patients who underwent both wrist MRI and EDX within a 90-day interval between January 2000 and December 2022. Median nerve CSA was measured on axial T2-weighted images at 3 levels: proximal to the carpal tunnel inlet (the distal radioulnar joint [DRUJ]), the inlet, and the outlet. Continuous variables are presented as means $\pm$ standard deviations. A logistic regression model was constructed to evaluate the diagnostic accuracy of median nerve CSA, at the 3 anatomical levels, in identifying CTS. Empirical cut point estimation determined optimal cutoffs and corresponding areas under the receiver operating characteristic curve (AUCs).

RESULTS: Sixty-eight patients (76 wrists; mean age, 51.4 ± 14.2 years; male-to-female ratio, 26 to 50; 59 White patients, 8 Hispanic patients, and 1 Asian patient) were included. The mean median nerve CSA in the EDX- [...]

[PubMed](#) · [DOI](#)

Oocyte Cryopreservation Experiences and Attitudes Among Female Orthopaedic Surgeons.

Chen GQ, Seidenstein AH, LaPorte DM

BACKGROUND: Female surgeons more commonly delay childbearing and experience higher rates of infertility than women in the general population. More women are entering orthopaedic surgery but face unique challenges in family building. The accessibility of fertility preservation strategies among female orthopaedic surgeons remains underexplored. We aimed to investigate facilitators of and barriers to oocyte cryopreservation, especially during orthopaedic residency.

METHODS: A survey regarding family planning and experiences surrounding oocyte cryopreservation was designed and distributed via residency program directors, regional orthopaedic societies, and the Ruth Jackson Orthopaedic Society. Female orthopaedic surgery residents, fellows, and attending physicians were invited to participate.

RESULTS: Of 169 participants, 107 (63%) reported intentionally delaying childbearing and 58 (34%) reported delaying or planning to delay for ≥ 4 years. Although 91 respondents (54%) had considered oocyte cryopreservation, only 36 (21%) had undergone or planned to undergo at least 1 cycle and 55 (33%) ultimately decided not to undergo it. Inflexible scheduling of work, lack of insurance coverage, and restrictive leave policy were the most important barriers to oocyte cryopreservation during residency. Among current residents and fellows, 30 (38%) would not have been comfortable telling program [...]

[PubMed](#) · [DOI](#)



The Journal of Arthroplasty

Volume 41, Issue 8 · August 2026 · 53 new articles

Letter to the Editor on "Patient-Specific Instrumentation versus Standard of Care in Total Knee Arthroplasty in an Obese Population: Minimum Five-Year Follow-Up".

Kocak S

[PubMed](#) · [DOI](#)

Response to Letter to the Editor on "Patient-Specific Instrumentation versus Standard of Care in Total Knee Arthroplasty in an Obese Population: Minimum 5-year Follow-Up".

Braun S, Allen J, Somerville L, Howard JL, Lanting BA, Vasarhelyi EM

[PubMed](#) · [DOI](#)

Letter regarding "Optimizing Postoperative Pain Management in Total Knee Arthroplasty With Preoperative Methadone: A Prospective Randomized Study".

Maricar MA, Tariq M

[PubMed](#) · [DOI](#)

Reply to Letter to the Editor on "Optimizing Postoperative Pain Management in Total Knee Arthroplasty With Preoperative Methadone: A Prospective Randomized Study".

Whitaker JE, Malkani AL

[PubMed](#) · [DOI](#)

Reply to Letter to the Editor On "Preserving Scientific Integrity in Academic Publishing: Navigating Artificial Intelligence, Journal Policies, and the Impact Factor as a Quality

Indicator".

Matsubara S

[PubMed](#) · [DOI](#)

Reply to Letter to the Editor On "Preserving Scientific Integrity in Academic Publishing: Navigating Artificial Intelligence, Journal Policies, and the Impact Factor as a Quality Indicator".

Mont MA, Smitterberg CW, Ayeni O, Becker R, Herbst E, Heyse T, Hirschmann MT, Karlsson J et al.

[PubMed](#) · [DOI](#)

Letter to the Editor Regarding: "Synovial Absolute Neutrophil Count as a Superior Diagnostic Biomarker for Acute Periprosthetic Joint Infection in Hip and Knee Arthroplasty With Low-Virulence Organisms".

Zhang L

[PubMed](#) · [DOI](#)

Reply to Letter to the Editor on "Synovial Absolute Neutrophil Count as a Superior Diagnostic Biomarker for Acute Periprosthetic Joint Infection in Hip and Knee Arthroplasty With Low-Virulence Organisms".

Karlidag T, Citak M

[PubMed](#) · [DOI](#)

Reconsidering Centers for Medicare & Medicaid Services Functional Status Measures for Total Hip and Knee Arthroplasty: From Process Compliance to Meaningful Outcome Assessment.

Chen AF, Della Valle CJ, Bostrom MPG, Mont MA

[PubMed](#) · [DOI](#)

American Association of Hip and Knee Surgeons Position Statement on Opioid Use for the Treatment of Osteoarthritis of the Hip and Knee.

Nandi S, Srivastava AK, Hannon CP

[PubMed](#) · [DOI](#)

The 2026 Centers for Medicare & Medicaid Services Physician Fee Schedule Final Rule: Understanding the 7% Cut to Procedural Reimbursement.

LeBrun DG, Bradley AT, Frisch NB, Rana AJ

[PubMed](#) · [DOI](#)

Androgen Awareness: Navigating Testosterone Replacement Therapy in Total Joint Arthroplasty.

Ganz MP, Mont MA, Scuderi GR

[PubMed](#) · [DOI](#)

Preoperative Testosterone Replacement Therapy Is Associated With Increased Complication Risk After Total Hip Arthroplasty: A Propensity-Matched Analysis of Real-World Data.

Omurzakov AM, Omurzakov A, Bhatti P, Debbi EM, Gausden EB, Chalmers BP

BACKGROUND: While testosterone replacement therapy (TRT) is known to affect cardiovascular physiology, its impact on outcomes following total hip arthroplasty (THA) remains unclear. This study aimed to assess whether preoperative TRT use is associated with increased complications following THA.

METHODS: A retrospective cohort study using a large national database was performed. Adult patients undergoing primary THA before February 2020 with 5-year follow-up were stratified based on preoperative TRT use within one year of surgery. Patients who had a history of septic arthritis, osteonecrosis, or pathologic fractures were excluded. Propensity score matching (1:1) was used to balance cohorts. Outcomes included medical complications at 90 days and one year and prosthetic complications up to five years postoperatively. Following matching, 3,953 patients were included in each cohort.

RESULTS: At 90 days, TRT use was associated with higher rates of deep vein thrombosis (2.8 versus 2.0%; $P = 0.023$), pneumonia (3.2 versus 1.7%; $P < 0.001$), and sepsis (4.2 versus 0.9%; $P < 0.001$). At one year, TRT users had increased rates of deep vein thrombosis (4.3 versus 3.0%; $P = 0.002$), cardiac events (2.9 versus 1.7%; $P < 0.001$), pneumonia (6.1 versus 3.4%; $P < 0.001$), acute kidney injury (7.6 versus 5.6%; $P < 0.001$), and sepsis (3.8 versus 1.9%; $P < 0.001$). At five years, TRT users had higher r [...]

[PubMed](#) · [DOI](#)

Surgeon Volume and Experience Both Independently Drive Value in Primary Total Joint Arthroplasty: Findings From a Patient-Level Value Analysis of 4,989 Procedures With Time-Driven Activity-Based Costing.

Sauder N, Peterson SL, Sharma A, Lim PL, Blank DH, Bedair HS, Melnic CM

BACKGROUND: Surgeon and surgeon experience may influence clinical outcomes in both total knee arthroplasty (TKA) and total hip arthroplasty (THA). Clinical outcomes can be integrated with costs to calculate value. We aimed to determine if surgeon experience and volume were associated with increased value in arthroplasty.

METHODS: We retrospectively sourced 2,786 primary TKAs and 2,203 primary THAs from a prospectively maintained multi-institutional database. A total of 37 surgeons performed the procedures. Knee Injury and Hip Disability Osteoarthritis Outcome Score-Physical Function Short Forms (HOOS-PS) were collected. Costs of care were determined using Time-Driven Activity-Based Costing and converted from US dollars to arbitrary cost units (CUs). Value was calculated as the quotient of the 1-year change in HOOS-PS or Knee Injury Osteoarthritis Outcome Score-Physical Function Short Forms and the costs of care. We determined if surgeon volume and experience correlated with value after adjusting for confounders. Stratum-specific likelihood ratio analysis determined meaningful thresholds in any associations.

RESULTS: In THA, increased Value HOOS-PS was independently associated with both surgeon annual THA volume ($P = 0.001$) and surgeon years of experience ($P < 0.001$). The stratum-specific likelihood ratio analysis revealed thresholds in volume (≥ 61 annual THAs) and experience [...]

[PubMed](#) · [DOI](#)

Medicare Reimbursement Trends for Common Knee Arthroplasty Procedures: A Generational Perspective, 2006 to 2024.

Anoushiravani AA, Hoveidaei AH, Borazjani R, Katanbaf RM, Misch M, Feng JE, O'Connor CM, Nace J et al.

BACKGROUND: The primary aim of this study was to assess the trends in Medicare reimbursement for knee arthroplasty procedures between 2006 and 2024.

METHODS: This study examined payment trends for common knee arthroplasty procedures by analyzing data from the Centers for Medicare and Medicaid Services Physician Fee Schedule between 2006 and 2024. Medicare reimbursement was determined by multiplying relative value units by the Centers for Medicare and Medicaid Services conversion factor, with adjustments made for inflation to 2024 US dollars using the Consumer Price Index inflation calculator provided by the US Department of Labor. Future trend projections were generated based on average values from 2020 to 2024.

RESULTS: Between 2006 and 2024, total reimbursements for all knee procedures dropped by 2.7%, whereas the Consumer Price Index increased by 55%, leading to an inflation-adjusted average reimbursement drop of 37.2%. Primary total knee arthroplasty had the most significant drop, with reimbursements decreasing by 45.1%. Unadjusted work reimbursements across all procedures decreased by 8.5%, translating to a 40.9%

inflation-adjusted reduction. Considering the current trend, projections for 2029 indicate a further projected reduction in total surgeon reimbursement of 18.5 to 54.7%, whereas work reimbursements could decrease by 23.2 to 63.6%.

CONCLUSIONS: From 2006 to 2024 [...]

[PubMed](#) · [DOI](#)

Greater Resource Utilization for Patient-Reported Outcome Measures Collection in Medicare Inpatient Versus Outpatient Total Knee Arthroplasty: Implications for the New Centers for Medicare and Medicaid Services Patient-Reported Outcomes-Based Performance Measure Policy.

Elmenawi KA, Emara AK, Khan ST, Gudapati LS, Zhang C, Deren ME, Cleveland Clinic Adult Reconstruction Research Group, Piuze NS

BACKGROUND: The Centers for Medicare and Medicaid Services now mandates the collection and reporting of patient-reported outcome measures (PROMs) for total joint arthroplasty. However, ensuring adequate follow-up remains a challenge. Furthermore, differences in PROMs capture rates and collection strategies between inpatient and outpatient Medicare total knee arthroplasty (TKA) populations are not well understood. We aimed to (1) evaluate differences in PROMs collection processes and success rates between inpatient and outpatient Medicare TKA patients and (2) identify factors associated with requiring active follow-up to obtain 1-year PROMs.

METHODS: A prospective analysis was conducted on 7,926 Medicare patients aged ≥ 65 years who underwent elective TKA from 2016 to 2021 at a single institution. Patients were stratified as inpatient (length of stay > 1 day) or outpatient (length of stay ≤ 1 day). The PROMs at one year were collected using passive methods (email, text message, patient portal) and active methods (phone calls, personalized communication). Rates and predictors of active follow-up were compared.

RESULTS: Inpatients had lower overall 1-year PROMs completion rates than outpatients (77 versus 84%, $P < 0.001$). Passive follow-up was more successful for outpatients (52.1 versus 43.8%, $P < 0.001$), while inpatients more often required active follow-up (56.2 versus 47 [...])

[PubMed](#) · [DOI](#)

Optimizing Postoperative Care: The Role of a Total Joint Hotline in Reducing Unnecessary Emergency Department Visits.

Jaramillo M, Desai B, Daher J, Velasco-Gonzalez C, Moore WA, Chimento GF

BACKGROUND: Postoperative concerns after total joint arthroplasty (TJA) frequently lead to emergency department (ED) visits. To reduce unnecessary ED use, our institution implemented a Total Joint Hotline (TJH) to triage patient concerns. The purpose of this study was to determine if the TJH effectively reduced avoidable ED visits.

METHODS: The TJH was launched in March 2021. Patients received a wristband with a phone number, providing direct access to the adult reconstruction care team. They were instructed to call with any postoperative concerns, which were addressed by reassurance, follow-up scheduling, or ED referral as needed. We retrospectively reviewed 1,410 TJA patients between March 2020 and February 2022 (642 pre-TJH; 768 post-TJH). Primary endpoints included ED visits and 30-day readmissions. Statistical analyses included Chi-square, t-tests, and logistic regressions with Firth correction (significance set at $P < 0.05$).

RESULTS: Visit rates to the ED decreased from 5.8% pre-TJH to 3.8% post-TJH (odds ratio (OR) 0.64; 95% confidence interval (CI) 0.39 to 1.06; $P = 0.08$). Overall, 30-day readmission rates were low (0.6% pre-TJH versus 1.3% post-TJH, $P = 0.3$). Among ED presenters, readmission was more frequent after TJH implementation (34.5 versus 10.8%; adjusted OR 5.0; 95% CI 1.19 to 20.95; $P = 0.03$).

CONCLUSIONS: The implementation of the TJH was associated with a [...]

[PubMed](#) · [DOI](#)

Stairway to Heavenly Results: Do Stairs Affect Postoperative Recovery After Total Joint Arthroplasty?

Ingawa HS, Rainey JP, Kapron CR, Blackburn BE, Pelt CE, Peters CL, Gililand JM, Anderson LA et al.

BACKGROUND: A common concern for patients undergoing total hip arthroplasty (THA) or knee arthroplasty (TKA) is the presence of stairs in their home. However, there is currently no evidence to suggest how stairs affect postoperative recovery. The aim of this study was to evaluate how stairs impact patient-reported outcomes (PROs) and physical activity metrics following TKA or THA in relatively young, active patients.

METHODS: A secondary analysis of data from a multicenter, prospective study evaluated steps-per-day, flights of stairs, gait speed, and percent walking asymmetry along with the EuroQol-5 Dimension (EQ-5D) and either Hip dysfunction and Osteoarthritis Outcome Score for Joint Replacement (HOOS JR) or Knee injury and Osteoarthritis Outcome Score for Joint Replacement (KOOS JR) preoperatively and at one, three, six, and 12 months postoperatively. Generalized estimating equations were used to evaluate PROs and physical activity metrics over time in the presence or absence of stairs in the household. In total, 3,965 patients were included in the study, 1,609 undergoing THA (61% with stairs), and 2,356 undergoing TKA (53% with stairs).

RESULTS: Patients who had stairs at home climbed significantly more daily flights of stairs compared to patients who did not have stairs ($P < 0.01$). There was no difference in steps-per-day, gait speed, or percent walking asymmetry ($P > 0$ [...])

[PubMed](#) · [DOI](#)

Conversion Total Knee and Hip Arthroplasty Requires Increased Energy Expenditure Compared to Primary Knee Arthroplasty.

Burnett RA, Matheson DJ, Kapron CR, Blackburn BE, Gililand JM, Anderson LA, Pelt CE, Peters CL et al.

BACKGROUND: Conversion total knee arthroplasty (TKA) requires increased preoperative planning, surgical time, and perioperative resources; yet, there is currently no separate procedural code for conversion TKA as there is for conversion total hip arthroplasty (THA). The purpose of this study was to compare energy expenditure for the surgeon when performing primary, conversion, and revision TKA/THA.

METHODS: Surgery was performed on 179 arthroplasty patients prospectively by five surgeons. There were 77 primary TKA, 10 conversion TKA, 18 revision TKA, 58 primary THA, four conversion THA, and 12 revision THA patients. Energy expenditure of the surgeon's institution was recorded via a smart garment. Demographic variables and energy expenditure were compared between cohorts using analysis of variance tests.

RESULTS: Overall, TKA patients were similar in age, sex, and laterality. Body mass index varied among TKA groups, with conversion and primary being lower than revision TKA ($P = 0.018$). There were no differences in body mass index and sex in THA patients. The conversion THA cohort was younger than the primary and revision THA cohorts ($P = 0.015$). Conversion TKA (467.2 ± 219.4 calories) required 152% higher energy expenditure than primary TKA (308.0 ± 109.2 cal), but less than revision TKA (703.8 ± 368.0 cal; $P < 0.001$). Conversion THA (523.8 ± 117.9 cal) required 143% higher energy [...]

[PubMed](#) · [DOI](#)

The Application of Flowable Gelatin Without Thrombin in Total Joint Arthroplasty: A Propensity Score-Matched Comparative Study.

Zhang Y, Yuan J, Zhang Y, Tong P, Wu Z, He B

BACKGROUND: Flowable gelatin (FG) without thrombin is a widely utilized intraoperative hemostatic agent, yet its efficacy in total joint arthroplasty remains debated. This study aimed to assess its clinical impact on total knee arthroplasty (TKA) and total hip arthroplasty (THA).

METHODS: We conducted a retrospective analysis of patients who underwent primary TKA and THA at a single institution from 2018 to 2024, categorizing them based on intraoperative FG usage. Patients who received FG were propensity score-matched in a 2:1 ratio to those who did not, based on demographic characteristics, comorbidities, and surgical factors. Primary outcomes included blood loss, transfusion rate, thromboembolic events, and other perioperative complications. Regression analyses were conducted to assess the impact of intraoperative FG use on total blood loss in both TKA and THA. A total of 612 TKA patients (non-FG: 233; FG: 117) and 844 THA patients (non-FG: 391; FG: 203) were included, with a mean follow-up exceeding two years (up to six years).

RESULT: Both total blood loss (TKA: 899.0 versus 1,122.1 mL; THA: 1,021.8 versus 1,328.4 mL) and hidden blood loss (TKA: 789.1 versus 1,010.8 mL; THA: 858.6 versus 1,169.9 mL) were significantly reduced in the FG group, accompanied by lower transfusion rates (TKA: 4.3 versus 12.9%; THA: 8.87 versus 19.69%). The incidence of thromboembolic events, in [...]

[PubMed](#) · [DOI](#)

Trends in Utilization of Technology-Assisted Navigation in Unicompartmental and Total Knee Arthroplasty Between 2009 and 2022.

Worden JA, Fuqua A, Ross BJ, Gillis J, Heo KY, Wilson JM, Premkumar A

BACKGROUND: In cases of isolated, single-compartment osteoarthritis of the knee, both unicompartmental knee arthroplasty (UKA) and total knee arthroplasty (TKA) can be effective treatment options. This study evaluated rates of technology-assisted utilization in UKA (TA-UKA) using a national claims database over a contemporary 13-year period to determine the trends in use of TA-UKA relative to TKA in the United States.

METHODS: Patients undergoing primary UKA and TKA from 2009 to 2022 were identified using a large national claims database. Rates of computer and robotic assistance were then determined to ascertain trends in technology-assisted TKA (TA-TKA) and TA-UKA over the study period. Technology assistance was defined as computer navigation or robotics. Utilization trends were also stratified by sex, region, and age, and significance was determined using Mann-Kendall statistical tests.

RESULTS: Between 2009 and 2022, 71,396 patients underwent UKA, while 869,794 patients underwent primary TKA. The rate of TA-UKA cases increased by 17.3% from 2009 to 2022 ($P < 0.001$), while rates of TA-TKA procedures increased by 10.8% ($P < 0.001$). Higher volumes of TA-UKA cases were performed in urban areas and on women. By 2022, 20.6% of all UKA cases and 17.6% of all TKA cases were performed with technology.

CONCLUSIONS: While a significant increase in the rates of both TA-UKA and TA-TKA [...]

[PubMed](#) · [DOI](#)

Age-Related Risk of Conversion to Total Knee Arthroplasty After Mobile-Bearing Unicompartmental Knee Arthroplasty: A Survival and Competing Risk Analysis.

Tai TW, Restrepo DJ, Guarin Perez SF, Good KM, Sierra RJ

BACKGROUND: Mobile-bearing unicompartmental knee arthroplasty (UKA) is a well-established treatment for isolated compartment osteoarthritis. However, long-term outcomes may vary with the age of the patient at arthroplasty. This study evaluated the association between age and the risk of conversion to total knee arthroplasty (TKA) and the ultimate outcome for patients undergoing conversion from UKA to TKA.

METHODS: A consecutive series of 468 patients who underwent mobile-bearing UKA between December 2006 and June 2023 were prospectively followed. Patients were stratified by age: ≤ 60 years ($n = 180$) and > 60 years ($n = 288$). The mean follow-up was 10.7 years (range, two to 18). Survivorship free from TKA conversion was analyzed using competing risk methods, with death as a competing event. Multivariate Cox regressions identified independent predictors of conversion. Outcomes following conversion to TKA were also evaluated.

RESULTS: The overall rate of conversion to TKA was 7.5% for the entire cohort. Competing risk analysis revealed cumulative incidence of TKA conversion at two, five, 10, and 15 years was 4.4, 5.6, 10.2, and 15.0% in the younger group, compared to 1.0, 3.2, 5.1, and 6.0% in the older age group ($P = 0.009$). Multivariate analyses identified younger age (hazard ratio 2.2; 95% confidence interval, 1.1 to 4.6; $P = 0.029$) and operative time (hazard ratio 1.01; $P = [...]$

[PubMed](#) · [DOI](#)

Development and Validation of a Nomogram for Predicting High-Risk Hidden Blood Loss After Total Knee Arthroplasty.

Huang Y, Weng J, Lin R, Liu Q, Ye X, Liang P, Luo X, Zhan C et al.

BACKGROUND: Hidden blood loss (HBL) accounts for 49% of total postoperative blood loss after total knee arthroplasty (TKA), predisposing patients to anemia, infection, and delayed functional recovery. Despite partial elucidation of its pathophysiology, standardized predictive models for HBL remain lacking, rendering current

clinical management reliant on subjective judgment and precluding individualized risk stratification. This study aimed to identify predictors of high-risk HBL after TKA and develop a validated visual prediction model integrating multiple clinical parameters.

METHODS: Clinical data from 400 TKA patients were analyzed. The dataset was randomly divided into a training group (70%) and a validation group (30%). Univariate analysis initially screened potential predictors, followed by least absolute shrinkage and selection operator (LASSO) regression for dimensionality reduction and key feature extraction. Multivariate logistic regression identified independent predictors of HBL. A nomogram was constructed and validated through receiver operating characteristic curves, calibration curves, and decision curve analysis to assess discrimination, calibration, and clinical utility.

RESULTS: Univariate analyses identified 18 significant variables associated with high-risk HBL ($P < 0.05$). A least absolute shrinkage and selection operator regression refined these to eight [...]

[PubMed](#) · [DOI](#)

Single Functioning Kidney Patients Face Increased Odds for Defined Complications Following Total Knee Arthroplasty.

Kim LY, Lee AH, Day W, Rubin LE, Grauer JN

BACKGROUND: Total knee arthroplasty (TKA) patients may present with a single functioning kidney (SFK) for reasons including congenital agenesis, living donation, or resection due to malignancy. While many SFK patients are asymptomatic, this may predispose them to adverse events following TKA. The present study investigated the correlation of SFK and its various etiologies on postoperative outcomes following TKA.

METHODS: A national database from 2010 to 2022 was used to identify 1,089,735 adult patients undergoing primary TKA. Then, 5,821 patients who had SFK were identified and matched 1:4 with control patients based on age, sex, Elixhauser comorbidity index (ECI), year of surgery, and history of chronic kidney disease (CKD). The 90-day adverse events were abstracted and compared by multivariable regression. The 5-year revision rates were compared using Cox proportional hazards regressions. The relative odds ratio (OR) of adverse events for different etiologies of SFK were compared using multivariable regressions.

RESULTS: On multivariable analyses, SFK patients had higher odds of aggregated adverse events driven by sepsis (OR [odds ratio] 1.75), urinary tract infection (OR 1.51), acute kidney injury (OR 1.30), and pneumonia (OR 1.27) ($P < 0.001$ for each). The SFK patients had equivalent rates of revision surgery within five year compared to controls (hazard ratio 0.93, $P =$ [...])

[PubMed](#) · [DOI](#)

Ten-Year Outcomes of Hypoallergenic Coated and Standard Implants in Total Knee Arthroplasty: A Randomized Controlled Trial.

Roitzsch C, Beyer F, Lützner C, Postler AE, Lützner J

BACKGROUND: Patients who have metal allergies frequently request hypoallergenic implants. In total knee arthroplasty (TKA), coating of standard implants is commonly used, with almost 10% of all primary TKAs in Germany in 2023. Registry data, however, have shown increased revision rates for coated implants. Whether this is due to the coating itself or the characteristics of allergy-prone patients remains unclear. The aim of this study was to compare revision rates, patient-reported outcomes, and serum metal levels in patients receiving coated versus standard TKA.

METHODS: In this randomized controlled trial, patients undergoing primary, unconstrained TKA were enrolled. Exclusion criteria included other metal implants, the need for constrained prosthesis, or known implant material allergy. A total of 118 patients were randomized (59 per group) to receive a coated or standard TKA. Serum metal levels were assessed up to five years. At the 10-year follow-up, 20 patients had died, and two had undergone revision. Of the remaining 96, 79 were available for clinical evaluation. Outcomes included the Oxford Knee Score, Short Form-36, and patient satisfaction.

RESULTS: Overall, 10-year implant survival was 98% for both groups. A revision occurred in each group. After 10 years, both groups showed good functional outcomes, health-related quality of life, and patient satisfaction, with no [...]

[PubMed](#) · [DOI](#)

Total Knee Arthroplasty in the Stiff and Ankylosed Knee at a Mean Follow-Up of 16 Years.

Kk M, Aggarwal K, Rajgopal A

BACKGROUND: Total knee arthroplasty (TKA) in patients who have stiff or ankylosed knees (range of motion [ROM] < 20°) presents technical challenges. Outcomes of TKA in this group of patients at follow-up greater than 12 years are under-reported. This study assesses the outcomes of TKA in this cohort at a mean follow-up of 16.3 years.

METHODS: This retrospective review of 109 patients (135 knees) who underwent TKA for stiff or ankylosed knees between 2000 and 2014 included cases from our previously published series, with a minimum follow-up of 10 years. There were 61 knees that were stiff in extension and 74 in flexion (SF). The mean follow-up duration was 16.3 years (range, 10 to 23). We assessed and compared Knee Society Score, Knee Society Functional Score, ROM, complications, revision rates, and implant survivorship between the two cohorts.

RESULTS: At final follow-up, mean Knee Society Score improved from 15.8 ± 6.3 to 72.8 ± 5.1 , Knee Society Functional Score from 9.4 ± 3.6 to 62.9 ± 3.9 , and ROM from $11.4^\circ \pm 5.9$ to $89.2^\circ \pm 14.1$ ($P < 0.001$). Complications occurred in 33 knees, with 17 requiring revision. The 10 and 15 years implant survival was 88.9, and 81.7% respectively. Postoperative ROM and maximum flexion did not differ significantly between the two groups. There was no significant difference in complications between the two groups, although residual flexion contra [...]

[PubMed](#) · [DOI](#)

Complications Are Elevated in Patients Who Have Decubitus Ulcers Undergoing Hemiarthroplasty, but There Are No Significant Differences in Two-Year Outcomes.

Wang JY, Harris AB, Schmerler J, Golladay GJ, Thakkar SC, Rao S

BACKGROUND: Patients undergoing hemiarthroplasty for femoral neck fracture are generally frail and prone to complications. Decubitus ulcers, associated with poor mobility, may indicate worse outcomes, though this relationship remains understudied. We hypothesized that preoperative decubitus ulcers increase the risk of postoperative complications at 90 days and two years following hemiarthroplasty.

METHODS: A national administrative claims database was queried for patients undergoing hemiarthroplasty for femoral neck fracture from 2010 to 2021. Patients were stratified by the presence of absence of a preoperative decubitus ulcer. Multivariable logistic regressions were used to evaluate the association between preoperative decubitus ulcers and postoperative complications occurring within 90 days and two years, including revision surgery, periprosthetic joint infection, and periprosthetic fracture. Odds ratios (ORs) with 95% confidence intervals (CIs) were reported, with statistical significance set at $P < 0.05$. A total of 76,539 patients were included, of whom 1,288 (1.7%) had a diagnosed decubitus ulcer prior to surgery. The mean age was 72 years (range, 50 to 89).

RESULTS: Within the first 90 days postoperatively, patients who had decubitus ulcers had an increased risk of 90-day readmission (OR: 1.4, 95% CI: 1.2 to 1.6), surgical site infection (OR: 1.6, 95% CI: 1.2 to 2.2), [...]

[PubMed](#) · [DOI](#)

Outcomes and Complications of Total Hip Arthroplasty for Crowe Type IV Dysplasia of the Hip: An International Multicenter Study.

Lari A, Ayyad SH, Khan T, Alemairi T, El-Bakoury A, Hammad AS

BACKGROUND: Crowe type IV developmental dysplasia of the hip (DDH) represents the most severe form of hip dislocation and poses major technical challenges for total hip arthroplasty (THA). This study aimed to evaluate outcomes and complications of THA in Crowe type IV hips across multiple international centers, highlighting outcomes that may be relevant across practices in technique and implant selection.

METHODS: A retrospective review of 37 patients (49 hips) who had Crowe type IV DDH who underwent THA between 2013 and 2022 at four centers in three countries was conducted. Clinical and radiographic data included Harris Hip Score (HHS), Oxford Hip Score (OHS), limb-length discrepancy (LLD), component positioning, and complications. Subtrochanteric osteotomy, surgical approach, and implant type were documented. The mean follow-up was 7.8 years (range, three to 12).

RESULTS: The LLD improved from 4.10 ± 1.20 cm preoperatively to 0.70 ± 0.42 cm postoperatively ($P < 0.001$). The HHS improved from 36.7 to 80.1 ($P < 0.001$) and OHS from 14.0 to 40.2 ($P < 0.001$). Abductor lurch

decreased from 100 to 29.7%. Return to independent mobility was achieved in 81% of patients. Shortening osteotomy was performed in 64.9% of cases. Complications occurred in 21% of patients (16% of hips); the most common was periprosthetic fracture (n = five, 10%), and osteotomy nonunion occurred in two cases ([...])

[PubMed](#) · [DOI](#)

Patient Perception of Leg-Length Discrepancy and Satisfaction Following Primary Total Hip Arthroplasty.

Llanes PD, Lee YK, Park SY, Park JW, Koo KH

BACKGROUND: Leg-length discrepancy (LLD) is a common complication following total hip arthroplasty (THA), which might affect patient function and satisfaction. It is unclear how a patient's perception of LLD changes over time. The purpose of our study was to evaluate the incidence, affecting factors, and lapse of perceived LLD after primary THA.

METHODS: This was a prospective, single-center, cohort study of primary THAs performed on 126 patients from 2019 to 2021 by a single surgeon via a postero-lateral approach. Pre-, intra-, and postoperative variables, including demographic data and LLD, were assessed. Patients were grouped according to the presence of perceived LLD, and the satisfaction questionnaire was used for the evaluation at six weeks, three months, six months, and one year postoperatively.

RESULTS: There was no statistical difference in the pre-, intra-, and postoperative variables assessed in both groups. Of the 126 patients, 62 (49%) had perceived postoperative LLD, and the remainder were asymptomatic. After one year, 73% (45 of 62) of symptomatic patients reported complete resolution of symptoms, and 15 of 17 symptomatic patients who had persistent perceived LLD reported favorable patient satisfaction scores even without surgical intervention.

CONCLUSIONS: The majority of patients (73%) who had a perception of LLD following THA observed their symptoms resolve [...]

[PubMed](#) · [DOI](#)

Should Total Hip Arthroplasty Surgeons Be Concerned that a Delay Between the Date of Surgical Booking and Surgery Influences Patient Body Mass Index and Short-Term Outcomes?

Di Pauli von Treuheim T, Sarfraz A, Ruff G, Saba BV, Schwarzkopf R, Rozell JC, Aggarwal VK

BACKGROUND: Obesity is a known risk factor for complications after total hip arthroplasty (THA), with societal guidelines recommending surgical delay for patients above body mass index (BMI) targets. Consequently, patients are motivated to reach BMI targets before the office visit discussing surgical booking. Our study investigates BMI fluctuations between surgical booking and the surgery date and whether these fluctuations have implications for perioperative and postoperative outcomes.

METHODS: We retrospectively reviewed our institutional database for elective, primary, unilateral THA from 2015 to 2024 with a minimum 90-day follow-up. The cohort was stratified into three groups by percent BMI change from booking date to THA date: group 1, decrease in BMI; group 2, 0 to 5% increase in BMI; and group 3, > 5% increase in BMI. Baseline demographic factors, as well as perioperative and postoperative outcomes, were compared. A multivariate regression analysis evaluated risk factors for interval change in BMI. We reviewed 10,400 THA patients who had an average 62-day delay between booking and surgery dates, where 44.2, 42.2, and 13.6% were allocated to groups 1, 2, and 3, respectively.

RESULTS: Operative time, length of stay, and discharge to a rehab facility were significantly higher in group 3 compared to the others. There were no significant differences seen in 90-day outcomes, [...]

[PubMed](#) · [DOI](#)

Admission Lymphocyte-to-C-Reactive Protein Ratio Predicts 90-Day Adverse Outcomes Following Aseptic Revision Total Hip Arthroplasty.

Karlidag T, Zanna L, Mundhe A, Gehrke T, Citak M

BACKGROUND: Revision total hip arthroplasty (rTHA) is linked to a greater risk of adverse postoperative outcomes in comparison to primary THA, owing to heightened surgical complexity and the burden of patient comorbidities. Consequently, there is an imperative need for an easily accessible predictor to facilitate preoperative risk stratification before rTHA. This study was conducted to evaluate the predictive utility of the admission lymphocyte-to-C-reactive protein ratio (LCR) concerning adverse events within 90 days postoperatively following aseptic rTHA.

METHODS: A retrospective cohort analysis was conducted involving 5,596 patients who underwent aseptic rTHA from 2010 to 2022 at a tertiary arthroplasty center. Patient demographics, clinical parameters, and perioperative data were collected. Adverse outcomes within 90 days postoperatively, including major and minor complications, unplanned readmissions, and mortality, were analyzed. Logistic regression analysis was performed to estimate the predictive value of preoperative LCR in conjunction with other potential risk factors, whereas receiver operating characteristic curve analysis was used to establish the most appropriate LCR cutoff values.

RESULTS: A total of 716 patients (12.8%) experienced postoperative adverse events. The admission LCR was significantly lower among these patients compared to controls. Multivariate an [...]

[PubMed](#) · [DOI](#)

Contemporary Cemented Revision Stems Demonstrate Excellent Survivorship in Revision Total Hip Arthroplasty up to 13-Year Follow-Up.

Baylac A, Guiraud T, Cardoso M, Abi Abboud E, Mathieu J, Dagneaux L

BACKGROUND: An extended diaphyseal femoral fixation can be mandatory in complex femoral revision total hip arthroplasty (THA). While outcomes of uncemented revision components are well documented in this context, little data are available about cemented revision stems with different options. This study aimed to evaluate survivorship, risk factors for re-revision, and radiographic and clinical outcomes of a series of one contemporary monoblock cemented revision stem implanted for femoral revision.

METHODS: Between 2008 and 2022, 230 cemented revision femoral components were implanted in revision THAs in 227 patients. The mean age was 77 years, with 59% being women. The Kaplan-Meier method was used to estimate survival with its confidence interval (95% CI). Risk analysis used the Cox proportional hazard method. Preoperative bone stock deficiency and stem fixation were assessed. Harris Hip Scores (HHSs) were collected at the last follow-up. The mean follow-up was five years (range, two to 13).

RESULTS: The 10-year survivorships free from aseptic femoral loosening and from any femoral revision were 98 (95% CI [confidence interval]: 92 to 99) and 84% (95% CI: 56 to 95), respectively. There were 12 femoral re-revisions: nine chronic infections and three cases of femoral loosening. Vancouver C femoral fractures were the main cause for reoperation (eight cases, 3% of the cohort). Dem [...]

[PubMed](#) · [DOI](#)

Synovial Absolute Neutrophil Count as a Superior Diagnostic Biomarker for Acute Periprosthetic Joint Infection in Hip and Knee Arthroplasty With Low-Virulence Organisms.

Karlidag T, Mundhe A, Zanna L, Gehrke T, Citak M

BACKGROUND: Diagnosing acute periprosthetic joint infection (PJI) soon after surgery presents a challenge because the inflammation caused by the procedure can mimic that of an infection. Key diagnostic measures include synovial white blood cell (WBC) count, polymorphonuclear percentage (PMN%), and alpha-defensin (AD). Nevertheless, the precise thresholds, particularly for infections with low virulence, remain ambiguous. This research assessed the effectiveness of synovial absolute neutrophil count (ANC) in diagnosing acute PJI in hip and knee joints and established specific thresholds based on the procedure and the organism involved.

METHODS: An examination was carried out on 502 patients who underwent joint aspiration within a 90-day period following their primary total joint arthroplasty, conducted from March 2011 to August 2022 at a tertiary arthroplasty center. Of the 502 patients, 123 (24.5%) had an acute PJI. The diagnosis of PJI was established according to the criteria set by the 2018 International Consensus Meeting. Receiver operating characteristic curves were used to assess synovial fluid WBC, PMN%, ANC, and AD, aiming to determine the optimal cutoff values for total knee arthroplasty and total hip arthroplasty, with differentiation according to the virulence of the organism.

RESULTS: For total knee arthroplasty, optimal thresholds were WBC 5,162 cells/ μ L, PMN% 88. [...]

Use of Cementless or Cemented Stems in One-Stage Exchange for Periprosthetic Joint Infection in Total Hip Arthroplasty Does Not Affect Outcomes: A Propensity-Matched Cohort Study from an Institutional Registry With a Mean Follow-Up of Six Years.

Bouché PA, Hallé A, Mouton A, Aubert T, Zeller V, Auburger G, Graff W, Marmor S

BACKGROUND: Periprosthetic joint infection (PJI) is the most dramatic complication of total hip arthroplasty (THA). We investigated whether a one-stage arthroplasty using a cemented or uncemented stem was associated with better survival in terms of infection and mechanical complications.

METHODS: Patients were selected from a PJI institutional registry of THA between 2003 and 2019. After matching the two groups by propensity score at a 1:3 ratio, 241 patients were finally included: 62 in the cemented group and 179 in the cementless group. The mean age of the population was 68 years (range, 25.1 to 91.4), and 45.2% (109 of 241) of the patients were women, with a mean body mass index of 26.9 (range, 15.6 to 46.4). The mean follow-up was 6.6 years (range, 0.1 to 18.1). To compare the overall survival rate between groups, Kaplan-Meier analysis was performed.

RESULTS: The overall survival rate in our series was 93.3% [88.2 to 98.7] at 10 years. At the last follow-up, nine THAs had been reinfected: four relapses and five new infections. Regarding the reinfection-free survival rate, there was no significant difference ($P = 0.62$) between the cemented group (95.8% [90.2 to 100.0]) and the cementless group (94.1% [87.8 to 100.0]) at 10 years. At the last follow-up, 34 (14.1%) THAs had aseptic complications that required 30 revisions. Regarding the aseptic survival rate, there was no si [...]

PubMed · DOI

The In Vitro Performance of Surgical Irrigation Solutions in Preventing Biofilm Formation on Implants.

Tabatabaei F, McMahon R, Estlack L, Sanchez TF, Medina S, Bionda N

BACKGROUND: Total joint arthroplasties are widely performed in the United States. Although various intraoperative irrigation solutions have been studied, comprehensive in vitro comparisons of their efficacy in preventing biofilm formation across multiple implant materials remain limited. This study evaluated the effectiveness of nine commercially available irrigation solutions in preventing biofilm formation by *Staphylococcus aureus* and *Pseudomonas aeruginosa* on four orthopaedic implant materials under clinically relevant conditions. Cytotoxicity was also assessed by evaluating human dermal fibroblast recovery following exposure.

METHODS: The solutions tested included a polyhexamethylene biguanide (PHMB)-synergy solution, a benzalkonium chloride (BZK) solution, a PHMB-betaine solution, a chlorhexidine gluconate solution, a povidone-iodine solution, a hypochlorous acid solution, a citric acid solution, a triple antibiotic solution (bacitracin, cefazolin, gentamicin), and normal saline. The testing laboratory was blinded to solution identities. Stainless steel, cobalt-chromium alloy, titanium alloy, and ultrahigh molecular weight polyethylene coupons ($n =$ four per group) were pretreated with each solution for two minutes. After aspiration, bacteria in simulated synovial fluid were added. Following 24-hour incubation at 37° C, adherent bacteria were recovered and quantified. For [...]

PubMed · DOI

The Influence of Total Knee Arthroplasty Surgical Approach on Range of Motion, Patient-Reported Outcome Measures, and Reoperation Rates: A Systematic Review of Randomized Controlled Trials.

Rainey JP, Radtke LE, Gililand JM, Mont MA

BACKGROUND: Various surgical approaches may be used to perform total knee arthroplasty (TKA). This systematic review assessed randomized controlled trials (RCTs) evaluating various TKA approaches and their influence on knee range of motion (ROM), patient-reported outcome measures (PROMs), and reoperation rates.

METHODS: A search of PubMed and Embase databases was performed for manuscripts published between January 1, 2010, and December 31, 2024. Inclusion criteria consisted of RCTs evaluating knee ROM, PROMs, and reoperation rates of various TKA surgical approaches. Modified Coleman methodology scoring and level of evidence were assigned to each study. Non-English manuscripts, studies with fewer than 30 patients in each

group, or studies with a mean follow-up of less than three months were excluded. A total of 27 manuscripts met final inclusion, and all were level 1 evidence with fair or better modified Coleman scores (mean 74.9, range 64 to 89).

RESULTS: Among the 25 RCTs that evaluated postoperative knee ROM, three studies supported the subvastus approach, two supported the midvastus approach, and two supported the mini medial parapatellar (MPP) approach when compared to a standard MPP. The remaining 18 studies found no difference in ROM between surgical approaches. Various PROMs were evaluated in 26 studies. Of these, three studies favored the subvastus approach, two favor [...]

[PubMed](#) · [DOI](#)

Femoral Valgus Leads to Earlier Total Hip Arthroplasty Independent of Acetabular Coverage, While Varus Deformity Has a Protective Effect.

Schwarz GM, Simon S, Mitterer JA, Huber S, Leder-Berg S, Hofstaetter JG

BACKGROUND: The presence of acetabular dysplasia or femoro-acetabular impingement increases the chance of an earlier total hip arthroplasty (THA). Surprisingly, the influence of femoral and acetabular parameters on the age at which THA is required remains poorly investigated. The aim of this study was to evaluate the radiographic hip morphotype at the time of THA using an artificial intelligence-based analysis to assess potential differences.

METHODS: Overall, 6,767 patients (mean age 66 years [range, 16 to 99]; women: 60.6%; men: 39.4%) from a single orthopaedic center were included in the final analysis. Images were analyzed using artificial intelligence-powered software. The lateral-center-edge angle and the neck-shaft angle (caput-collum-diaphyseal) were used to determine the nine different hip morphotypes based on the coronal plane alignment of the hip (CPAH, I to IX).

RESULTS: Patients who had valgus-aligned femoral necks (CPAH, I, IV, and VII) were significantly younger at the time of THA compared to patients who had femoral varus necks (CPAH, III, VI, and IX), suggesting a potential negative effect of valgus and a protective effect of varus alignment independent of acetabular coverage. Type I hips, defined as acetabular undercoverage in combination with femoral valgus alignment, were on average 16.2 years younger at the time of THA compared to type IX hips with acetab [...]

[PubMed](#) · [DOI](#)

Glucagon-Like Peptide-1 Receptor Agonists, Readmission, and Postoperative Complications in Arthroplasty: A Systematic Review and Meta-Analysis.

Chan YC, Chuang SH, Wu LC, Kuo YJ, Lian YZ, Chen YP

BACKGROUND: Agonists targeting the glucagon-like peptide-1 receptor have gained prominence in treating individuals who have type 2 diabetes and obesity. However, their influence on postoperative medical complications following total joint arthroplasty remains poorly characterized.

METHODS: A systematic review and meta-analysis were conducted using data from eight retrospective cohort studies encompassing 76,091 patients who underwent total hip or knee arthroplasty. The primary outcome was hospital readmission within 90 days. Additional postoperative events assessed included deep vein thrombosis, pulmonary embolism, pneumonia, acute kidney injury, surgical site infection, cerebrovascular accident, urinary tract infection, and hypoglycemic event. Pooled odds ratios were calculated using a random-effects model to account for interstudy variability.

RESULTS: Patients who received glucagon-like peptide-1 receptor agonists (GLP-1 RAs) before surgery exhibited a significantly lower likelihood of 90-day readmission (odds ratio 0.73; 95% confidence interval: 0.64 to 0.82). There were no statistically significant differences observed in other postoperative medical complications. Sensitivity and trim-and-fill analyses revealed modest variability in select outcomes, but did not substantially alter overall conclusions.

CONCLUSIONS: Preoperative glucagon-like peptide-1 receptor agonist use [...]

[PubMed](#) · [DOI](#)

Does Testosterone Replacement Therapy Increase Risk of Thrombosis After Total Hip or Knee Arthroplasty?

Ni A, Cal E, Martinez V, Gerhart CR, Moore CC, Buttacavoli F

BACKGROUND: Testosterone is a prothrombotic hormone, but the effect of exogenous testosterone on venous thromboembolic events (VTEs) following total joint arthroplasty (TJA) remains unclear. This study investigated the association between testosterone replacement therapy (TRT) and postoperative risk of VTE following primary total hip arthroplasty (THA) and total knee arthroplasty (TKA).

METHODS: We conducted a retrospective, matched cohort study using a national registry to identify primary THA or primary TKA patients from the years 2005 to 2025. Patients were stratified by TRT within three months of the TJA. There were 640,219 patients who had a THA (265,477) or a TKA (374,742) that were included. There were 1,469 THA (0.6%) and 2,032 (0.5%) TKA patients who were prescribed testosterone within three months before or after their TJA. A matched cohort was created after balancing demographics and comorbidities. Primary outcomes were DVT or pulmonary embolism (PE) within three months after their TJA. Measures of association models were used to calculate odds ratios (ORs).

RESULTS: We found a statistically significant increase in the rate of PE in TKA patients on TRT compared to those who were not (OR = 1.45; 95% confidence interval (CI), 1.04 to 2.02; P = 0.029). Rates of DVT in TKA patients on TRT were similar to those not on TRT (OR = 1.27; 95% CI, 0.98 to 1.65; P = 0.073). Si [...]

[PubMed](#) · [DOI](#)

Variation and Slower-Than-Recommended Recovery of Daily Life Activities Following Knee Arthroplasty With a Personalized eHealth Program: Results of the ACTIVE Trial Intervention Cohort.

Straat AC, Kuijer PPF, Maarleveld JM, van den Berg XT, Visser SV, Huirne JA, Kerkhoffs GM, Anema JR et al.

BACKGROUND: The rising prevalence of knee arthroplasty (KA) affects daily life, especially for working-age patients who have high expectations for resuming activities. Variability in guidelines and lack of personalized recovery recommendations highlight the need for improved guidance, such as through eHealth solutions. This study identifies the most important daily activities chosen in an eHealth application and evaluates patients' recovery times compared to expert recommendations.

METHODS: Data from the intervention group of the Dutch ACTIVE trial were analyzed (n = 145, mean age 58 years [range, 46 to 66]; 54% women); 133 of these patients started follow-up. Patients used an app, which provided expert-based personalized recommendations for 27 activities, all with three recovery speeds, for unicompartmental and total KA: slow, average, and fast. We analyzed activity choices and compared actual resumption times with the expert recommendations.

RESULTS: The top activities included cycling (92%), driving (88%), and lifting/carrying (73%). In 79% of cases, actual median recovery time exceeded recommendations. For total KA patients in average and fast recovery groups, climbing stairs (52 and 46%) and walking on even terrain (50 and 53%) most often matched recommendations.

CONCLUSION: Generally, recovery was slower than the provided recommendations, with 8 out of 10 activities no [...]

[PubMed](#) · [DOI](#)

An Asymmetric Medio-Lateral Flexion Laxity Target Yields Excellent Results in Robotic-Assisted Total Knee Arthroplasty With Functional Knee Positioning.

Vermue H, Zambianchi F, Clemenza S, Clò M, Victor J, Catani F

BACKGROUND: Functional knee positioning (FKP) in robotic-assisted (RA) total knee arthroplasty (TKA) aims to optimize soft-tissue balance through patient-specific implant fine-tuning. However, it remains unclear (1) how intraoperative ligament laxity affects joint awareness and (2) whether the discrepancy between intraoperative planned and achieved laxity is affected by preoperative coronal limb alignment or fixed flexion deformity (FFD).

METHODS: This retrospective study analyzed 130 patients undergoing image-based RA TKA with FKP using a cementless cruciate-retaining implant. Intraoperative adjustments were made to balance the flexion-extension laxity, targeting one to two mm larger lateral flexion laxity. Medial and lateral gaps before and after implant positioning were recorded at near extension and 90° flexion. Patient-reported outcome measures (PROMs), including the Forgotten Joint Score (FJS-12) and a 5-point Likert scale for satisfaction, were assessed at a

minimum of one year postoperatively to investigate correlations between PROMs and final intraoperative laxities. The impact of preoperative coronal limb alignment, measured on long-leg radiographs, and FFD on final laxities was also evaluated.

RESULTS: Final laxities were: median medial extension laxity 1.5 mm (1.0 to 2.0), lateral extension laxity 1.5 mm (1.1 to 2.0), medial flexion laxity 1.0 mm (0.5 to 1.5), and [...]

[PubMed](#) · [DOI](#)

Predicting Outcomes of Early and Late Acute Periprosthetic Joint Infections Treated With Debridement, Irrigation, and Implant Retention: An External Validation of the CRIME80 (COPD, C-Reactive Protein, Rheumatoid Arthritis, Index Surgery, Male, Exchange of Mobile Components, Age Greater than 80 years) Score.

Naufal ER, Babazadeh S, Aboltins C, Choong PF, Fillingham YA, Higuera CA, Otero JE, Piuze NS et al.

BACKGROUND: Debridement, antibiotics, and implant retention (DAIR) requires careful patient selection. The CRIME80 (COPD, C-reactive protein, Rheumatoid arthritis, Index surgery, Male, Exchange of mobile components, age greater than 80 years) score is a tool designed to predict treatment failure in patients who have late-acute PJI treated with DAIR. We aimed to assess the predictive performance of the CRIME80 score in patients who have early and late acute PJI.

METHODS: We conducted an international multicenter validation study using retrospectively collected data from patients who had acute infections of the hip or knee treated with DAIR. Failure was defined as: (i) removal of the prosthesis; (ii) the need for suppressive antibiotics one or more years after surgery for PJI; (iii) relapse of infection as determined by the treating clinician; (iv) repeated DAIR more than seven days after the initial procedure; or (v) death. Nonparametric estimation of the area under the receiver operating curve (AUC) was used to establish the discriminative performance of the CRIME80 score. The cohort included 281 knee and 99 hip infections treated with DAIR.

RESULTS: Although the CRIME80 score showed a linear trend with failure, the score only had modest discriminative performance (AUC in hip PJI: 0.63, 95% confidence interval (CI) 0.51 to 0.74; AUC in knee PJI: 0.57, 95% CI 0.51 to 0.64). Th [...]

[PubMed](#) · [DOI](#)

Return to Sports After Total Hip Arthroplasty: Patterns of Participation and Sport-Specific Outcomes.

Lin YJ, Terner B, Piergrossi D, Rozell J, Schwarzkopf R, Arshi A

BACKGROUND: Return to sports is a concern for many patients undergoing total hip arthroplasty (THA). As younger, active patients increasingly undergo THA, identifying factors that influence athletic recovery is critical. Although prior work has reported favorable outcomes, large-scale studies comparing return rates across both surgical approaches and sport types remain limited.

METHODS: We retrospectively analyzed 1,115 athletically participant THA patients and compared rates of returning to sports up to 1 year postoperatively. Survey data captured sport type, participation at four time points, frequency, exertion, and perceived recovery. Patients were stratified by surgical approach: posterior approach (PA) (n = 519), anterior approach (AA) (n = 556), and lateral approach (LA) (n = 50).

RESULTS: Preoperatively, 45.2% of patients were active in at least one sport versus 43.7% at one year postoperatively; 73.2% of preoperative athletes returned, and 23.3% of those previously inactive took up sports. Return-to-sport rates were 68.1, 77.0, and 81.8% for PA, AA, and LA, respectively (Chi-square = 3.42, P = 0.181). There were no significant differences between AA and PA, and the LA was not statistically comparable due to a smaller sample size. Low-impact sports had significantly higher return rates than high-impact sports (72.4 versus 50.0%; P < 0.001). At survey completion (> one [...])

[PubMed](#) · [DOI](#)

Disease-Modifying Antirheumatic Drugs May Reduce the Risk of Manipulation Under Anesthesia or Lysis of Adhesions Following Primary Total Knee Arthroplasty.

Goh GS, Lee S, Kim MT, Bortman JM, Paré DW, Wang CY, Aurigemma PH

BACKGROUND: Stiffness is a debilitating complication following total knee arthroplasty (TKA), often necessitating manipulation under anesthesia (MUA) or arthroscopic lysis of adhesions (LOA). While nonsteroidal anti-inflammatory drugs and corticosteroids have been shown to reduce the risk of stiffness, the effect of disease-modifying antirheumatic drugs (DMARDs) remains unknown. This study aimed to evaluate whether DMARD use was associated with reduced rates of MUA or LOA after TKA.

METHODS: A national health network was queried for patients who had rheumatoid arthritis and underwent primary TKA between 2013 and 2023. Patients were stratified into traditional, biologic, and dual DMARD groups based on prescriptions within one year before surgery. Each DMARD group was matched with non-DMARD controls using propensity score matching (1:1), controlling for demographics, comorbidities, and perioperative nonsteroidal anti-inflammatory drug and corticosteroid use. The primary outcome was MUA or LOA at 90 days, six months, one year, and two years. Other surgical complications and inflammatory markers were also recorded. A total of 16,596 patients were included (traditional: 5,507; biologic: 929; dual: 1,862).

RESULTS: Traditional DMARD use was associated with significantly reduced MUA or LOA risk at 90 days (odds ratio (OR) 0.73), six months (OR 0.66), one year (OR 0.69), and two year [...]

[PubMed](#) · [DOI](#)

Diabetic Neuropathy Is a Strong Risk Factor for Poor Orthopaedic Outcomes Following Total Knee Arthroplasty.

Kotzur T, Barnes M, Peterson B, Jodoin Z, Buttacavoli F, Moore C

BACKGROUND: The aim of this study was to investigate the impact of diabetic peripheral neuropathy (DPN) on total knee arthroplasty (TKA) outcomes when compared with both nondiabetic and diabetic patients who did not have neuropathy (DM).

METHODS: This retrospective cohort study utilized a global clinical research network, comprising data collected between 2005 and 2022. Patients undergoing TKA were identified using international classification of diseases, 10th revision and Current Procedural Terminology codes. Patients were stratified based on diabetes status and the presence of diabetic neuropathy. Orthopaedic complications were assessed over a 2-year period postoperatively, while medical and hospital complications were evaluated within 90 days. Propensity score matching (1:1) was performed to control for baseline covariates, including age, sex, body mass index, Charlson Comorbidity Index, and hemoglobin A1c where applicable. Postmatching logistic regression was used to calculate risk ratios (RRs) with 95% confidence intervals. Statistical analyses were conducted with significance set at $P < 0.05$.

RESULTS: This study included 318,851 patients who underwent TKA: 227,597 (71.4%) patients who did not have diabetes, 68,694 (21.5%) patients who had diabetes, but did not have diabetic neuropathy, and 22,560 (7.1%) patients who had diabetic neuropathy. When compared to patients wh [...]

[PubMed](#) · [DOI](#)

Liposomal Bupivacaine Versus Ropivacaine With Perineural Dexamethasone in Adductor Canal Block for Total Knee Arthroplasty: A Randomized Clinical Trial.

Liao C, Li J, Hu X, Sun Y, Zheng K, Pan X, Wang L

BACKGROUND: Adductor canal block (ACB) provides effective analgesia for knee surgery while preserving quadriceps muscle strength. However, conventional local anesthetics yield only a short-lasting effect. To extend analgesia duration and reduce opioid usage, the administration of liposomal bupivacaine and perineural dexamethasone has been proposed. The present study assessed and compared the analgesic effectiveness and duration between liposomal bupivacaine and ropivacaine with perineural dexamethasone in patients undergoing total knee arthroplasty (TKA).

METHODS: A total of 72 patients scheduled for primary unilateral TKA were prospectively randomized to receive a single-injection ACB with either liposomal bupivacaine combined with bupivacaine hydrochloride ($n = 36$) or ropivacaine with perineural dexamethasone (control group, $n = 36$). The primary endpoint was the incidence of moderate-to-severe pain with movement at 48 hours postoperatively. The secondary outcomes included the incidence of moderate-to-severe pain both at rest and with movement at other time points, numerical rating scale pain score, opioid consumption, quality of postoperative recovery, and adverse events.

RESULTS: At 48 hours, the incidence of moderate-to-severe pain with movement did not differ between the liposomal bupivacaine group and the control group (47.2 versus 63.9%; relative risk 0.71; 95% confidence interval 0.41 to 1.24).

[PubMed](#) · [DOI](#)

Do Implant-Specific Associations Exist Between Patellar Resurfacing and All-Cause Revision Risk After Primary Total Knee Arthroplasty? A Retrospective Analysis of Two National Joint Registries.

Ross BJ, Arellano E, McCall M, Manz W, Armstrong C, Khan I, Debbi EM, Dickenson E et al.

BACKGROUND: The role of patellar resurfacing during primary total knee arthroplasty (TKA) remains controversial. It is unclear if patellar resurfacing impacts revision risk differently across implant systems. The purpose of this study was to compare estimated revision rates for highly utilized TKA implants based on the presence or absence of primary patellar resurfacing.

METHODS: Over 20 national joint registries were screened. Only the National Joint Registry (NJR) Annual Report 2023 and the Endoprothesenregister Deutschland (EPRD) Annual Report 2023 reported implant-specific estimated all-cause revision rates stratified by primary patellar resurfacing status. Individual implants at or above the 50th percentile with respect to implant utilization were analyzed. Kaplan-Meier estimated all-cause revision rates at three and 5 years were compared for individual implants among patients who had versus those who did not have patellar resurfacing during primary TKA using multivariable logistic regression.

RESULTS: There were 13 unique implant systems from EPRD and 46 implants from NJR that were analyzed, yielding 1,379,573 primary TKAs in total. In the EPRD, estimated all-cause revision rates were significantly higher without primary patellar resurfacing for 58.3% of implants (all odds ratios (ORs) > 1.00, $P < 0.05$), comparable for 34.0% of implants ($P > 0.05$), and significantly low [...]

[PubMed](#) · [DOI](#)

Validity of Substantial Clinical Benefit Estimates in Knee Arthroplasty Is Weak Based on Comparisons to a Gold Standard of Meaningful Improvement.

Riddle DL, Dumenci L

BACKGROUND: Estimates of the substantial clinical benefit (SCB) for knee arthroplasty have been endorsed by the Centers for Medicare and Medicaid Services but have not been validated. We examined validity by comparing SCB estimates to an externally validated gold standard of meaningful improvement.

METHODS: There were two datasets, the Knee Arthroplasty pain coping Skills Training (KASTPain) no-effect randomized clinical trial and the Osteoarthritis Initiative (OAI) data on patients who had a knee arthroplasty. Both studies captured preoperative and multiple postoperative Western Ontario and McMaster Universities Osteoarthritis Index Physical Function scores. A total of 396 participants were included from the OAI, and 384 participants were from KASTPain.

RESULTS: The validity of Latent Class Analysis estimates was supported. Compared to the gold standard, KASTPain participants meeting/exceeding the SCB generally ruled in a good outcome, while change scores less than the SCB did not rule in a poor outcome. The diagnostic validity of SCB estimates in OAI was weak.

CONCLUSIONS: We found evidence of weak validity for SCB estimates, particularly for the subset of participants not meeting the SCB. Future work should explore the validity of baseline quartile-adjusted SCB estimates obtained at multiple sites.

LEVEL OF EVIDENCE: Level 1 Diagnostic test study.

[PubMed](#) · [DOI](#)

Preoperative Hemoglobin as a Continuous Predictor of Postoperative Complications Following Elective Total Hip Arthroplasty.

Palmer RC, Lim MA, Telang SS, Iyer A, Culler MW, Yun AG, Lieberman JR, Heckmann ND

BACKGROUND: Preoperative anemia is associated with adverse outcomes following total hip arthroplasty (THA). However, discrete hemoglobin thresholds may obscure potentially meaningful relationships between anemia severity and postoperative risk. This study assessed preoperative hemoglobin as a continuous variable to predict

periprosthetic joint infection (PJI), medical, and surgical complications following elective THA.

METHODS: A health care database was utilized to identify all primary THA patients between 2016 and 2023 with a hemoglobin value available within 28 days preoperatively. The primary outcome was PJI within 90 days of surgery. The secondary outcomes included 90-day aggregate medical and surgical complications. Multivariable logistic regressions with restricted cubic splines (RCS) evaluated hemoglobin as a continuous predictor of postoperative risk. Change-points were identified using a Metropolis-Hastings bootstrap simulation. In total, 25,654 THAs were included, of which 126 (0.5%) developed PJI within 90 days.

RESULTS: The RCS analysis identified hemoglobin change-points of 13.5 g/dL for PJI and aggregate medical complications and 13.3 g/dL for aggregate surgical complications. Compared to 13.5 g/dL, patients who had hemoglobin values of 11.0 and 8.0 g/dL had a 1.91-fold (95% confidence interval (CI): 1.37 to 2.68) and 4.40-fold (95% CI: 1.91 to 10.14) increased risk [...]

[PubMed](#) · [DOI](#)

International Classification of Diseases-10-Based Risk Stratification and Implant Selection for Knee Arthroplasty Among Patients Who Have Psychiatric and Neurologic Comorbidities: A Registry-Based Analysis of 549,234 Cases.

Wagner N, Wu Y, Grimberg A, Gwinner C, Perka C, Hardt S

BACKGROUND: Psychiatric and neurologic comorbidities may worsen outcomes after total knee arthroplasty (TKA) or unicompartmental knee arthroplasty (UKA). We compared risks across eight International Classification of Diseases, Tenth Revision, German Modification clusters-psychiatric F00 to F39 and neurologic G20 to G64-and by implant constraint (unconstrained, constrained, and unicompartmental).

METHODS: We analyzed 549,234 primary knee arthroplasties (2012 to 2024) from a national arthroplasty registry. Patients who had diagnoses were matched one-to-one to controls without F00 to F99 or G00 to G99 codes, adjusting for age, sex, body mass index (BMI), Elixhauser index, fixation, and implant type. Kaplan-Meier with 95% confidence intervals (CIs) estimated 8-year incidence; multivariable Cox models estimated hazard ratios (HRs) by diagnoses and implant.

RESULTS: Complication patterns varied by cluster. The revision risk was highest in F20 to F29 (schizophrenia spectrum; HR 1.92, $P = 0.001$) and G20 to G26 (extrapyramidal/parkinsonian; HR 1.66, $P < 0.001$). Infection peaked in F00 to F09 (organic/cognitive; HR 2.34, $P < 0.001$) and G60 to G64 (polyneuropathy; HR 1.52, $P < 0.001$). Loosening increased from F30 to F39 (affective; HR 1.50) and from G20 to G26 (extrapyramidal/Parkinsonian; HR 1.92, $P < 0.001$). All-cause mortality was highest in F20 to F29 (schizophrenia spectrum; HR 3.0 [...])

[PubMed](#) · [DOI](#)

Defining Preoperative Hemoglobin Thresholds Can Help Avoid Transfusion or Adverse Events for Revision Total Joint Arthroplasty.

Sutton R, Sherman MB, Leipman JH, Linton A, Martinazzi BJ, Fillingham YA

BACKGROUND: Anemia optimization before revision total joint arthroplasty (rTJA) could improve outcomes; however, there is little literature on hemoglobin optimization efforts before rTJA. Simulation models in orthopaedic literature help define the likelihood of successful surgery based on thresholds of preoperative parameters. We evaluated whether certain preoperative hemoglobin thresholds could be defined for revision total knee arthroplasty (rTKA) and revision total hip arthroplasty (rTHA) in order to predict reduced likelihood of intraoperative or postoperative transfusion or perioperative adverse events.

METHODS: We retrospectively reviewed consecutive rTKA and rTHA patients from 2019 to 2024. Exclusion criteria were primary arthroplasty, no preoperative hemoglobin, and incomplete 1-year follow-up. Preoperative hemoglobin values were used to determine the likelihood of transfusion (intraoperative and postoperative) and perioperative adverse outcome (defined as medical complication, surgical complication, readmission, or re-revision).

RESULTS: Preoperative hemoglobin was more sensitive and specific at defining the need for perioperative transfusion compared to perioperative adverse events. For both rTKA and rTHA, men required a higher preoperative hemoglobin threshold than women when predicting perioperative transfusion (rTKA: men 11.7 g/dL;

area under the curve (AUC) 0.86 [...]

[PubMed](#) · [DOI](#)

Chronic Kidney Disease and Outcomes of Total Hip Arthroplasty Patients Who Have Periprosthetic Hip Fractures.

Juels ML, Galoustian NA, Balian J, Cho NY, Benharash P, Stavrakis AI

BACKGROUND: Patients who have end-stage renal disease (ESRD) and chronic kidney disease (CKD) have an increased risk of mortality and perioperative complications after primary total hip arthroplasty. The association between increasing severity of renal disease with clinical and financial end points in patients treated for hip periprosthetic fracture (PPF) is underexplored in the literature.

METHODS: Hip PPF hospitalizations were identified using a large national database and grouped into non-CKD, early-stage CKD (CKD 1 to 3), advanced-stage CKD (CKD 4 to 5), and ESRD. The primary outcome of the study was in-hospital mortality. The secondary outcomes included perioperative complications, hospitalization length of stay, costs, non-home discharges, and 30-day readmissions. Nonparametric tests were used to assess temporal trends. Of 108,836 patients treated for PPF, 17,115 (15.7%) had CKD. The overall prevalence of renal disease in patients who had PPF increased significantly from 2016 to 2020 (2020: 16.3% versus 2016: 14.6%, $P < 0.001$), whereas those who had ESRD decreased (2020: 1.3% versus 2016: 1.8%, $P = 0.001$). Multivariable logistic and linear regression models were used to evaluate the association of renal disease severity with outcomes of interest.

RESULTS: Chronic kidney disease 1 to 3 (adjusted odds ratio [AOR] 1.28, 95% CI 1.05 to 1.55), CKD 4 to 5 (AOR 2.00, 95% CI 1.4 [...]

[PubMed](#) · [DOI](#)

Etiology of Failure in Revision Total Knee Arthroplasty: A Systematic Review and Meta-Analysis.

Lakpriya S, De C, Tahir M, Sanka SK, Pierce TP, Gwam C

BACKGROUND: Revision total knee arthroplasty (TKA) remains a burden and will continue to increase in incidence. The purpose of this systematic review and meta-analysis was to analyze the current revision rates and most common etiologies of failure of revision TKA.

METHODS: A comprehensive search of multiple electronic databases (PubMed, CINAHL Plus, EMBASE, and SCOPUS) was performed for all articles pertaining to this topic from 2014 to 2025. A total of 16 studies were included in this study. Study characteristics, patient demographics, rate of re-revision, and etiology for revision failures were extracted from each study.

RESULTS: Among 39,723 revision arthroplasties, the re-revision rate for this cohort was 13.5% ($n = 5,350$ revisions). The most common etiologies for revision failure were periprosthetic joint infection (27.5%; 95% confidence interval [CI], 26.4 to 28.8), instability (13.6%; 95% CI, 12.7 to 14.5), aseptic loosening (12.8%; 95% CI, 11.9 to 13.7), arthrofibrosis (9.4%; 95% CI, 8.5 to 10.4), unexplained pain (8.0%; 95% CI, 7.1 to 8.9), and periprosthetic fracture (7.6%; 95% CI, 6.8 to 8.5). There was substantial heterogeneity found among the studies, with very few showing publication bias.

CONCLUSIONS: Our revision failure rate of 13.5% is similar to other studies. The most common etiologies for revision TKA failure were infection, instability, and aseptic loos [...]

[PubMed](#) · [DOI](#)

Fractures of the coronoid process: state of the art.*Marinelli A, Riva M, Coliva F, Minerba M, Carbone G, Guerra E*

Coronoid fractures are rarely isolated and are much more frequently associated with other osseous or ligamentous structures injuries. On the basis of the coronoid fracture patterns, described by the O'Driscoll classification, it is possible to recognize three main patterns of injury that differ on traumatic mechanism and on associated lesions: posterolateral rotatory instability, posteromedial rotatory instability, and axial load injuries. The management of coronoid fractures is challenging and varies according to characteristics of the fracture, associated lesions, and amount of elbow instability. In general, operative treatment is indicated in every case the fracture is at least 50% of the whole coronoid, whether the sublime tubercle is involved, and whether the ulno-humeral joint is not perfectly reduced. In conclusion, the correct management of the coronoid, especially in the setting of complex elbow instability, represents a predictive factor for patient outcomes and functional results. The stability of the elbow, rather than the size of the coronoid fragment, is the main parameter for surgical indication, aimed to fix the coronoid and/or repair the associated lesions.

[PubMed](#) · [DOI](#)**Total hip arthroplasty in Italy: an observational, population-based study on surgical volume growth from 2001 to 2023 and forecasts until 2050 with six different statistical models.***Ciminello E, Cuccu A, Romanini E, Venosa M, Cazzato G, Tucci G, Boniforti F, Carpanese L et al.*

BACKGROUND: The number of total hip arthroplasty (THA) procedures has been steadily increasing worldwide, driven by aging population, improvements in surgical techniques and implant design. This study aimed to analyze the temporal trends of elective THA in Italy since 2001-2023 and forecast THA volumes up to 2050 to provide insights for healthcare planning.

MATERIALS AND METHODS: International Classification of Diseases, 9th Revision, Clinical Modification (ICD9-CM) coding system was used to extract records of interest (elective THA) from the Italian National Hospital Discharge Record database. Six statistical models were applied to forecast future THA volumes: logistic regression; Poisson regression; logarithmic regression; inverse/power regression; Poisson log-normal regression; and hierarchical Poisson regression with temporal effects (HPTE). Model performances were assessed by using error metrics and internal validation on the basis of a rolling-origin approach. An out-of-sample validation was conducted to ensure a robust assessment of forecasting reliability. THA volume forecasts were provided with 95% prediction intervals.

RESULTS: A total of 1,318,400 records for primary elective THAs performed in Italy since 2001-2023 were analyzed. The number of THAs increased by approximately 80%, rising from 68.270 in 2001 to 122.777 in 2023. Among the tested models, HPTE generally [...]

[PubMed](#) · [DOI](#)**Clinical relevance of patient-reported outcome measures in the surgical management of focal chondral defects of the knee: a systematic review.***Migliorini F, Maffulli N, Memminger MK, Hofmann UK*

INTRODUCTION: To evaluate clinical outcome following surgical management of focal chondral defects in the knee, patient-reported outcome measures (PROMs) are used. To give these measures meaning, parameters such as the minimal clinically important difference (MCID), patient-acceptable symptom state (PASS), minimally detectable change (MDC), clinically important difference (CID) and substantial clinical benefit (SCB) have been introduced. This systematic review investigated the MCID, SCB, CID, PASS and MDC of the most commonly used PROMs for assessing patients following surgical repair of focal chondral defects of the knee.

METHODS: This systematic review was conducted in accordance with the 2020 Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines and the recommendations of the Cochrane

Handbook for Systematic Reviews of Interventions. All clinical studies investigating tools to assess the clinical relevance of PROMs in the surgical repair of focal chondral defects of the knee were reviewed. In April 2025, the following databases were accessed: PubMed, Web of Science and Embase. The PROMs of interest included: the International Knee Documentation Committee (IKDC) questionnaire, the Knee Injury and Osteoarthritis Outcome Score (KOOS) and its related subscales activities of daily living (ADL), pain, quality of life (QoL), sports and recreational [...]

[PubMed](#) · [DOI](#)

Simultaneous versus staged bilateral total hip arthroplasty: a meta-analysis.

Soler F, Hernández J, Lamo-Espinosa JM, Benlloch M, Mariscal G

BACKGROUND: There is debate regarding the optimal timing of bilateral total hip arthroplasty (THA), with simultaneous or staged approaches considered. Cost-effectiveness is an important factor that influences resource allocation. The main aim of this study was to assess the cost implications of bilateral simultaneous (sim-THA) versus staged (st-THA) total hip arthroplasty. The secondary objective was to evaluate the efficacy and safety of sim-THA compared those with of st-THA.

METHODS: A systematic review and meta-analysis were conducted according to the Preferred Reporting Items for Systematic reviews and Meta-Analyses (PRISMA) guidelines. Studies comparing simultaneous and staged bilateral THA in terms of cost, complications, length of hospital stay, and patient outcomes were eligible. Odds ratios (ORs), mean differences (MD), and standard mean differences (SMD) with 95% confidence intervals (CIs) were calculated. The risk of bias was assessed using the Methodological Index for Non-randomised Studies (MINORS). Meta-analyses were performed using Review Manager version 5.4 (Cochrane, Oxford, United Kingdom).

RESULTS: A total of 20 observational studies including 13,984 patients were included. Sim-THA was associated with significantly lower total costs (SMD -0.54, 95% CI -0.92 to -0.16; $p = 0.005$) and shorter hospital stay (MD -2.90, 95% CI -4.38 to -1.42; $p = 0.0001$) than tho [...]

[PubMed](#) · [DOI](#)

Long-term outcomes of the modified Dunn procedure in moderate and severe slipped capital femoral epiphysis: a prospective case series with 7-year follow-up.

Fahmy M, Abdelazeem AH, Shawky MA

INTRODUCTION: Slipped capital femoral epiphysis (SCFE) is the most common hip disorder in adolescents and may lead to femoroacetabular impingement, early osteoarthritis, and long-term functional disability if inadequately treated. While in situ pinning remains the standard treatment for mild slips, it fails to correct the deformity in moderate and severe cases, potentially predisposing to degenerative changes. The modified Dunn procedure (MDP) was developed to restore proximal femoral anatomy through surgical hip dislocation while preserving vascular supply. The aim of the study is to evaluate the long-term radiological and functional outcomes of the MDP in patients with moderate (14 cases) and severe (10 cases) SCFE, and to assess the incidence of avascular necrosis (AVN), osteoarthritis, and other complications.

METHODS: A prospective case series was conducted between August 2015 and January 2019 at a single tertiary institution. A total of 24 hips with moderate-to-severe SCFE and open physis were treated using the MDP via surgical hip dislocation. Mild and acute-only slips were excluded. MDP was used as a primary procedure, performed early in severe slips and in selected moderate slips after clinical assessment. Patients were followed clinically and radiologically for a mean duration of 84 ± 2.6 months (range 80-88 months). Functional outcomes were assessed using the Harris [...]

[PubMed](#) · [DOI](#)

The presence of a positive blood culture should be regarded a risk factor for DAIR failure in cases of haematogenous total knee arthroplasty infection.

Pedemonte-Parramón G, Reynaga E, Molinos S, de Los Ríos JD, Vivero A, García-Oltra E, López-Pérez V, Paredes R et al.

PURPOSE: Late acute haematogenous total knee arthroplasty (LAH TKA) infections represent approximately 20% of all prosthetic infections. Debridement, antibiotics, and implant retention (DAIR) is a widely accepted treatment; however, it carries a significant risk of failure. The study aims to identify risk factors for DAIR failure in LAH TKA infections, focusing on blood cultures (BC) as a potential contributor.

METHODS: A retrospective cohort study was conducted on 37 cases of LAH TKA infections from 2015 to 2023. Patients were divided into two groups: 20 in the success group (SG) and 17 in the failure group (FG). The study analyzes various factors, including demographics, comorbidities, serum and joint fluid biomarkers, blood and intraoperative cultures, prior infection or antibiotic use, and surgical and post-surgical variables.

RESULTS: Positive BC were more frequent in the FG compared with the SG ($P = 0.03$), and were also more common in patients with a history of prior infection ($P = 0.03$). Logistic regression identified positive BC as the only significant predictor of DAIR failure (odds ratio (OR) 12, 95% confidence interval (CI) 1.1-18, $P = 0.04$), even after adjusting for other variables. Positive intraoperative cultures were more frequent in the FG, particularly in those with a prior infection history ($P = 0.08$), even if they were receiving antibiotics ($P = 0.05$).

CON [...]

[PubMed](#) · [DOI](#)

Iatrogenic femoral neck fractures or separation of the proximal femoral epiphysis during closed reduction of irreducible femoral head fracture-dislocations in children: a review of 12 cases.

Lu Y, Xue C, Canavese F, De Salvo S, Yao H, Huang D, Lin R, Zheng J et al.

OBJECTIVES: This study aimed to analyze the radiologic characteristics of irreducible pediatric femoral head fracture-dislocations to prevent potential iatrogenic femoral neck fractures (FNF) or separation of the proximal femoral epiphysis (SPFE).

METHODS: This is a retrospective review of patients who were skeletally immature and diagnosed with traumatic hip dislocations combined with femoral head fractures. The collected data included patient demographics, fracture classification, fragment ratio, combined injuries, urgent reductions, treatment strategies, complications, and final outcomes.

RESULTS: We treated 12 patients with femoral head fractures and dislocations; 11 out of 12 patients underwent urgent closed reduction (91.7%). Six of the patients failed closed reduction and experienced FNF ($n = 2$, 33.3%) or SPFE ($n = 4$, 66.7%). Five patients presented with avascular necrosis of the femoral head after open reduction with internal fixation via a surgical hip dislocation approach, and two patients required further surgical treatment. Analysis of radiographs and computed tomography (CT) scans of irreducible femoral head fracture-dislocations revealed that the fractured femoral head was perched on the sharp angle of the posterior wall of the acetabulum, with a fragment ratio of 18-27%. After recognizing the irreducibility, one case with a fragment ratio of 26% underwent immed [...]

[PubMed](#) · [DOI](#)

Orthopaedic publications receive fewer citations than general medicine articles: a scientometric analysis of temporal trends.

Migliorini F, Vaishya R, Maffulli N, Eschweiler J, Kobbe P, Betsch M, Schäfer L, Oliva F et al.

BACKGROUND: Orthopaedics, as a surgical speciality with a structurally narrower citing audience and lower citation density than general medicine, is systematically disadvantaged in any cross-disciplinary comparison relying on absolute citation counts. Whether this disadvantage is stable or has changed over time has not been previously quantified.

METHODS: The 100 most-cited articles indexed in the Web of Science Core Collection were identified for medicine and orthopaedics in 2014 and 2024. Citation data were extracted from a single snapshot taken on 19 January 2026. Total citations and citation velocity, defined as citations per year since publication, were compared between fields and across time points using Mann-Whitney U tests. Ratio-based analyses and distributional characteristics were also assessed.

RESULTS: Median citation counts were substantially higher in medicine than in orthopaedics in both years, reaching 5055 versus 288 in 2014 and 720.5 versus 34.0 in 2024, corresponding to 17.6- and 21.2-fold differences, respectively. Citation velocity followed the same pattern, with medicine maintaining an approximately 17- to 21-fold advantage at both time points. Within-field temporal analysis showed that citation velocity remained broadly stable in medicine between cohorts, whilst orthopaedics demonstrated reduced early citation accrual in the more recent cohort. Citatio [...]

[PubMed](#) · [DOI](#)

Sleep quality and nocturnal pain in patients with elbow disorders: a prospective multicenter study.

Schneider MM, Nietschke R, Migliorini F, Schoch C, Burkhart K, Hollinger B, Kimmeyer M, Lehmann L et al.

BACKGROUND: Mental health disorders have become a growing global concern. Sleep quality is known to have a crucial impact on mental well-being. Poor sleep contributes to increasing economic costs owing to reduced productivity as well as the development of mental disorders. Conversely, anxiety and depression often lead to sleep disturbances. Pain is a key factor affecting sleep quality. Studies have shown that musculoskeletal disorders, especially in the shoulder, may cause sleep disturbances. Research on the relationship between elbow pain and sleep quality remains limited. This study aims to evaluate the impact of common elbow pathologies on sleep quality.

METHODS: This prospective multicenter cohort study used self-reported outcome measures in patients with elbow disorders, including the Pittsburgh Sleep Quality Index (PSQI), to assess sleep quality. The PSQI was compared with other outcome measures.

RESULTS: A total of 664 patients (284 female, 380 male) were included, of whom 81% reported nocturnal pain. The mean age was 47.9 ± 14.2 years, body mass index (BMI) 26.9 ± 6.7 kg/m² and symptom duration 15.8 ± 7.3 months. The most common diagnoses were lateral epicondylar pain (66%), medial epicondylar pain, arthritis, ulnar nerve entrapment and biceps/triceps tendinopathies. The overall PSQI was 7.3 ± 3.8 , disabilities of arm, shoulder and hand questionnaire (DASH) 35.0 ± 19 . [...]

[PubMed](#) · [DOI](#)

A novel figure-of-eight polymeric cerclage wiring technique for greater trochanter fixation in revision arthroplasty and conversion of failed proximal femur fracture fixations to total hip arthroplasty.

Leggieri F, Martín Cocilova FN, Secci G, Stimolo D, Zanna L, Civinini R, Innocenti M

INTRODUCTION: Periprosthetic femoral fractures occur in 0.8-10% of primary total hip arthroplasties (THA) and up to 17.5% for revision THA (revTHA), with isolated greater trochanter (GT) fractures being the most predominant pattern. This study aimed to evaluate a novel figure-of-eight cerclage wiring technique for greater trochanter fixation in hip replacement cases.

METHODS: These retrospectively collected data included 36 hips (36 patients) undergoing revision THA or conversion of failed intertrochanteric fracture to THA with figure-of-eight cerclage wiring between April 2021 and June 2023. Patients under 18 years, with incomplete records, or lost to follow-up before 24 months were excluded. Primary outcomes included Trendelenburg sign, periprosthetic infections, and hip dislocation rates. Secondary outcomes evaluated THA stem subsidence and trochanteric migration using calibrated digital imaging. Statistical analysis included descriptive statistics, independent t-tests, linear regression, and correlation analyses with significance set at $p < 0.05$.

RESULTS: Mean follow-up was 404.76 ± 22.51 days. Patient demographics showed mean age 76.5 ± 12.3 years, body mass index (BMI) 25.8 ± 4.5 kg/m², with 73.8% female patients. At final follow-up, one patient (2.8%) experienced post-surgical infection, one patient (2.8%) had dislocation, and three patients (8.3%) exhibited positive T [...]

[PubMed](#) · [DOI](#)

Minimal clinically important difference (MCID), patient-acceptable state (PASS), minimally detectable change (MDC), and substantial clinical benefit (SCB) in patients who have

undergone arthroscopic surgery for femoroacetabular impingement: a systematic review.

Migliorini F, Maffulli N, Schäfer L, Jeyaraman M, Betsch M, Mahmoud M

INTRODUCTION: Hip arthroscopy has been established as a successful management for femoroacetabular impingement (FAI) syndrome. Patient-reported outcome measures (PROMs) are frequently used to assess treatment results in patients. However, clinically important outcome values have recently been calculated to better understand and interpret PROMs and define successful treatment. This systematic review aimed to investigate the minimal clinically important difference (MCID), patient-acceptable state (PASS), substantial clinical benefit (SCB), and minimally detectable change (MIC) of the most commonly used PROMs for assessing patients who have undergone arthroscopic surgery for FAI.

MATERIAL AND METHODS: This study was conducted according to the Preferred Reporting Items for Systematic Reviews and Meta-Analyses: the 2020 PRISMA statement. In October 2025, the following databases were accessed: PubMed, Web of Science, and Embase. All clinical studies investigating tools to assess the clinical relevance of PROMs used to evaluate patients who have undergone arthroscopic surgery for FAI were accessed. Only studies that assessed the minimal MCID, PASS, SCB, and/or MIC were considered.

RESULTS: The methodological quality of the 21 included studies was assessed, and the overall risk of bias was found to be low-to-moderate. The main characteristics of the included studies were analyzed and [...]

[PubMed](#) · [DOI](#)

Efficacy and safety of MIPO versus ORIF in distal tibial fractures: a systematic review and meta-analysis.

Hassaan AM, Hassan AA, Abdelkhalik WS, Saad AM, Hosny AS, Elfeky BAEH, Mandour I, Ragheb MG et al.

BACKGROUND: Distal tibial fractures are challenging to manage owing to limited soft-tissue coverage and compromised blood supply. Minimally invasive plate osteosynthesis (MIPO) and open reduction and internal fixation (ORIF) are among the widely used surgical options, but the optimal approach remains unclear. MIPO preserves periosteal blood flow with smaller incisions, while ORIF offers direct visualisation but increases soft-tissue damage. With growing comparative evidence, an updated evaluation of these techniques is necessary.

AIM: To compare the efficacy and safety of MIPO versus ORIF in adult patients with distal tibial fractures.

METHODS: Following PRISMA guidelines and the Cochrane Handbook, we searched in PubMed, Embase, Cochrane, Scopus, and Web of Science till November 2025. Randomised trials and cohort studies comparing MIPO and ORIF were included. Data extraction was performed independently by two reviewers. Risk of bias was assessed using RoB 2 for RCTs and the Newcastle-Ottawa Scale for observational studies. Random-effects models were used to estimate pooled mean differences and odds ratios, and heterogeneity was assessed using I². Sensitivity and subgroup analyses were conducted by study design.

RESULTS: Nine studies involving 530 patients met our inclusion criteria. No significant differences were found between MIPO and ORIF in AOFAS scores, union time, oper [...]

[PubMed](#) · [DOI](#)

Comparative efficacy and safety of aspirin, low-molecular-weight heparin, and rivaroxaban for deep venous thrombosis prophylaxis after hip arthroplasty for femoral neck fracture: a single-center randomized controlled trial.

Miao S, Yue Q, Zhang X, Lu J, Zhang Y, Lu N, Wei L

OBJECTIVE: To evaluate the efficacy and safety of aspirin, low-molecular-weight heparin (LMWH), and rivaroxaban for deep vein thrombosis (DVT) prophylaxis after total hip arthroplasty (THA) or bipolar hemiarthroplasty (BHA) for femoral neck fracture, with subgroup analysis stratified by surgical type.

METHODS: This single-center randomized controlled trial (RCT) was conducted between June 2024 and June 2025. A total of 217 consecutive patients with femoral neck fracture undergoing hip arthroplasty were initially screened, and 161 eligible patients were finally enrolled and randomly assigned to the aspirin group, LMWH group, or rivaroxaban group at a 1:1:1 ratio. All patients received standardized intermittent pneumatic compression (IPC) and graduated compression stockings for mechanical DVT prophylaxis, as well as a unified postoperative rehabilitation protocol. The primary outcome was the incidence of lower extremity DVT within 12 weeks

postoperatively, confirmed by color Doppler ultrasound. The secondary outcome was the incidence of bleeding-related complications graded by the International Society on Thrombosis and Hemostasis (ISTH) criteria. All patients were followed up at 6, 8, and 12 weeks postoperatively, with routine bilateral lower extremity ultrasound performed at 12 weeks regardless of symptoms. Subgroup analysis was performed on the basis of surgical type (THA ver [...])

[PubMed](#) · [DOI](#)

Risser growth plate injury in unstable paediatric pelvic fractures: a multicentre retrospective study.

Oransky M, Aulisa AG, Roncoroni A, Zoppi AR, Mata M, Rohayem MA, Falciglia F, Toniolo RM

INTRODUCTION: Unstable pelvic fractures in children are rare but severe injuries and are associated with high-energy trauma. Unlike adults, children have open growth plates, particularly the Risser growth plate (RGP), which is vulnerable to injury. The presence of growth plates, such as the triradiate cartilage and Risser's plate, represents point of vulnerability that determines distinctive fracture patterns, similar to those seen with Salter-Harris growth plate injuries. Lateral compression fracture is the most common mechanism and results in an irreducible crushing of the sacral alae, causing a rotational deformity. Both vertical and lateral compression displaced pelvic fractures require careful evaluation of the posterior iliac apophysis-associated injury. Failure to treat Risser growth plate injury (RGPI) can lead to severe long-term problems such as limb length discrepancy and scoliosis. This study aimed to determine how commonly RGPI occurs in unstable paediatric pelvic fractures, link it to specific fracture patterns and propose a subgroup type to improve treatment.

MATERIALS AND METHODS: This multicentre retrospective study included 40 children aged up to 12 years with unstable pelvic fractures (AO/OTA 61B and 61C). The patients were treated between 1987 and 2022 at trauma centres in Italy, Argentina and Venezuela. We reviewed patient demographics, injury mechanisms a [...]

[PubMed](#) · [DOI](#)

Total femur replacement: function, quality of life, and gait: a multicentric EMSOS study.

Valentini M, Svehlik M, Leithner A, Bergovec M, EMSOS TFR Study Group, Smolle MA

BACKGROUND: Total femur replacements (TFRs) involve the replacement of the whole femoral bone, reconstruction of the hip and knee joints, and soft tissue reconstruction for function and joint stability. The aim of this multicentric European Musculo-Skeletal Oncology Society (EMSOS) study is to report on the functional outcome, quality of life, as well as gait analysis in patients with TFR.

MATERIALS AND METHODS: Retrospective data from 12 international participating centers were collected. In total, 65 patients [median (IQR) age at surgery 40 (18-56) years, 60% male] who received a TFR owing to oncologic or nononcologic indications between 1 January 1990 and 31 March 2024 were included. Patient-related outcome measures (PROMs), range of motion (ROM) of the hip and knee, and gait videos were collected.

RESULTS: At a median (IQR) follow-up of 28 (14-126) months, the median (IQR) percentage Musculoskeletal Tumor Society (MSTS) score was 63% (43-73%), 12-Item Short-Form Health Survey (SF-12) was 70% (55-79%), Harris Hip Score (HHS) was 63% (52-77%), and Oxford Knee Score (OKS) was 63% (52-79%); the median (IQR) extension-to-flexion ROM were 80° (70-97.5°, hip) and 70° (40-90°, knee). Gait videos were available for 24/65 patients (37%). The median (IQR) total Edinburgh Visual Gait Score of the operated side was 8.5 (6-13), with deviations from norm in all patients (24/24). Compens [...]

[PubMed](#) · [DOI](#)

Vascularized versus nonvascularized free fibular grafts in reconstruction of post-traumatic critical long bone defects, a comparative study.

Fouaad AA, Massoud AH, Shaban M, Abdel-Naby M, Abulsoud MI

BACKGROUND: Reconstruction of critical-sized long bone defects is a complex orthopedic challenge. Free fibular grafts, vascularized (FVFG) and nonvascularized (NVFG), are established reconstructive options, but comparative clinical outcomes remain uncertain.

PURPOSE: To compare clinical and radiological outcomes of FVFG and NVFG in patients with post-traumatic critical bone defects more than 10 cm.

METHODS: A randomized controlled trial was conducted with 50 patients assigned equally to FVFG or NVFG groups. The primary outcome was time to union, while secondary outcomes included graft hypertrophy, functional scores (DASH and LEFS), complication rates, and donor site morbidity.

RESULTS: The mean time to union was 5.78 months in the FVFG group and 6.17 months in the NVFG group, showing no statistically significant difference ($p = 0.447$). Rates of graft hypertrophy, functional recovery, and complications were comparable between the groups.

CONCLUSIONS: Both FVFG and NVFG provide effective reconstruction for critical bone defects, with nearly similar healing times and functional outcomes. NVFG may represent a less complex alternative in selected cases.

[PubMed](#) · [DOI](#)

Five-year outcomes of medial open-wedge high tibial osteotomy with lateral supplemental lag screw in patients with obesity.

Tian YH, Lee KH, Huang CW, Lin SY, Yang JC

BACKGROUND: Obesity contributes to the accelerated progression of knee osteoarthritis. Medial open-wedge high tibial osteotomy (MOWHTO) is a joint-preserving surgical intervention for unicompartmental knee osteoarthritis; however, its efficacy in patients with obesity remains controversial. This study aimed to evaluate the 5-year clinical and radiographic outcomes of MOWHTO with lateral supplemental lag screw fixation in patients with obesity.

MATERIALS AND METHODS: This retrospective cohort study included patients with obesity who underwent MOWHTO between 2017 and 2020, with a minimum follow-up of 5 years. All procedures were performed by a single surgeon using three-dimensional printed, patient-specific instrumentation, locking plates, and a lateral supplemental lag screw. Clinical outcomes were assessed using the visual analog scale (VAS) and Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC). Radiographic parameters, including the weight-bearing line percentage (WBL%) and medial proximal tibial angle (MPTA), were evaluated.

RESULTS: Significant improvements in VAS and WOMAC scores were observed at all postoperative time points ($p < 0.001$) and were accompanied by improved radiographic alignment, with WBL% shifting toward the Fujisawa point and increased MPTA values. At 5 years, mild regression in alignment was noted; however, the overall correction was [...]

[PubMed](#) · [DOI](#)

Outcomes of surgical fixation of osteochondral fracture: a systematic review.

Pang AS, Oviya R, Tan JJ, Lim KSA, Hui JHP, Tan SHS

BACKGROUND: Dealing with knee osteochondral defects presents a substantial clinical challenge due to the variable nature of repair outcomes and intricate biomechanical complexities inherent to the knee joint. Numerous surgical alternatives exist for addressing knee osteochondral lesions, yet there is limited evidence available for comparing their respective outcomes.

METHODS: According to the Preferred Reporting Items for Systematic Review and Meta-analyses (PRISMA) guidelines, a systematic literature search was conducted to identify publications from inception to September 2023 on PubMed, Cochrane, and Medline academic search engines with the MeSH terms "osteochondral" AND ("patella" OR "patellar" OR "patellofemoral" OR "femoral condyle lesions" OR "trochlear lesions"). Inclusion criteria were clinical human studies, English language, subjects who underwent surgical management (fixation, reconstruction, or debridement) for knee osteochondral lesions with reported treatment outcomes, and a minimum sample size of ten patients. The methodological index for non-randomized studies (MINORS) was used to evaluate the non-randomized studies' methodological quality. Main outcome measurements Pediatric International Knee Documentation Committee (Pedi IKDC) scale, Tegner-Lysholm Scoring Scale, Tegner Activity Score, Visual Analog Scale, Knee Society Score (KSS) Score, Kujala Score, and F [...]

[PubMed](#) · [DOI](#)

Do modular tapered long stems differ in stability and function? A prospective comparison of two stems in complex femoral revision THA.

Fahmy M, Shawky MA

BACKGROUND: Revision total hip arthroplasty (rTHA) in the setting of substantial proximal femoral bone loss is technically challenging. Modular tapered fluted stems provide predictable diaphyseal fixation while allowing independent adjustment of version, offset, and limb length. Among these, two commonly used systems-modular tapered fluted stem Type A (Revitan™) and Type B (LIMA Modulus™)-have limited direct comparative evidence. This study aimed to prospectively compare radiographic stem subsidence (primary outcome), as well as functional outcomes, complications, survivorship, and secondary outcomes, between Type A and Type B modular long stems in femoral rTHA.

METHODS: In this single-center randomized prospective study, 110 patients undergoing femoral revision rTHA were randomly assigned to receive either Type A (n = 55) or Type B (n = 55) stems. All procedures were performed by experienced revision surgeons under standardized perioperative and rehabilitation protocols. Radiographs were analyzed for stem subsidence, osseointegration, and limb-length restoration. Functional outcomes were assessed using the Harris Hip Score (HHS), Oxford Hip Score (OHS), and European Quality of Life Visual Analogue Scale (EQ-VAS) at baseline and final follow-up (mean 61.4 months). Complications and stem survivorship were recorded prospectively. Statistical analysis included paired and unpaired [...]

[PubMed](#) · [DOI](#)

Clinical comparison of single posterolateral plate with medial-cannulated-screw fixation and double-plate fixation for extra-articular distal humerus fractures.

Jung HS, Chu MS, Lee JS

BACKGROUND: Double-plate fixation is the gold standard for extra-articular distal humerus fractures, but it carries a substantial risk of postoperative ulnar neuropathy. Fixation using a single posterolateral plate with a medial cannulated screw may reduce ulnar neuropathy while maintaining fracture stability. This study aimed to compare the clinical outcomes of single-plate-with-medial-screw fixation versus double-plate fixation for extra-articular distal humerus fractures.

MATERIALS AND METHODS: Fifty-six patients who underwent surgery for extra-articular distal humerus fractures (Arbeitsgemeinschaft für Osteosynthesefragen/Orthopaedic Trauma Association [AO/OTA] classification A2 or A3) between January 2018 and August 2024 were divided into a double-plate group and a single-plate-with-medial-screw group. We conducted a retrospective, nonrandomized comparative study. The double-plate fixation was used in 30 patients from January 2018 to October 2021, while the single-plate fixation with a medial screw was used in 26 patients from November 2021 to August 2024. All surgeries were performed using a posterior paratricipital approach. Bony union, radiographic healing, and loss of reduction were evaluated. Postoperative pain scores (visual analog scale at 2 days after the operation), operative time (minutes), elbow range of motion, elbow function (Mayo Elbow Performance Score [MEP] [...])

[PubMed](#) · [DOI](#)

A study on the association between tibial plateau fractures and intra-articular soft-tissue injuries under valgus injury mechanisms.

Duan S, Wang S, Zhu T, Li Q, Zhang M

BACKGROUND: While recent investigations have focused on injury mechanism classifications of tibial plateau fractures (TPFs), the association between valgus TPFs and concomitant soft-tissue damage involving menisci and ligaments remains insufficiently elucidated. This study aimed to characterize intra-articular soft-tissue injuries associated with various valgus TPFs and assess the predictive value of lateral plateau depression (LPD) and widening (LPW). Additionally, the analysis extended to other injury mechanisms.

MATERIALS AND METHODS: This study included adult patients with acute tibial plateau fractures who had complete imaging data, excluding patients with open fractures and multiple fractures throughout the body. Imaging examinations were used to assess the fracture mechanism and intra-articular soft-tissue damage.

RESULTS: The study retrospectively analyzed the clinical data of 138 patients with valgus injury TPFs in our hospital. The study compared the incidence of TPFs with intra-articular soft-tissue damage under different injury mechanisms. The result demonstrated that the incidence of valgus hyperextension TPFs combined with medial collateral ligament injuries was relatively high (61.5%) compared with TPFs with other injury mechanism. Multivariable logistic regression and smooth curve fitting revealed significant dose-response relationships of LPD (odds ratio [OR] [...])

The coronoid as the key fragment of trans-ulnar fracture-dislocations of the elbow: Insights from a retrospective cohort comparison using the coronoid-centric Mayo classification system.

Henssler L, Kerschbaum M, Zanklmaier M, Wieczorek LA, Alt V, Klute L

BACKGROUND: Trans-ulnar fracture-dislocations of the elbow are rare injuries with complex fracture patterns and variable outcomes. Traditional classification systems offer limited prognostic value. A recently introduced coronoid-centric Mayo classification distinguishes injury subtypes based on coronoid attachment and identifies trans-ulnar basal coronoid (TUBC) fractures as a particularly challenging entity. This study evaluated outcomes across Mayo fracture types and explored factors associated with inferior results in TUBC injuries.

MATERIALS AND METHODS: In this retrospective cohort study, surgically treated trans-ulnar elbow fracture-dislocations managed at a level I trauma center between 2010 and 2022 were identified and classified according to the Mayo system. Demographic data, injury characteristics, surgical management, radiographic outcomes, and complications were recorded. Functional outcomes were assessed after a minimum follow-up of 12 months using the Mayo Elbow Performance Score (MEPS); Oxford Elbow Score (OES); Quick Disabilities of Arm, Shoulder and Hand Questionnaire (QuickDASH); European Quality of Life Five-Dimension, Five-Level Version (EQ-5D-5L); and range-of-motion measurements. Radiographs were analyzed for union, instability, heterotopic ossification, and post-traumatic osteoarthritis (OA).

RESULTS: A total of 52 patients were included (14 trans-olecr [...]

PubMed · DOI

Feasibility study of a robot-assisted endoscopic technique for plate osteosyntheses of the acetabulum and anterior pelvic ring.

Hinz N, Althoff G, Thomaschewski M, Bonke LA, Eberenz A, Münch M, Behrends L, Männel G et al.

BACKGROUND: Minimally invasive stabilization of acetabular and pelvic ring fractures using endoscopic techniques has become increasingly important. The logical advancement of conventional endoscopic techniques is a robot-assisted approach, which benefits from the advantages of robot-assistance systems (e.g., more degrees of freedom for instruments and improved visualization). The aim was therefore to investigate the feasibility of a robot-assisted endoscopic technique for plate osteosyntheses of the acetabulum and anterior pelvic ring.

MATERIALS AND METHODS: In this two-part feasibility study, four different plate osteosyntheses were performed endoscopically using the Hugo™ robot-assisted surgery system, first on ten synthetic pelvic models and then on ten human cadavers. During robot-assisted dissection for the preperitoneal approach, identification of ten relevant anatomical landmarks was also assessed. In both parts, the success, number of drilling errors, and time for each plate were described as learning curves and analyzed using linear regression.

RESULTS: The infrapectineal plate could be successfully performed in 100% of synthetic models, the posterior column plate in 100%, the suprapectineal plate in 90%, and the superior pubic ramus plate in 80%. Learning curves could be observed for the number of drilling errors per plate (e.g., 0.67 to 0, $p = 0.009$) and the time t [...]

PubMed · DOI

Favorable subjective clinical outcomes after revision hip arthroscopy for femoroacetabular impingement syndrome despite a high rate of capsular defects and increased capsular thickness.

Cao Z, Gao G, Lin W, Zhu Y, Zhou X, Wang J, Xu Y

BACKGROUND: The capsular healing status and capsular thickness changes in patients with femoroacetabular impingement syndrome (FAIS) following revision hip arthroscopy are poorly documented, and their relationship with subjective clinical outcomes remains unclear. The purpose of this work is to evaluate the incidence of capsular defects and changes in capsular thickness using magnetic resonance imaging (MRI) following hip arthroscopy in patients with FAIS, and to compare the subjective clinical outcomes between patients with and without capsular healing after revision hip arthroscopy.

PATIENTS AND METHODS: Consecutive patients with FAIS who underwent revision hip arthroscopy between 2013 and 2023 were included. Patients were categorized into two groups on the basis of capsular healing status after revision hip arthroscopy. Patient-reported outcomes (PROs) were collected preoperatively and at minimum 2-year follow-up, including the modified Harris Hip Score (mHHS), International Hip Outcome Tool-12 (iHOT-12), Hip Outcome Score-Activity of Daily Living Scale (HOS-ADL), Hip Outcome Score-Sport Specific Subscale (HOS-SSS), and visual analog scale (VAS). Achievements in minimal clinically important difference (MCID) and patient acceptable symptom state (PASS), patient satisfaction, and re-revision rates were evaluated and compared between groups. Capsular thickness was assessed by [...]

[PubMed](#) · [DOI](#)

From bench to bedside: a novel suture-augmented prosthesis for greater trochanteric fixation in osteoporotic hip arthroplasty.

Yang M, Yang J, Wang X, Wu Z, Bian S, Sheng R, Liao F, Qin K et al.

BACKGROUND: Stable fixation of the greater trochanter during hip arthroplasty for unstable osteoporotic intertrochanteric fractures remains a significant challenge. Conventional techniques depend on securing the bone-implant interface, which is particularly compromised in osteoporotic bone. Here, we propose a novel conceptual approach that establishes a direct mechanical bridge from the abductor tendon to the prosthesis, thereby reducing reliance on the fragile bone-implant interface.

METHODS: In this two-stage translational study, we progressed from biomechanical validation to clinical application. First, using a decalcified caprine femoral model simulating osteoporosis, three fixation constructs were compared: locking plate (LP), suture-augmented locking plate (LPSA), and Kirschner-wire tension band (KWTB). Ultimate load and construct stability were evaluated. Biomechanical testing confirmed the principle of suture-mediated load sharing and highlighted the intrinsic weakness of screw fixation in osteoporotic bone. Guided by these results, we designed a novel femoral prosthesis that eliminates screw fixation, employs sutures as the primary load-bearing element, and incorporates integrated suture anchor tunnels. This prosthesis was then assessed in a retrospective series of 15 consecutive elderly patients with osteoporotic intertrochanteric fractures. Clinical outcomes were ev [...]

[PubMed](#) · [DOI](#)

Updating radiographic parameters for the healthy growing hip: are current parameters still valid?

Hütter K, Smolle MA, Butter S, Schroedter R, Tschauner S, Kraus T

BACKGROUND: This study aimed to evaluate whether the acetabular angle (AA) by Tönnis and lateral center-edge angle (LCEA) by Wiberg-key radiographic parameters in pediatric hip assessment-have changed in a contemporary pediatric population compared with historical reference values, considering trends in earlier skeletal maturation and body composition.

MATERIALS AND METHODS: This retrospective diagnostic accuracy study assessed changes in AA and center-edge angle (CE) angles in a contemporary cohort. A total of 1774 anteroposterior pelvic radiographs (3548 hips) from patients aged 0-18 years without hip pathology were analyzed. Radiographs were obtained between 2006 and 2018 and measured using the Supervisely digital platform. Standardized anatomical landmarks were applied. Data were stratified by age and sex and compared with historical values using two-sample t-tests ($p < 0.05$). To minimize measurement bias, two independent observers conducted all assessments using standardized digital protocols.

RESULTS: A total of 1774 patients aged 0-18 years (mean age, 8.30 ± 5.20 years; 666 females, 1108 males) were evaluated. AA values did not show statistically significant differences in 11 of 16 age groups compared with historical data ($p > 0.05$). In contrast, LCEAs were significantly higher in the contemporary cohort, especially in the 5-8, 9-12, and 12-16 age groups (all $p < 0.01$) [...]

[PubMed](#) · [DOI](#)

In-hospital deep vein thrombosis after tibial plateau fractures: incidence, laterality/anatomy, and risk factors in a multicenter retrospective cohort of 3366 patients.

Yang S, Li G, Li Y, Long Y, Zhang J, Liu L, Zhao Z, Sun C et al.

BACKGROUND: We aimed to determine the cumulative in-hospital incidence, anatomic distribution, and predictors of duplex ultrasonography-detected lower-extremity deep vein thrombosis (DVT) in patients hospitalized with tibial plateau fractures under routine thromboprophylaxis and protocolized surveillance.

METHODS: We retrospectively analyzed consecutive adults (≥ 18 years) admitted to two level-I trauma centers (November 2014-January 2024). Bilateral duplex ultrasonography was performed on admission/preoperatively (as soon as feasible after arrival and prior to surgery) and repeated serially during hospitalization (approximately every 5-7 days and when clinically suspected). The primary outcome was cumulative in-hospital DVT from admission to discharge; isolated intermuscular calf-muscle vein thrombosis (soleal/gastrocnemius) was recorded descriptively but excluded from the primary endpoint. Multiple injuries (polytrauma) were defined as ≥ 1 additional acute traumatic lesion beyond the index tibial plateau fracture documented on admission. Pulmonary embolism (PE) was assessed only when clinically suspected and confirmed by computed tomography pulmonary angiography (CTPA). Multivariable logistic regression identified independent predictors; receiver operating characteristic (ROC) analysis evaluated discrimination for key continuous predictors.

RESULTS: Among 3366 patients, 675 [...]

[PubMed](#) · [DOI](#)

Retained articulating spacers (1.5 stage) in chronic knee periprosthetic joint infections: an analysis of associate factors and outcomes in a single center cohort of patients.

Balato G, Ascione T, Festa E, De Mauro D, Di Gennaro D, Rosa D, Mariconda M

BACKGROUND: Two-stage revision remains the gold standard for chronic periprosthetic joint infection (PJI), but it can significantly impact patients' quality of life and functional recovery. The 1.5-stage revision, involving indefinite spacer retention, has emerged as a potential alternative in selected high-risk patients. Our questions were: (1) What are the early functional outcomes, including their infection-free survival? (2) Which pre- and perioperative factors are associated with spacer retention? and (3) What is the mechanical survivorship of retained spacers?

MATERIALS AND METHODS: This was a retrospective observational study including 108 consecutive patients with chronic PJI between 2016 and 2020 who were followed prospectively for clinical outcomes. Functional status and quality of life were assessed using the EuroQol-5-Dimension-5-Level (EQ-5D-5L), Western Ontario and McMaster Universities Osteoarthritis Index (WOMAC), and Knee Society Score (KSS) before spacer placement and prior to scheduled reimplantation. Infection-free status was defined as the absence of clinical, radiographic, and laboratory signs of infection recurrence over a 24-month follow-up. Univariate and multivariate logistic regression were used to identify factors associated with spacer retention.

RESULTS: All patients initially underwent articulating spacer implantation as part of a planned two-st [...]

[PubMed](#) · [DOI](#)

Moderate leg length discrepancy: a long-term risk factor for hip osteoarthritis?

Aprato A, Donis A, Via RG, Scandurra A, Tuè F, Audisio A, Fusini F, Massè A

BACKGROUND: Leg length discrepancy (LLD) has been implicated as a biomechanical factor contributing to hip osteoarthritis (OA), yet the extent of its influence remains unclear. This study examines the correlation between moderate LLD (≥ 10 mm) and hip OA progression, focusing on asymmetrical OA distribution in patients over 65 years old.

MATERIALS AND METHODS: A retrospective analysis was conducted from a database of 1672 full-length standing X-rays. Patients under 65 years, with deformities, unilateral limb issues, or prosthetics, were excluded; therefore, the study group was composed of 220 patients. Tibial and femoral lengths were measured bilaterally, and hip OA severity was assessed using the Tönnis classification. Statistical analyses included Pearson's Chi-squared test and linear regression to explore correlations between LLD and OA distribution.

RESULTS: Among the sample, 18% showed an LLD ≥ 10 mm. A significant correlation was found between LLD and the asymmetrical distribution of hip OA ($p = 0.002$), with higher OA severity observed in the hypometric limb. Linear regression analysis suggested that each millimeter of LLD corresponded to a 0.74-unit change in OA severity difference between hips.

CONCLUSIONS: This study highlights a significant association between moderate LLD and contralateral hip OA in the elderly, emphasizing the biomechanical impact of asymmetrical [...]

[PubMed](#) · [DOI](#)

Authorship, titles and open access as drivers of citation performance in orthopaedics: a scientometric analysis.

Migliorini F, Vaishya R, Rivera F, Eschweiler J, Kobbe P, Betsch M, Oliva F, Maffulli N

BACKGROUND: Bibliometric analyses are increasingly used to explore how scientific knowledge is created, disseminated, and perceived. In orthopaedics, research output has expanded rapidly over the past decade, yet the factors determining whether an article achieves wide visibility and scholarly impact remain poorly understood. Beyond the inherent quality of a study, elements such as authorship patterns, title construction, and open access (OA) availability may play an essential role in shaping citation performance. However, evidence in this field is still limited and sometimes contradictory, highlighting the need for large-scale, field-specific analyses.

METHODS: Orthopaedic publications from 2010 to 2020 were identified in Scopus using the keyword 'orthopaedic'. After duplicate removal, 97,806 unique articles were included with complete data on authorship, titles, citation counts, study design, and OA status. Citation rates were normalised per year since publication. Associations between bibliographic features and citation performance were assessed using multiple linear regression, while differences across title styles and study designs were evaluated with comparative statistical testing. Exploratory modelling was performed to identify combinations of authorship and title characteristics linked to the highest predicted citation rates.

RESULTS: Larger author teams were associa [...]

[PubMed](#) · [DOI](#)

Interpreting nonsignificant findings: Reassessing the safety of self-locking standalone cages in octogenarian patients undergoing ACDF.

Abudayeh A, Fishchenko I

[PubMed](#) · [DOI](#)

A novel, simple, and affordable technique for arthroscopic soft-tissue tenodesis of the long head of the biceps tendon.

Fossati C, Ioppoli A, Ravaglia R, Vaja F, Menon A, Randelli PS

BACKGROUND: The purpose of this study was to describe a novel, simple, and implant-free arthroscopic soft-tissue tenodesis technique of the long head of the biceps tendon (LHBT) to the rotator cuff using an absorbable monofilament suture and to evaluate its clinical and functional outcomes at a minimum 1-year follow-up.

METHODS: A retrospective case series of 23 patients (mean age 58.0 ± 7.8 years) who underwent arthroscopic rotator cuff repair with concomitant LHBT soft-tissue tenodesis between June 2021 and June 2023 was analyzed. Functional outcomes were assessed using Constant-Murley Score (CMS), American Shoulder and Elbow Surgeons (ASES) score, Single Assessment Numeric Evaluation (SANE), visual analog scale (VAS) for pain, and the Long Head of Biceps (LHB) score. Supination strength was measured with a handheld dynamometer and compared with the contralateral side. Incidence of Popeye deformity and tenderness over the bicipital groove were recorded.

RESULTS: Objective Popeye deformity was observed in 13% of patients, with subjective concern reported by only 4.3%. Supination strength and LHB scores were similar to the contralateral side (means of 104.65 versus 104.64 N; LHB 93.7 versus 94.6 points). The mean CMS and ASES scores were 90.3 ± 12.4 and 89.6 ± 15.9 points, respectively. The SANE score averaged 87.4 ± 20.9 , and the VAS for pain was low (1.65 ± 2.59 cm).

CONCL [...]

[PubMed](#) · [DOI](#)

ORIF of three- and four-part proximal humeral fractures: a retrospective study of long-term clinical and complication outcomes of two-plate fixation methods.

Conteduca J, Gasbarra E, Rausa I, Casto A, Valentino L, Solarino G, Meccariello L, Rollo G

BACKGROUND: The aim of this study was to compare clinical outcomes and complication rates during long-term follow-up (≥ 6 years) after treatment for three- and four-part proximal humeral fractures using two different types of locking plates.

MATERIALS AND METHODS: A total of 113 patients with three- and four-part proximal humeral fractures who underwent surgery between September 2012 and January 2019 were enrolled retrospectively. Data for 49 patients [9 males, 40 females; mean age [standard deviation] (SD) 68.9 (5.8) years] treated with a PGR (intrauma) plate (group A) and 51 patients [10 males, 41 females; mean age (SD): 66.0 (7.5) years] treated with a PHILOS humeral plate (PHP, group B) were available at the last follow-up. The mean follow-up periods in groups A and B were 10.8 and 10.2 years, respectively, with a minimum follow-up of 6 years. At the final follow-up evaluation, functional outcomes were assessed using the Oxford Shoulder Scale (OSS), Simple Shoulder Test (SST), and Disabilities of the Arm, Shoulder and Hand (DASH) questionnaires. X-ray evaluation was also performed, and complications were recorded.

RESULTS: The mean OSS score in the PGR and PHP groups at follow-up was 42.9 and 39.1, respectively ($p > 0.05$). The SST score (PGR: 7.5 ± 2.0 ; PHP: 7.3 ± 4.0) and DASH score (PGR: 22.1 ± 5.5 ; PHP: 20.5 ± 4.0) were similar in both groups ($p > 0.05$). In our serie [...]

[PubMed](#) · [DOI](#)

Incidence rates and treatment of the transcervical fracture of the neck of femur in Italy: is total hip arthroplasty an increasingly preferred approach? A population study on trends between 2001 and 2023 based on 1,120,770 hospital discharge records.

Ciminello E, Romanini E, Venosa M, Cazzato G, Tucci G, Boniforti F, Carpanese L, Cuccu A et al.

INTRODUCTION: Transcervical femoral neck fractures (TFNFs) are among the most devastating fragility fractures in the elderly. TFNF are associated with excess 1-year mortality rates ranging from 15% to 30%. Treatments include conservative methods, internal fixation, and arthroplasty (partial or total hip arthroplasty). This study aims to analyze the changes in incidences of TFNF in the Italian population between 2001 and 2023 and the evolution of the choices of treatment.

MATERIALS AND METHODS: Using hospital discharge record (HDR) data from 2001 to 2023, records with ICD9-CM codes for femoral neck fractures (820.0 and 820.1) among diagnoses were selected and categorized into four treatment groups: total arthroplasty, partial arthroplasty, fixation, and conservative. Time series were analyzed with stratification by sex and age.

RESULTS: The extracted data included 1,120,724 records of TFNFs, with 871,161 cases treated surgically (total or partial arthroplasty or internal fixation) and 249,563 treated conservatively; the average patient age was 79.1 years, with a higher proportion of women (72.8%). Partial hip arthroplasty was the preferred treatment overall. For younger patients, in the age classes < 45 and 45-54 years, fixation was the most chosen treatment. Over time, the use of the conservative treatment decreased from 27.5% in 2001 to 14.6% of cases in 2023. The use of par [...]

[PubMed](#) · [DOI](#)

Does latissimus dorsi tendon transfer provide a normal scapular rhythm and external rotation strength in posterosuperior massive irreparable rotator cuff tears? A kinematic analysis in a retrospective cohort.

Calvo C, Fiumana G, Brigo A, Ruggiero ED, Bonfatti R, Donà A, Micheloni GM, Giorgini A et al.

BACKGROUND: Posterosuperior massive irreparable rotator cuff tears (PMIRT) are rare and disabling conditions. When conservative treatment fails, latissimus dorsi tendon transfer (LDTT) is a viable surgical option for symptom relief in carefully selected patients. However, its effectiveness in restoring glenohumeral function and its influence on scapulothoracic rhythm remain subjects of ongoing debate.

PURPOSE: The purpose of this study was to evaluate the clinical outcomes of LDTT and assess its impact on scapulothoracic rhythm using kinematic and electromyographic analysis.

MATERIAL AND METHODS: A total of 18 patients with PMIRT underwent LDTT. Functional scores consisted of Constant-Murley score (CMS), America Shoulder and Elbow Surgeons (ASES) score, and Quick Disabilities of the Arm, Shoulder, and Hand (QuickDASH) score. Electromyography (EMG) activity of the latissimus dorsi muscle was

evaluated in conjunction with three-dimensional (3D) kinematic tracking system. Differences in external rotation strength were compared with and without active adduction.

RESULTS: The mean age was 55.9 years (range 40-69 years), with a mean follow-up of 20.4 months (range 6-39 months). At final follow-up, the mean CMS was 68.2 (95% CI 64.9-71.5), ASES was 76.9 (95% CI 66.1-87.6), and QuickDASH was 17.6 (95% CI 9.5-25.6). A significant difference in external rotation strength was observed [...]

[PubMed](#) · [DOI](#)

Is there a need for exploration in pulseless supracondylar fractures of the humerus? A systematic review and individual patient data meta-analysis.

Haoyang C, Chen JG, Lai H, Lim AKS, Tan SHS, Hui JHP

BACKGROUND: Different approaches have been proposed to treat patients with pulseless supracondylar humeral fractures (SHF). We aim to analyze the current outcomes of patients who have undergone different treatment options for pulseless SHF, namely watchful waiting versus urgent surgical exploration.

METHODS: Electronic databases including PubMed, Embase, and Cochrane Library were explored. Our study included patients from all age groups but only included English-language articles. Single case reports and case studies with adequate description of patient population, injury, and outcomes were included. An individual patient data meta-analysis was done to evaluate key outcomes of surgical exploration versus no surgical exploration in pulseless patients with SHF.

RESULTS: Overall, the data for 1070 individual patients from a total of 48 studies were included. Patients with pulseless SHF with open fractures have a higher probability of requiring vascular intervention ($p < 0.001$) as they are more likely to have disrupted arteries ($p = 0.050$) and more likely to require vascular repairs ($p < 0.001$) than patients with closed fractures. Similarly, patients with pulseless SHF with pucker sign ($p = 0.003$) and ecchymosis ($p = 0.002$) were more likely to undergo surgical exploration. However, the neurological status had no relation to them undergoing surgical exploration ($p = 0.382$), and al [...]

[PubMed](#) · [DOI](#)

Increasing complexity and plateauing outcomes in AO type C distal humerus fractures: a 50-year institutional case series.

Hönck K, Wenzelburger P, Eibinger N, Sadoghi P, Puchwein P

BACKGROUND: Distal humerus fractures in adults are rare but complex injuries, particularly AO type C3 fractures, often involving comminution, instability, and osteoporotic bone. Demographic trends indicate rising incidence, especially among elderly patients, with increasing functional demands challenging modern osteosynthesis. Open reduction and internal fixation (ORIF) with bicolumnar plating remains the preferred reconstructive approach, although complications remain substantial. This study aimed to evaluate trends in fracture complexity, patient demographics, and functional outcomes over five decades, focusing on the most recent cohort (2009-2018, series E) and comparing with historical cohorts (series A-D, 1969-2008).

METHODS: This retrospective analysis included five consecutive 10-year institutional cohorts (series A: $n = 43$; B: $n = 29$; C: $n = 47$; D: $n = 58$) of 233 patients with 235 supracondylar distal humerus fractures (AO type C1-3) treated with ORIF. Functional outcome parameters (Jupiter and Cassebaum Scores) and demographic data were available for all cohorts; additional parameters (range of motion (ROM), QuickDASH Score) were available for series D and E, Mayo Elbow Performance Score (MEPS), Short Form (SF)-36, and strength were assessed specifically in series E ($n = 56$; follow-up $n = 21$). Cross-cohort comparisons of categorical variables used chi-square tests; [...]

[PubMed](#) · [DOI](#)

Magnetic resonance imaging-based fat infiltration grading improves reparability prediction in massive rotator cuff tears.

Hu Q, Zhang G, Wang D, Tung TH, Ding J, Zhou X

PURPOSE: This study aims to investigate the extent of fat infiltration (FI) in the rotator cuff (RC) muscles in more medial slices from the Y-shaped view (Y-view) on shoulder magnetic resonance imaging (MRI) and assess whether these slices provide a more accurate prediction of the reparability of massive RC tears (massive RCTs).

METHODS: The retrospective study included 57 patients with massive RCTs who successfully underwent arthroscopic surgery between 1 January 2023 and 31 December 2023. Patients were categorized into two groups: the irreparable group and the repairable group. All patients underwent shoulder MRI covering the region from the acromion to the medial border of the scapula in oblique coronal and oblique sagittal orientations. The FI stage of the RC was assessed across three different views to determine which view most effectively predicts the reparability of patients with massive RCTs.

RESULTS: The FI stage of the infraspinatus (IS) muscle across three distinct views demonstrated a significant correlation with the reparability of massive RCTs. The FI stage was more severe in the irreparable group. Receiver operating characteristic (ROC) analysis revealed that the area under the curve for the 1/2 scapula view (0.737) and the most severe view (0.745) exceeded that of the traditional Y-view (0.644). Paired-sample ROC curve analysis revealed a significant difference [...]

[PubMed](#) · [DOI](#)

Mid-term clinical and functional outcomes after reverse shoulder arthroplasty with latissimus dorsi transfer.

Colombini AG, Rab P, Macken AA, Soares MN, Kimmeyer M, Shirinskiy IJ, Popescu IA, Lafosse L et al.

BACKGROUND: Although reverse total shoulder arthroplasty (rTSA) with concomitant latissimus dorsi transfer (LDT) has been shown to effectively treat external rotation (ER) deficits, there are limited data regarding its outcomes with modern implants and its impact on activities of daily living (ADLs) requiring ER. The purpose of this study was to assess the mid-term clinical and radiographic outcomes of rTSA with concomitant isolated LDT in patients with an ER lag sign and posterior rotator cuff deficiency.

METHODS: This retrospective cohort study with prospective follow-up included consecutive patients who underwent rTSA with concomitant isolated LDT between 2010 and 2022 with a minimum follow-up of 2 years. Primary outcomes included the resolution of ER lag sign, active ER, and the Activities of Daily Living and External Rotation (ADLER) score. Secondary outcomes included the Auto-Constant Score (CS), Subjective Shoulder Value (SSV), activities of daily living requiring internal rotation (ADLIR) score, visual analog scale (VAS) for pain, and radiographic analysis of standardized radiographs.

RESULTS: In total, 32 procedures in 32 patients were identified. Of these, 22 procedures in 22 patients (68% female, 72.9 ± 8.4 years at surgery) were available for follow-up at 4.8 ± 2.2 years postoperatively (response rate 73%). The ER lag sign resolved in 95.5% of patients, the active [...]

[PubMed](#) · [DOI](#)

Partial versus full revision in cementless single-stage exchange for hip periprosthetic joint infection: a multicenter comparative study with 10-year survivorship analysis.

Mu W, Jarusriwanna A, Xu B, Brown SA, Zhang X, Dawes A, Parvizi J, Cao L

BACKGROUND: Periprosthetic joint infection (PJI) is one of the most challenging complications following total hip arthroplasty (THA). Traditional management typically involves full revision to ensure comprehensive infection eradication. However, for patients with well-fixed implants, partial revision in a single-stage cementless approach may provide a viable alternative, potentially preserving bone stock and reducing operative time. The relative efficacy of this approach compared with full revision remains to be fully explored. This multicenter study aims to determine whether partial revision in cementless single-stage exchange surgery offers comparable infection control outcomes to full revision for select patients with well-fixed implants.

METHODS: We conducted a retrospective, multicenter cohort study involving 226 patients who underwent cementless single-stage exchange hip arthroplasty for PJI between 1 October 2013 and 31 July 2022. Patients were divided into partial revision (n = 61) and full revision (n = 165) groups. The primary outcome was treatment success, defined as the absence of clinical symptoms and signs of infection at a minimum follow-up of 2 years.

RESULTS: The success rates were 77.0% for the partial revision group and 80.0% for the full revision group, with no significant difference (p = 0.629). Both groups showed comparable 10-year survival rates for over [...]

[PubMed](#) · [DOI](#)

AC Joint Reconstruction Using Cortical Suture Button and Semitendinosus Allograft.

Egol AJ, Lowe DT, Egol KA

Acromioclavicular joint injury reconstructive options include a wide number of described techniques. Although no single technique has consistently demonstrated superior clinical outcomes in the literature, there are theoretical biomechanical benefits to a combined reconstructive technique using cortical suture button and semitendinosus allograft. This review describes an acromioclavicular joint reconstruction using a cortical suture button and semitendinosus allograft.

[PubMed](#) · [DOI](#)

Introduction.

Jeray KJ

[PubMed](#) · [DOI](#)

Humerus Shaft-Antegrade Intramedullary Nailing.

Karsmarski O, Stroh DA

This review describes intramedullary nailing of humeral shaft fractures through antegrade technique. The case involved a high-energy injury to the humerus in a patient with relevant comorbidities, and discusses the rationale for both surgery in general and nailing in specific. Indications for surgery, setup considerations unique to the lateral decubitus position, and intraoperative methods for avoiding complications are all discussed. A brief review of current relevant literature is included.

[PubMed](#) · [DOI](#)

Intraarticular Distal Humerus With Anconeus Pedicle and Olecranon Osteotomy.

Cerasani MN, Zhang G, Konda S, Ganta A

This review details an open reduction internal fixation technique of a comminuted, intraarticular Y-type bicolumnar distal humerus fracture using supplemental minifragment plating in addition to dual locking plates. Owing to significant articular comminution, an olecranon osteotomy with an anconeus pedicle flap was used to facilitate exposure to achieve an accurate articular reduction to restore elbow function.

[PubMed](#) · [DOI](#)

Carpal Tunnel Release: Technical Considerations for the Orthopaedic Trauma Surgeon.

Coscia A, Weinerman J, Hake M, Lawton J

Emergent carpal tunnel release is indicated for acute carpal tunnel syndrome, most commonly secondary to distal radius fracture and may be performed in the setting of forearm/hand compartment syndrome along with forearm and/or hand fasciotomies if median nerve compression is suspected. Carpal tunnel release can safely and efficiently be performed concurrent to fracture fixation and/or compartment release, generally with excellent results. This abstract is intended to serve as a succinct review for orthopaedic traumatologists who do not routinely perform carpal tunnel release on an elective basis, describing the relevant anatomy, structures at risk, and an incision design required to successfully perform open carpal tunnel release.

[PubMed](#) · [DOI](#)

Transradial Amputation With Nerve Management: Case Presentation and Surgical Technique.

Moll S, Dewing J, Bouz G, Hacquebord J, Ayalon O

Transradial amputation of appropriate length preserves forearm rotation and facilitates prosthetic use; however, high rates of phantom limb pain and neuropathic medication use persist. This case describes a multidisciplinary approach to transradial amputation that integrates targeted muscle reinnervation, regenerative peripheral nerve interfaces, and bone and soft tissue management to optimize pain control and prosthetic success. A 29-year-old man with a severe ballistic injury and complex regional pain syndrome underwent transradial amputation with targeted muscle reinnervation and regenerative peripheral nerve interface. Limb length was preserved in accordance with prosthetic guidelines, and myodesis was performed to improve residual limb stability and myoelectric signal quality. This case highlights how nerve management and prosthetic-informed surgical planning optimizes functional outcomes and quality of life after traumatic transradial amputation.

[PubMed](#) · [DOI](#)

The Modified Hardinge Approach to Hip Hemiarthroplasty: An Alternative in High-Risk Patients.

Beck A, Winzenried A, Beilfuss M, Dunahoe J

This review describes an alternative hemiarthroplasty technique for high-risk patients using a direct lateral approach through the anterior one-third of the gluteus medius and vastus lateralis.

[PubMed](#) · [DOI](#)

Early Open Repair of a Multiligamentous Knee Injury.

Curran CC, Davis ME

This review discusses the utility of early open repair of multiligamentous knee injuries in patients with high-energy mechanisms. The surgical technique for anterior cruciate ligament, posterior cruciate ligament, and medial collateral ligament repair was demonstrated on a polytraumatized patient using the medial parapatellar approach. The posterior cruciate ligament was repaired first with suture through a drill tunnel in the femoral insertion followed by the anterior cruciate ligament in a similar fashion. The medial collateral ligament was repaired last with a staple. With some exceptions, weight bearing and knee range of motion are permitted immediately postoperatively. In the right patient population, this procedure may expedite patients' recovery and return to daily life.

[PubMed](#) · [DOI](#)

Articular Reduction of a Bicondylar Tibial Plateau Fracture Using a Lateral Tibial Osteotomy.

Arnold JC, Druten E, Soni C, Suryavanshi J, Wynn M, Jang Y, Lopas LA

Tibial plateau fractures with significant articular depression can present challenges in achieving anatomic reduction, especially without an associated cortical "split" to access the depression. This case involved a 23-year-old woman with a valgus-unstable bicondylar tibial plateau fracture and substantial lateral joint depression. This review describes the use of a lateral tibial osteotomy to directly visualize and elevate a depressed articular surface during open reduction and internal fixation, allowing for restoration of alignment and stable fixation.

[PubMed](#) · [DOI](#)

Posterior Tibial Tendon Transfer and Achilles Tendon Lengthening for Foot Drop.

Mercer NP, Lezak BA, Lowe DT, Egol KA

This review describes a case of chronic post-traumatic foot drop managed with posterior tibial tendon transfer through the interosseous membrane and percutaneous Achilles tendon lengthening. Key technical considerations and postoperative rehabilitation principles are highlighted. This approach provides a durable solution for restoring ankle dorsiflexion and improving ambulation in patients with irreversible peroneal nerve injury.

[PubMed](#) · [DOI](#)

Ankle Syndesmotic Injury With Posterior Tibial Tendon Rupture.

Lezak BA, Mercer NP, Lowe DT, Egol KA

Isolated ankle syndesmotic injuries are uncommon and are typically discussed in the context of athletic high ankle sprains; however, concomitant rupture of the posterior tibial tendon represents an exceedingly rare injury pattern.

This case highlights the importance of maintaining a high index of suspicion for associated soft-tissue pathology in patients presenting with syndesmotic instability in the absence of fracture. This case underscores the need for comprehensive clinical and imaging evaluation of isolated syndesmotic injuries and supports primary tendon repair when rupture is identified.

[PubMed](#) · [DOI](#)

Supination Adduction Ankle Fracture Treated With Fibular Intramedullary Screw Fixation.

Di Pauli von Treuheim T, Cerasani MN, Konda S, Ganta A

This abstract describes an open reduction and internal fixation technique used for a supination adduction ankle injury. This technique demonstrates medial plafond articular disimpaction with backfilling of the bony void with cancellous allograft. The medial malleolus was then fixed with a mini fragment plate in antiglide fashion with subchondral rafting screws. The lateral malleolus fracture was addressed with use of a retrograde screw rather than a fibular plate. This technique emphasizes the importance of both recognizing and addressing the anteromedial plafond impaction of supination adduction ankle fractures.

[PubMed](#) · [DOI](#)

Reduction Techniques for Calcaneus Fractures-Using a Femoral Distractor.

Arnold JC, Druten E, Landis T, Lopas LA, Jang Y

Displaced intra-articular calcaneal fractures disrupt hindfoot alignment and are associated with long-term morbidity. Although minimally invasive approaches reduce wound-related complications, they can limit reduction of a displaced calcaneal tuberosity. This report describes a 29-year-old man with a displaced intra-articular calcaneal fracture treated using a laterally applied femoral distractor to facilitate correction of coronal, sagittal, and axial malalignment while limiting soft-tissue disruption.

[PubMed](#) · [DOI](#)

Sinus Tarsi Approach With Distractor-Aided Reduction of Displaced Intra-articular Calcaneus Fracture.

Judson WR, Ricci G

This review describes the sinus tarsi approach for operative fixation of intra-articular calcaneus fractures, using a fixed distractor for assistance. The sinus tarsi approach has been shown to decrease wound complication rates and allow for earlier surgical intervention compared with the extensile lateral approach, while still achieving satisfactory anatomical restoration in the operative management of calcaneus fractures. The case presented involves a 46-year-old man who sustained a displaced intra-articular calcaneus fracture after a fall from height. The discussion includes operative indications, relevant imaging studies, patient positioning, and a detailed description of the sinus tarsi technique, as well as intraoperative observations and fracture fixation procedures.

[PubMed](#) · [DOI](#)

A comparative analysis of three knee arthrodesis techniques to manage failed total knee arthroplasty due to infection: intramedullary nailing, LRS external fixator, and compression plating.

Rajasekaran RB, Ponniah HS, Palanisami DR, Sundaram VP, Palanivelayutham SS, Jayaramaraju D, Dhanasekaran S, Rajasekaran S

INTRODUCTION: This study compared external fixation (EF), intramedullary nailing (IMN), and compression plating (CP) for knee arthrodesis (KA) following failed total knee arthroplasty (TKA) due to periprosthetic joint infection, focusing on complications, radiological, and functional outcomes.

METHODS: Patients aged ≥ 18 who underwent knee arthrodesis following failed TKA due to infection from 2008 to 2023 were included. Four surgeons used three techniques: EF, IMN, and CP. Retrospective data on demographics, surgical techniques, complications, and outcomes were collected. Postoperative patient-reported outcome measures (PROMs), pain, and functional status were assessed. Radiographic data was reviewed for limb shortening and union status.

RESULTS: 46 patients (mean age 61.9 years, BMI 28.2), predominantly male (84.8%) were involved in this study. Twenty patients underwent EF, 16 underwent CP, and 10 underwent IMN, all for infected primary TKA. *Pseudomonas* was the most common infecting organism (15.2%). Average follow-up was 29.1 months. Bony fusion was achieved in 41 knees (89.1%), with five requiring additional surgery; four eventually required amputation due to non-fusion. EF group had the shortest time to union and no major complications or amputations. Time to union was significantly shorter in the CP group compared to the IMN group ($p < 0.001$), and EF group also had a si [...]

[PubMed](#) · [DOI](#)

Accuracy of iliac crest reference pin placement in total hip arthroplasty: a CT-based analysis of risk factors for cortical perforation.

Masada Y, Tetsunaga T, Yamada K, Inoue T, Okuda R, Tetsunaga T, Okazaki Y, Ozaki T

INTRODUCTION: Accurate iliac crest reference pin placement is essential for safe navigation-assisted total hip arthroplasty (THA). Although complications are uncommon, inaccurate placement may cause neurovascular injury. This study aimed to evaluate pin insertion accuracy using postoperative computed tomography (CT), identify risk factors for cortical perforation, and determine whether CT-detected cortical perforation was associated with postoperative acetabular cup positioning accuracy.

MATERIALS AND METHODS: We retrospectively analyzed 200 primary THAs performed using portable or CT-based navigation between December 2022 and July 2025. Pin position was assessed on CT 1 week postoperatively and classified as complete intraosseous insertion or cortical perforation. Risk factors were examined using univariable and multivariable logistic regression analyses. Perforation direction was evaluated using Firth's penalized logistic regression. Postoperative cup positioning accuracy was assessed using CT-based radiographic inclination (RI), radiographic anteversion (RA), and absolute errors from target angles of 40° for RI and 15° for RA.

RESULTS: Cortical perforation occurred in 70 of 200 hips (35.0%): 32 medial and 38 lateral. On a pin-by-pin basis, 113 of 400 pins (28.3%) demonstrated perforation. In multivariable analysis, smaller ASIS width, lateral decubitus positioning, and 4-m [...]

[PubMed](#) · [DOI](#)

Long-term outcomes of arthroscopic single-anchor fixation versus microfracture for osteochondral fractures of the knee weight-bearing area in adolescents and young adults.

Yang W, Pan Q, Wang H, Peng Y, Meng C, Huang W

PURPOSE: Osteochondral fractures involving the weight-bearing area of the knee in young patients require stable restoration of the native osteochondral fragment whenever possible. This study aimed to evaluate the long-term clinical and radiological outcomes of arthroscopic single-anchor fixation and to compare them with those of microfracture.

METHODS: This retrospective comparative study included 24 adolescents and young adults with osteochondral fractures of the femoral condyle or trochlear region treated between 2016 and 2017. Twelve patients underwent arthroscopic single-anchor in situ fixation, and 12 underwent microfracture. The fixation technique used one double-loaded suture anchor to obtain multi-point compression of the osteochondral fragment while preserving the subchondral plate. Clinical outcomes were assessed using the Lysholm, International Knee Documentation Committee, and Tegner scores. Magnetic resonance imaging was evaluated using the MOCART 2.0 score. The mean follow-up was 93.7 months.

RESULTS: The single-anchor group showed a significantly higher intraoperative contact area ratio than the microfracture group ($96.5\% \pm 3.2\%$ vs. $85.8\% \pm 5.6\%$, $P < 0.001$). At final follow-up, the single-anchor group had superior Lysholm scores (91 ± 5 vs. 83 ± 8 , $P = 0.009$), IKDC scores (86 ± 6 vs. 78 ± 7 , $P = 0.010$), and Tegner activity levels (5.0 ± 1.1 vs. 3.9 ± 0.9 , $P = 0$ [...])

[PubMed](#) · [DOI](#)

Isometric points in cerclage for patellar tendon repair: a 3-dimensional, weightbearing computed tomography kinematic simulation.

Zindel C, Hodel S, Luo Y, Meisterhans M, Fucentese SF, Vlachopoulos L

PURPOSE: The gold standard for the management of ruptured patellar tendon is the surgical repair with nonabsorbable sutures and additional reinforcement e.g. cerclage wires as described by McLaughlin in 1947. Despite the early description, the exact placement of such cerclage wires remains unclear. The aim of the study was (1) to analyze how the cerclage elongates during knee flexion and (2) to define the insertion points with the most isometric behavior at the patella and tibial tuberosity (TT).

METHODS: 3D simulations of 10 healthy knees in the range from 0° to 120° of flexion using weightbearing computed tomography scans were generated. 12 insertion points were defined at the patella in equal distances along the lateral and medial border from the proximal to the distal pole. 42 insertion points were defined at the tibial tuberosity (TT) in a 3.0×3.0 cm large grid around its center. An automatic string simulation algorithm was used to calculate the length for all possible combinations of the cerclage from 0° to 120° flexion. The most isometric combination of insertion points was found using the isometric score.

RESULTS: The most isometric insertion at the TT is distally ($p < 0.01$) and towards the center of the TT ($p < 0.01$). The most isometric points at the TT is located 61 mm distal to the joint line and 5 mm from the anterior border of the TT in a true lateral view fluo [...]

[PubMed](#) · [DOI](#)

Patient mortality exceeds implant revision risk with equivalent survival following unicompartmental versus total knee arthroplasty in octogenarians.

Innocenti M, Leggieri F, Massenzi M, Stimolo D, Akkaya M, Matassi F, Civinini R

BACKGROUND: While traditional surgical decision-making emphasizes long-term implant survivorship for octogenarians knee reconstruction, this paradigm may be inappropriate when patient mortality risk substantially exceeds revision risk. This study compared mid-term mortality between unicompartmental knee arthroplasty (UKA) and total knee arthroplasty (TKA) in octogenarians, hypothesizing that implant type would not significantly influence patient survival.

METHODS: A retrospective cohort study was conducted on 238 consecutive octogenarian patients (age ≥ 80 years) who underwent primary knee arthroplasty at a single tertiary orthopaedic center between January 2018 and December 2020. Patients with revision arthroplasty, fracture, tumor, inflammatory arthropathy, or incomplete data were excluded. The primary outcome was all-cause mortality at minimum 5-year follow-up. Secondary outcomes included age-stratified mortality (80-84, 85-89, ≥ 90 years) and identification of independent mortality predictors. Statistical analyses included independent t-tests, chi-square/Fisher's exact tests, standardized mean differences, and multivariable logistic regression adjusting for age, follow-up duration, and registry year.

RESULTS: Of 238 patients, 185 (77.7%) underwent TKA and 53 (22.3%) underwent UKA. Overall mortality was 35.3% (84/238): 33.5% in TKA versus 41.5% in UKA (OR = 1.41, 95% CI: 0 [...])

[PubMed](#) · [DOI](#)

Short- to mid-term clinical outcomes and return to sport and work after arthroscopic debridement of the extensor carpi radialis brevis in patients with chronic lateral epicondylopathy.

Siebenlist S, Doucas A, Hatt FL, Lappen S, Vieider RP, Lacheta L, Kadantsev P

BACKGROUND: This study aimed to evaluate the short- to mid-term clinical outcomes following arthroscopic extensor carpi radialis brevis (ECRB) debridement in patients with chronic epicondylopathy humeri radialis (cEHR). We hypothesized that arthroscopic ECRB debridement would result in favorable short- to mid-term patient-reported outcomes, improving pain levels, strength, range of motion (ROM) and patient satisfaction.

METHODS: A retrospective case series was conducted on patients who underwent arthroscopic ECRB debridement for cEHR with symptoms > 6 months. Patients were assessed at a minimum of 12 months postoperatively using the Mayo Elbow Performance Score (MEPS), Patient-Rated Tennis Elbow Evaluation (PRTEE), Disabilities of the Arm, Shoulder, and Hand (DASH) score, and Visual Analog Scale (VAS) for pain. Pre- and postoperative scores were compared using DASH and VAS. Maximum isometric wrist extension strength and postoperative ROM were measured bilaterally. Return to sport and work, patient satisfaction, and complications were documented.

RESULTS: Of the 35 eligible patients, 30 completed follow-up (86% follow-up rate; mean age 48.4 ± 6.6 years; range: 36-65 years; 47% male). The mean duration of symptoms before surgery was 12 months (IQR 8-20, range 6-62). The mean follow-up duration was 28.13 ± 10.9 months (range 12-45). At final follow-up, median outcome scores were [...]

[PubMed](#) · [DOI](#)

Complications and outcomes after operative treatment of low-energy pelvic ring and acetabular fractures.

Salonpää KB, Nurkkala J, Lantto I, Kakko S, Kaakinen T, Aitavaara-Anttila M, Liisanantti J

BACKGROUND: Fragility fracture of the pelvis (FFP) with a low-energy mechanism of injury is relatively common among the growing geriatric population. The incidence of postoperative complications after surgically treated FFP is not well documented. The aim of this study was to report complications after operatively treated FFP and to determine whether these complications would have a detrimental effect on outcomes.

METHODS: Retrospective study of 200 consecutive patients who underwent operative treatment at Oulu University Hospital, Oulu, Finland, between 1.1.2010 and 15.4.2021 for FFP with a low-energy injury mechanism. Postoperative complications were recorded, and associations with outcomes were depicted.

RESULTS: Postoperative complications were recorded in 31 (15.5%) patients. Patients with complications had higher intraoperative bleeding (OR 2.8 (1.0-7.9), $p = 0.048$) and higher ASA classification 3.4 (1.3-8.4), $P = 0.009$). Patients with complications had longer hospital stays, but no clear association between complications and mortality or discharge location was observed.

CONCLUSIONS: The incidence of postoperative complications was comparable with that reported in previous literature. Patients who experienced complications had longer hospital stays; however, no definitive association between complications and outcomes was identified.

[PubMed](#) · [DOI](#)

Minced cartilage implantation for cartilage regeneration: a survey of current clinical practices.

Yilmaz T, Leutheuser S, Niemeyer P, Behrendt P, Roessler P, Gille J, Mumme M, Vogt S

INTRODUCTION: The objective of this study is to evaluate how minced cartilage implantation (MCI) is currently applied in clinical settings, focusing on the selection criteria, procedural execution, handling of concomitant joint pathologies, and postoperative management.

MATERIAL AND METHOD: An online survey was conducted among all 4,915 members of the German Society for Arthroscopy and Joint Surgery (AGA). Invitations with a survey link were emailed, and the questionnaire was accessible anonymously from January 10 to February 20, 2024. A total of 927 responses were received and analyzed, resulting in a 19% response rate. Demographic data were described descriptively; survey answers were reported as percentages. The 25-item questionnaire, developed by AGA and the Quality Circle for Cartilage Repair & Joint Preservation (QKG e.V.), included single- and multiple-choice questions on experience with cartilage therapy, minced cartilage procedure application and postoperative care, and the overall assessment of this technique within cartilage repair.

RESULTS: MCI has emerged as one of the three most commonly used cartilage repair techniques in the knee joint. Defect sizes up to 4 cm² represent the largest treated group. The application of the procedure is quite heterogeneous, with some practitioners using a shaver and others manually mincing with a scalpel. Fixation methods also var [...]

[PubMed](#) · [DOI](#)

Variability in United States online rehabilitation protocols after open and endoscopic hip abductor repair.

Anwunah C, Rosenberg S, Hartwell M

PURPOSE: The purpose of this study was to evaluate readily accessible U.S. based online physical therapy protocols for open and endoscopic hip abductor repairs and to assess the variability in post-operative rehabilitation protocols.

METHODS: Online physical therapy protocols from U.S. based orthopaedic institutions for gluteus medius/minimus repair were collected using a web-based query utilizing search terms "hip abductor repair rehabilitation protocol" and "gluteus repair rehabilitation protocol." Published protocols from both Electronic Residency Application Service (ERAS) academic orthopaedic surgery programs and private groups were collected. All publicly available protocols were included. Protocols were analyzed using a standardized spreadsheet to assess rehabilitation components and timelines. Descriptive statistics (median values, percentages, ranges) were calculated.

RESULTS: 54 rehabilitation protocols were assessed, including 22 (41%) from ERAS-affiliated academic programs or associated physicians. Postoperative weight-bearing was addressed in 53/54 (98%) protocols, with 41 (77%) recommending toe-touch/touchdown/flat-foot/20-lb weightbearing. Median duration of restricted weight-bearing was 6 weeks (range, 2-8 weeks). Regarding range of motion restrictions, 37 (69%) protocols limited hip flexion to 90°, 42 (78%) advised against passive adduction past neutral, and [...]

[PubMed](#) · [DOI](#)

The incidence and epidemiology of patellar fractures following medial patellofemoral ligament reconstruction: a systematic review and meta-analysis.

Abelleyra Lastoria DA, Hunt A, Ross CK, Yemane M, Keynes S, Ghosh A, Hing CB

INTRODUCTION: Patellar instability can arise from a traumatic event with anatomical predisposing factors increasing the risk of dislocation. Medial patellofemoral ligament reconstruction (MPFLR) is commonly performed to treat patellar instability. The aim of this systematic review was to determine the incidence of and risk factors for patellar fractures after MPFLR.

MATERIALS AND METHODS: The protocol for this review was registered on PROSPERO with registration number CRD420251066107. The PRISMA 2020 checklist was followed. Electronic databases and conference proceedings were searched to the 15th of May 2025. Studies were eligible if they reported incidence of patellar fractures after MPFLR. A random-effects meta-analysis was performed. Included studies were appraised using tools respective of study design.

RESULTS: A total of 4,972 records were screened in the initial search. A total of 110 studies satisfied the inclusion criteria, evaluating 115,496 patients. The evidence was moderate to low in quality. The incidence of patellar fracture ranged from 0 to 9.50%, with a pooled incidence of 1.17% (95% confidence interval [CI] 0.76- 1.68%). There was no difference in incidence of patellar fractures between patients who did and did not undergo bone tunnel fixation (0.94%, 95% CI 0.69-1.24 vs. 1.00%, 95% CI 0.15-2.44%, respectively). There was no difference in fracture incidence [...]

Predictive accuracy of a general-purpose artificial intelligence model for cut-out following proximal femoral nailing.

Ben Arie G, Warschawski Y, Amzallag N, Gan-Or H, Ben-Tov T, Khoury A, Horev I, Graif N et al.

INTRODUCTION: Cut-out is the most consequential mechanical complication after proximal femoral nailing and requires prompt recognition. General-purpose artificial intelligence (AI) models with image-interpretation capability are increasingly accessible, yet their diagnostic performance for this task is unknown. We evaluated ChatGPT's accuracy for predicting cut-out following proximal femoral nailing and compared TFNA and Gamma nail subgroups.

MATERIALS AND METHODS: In this retrospective predictive-accuracy study, 989 patients (683 TFNA, 306 Gamma nail; mean age 83.8 years; cut-out prevalence 2.5%) were analysed. For each case, three baseline radiographs - the injury anteroposterior and lateral views and the immediate postoperative control radiograph, all obtained before any complication could be radiographically present - together with the full clinical data set (excluding complication-related information) were submitted to ChatGPT, which returned a binary prediction (yes/no) and probability estimate (0-100%). The follow-up radiographs on which cut-out becomes apparent were not provided to the model. Ground truth was established on subsequent follow-up radiographs by a departmental review board requiring agreement of two senior surgeons. Accuracy metrics were calculated with Wilson 95% confidence intervals (CI); exact McNemar's and Mann-Whitney U tests assessed directional bias [...]

PubMed · DOI

Staged management of a compound combined pelvi-acetabular injury with perineal degloving in a young female: a case report and literature review.

Dhankhar K, Kekatpure AL, Kekatpure AL

We present the case of a 30-year-old woman, the patient who sustained a compound combined pelvi-acetabular fracture with extensive perineal internal degloving after high-energy trauma. Computed tomography revealed an unstable pelvic ring injury (Tile type C1; anteroposterior compression type III) with right transverse acetabular fracture. The occurrence of extensive internal degloving was confirmed intraoperatively. The injury was treated using a graduated, principle-based approach incorporating early soft-tissue debridement, dead-space management and temporary pelvic stabilization, infection control, and delayed definitive internal fixation. An intercurrent pin-site infection was treated successfully without progression to deep infection. The patient demonstrated progressive radiological union at 4 months with a Harris Hip Score (HHS) of 80 and a Majeed pelvic score of 55. At one-year follow-up, complete union occurred and functional recovery was achieved, characterized by HHS of 90 and a Majeed score of 70. In this case, infection-guided staging as well as extremely aggressive soft tissue treatment may allow for safe delayed fixation even where pelvi-acetabular trauma has been compromised.

PubMed · DOI

Concomitant upper extremity fractures do not worsen outcomes in conservatively managed pelvic fractures: a cohort study of 1840 geriatric patients.

Graif N, Elbaz E, Kazum E, Rachevsky G, Gurel R, Factor S

INTRODUCTION: Pelvic fractures in elderly patients are associated with significant pain, functional decline, and prolonged rehabilitation, with the majority managed non-operatively. In the conservative management paradigm, intact upper extremity function is critical for mobilization with assistive devices; concurrent upper extremity (UE) fractures may therefore complicate recovery. The aim of this study was to evaluate the clinical outcomes associated with this injury combination in patients with conservatively managed pelvic fractures.

METHODS: A retrospective cohort study was conducted at a Level 1 trauma center, including all patients aged ≥ 65 years admitted with pelvic ring or acetabular fractures managed non-operatively between January 2010 and January 2024. Patients were stratified by the presence of a concomitant UE fracture sustained during the same traumatic event. Primary outcomes were hospital length of stay (LOS) and all-cause mortality at multiple time points. Secondary outcomes included readmission rates and systemic infection rates.

RESULTS: Of 1840 patients, 1693 (92.0%) sustained isolated pelvic fractures and 147 (8.0%) had concomitant UE fractures. Groups were comparable at baseline across all measured covariates. LOS did not differ significantly between groups (11.0 ± 14.2 vs. 11.8 ± 16.9 days; $p = 0.796$). Mortality was similar at all evaluated time points [...]

Single double-loaded anchor repair of isolated Lafosse III subscapularis tears in patients not involved in physically demanding activities provides satisfactory functional and radiologic outcomes at minimum 5-year follow-up.

Aral F, Oklaz EB, Ahmadov A, Asiret E, Delice C, Guler Oklaz EB, Akdulum I, Kanatli U

PURPOSE: To evaluate the outcomes of single-anchor repair for isolated Lafosse III subscapularis tears in patients not engaged in physically demanding activities.

METHODS: Patients with isolated subscapularis tears who underwent arthroscopic surgery between 2014 and 2020 were retrospectively evaluated. The inclusion criteria were patients who underwent single double-loaded anchor repair for Lafosse III tears and were not involved in physically demanding activities. Assessments included the American Shoulder and Elbow Surgeons (ASES) score, Constant-Murley Score (CMS), Visual Analog Scale (VAS), range of motion, specific clinical tests, and magnetic resonance imaging encompassing tendon integrity, cross-sectional area (CSA), vertical diameter, and transverse diameters.

RESULTS: 27 patients (mean age 52.4 ± 10.3 years; mean follow-up 80.8 ± 18.6 months) were included in the study. The final follow-up demonstrated significant improvements in the ASES (44.0 ± 12.4 to 82.4 ± 14.1), CMS (49.1 ± 15.3 to 85.3 ± 11.1), VAS (5.7 ± 2.1 to 1.3 ± 2.2), and internal rotation (L4 to T8) ($p < 0.001$ for all). The positivity of clinical tests decreased substantially: Bear-Hug, from 93% to 18%; Belly-Press, from 89% to 15%; and Lift-Off, from 89% to 18% ($p < 0.001$ for all). Radiological analyses indicated maintained tendon integrity in 93% of patients and increases in CSA (1.292 ± 453 to 1.544 [...])

PubMed · DOI

Shoulder injuries in rugby union: a systematic review and meta-analysis.

Doyle J, Bellamy M, Fowler K, Keiller H, Gunasekera R, Funk L

Shoulder injuries are common in Rugby Union, resulting in prolonged absence and considerable burden. Reported incidence, severity, and mechanisms vary widely, reflecting the multiple high-impact actions in the sport. Various management approaches exist to treat a wide range of pathologies. This systematic review and meta-analysis aimed to synthesise evidence on the epidemiology, mechanisms, and treatment of shoulder injuries in rugby. PUBMED, SCOPUS, and WEB OF SCIENCE were searched from database inception until 1st June 2025. From 1,099 abstracts screened, 37 studies were included. The pooled match incidence was 11.02 injuries/1,000 player-hours (95% CI 7.23-16.82), though heterogeneity was very high ($I^2 = 97\%$) and the 95% prediction interval was wide (2.11-57.61 injuries/1,000 player-hours), indicating the pooled rate should be read as an average across diverse populations rather than a single representative value. Higher rates were seen in male elite (13.74/1,000 h) and high school/university players (12.28/1,000 h). Training incidence was considerably lower (0.20/1,000 h). Substantial between-study heterogeneity was observed across all incidence analyses, reflecting genuine variation across playing levels, sex, and surveillance methods. Tackling was the predominant injury mechanism, most often affecting the tackler. Bankart, Latarjet, and Bristow procedures all produced fav [...]

PubMed · DOI

The pre-operative physical component subscale of the 12-item Short Form Health Survey was the only factor associated with 2-year clinical outcome and satisfaction after hip abductor tendon repair.

Ebert J, Edwards P, Klinken S, Janes G

INTRODUCTION: Surgical repair in the context of symptomatic hip abductor tears unresponsive to non-operative management has demonstrated encouraging outcomes. This study sought to investigate demographic, injury/management and post-operative factors associated with clinical outcome after hip abductor tendon repair.

METHODS: A total of 104 patients undergoing repair were included. Patients were assessed pre-surgery and 2-years post-operatively with the modified Harris Hip Score (mHHS), Oxford Hip Score (OHS), physical (PCS) and mental (MCS) component subscales of the 12-item Short Form Health Survey (SF-12), and satisfaction. Regression analysis assessed the contribution of pre-operative demographic (age, sex, body mass index, PCS, MCS), injury/treatment (symptom duration, prior corticosteroid injections, gluteus minimus/medius tear grade and

fatty infiltration) and post-operative (hip abduction strength symmetry) variables, to the 2-year mHHS and being 'very satisfied'.

RESULTS: Mean improvement in the mHHS and OHS from baseline to 2 years was 36.0 points (95% CI, 31.9 to 40.2; $p < 0.001$) and 17.5 points (95% CI, 15.8 to 19.3; $p < 0.001$), respectively. Seven (6.7%) patients presented with symptomatic re-tearing. Of the 97 patients with 2-year follow-up, 78 (80.4%) were 'very satisfied'. The pre-operative SF-12 PCS was the only factor associated with the 2-year mHHS ($B = 0.46$; [...])

[PubMed](#) · [DOI](#)

Temporising external fixation reduces loss of reduction compared with plaster splinting in ankle fracture-dislocations: a systematic review and meta-analysis of cohort studies.

Suravaram V, Solomon L, Bhat S, Wittauer M

PURPOSE: The optimal temporising strategy for ankle fracture-dislocations is a highly debated topic. These unstable injuries are commonly managed with staged fixation to allow for soft tissue recovery, using either plaster splinting or ankle-spanning external fixation. This study aimed to compare the efficacy and safety of these two approaches.

METHODS: A systematic search of MEDLINE, Embase, Scopus, Cochrane CENTRAL, and Google Scholar was conducted. Comparative studies evaluating temporary external fixation versus plaster splinting in skeletally mature patients with ankle fracture-dislocations were included. Meta-analyses were performed to assess differences in loss of reduction prior to definitive fixation, soft tissue complications, and time to definitive fixation. Risk of bias was assessed using ROBINS-I tool, and certainty of evidence was appraised using GRADE.

RESULTS: Nine retrospective cohort studies comprising 1638 patients were included (589 external fixation, 1049 splinting). Loss of reduction occurred in 3% of injuries in the external fixation cohort compared with 14.6% in those managed with plaster splinting (RR 0.22; 95% CI 0.09-0.52). Rates of overall soft-tissue complications were similar between cohorts. The data regarding time to definitive fixation were highly heterogeneous across studies. Overall certainty of evidence was moderate for LOR and very low for [...]

[PubMed](#) · [DOI](#)

Comprehensive ultrasound assessment of thumb metacarpophalangeal collateral ligament injuries reveals frequent concomitant abnormalities.

Bain O, Farkash U, Atzmon R, Shemesh S, Ohana N

INTRODUCTION: Ulnar collateral ligament (UCL) tears of the thumb are well-recognized injuries, while radial collateral ligament (RCL) and dorsal extensor-related abnormalities are less frequently studied. The pattern of concomitant ultrasound abnormalities in patients after thumb trauma is not well documented. The aim of this study was to characterize concomitant UCL, RCL, and dorsal extensor-related abnormalities within a selected cohort referred to thumb ultrasound after trauma, and to explore associations with injury mechanism.

MATERIALS AND METHODS: A retrospective review was performed of 42 consecutive adult patients who underwent high-resolution ultrasound of the thumb metacarpophalangeal joint between April 2014 and February 2025 for suspected ligamentous injury after acute trauma. Patients were included if ultrasound demonstrated pathologic findings in at least one stabilizing structure, UCL or RCL, and the extensor mechanism was systematically assessed. Collateral ligament injuries were graded using standard sonographic criteria, while dorsal extensor-related findings were recorded descriptively. Associations between injury severity and mechanism of injury were tested using chi-square or Fisher's exact analysis, as appropriate.

RESULTS: UCL pathology was present in 35/42 (83.3%) patients, RCL pathology in 23/42 (54.8%), and dorsal extensor-related abnormalities in 17 [...]

[PubMed](#) · [DOI](#)

Holding the line: predictors of coronal and sagittal loss of reduction after tibial plateau fracture fixation.

Hesmerg MK, Benner JL, Joosse P, Keijser LCM

INTRODUCTION: Tibial plateau fractures (TPFs) are challenging fractures that often suffer from secondary Loss of Reduction (LoR) after fracture fixation. In this study, we aimed to investigate the factors and characteristics associated with LoR after surgical fixation, using measurements of both coronal and sagittal displacement.

METHODS: A retrospective analysis of 213 TPF patients (79.3% female, mean age 57.5 ± 14.4 years) surgically treated between 2017 and 2024 in a level 1 trauma center was performed. Patients were included if they had a minimum of 100-days radiological follow-up. Loss of reduction was defined as late postoperative depression ≥ 3 mm, condylar widening ≥ 5 mm, varus/valgus or tibial slope collapse > 5 degrees, breakage, loosening, or conversion to total knee arthroplasty (TKA). Radiographic images were assessed for preoperative, direct postoperative, and late postoperative (up to 1 year) displacement. Univariable and multivariable logistic regression analyses were employed to study factors associated with LoR..

RESULTS: At a mean follow-up of 325 days (SD ± 161) 77 patients (36.2%) suffered from LoR. The adjusted multivariable logistic regression model revealed that patient age (OR 1.03, $p = 0.028$), involvement of the posterior column (OR 2.87, $p = 0.014$), preoperative step-off (OR 1.08, $p < 0.001$), postoperative step-off (OR 1.61, $p = 0.007$), and postope [...]

[PubMed](#) · [DOI](#)

Using an App based algorithm to determine axial rotation and pelvic tilt at the time of an AP radiograph: a sawbone study.

Boettner F, Premkumar A, Sterneder M, Aboaf C, Haralambiev L

INTRODUCTION: Axial rotation of the pelvis impacts inclination and anteversion measurements of the acetabular component on plain radiographs. Currently the axial rotation is not taken into account when providing acetabular component measurements on plain radiographs. The current study evaluates a novel App based algorithm to determine pelvic axial rotation at the time of an AP pelvis radiograph.

MATERIALS AND METHODS: AP and lateral radiographs were taken of the sawbone pelvis to calibrate the App. The sawbone was then rotated 11 times to the left (average 5.3° , range 0.2° - 10.5°) and 10 times to the right (average 5.2° , range 1.2° - 11.3°). The sawbone was then tilted forward 14 times (average 5.6° range 0.1° - 11.6°) and 13 times backward (average 5.9° , range 0.1° - 11.2°). The amount of axial rotation and tilt of the sawbone pelvis was determined using a digital caliper. AP pelvis radiographs were taken for each position. Using a novel App based calculation algorithm the amount of axial rotation and pelvic tilt was determined from the AP Pelvis radiograph.

RESULTS: The algorithm determined the axial rotation of the pelvis at the time of the AP pelvis radiograph with an average difference of 0.6° (range 0° - 1.8°) compared to the digital caliper measurement. For measuring pelvic tilt, the difference was 2.3° (range 0.4° - 4.5°) compared to the digital caliper measurement.

CONCLUSIONS [...]

[PubMed](#) · [DOI](#)

Risk factors for implant-associated infection after bone allograft reconstruction: a 20-year retrospective cohort study.

Scheurer F, Smolle M, Hauer G, Ruckenstein P, Valentin T, Leithner A

AIMS: Reconstruction of bone defects is a key challenge in orthopaedic surgery. While autologous bone grafts are considered the gold standard, osseous allografts are increasingly used as an alternative as they allow for treatment of large bone defects. However, they may be associated with an increased risk of postoperative infections. This study aims to analyze the frequency of and risk factors for implant-associated infections after bone allograft transplantation.

METHODS: All patients who underwent reconstruction with bone allografts at a single orthopaedic and trauma unit between 2003 and 2023 were retrospectively included. Data on demographics, disease-associated parameters, risk factors at index surgery, and development of implant-associated infections during follow-up were documented. Potential risk factors for development of implant-associated infections during follow-up were investigated with logistic regression models.

RESULTS: A total of 220 patients were included (46.8% males; median age at surgery 46.5 years [IQR: 22-67]; $n = 23$ with implant-associated infections during follow-up). Factors independently associated with subsequent

implant-associated infections were smoking ($p = 0.001$), BMI ($p = 0.015$) and radiotherapy ($p = 0.013$). Gender, age, preoperative chemotherapy, allograft material, preoperative C-reactive protein, hemoglobin and reason for surgery (tumor, n [...])

[PubMed](#) · [DOI](#)

Why it's better to believe in a larger definition of the diametaphyseal junction zone in pediatric distal radius fractures.

von Schrottenberg C, Beck SM, Schwert P, Fitze G, Schultz J

INTRODUCTION: Diametaphyseal radius fractures (DMRF) in children are characterized by special biomechanical behavior. They present with a stereotypical loss of reduction (radial translation and ulnar tilt) of the distal fragment when stabilized with elastic stable intramedullary nailing (ESIN), the standard osteosynthesis for pediatric diaphyseal forearm fractures. No consensus has been found on the definition of the diametaphyseal junction zone (DMJZ). Some authors claim that diaphyseal fractures immediately proximal to the metaphysis, as defined by the AO Pediatric Comprehensive Classification of Long Bone Fractures (AO-PCCF), can safely be stabilized with ESIN. They confine the DMJZ to a small part within the metaphysis. Other definitions of the DMJZ extend slightly more proximally, including a small part of the distal diaphysis. The forearm fracture index (FFI) calculates the ratio of the fracture's distance to the radius' growth plate over its width and defines DMRF to have an FFI between 1 and 2. The aim of this case series was to demonstrate that these particular fractures, which can be considered diaphyseal according to the AO-PCCF, still exhibit biomechanical characteristics typical of DMRF.

MATERIALS AND METHODS: Radiologic and clinical outcomes of 11 DMRF, that would be considered diaphyseal by the AO-PCCF classification but fall into the DMJZ when using the FFI, are [...]

[PubMed](#) · [DOI](#)

Optimizing initial biomechanical strength ("Time Zero") of posterior repair in total hip arthroplasty: a biomechanical comparison of suture materials and knot configurations.

Chen M, Liu X, Chen D, Li X, Du H, Sun Y, Wang H

INTRODUCTION: Posterior capsule repair (PCR) is commonly performed during total hip arthroplasty (THA) via the posterolateral approach to improve postoperative stability. Because biological healing of the repaired posterior structures requires time, the repair construct must provide sufficient immediate mechanical support after surgery. This study aimed to identify the suture material and knot configuration with favorable initial mechanical performance in a standardized in vitro model.

MATERIALS AND METHODS: This in vitro biomechanical study was conducted in two sequential phases. Phase 1 evaluated the baseline tensile properties of three commonly used suture materials: #2 Ethibond, #0 Vicryl, and 3 - 0 PGLA. Phase 2 used a standardized porcine dermal surrogate model to compare eight suture configurations, including single- and double-strand techniques. The primary biomechanical outcomes included maximum failure load, load at 2 mm gap formation, and construct stiffness during uniaxial quasi-static load-to-failure testing.

RESULTS: In Phase 1, non-absorbable #2 Ethibond showed significantly greater ultimate tensile strength (92.44 ± 8.72 N) and higher stiffness than the absorbable alternatives ($P < 0.001$). In Phase 2, double-strand configurations outperformed all single-strand techniques. Among them, the double-strand Nice knot achieved the highest maximum failure load (104.04 [...])

[PubMed](#) · [DOI](#)

Intravenous versus oral tranexamic acid in shoulder arthroplasty.

Parmar T, Adio A, Daher M, Khan A, Horneff J, Abboud J

PURPOSE: Tranexamic acid (TXA) is used in shoulder arthroplasty to reduce blood loss. While intravenous (IV) TXA is standard, oral TXA offers advantages in cost and administration. This study compares 90-day complications and postoperative laboratory values between IV and oral TXA in total shoulder arthroplasty (TSA).

METHODS: A retrospective cohort study using TriNetX identified adults undergoing primary TSA (2005-2025) receiving TXA on the day of surgery, stratified by route (IV vs. oral). One-to-one propensity score matching was performed. Ninety-day outcomes included transfusion, thromboembolic events, cardiac and renal complications, neurologic events, infection, wound dehiscence, readmissions, emergency department visits, opioid utilization, and

mortality. Postoperative laboratory outcomes included hemoglobin, erythrocyte count, hematocrit, and platelet count. Outcomes were compared using odds ratios (ORs) with 95% confidence intervals (CIs) and t tests where appropriate.

RESULTS: After matching, 5722 patients were included in each cohort. No significant differences were observed between IV and oral TXA in 90-day transfusion rates ($P = 0.873$) or thromboembolic events, including deep vein thrombosis and pulmonary embolism. Rates of myocardial infarction, stroke, cardiac ischemia, acute renal failure, infection, readmission, emergency department visits, opioid use, and mo [...]

[PubMed](#) · [DOI](#)

Robotic-arm assistance in revision total hip arthroplasty improves the accuracy of acetabular component positioning: a cadaver-based study.

Fitzgerald K, Borukhov I, Chandra V, Smitherberg C, Mont MA, LiArno S

INTRODUCTION: Acetabular component positioning is a key factor for stability, impingement, wear, and biomechanics in revision total hip arthroplasty (THA). However, anatomic distortion and previous implants may complicate positioning. Although evidence on computed-tomography (CT)-based robotic assistance in revision THA is limited, its planning features and precision may improve surgical plan execution. This study aimed to measure the precision and accuracy of acetabular component positioning of revision robotic-arm assisted total hip arthroplasty (RA-THA) compared with a published manual primary total hip arthroplasty (M-THA) benchmark for: (a) acetabular inclination; (b) acetabular version; and (c) center of rotation (COR), with comparison by surgeon case volume.

MATERIALS AND METHODS: There were seven cadavers (nine hips) that previously underwent primary THA revised with RA-THA by independent surgeons with varying primary RA-THA case volumes, surgical approaches, and acetabular component types. Due to surgical protocol deviations, two hips were excluded leaving seven hips from six cadavers. Pre- and postoperative CT scans were compared to calculate absolute error for inclination, version, and COR; normality was established with the Anderson-Darling test and two-variance tests assessed differences in standard deviations ($\alpha = 0.05$). Mean differences between cadaver-based rev [...]

[PubMed](#) · [DOI](#)

Correction: Modified rotational wedge distal metatarsal osteotomy versus chevron osteotomy for hallux valgus: long-term radiographic and clinical outcomes.

Aybar A, Gokkus K, Özden E, Çetin MH, Çetin MÜ

[PubMed](#) · [DOI](#)

Isolated compartment arthrosis of the knee: contemporary concepts, persistent controversies, and future directions in unicompartmental knee arthroplasty.

Akkaya M, Innocenti M

[PubMed](#) · [DOI](#)

Preoperative lower radiographic osteoarthritis severity in primary total knee arthroplasty increases revision risk by tenfold compared to severe osteoarthritis.

Stroobant L, Vermue H, Droogsmas T, Van Onsem S, Arnout N, van Hellemond G, Victor J, Heesterbeek PJC

BACKGROUND: Dissatisfaction following primary total knee arthroplasty (pTKA) is higher in patients with preoperative low-grade osteoarthritis (OA), but its impact on revision risk remains unclear.

QUESTIONS/PURPOSES: This study aimed to identify predictors for revision surgery within 10 years following pTKA, focusing on radiographic severity of OA using the Kellgren-Lawrence (KL) grading system, along with other demographic and radiographic factors.

PATIENTS AND METHODS: This retrospective case-control study was conducted at two European tertiary referral centers. The case group included 142 patients who underwent aseptic revision total knee arthroplasty (rTKA) between 2007 and 2023, within 10 years following pTKA. A 2:1 control group of 284 patients who had pTKA between 2011 and 2014, with no revision surgery within 10 years, was selected. Collected data included age, sex,

body mass index (BMI), side of surgery, and American Society of Anesthesiologists (ASA) classification. Radiographic data were collected, requiring a preoperative radiograph to assess the KL grade. Univariate analyses identified potential predictors for rTKA, and multivariable logistic regression assessed their relationship with revision risk.

RESULTS: Patients requiring rTKA were significantly younger at the time of pTKA. For each additional year of age, the revision risk decreased by 6% (OR 0.94, 95% CI [...])

[PubMed](#) · [DOI](#)

Correction: Non-tobacco nicotine dependence and postoperative complications following operative fixation of tibial shaft fractures: a retrospective cohort analysis.

Berk A, Bogart M, Eghtedari C, Good L, Florentino S, Moyal A, Burkhart R, Ochenjele G et al.

[PubMed](#) · [DOI](#)

Comparison of pediatric proximal femoral locking plates and dynamic hip screws in the fixation of pediatric intertrochanteric fractures, a biomechanical study.

Wang JH, Shih CA, Wang CH, Lee CC, Wu KW, Wang TM, Kuo KN, Hong CK

Treatment options for pediatric intertrochanteric fractures range from conservative casting in young children to surgical fixation in older patients, most commonly using dynamic hip screws (DHS) or pediatric proximal femoral locking plates (PF-LCP). As there is limited biomechanical evidence favoring one method over another, this study aimed to compare the biomechanical stability of DHS and PF-LCP for the treatment of pediatric intertrochanteric fractures using a synthetic bone model. Sixteen synthetic composite femurs were allocated into two groups stabilized with either a DHS or a PF-LCP. To simulate Delbet type IV intertrochanteric fractures, a vertical osteotomy was created in each specimen. Biomechanical evaluation was carried out using a universal testing machine, assessing axial stiffness, vertical, horizontal, and total displacement, as well as ultimate load to failure under standardized loading conditions. Comparative statistical analyses were then performed to determine differences in biomechanical performance between the two fixation methods. Group P demonstrated significantly smaller vertical and total displacements (0.15 ± 0.15 mm and 0.41 ± 0.31 mm) compared with the DHS group (0.59 ± 0.42 mm and 0.75 ± 0.33 mm) ($p = 0.018$ and $p = 0.045$), whereas horizontal displacement showed no difference ($p = 0.142$). Axial stiffness and ultimate failure load were comparable bet [...]

[PubMed](#) · [DOI](#)

Impact of severe varus deformity on the outcome of mobile-bearing unicompartmental knee arthroplasty: a comparison of patients with varus deformity of less than and more than 15 degrees using matched-pair analysis.

Mullaji AB, Gupta AK, George GK, Kazi R

INTRODUCTION: The outcome of patients undergoing UKA with larger preoperative deformity ($> 15^\circ$ varus) and lesser deformity is sparsely compared in the literature. This matched-pair retrospective study attempts to fill this research gap by comparing the effect of preoperative and postoperative residual varus on outcome.

METHODS: Patients undergoing UKA with preoperative varus deformity $> 15^\circ$ (HKA $> 15^\circ$ group) were selected and matched for age and gender with those with $< 15^\circ$ of preoperative varus deformity (HKA $< 15^\circ$ group). There were 72 age and gender-matched pairs knees. HKA was measured and Forgotten Joint Score, OKS, VAS and WOMAC scores were obtained at a minimum follow up of 5 years. Postoperative residual varus was classified into 3 groups based on postoperative HKA: varus $< 3^\circ$, varus 3 to 7° and varus $> 7^\circ$.

RESULTS: Mean correction of deformity was significantly more ($p < 0.001$) in the HKA $> 15^\circ$ group (10.5°) when compared to HKA $< 15^\circ$ group (4.5°). There was no significant difference in outcome measures OKS ($p = 0.21$), FJS ($p = 0.35$) and VAS ($p = 0.6$) between patients in the two groups. There was no significant correlation between the preoperative and postoperative HKA of the patient with postoperative OKS, FJS and VAS in the two groups. There was no statistically significant difference between FJS ($p = 0.73$) of the three postoperative varus groups. Excellent to good [...]

[PubMed](#) · [DOI](#)

A multimodal imaging approach to predict carpal tunnel syndrome after distal radius fractures.

Önüt M, Alkol N, Önaloğlu Y, Özer M, Apaydın E, Talmaç MA

OBJECTIVE: Distal radius fractures are commonly associated with the development of carpal tunnel syndrome (CTS), yet current evidence primarily focuses on radiographic alignment parameters. The role of carpal tunnel morphology, particularly total carpal tunnel cross-sectional area (CSA) measured on computed tomography (CT), remains insufficiently investigated. This study aimed to evaluate the relationship between CSA, volar tilt, and the development of CTS in conservatively managed distal radius fractures, and to assess the predictive performance of these parameters when used in combination.

METHODS: This retrospective case-control study included 150 patients, comprising 50 patients who developed CTS and 100 controls without CTS. CSA was measured on CT scans obtained at initial presentation, while volar tilt was assessed using standard radiographs. Additional variables, including the presence of ulna fracture and fracture type, were analyzed. Group comparisons were performed using nonparametric tests. Multivariable logistic regression analysis was conducted to identify independent predictors of CTS. Diagnostic performance was evaluated using receiver operating characteristic (ROC) curve analysis.

RESULTS: Patients who developed CTS had significantly lower CSA and higher volar tilt values (both $p < 0.001$). The presence of an ulna fracture was inversely associated with CTS ($p = [\dots]$)

[PubMed](#) · [DOI](#)

Unicompartmental knee arthroplasty and total knee arthroplasty in patients with 15 or more degrees of varus deformity: age- and gender-matched comparative study of radiographic and clinical outcomes.

Mullaji A, Chopperla S, George GK, Kazi R

INTRODUCTION: To date, few studies have directly compared matched cohorts of patients with severe varus deformity undergoing UKA versus TKA. The current study seeks to address this gap. The hypothesis was that, in patients with deformity of $> 15^\circ$, postoperative PROMs following UKA would be inferior to those achieved with TKA.

METHODS: Records of patients operated for UKA and TKA from January 2016 to December 2021 were evaluated, and the data for patients with deformity of $\geq 15^\circ$ were selected. The UKA patients were then matched for age and gender with TKA patients. 52 matched pairs of patients were obtained. The HKA, mLDFA, MPTA were measured. OKS, SF12, and FJS were collected. The magnitude of change in the mLDFA and MPTA was assessed using the Estimated Marginal Means.

RESULTS: The mean follow-up times for UKA and TKA were 3.7 years and 5.5 years, respectively. The mean deformity correction (Δ HKA) was significantly greater in TKA ($16.3^\circ \pm 4.5^\circ$) than UKA ($10.6^\circ \pm 3.1^\circ$) ($p < 0.001$), with significant changes in mLDFA and MPTA ($p < 0.001$). In UKAs, 68% of patients demonstrated a postoperative HKA between 171.2° and 175.7° , whereas in TKAs, values clustered between 174.5° and 179.8° . Mann-Whitney U test demonstrated no statistically significant differences between the UKA and TKA cohorts across all assessed PROMs, including OKS, SF-12, and FJS ($p > 0.05$ in all). 98% of the UKA an [...]

[PubMed](#) · [DOI](#)

Acetabular superior wall fractures: a distinct acetabular fracture entity.

Gänsslen A, Krappinger D, Liodakis E, Clausen JD, Omar-Pacha T, Lindtner RA, Sehmisch S, Osche DB

INTRODUCTION: A transitional "grey-zone" exists between typical locations of classical trapezoidal anterior wall and large-fragment posterior wall acetabular fractures. Part of this grey zone is the superior acetabular dome, which is commonly involved in acetabular fracture patterns, especially in geriatric patients. The aim of this study was to characterize isolated acetabular superior wall fractures and to assess their relevance as a distinct fracture pattern.

METHODS: A retrospective analysis of pelvic fracture databases from three tertiary trauma centers was performed. Acetabular fractures with superior dome involvement were screened, and cases with isolated superior articular involvement were included. Overall, seven patients were identified. Clinical and radiological data were analyzed regarding fracture morphology, associated fracture characteristics (including dislocation, comminution,

marginal impaction, intra-articular fragments, and femoral head injury), concomitant pelvic ring injuries, treatment, and clinical outcomes.

RESULTS: Isolated superior wall acetabular fractures occurred following both high-energy and low-energy mechanisms, including simple falls. The mean age was 48 years and four of seven patients were male. Hip dislocation or subluxation was observed in four cases (anterior-superior or purely superior). Surgical treatment was performed in six cases, p [...]

[PubMed](#) · [DOI](#)

A novel biomechanical model for reproducing and analysing the causal mechanisms of PFNA "cut-in" phenomenon.

Rasappan K, Joshua K, Chou SM, Shaw LKRM, Huang D, Yew A, Kwek EBK

INTRODUCTION: "Cut-in" is a rare but serious complication associated with cephalomedullary nail fixation for intertrochanteric hip fractures, particularly with Proximal Femoral Nail Antirotation (PFNA). This phenomenon involves paradoxical superomedial migration of the helical blade through the femoral head. Although increasingly recognised, the underlying mechanisms remain incompletely defined. This study aims to describe a new superolateral tensile loading set up in a bidirectional loading model, with the aim of incorporating possible mechanical deviations introduced by abductor weakness.

MATERIALS AND METHODS: This study was conducted in two stages. An initial stage with synthetic femurs (SYNBONE® 2420) was used to calibrate the set up. Within this stage, intramedullary canal diameter was compared (12 mm vs. 18 mm). In the final stage, 5 osteoporotic synthetic femurs (Sawbones® 3503) were used. In both stages, standardised AO/OTA 31-A1.1 intertrochanteric fracture were created in the synthetic femurs and fixed with PFNA-II implants. All femurs were subjected to cyclical bidirectional loading with a universal testing machine. Each cycle consisted of axial compression (720 N) and superolateral tensile loading (120 N) at 2 Hz. This configuration was designed to simulate altered hip biomechanics postulated with the trendelenburg gait. Testing was continued until mechanical fail [...]

[PubMed](#) · [DOI](#)

Routine intraoperative infection testing in presumed aseptic hip and knee revision: a matched cohort retrospective study.

Loppini M, Rocchi C, Di Maio M, Chiappetta K, Bulgarelli A, Grappiolo G

INTRODUCTION: Periprosthetic joint infections (PJIs) complicate 1-2% of primary and up to 4% of revision arthroplasties, and their diagnosis remains challenging due to the lack of a diagnostic gold standard. Differentiating PJI from aseptic failure is particularly difficult in low-grade or biofilm-related infections. Routine intraoperative microbiological screening is not recommended in presumed aseptic revisions. This study aimed to evaluate the association between routine intraoperative microbiological screening and implant survival in presumed aseptic revision arthroplasty. A secondary objective was to identify preoperative predictors of unexpected PJIs.

METHODS: Retrospectively collected data of patients undergoing presumed aseptic THA and TKA revisions between 2013 and 2019 were included. Two matched cohorts were compared: a screened group (2016-2019) undergoing routine intraoperative microbiological sampling, and an unscreened one (2013-2015) without systematic screening. Kaplan-Meier survival analysis with log-rank testing assessed overall and infection-free survival. Univariate conditional logistic regression and receiver operating characteristic (ROC) analyses evaluated potential associations between preoperative serum markers and unexpected PJI.

RESULTS: A total of 559 patients were included, of whom 295 underwent screening and 264 did not. Unexpected infections occ [...]

[PubMed](#) · [DOI](#)

Artificial intelligence advancements for orthopaedic clinical reasoning: longitudinal assessment of newer models (ChatGPT-5, Grok-3, Gemini 2.5 Flash) compared to clinicians.

Agharia S, Soroush S, Ameen D, Zhou Y

INTRODUCTION: This descriptive study aimed to longitudinally evaluate the performance of contemporary large language models - ChatGPT-5, Gemini 2.5 Flash, and Grok-3 - on orthopaedic clinical multiple-choice tasks,

benchmarked against pooled clinician consensus. A secondary aim was to assess whether recent advances in generative AI translated into improved alignment with clinician consensus compared with previous AI models.

MATERIALS AND METHODS: A total of 97 multiple-choice clinical cases spanning eight orthopaedic subspecialties were sourced from OrthoBullets and previously benchmarked against aggregated responses from thousands of practising clinicians. Using identical methodology to our 2023 study of ChatGPT-3.5, ChatGPT-4, and Bard, each model was prompted with standardised case stems and response options. The primary outcome was the proportion of AI responses matching the most popular clinician response; secondary analyses assessed agreement within 10% and 20% of clinician consensus, performance on 'controversial' (< 25% margin) questions, and inter-model concordance using Cohen's kappa coefficients.

RESULTS: Gemini 2.5 Flash achieved the highest alignment with clinician consensus (69.1%), followed by Grok-3 (66.0%) and ChatGPT-5 (58.8%). None of the LLMs refused to respond to any prompts, representing a reduction from 7.2% from our 2023 study. Subspecialty analysis de [...]

[PubMed](#) · [DOI](#)

Trajectory of urinary incontinence symptoms following total hip arthroplasty for hip osteoarthritis.

Kadowaki M, Fukutani S, Ushio K, Uchio Y

INTRODUCTION: Urinary incontinence (UI), a prevalent quality-of-life-limiting condition, frequently coexists with hip osteoarthritis, potentially through pelvic floor dysfunction related to obturator internus muscle involvement. Several reports suggest improvement after total hip arthroplasty (THA), although postoperative time-course changes remain unclear. This study aimed to clarify the temporal trends in UI after THA and examine association of postoperative UI trajectories with functional recovery.

MATERIALS AND METHODS: This retrospective study included 118 female patients who underwent THA for hip osteoarthritis between October 2021 and March 2024. UI, assessed using the International Consultation on Incontinence Questionnaire-Short Form (ICIQ-SF), was defined as ICIQ-SF ≥ 1 preoperatively and at 3, 6, 9, and 12 months postoperatively, and classified as stress (SUI), urge (UUI), or mixed (MUI). Hip-related patient-reported outcomes were evaluated using the Japanese Orthopaedic Association Hip Disease Evaluation Questionnaire (JHEQ). Associations with BMI, age, surgical approach (direct lateral/Bauer vs. modified Watson-Jones/OCM), and UI type were tested using Mann-Whitney U, chi-squared, and repeated-measures ANOVA tests.

RESULTS: Of 118 patients (mean age at surgery, 67 years; mean BMI, 24.3 kg/m²) preoperatively, 73 (61.9%) reported UI (mean ICIQ-SF 7.88), including S [...]

[PubMed](#) · [DOI](#)

The grand piano ratio: an adjunctive intraoperative screening tool for detecting excessive femoral component external rotation in total knee arthroplasty.

Avci Ö, Öztürk A, Güler BO, Bayrak HÇ, Kaya AÖ, Doğan Y

BACKGROUND: Accurate femoral component rotation is essential for optimal outcomes in total knee arthroplasty (TKA), yet reliable intraoperative assessment remains challenging. The "grand piano sign" has been described as a qualitative visual cue, but its quantitative clinical value has not been clearly established. This study aimed to evaluate the diagnostic performance of the grand piano ratio as an intraoperative tool for detecting excessive femoral component external rotation using advanced postoperative imaging as a reference standard.

METHODS: A retrospective analysis was conducted on 170 knees undergoing primary TKA. The intraoperative grand piano ratio was measured after completion of all femoral resections and before trial component implantation using the anterior femoral resection surface. Postoperative femoral component rotation was assessed using MAVRIC-sequence magnetic resonance imaging referenced to the surgical transepicondylar axis (sTEA). Excessive femoral component external rotation was defined as postoperative external rotation greater than 3° relative to the sTEA reference. Receiver operating characteristic (ROC) analysis was performed to evaluate diagnostic performance of the grand piano ratio and determine the optimal cutoff value, while multivariable logistic regression analysis was used to identify independent predictors of excessive external rotation. [...]

[PubMed](#) · [DOI](#)

The shoulder stiffness scale: a validated tool for monitoring stiffness in arthroscopic rotator cuff reconstructions.

Bouaicha S, Biner M, Selman F, Wieser K, Müller AM, Audigé L

BACKGROUND: Postoperative shoulder stiffness after arthroscopic rotator cuff repair is common and clinically heterogeneous. Established shoulder scores assess pain and function, but do not specifically isolate stiffness during follow-up. The aim of this study was to validate the newly developed Shoulder Stiffness Scale (SSS) as a pragmatic monitoring tool for postoperative shoulder stiffness after arthroscopic rotator cuff repair (ARCR).

METHODS: From a cohort of 973 primary arthroscopic rotator cuff reconstructions, patients without concomitant joint pathology of the same or contralateral arm were included. The new SSS was based on three items related to shoulder pain, subjective and objective ROM ranging from 0 to 10 points. It was collected before surgery, as well as 6 and 12 months postoperatively, along with other pain level parameters, Constant Score (CS), Subjective Shoulder Value (SSV), Oxford Shoulder Score (OSS). Changes of the SSS and its subscales over time, Cronbach's alpha for internal consistency, bottom and ceiling effects were reported. Validity was investigated through SSS associations with other parameters of similar dimension, and functional scores.

RESULTS: The selected cohort with 740 patients showed postoperative shoulder stiffness in 12.7 %. The SSS averaged 5.1 points (SD 2.5) preoperatively, and decreased significantly to mean 2.9 (SD 2.4) and 1.6 (S [...]

[PubMed](#) · [DOI](#)

Isolated screw fixation of posterior wall fractures.

Clausen JD, Özbek H, Omar-Pacha T, Sehmisch S, Gänsslen A

INTRODUCTION: Posterior wall (PW) fractures are common acetabular fractures and are most often treated by open reduction and internal fixation. The gold-standard of surgical stabilization is screw fixation of large wall fragments with additional buttress plating. Some reports focussed on screw fixation alone presenting adequate results.

MATERIAL AND METHODS: From a total of 208 PW-fracture treated between 1972 and 2008, 57 patients were identified with open reduction and internal fixation (ORIF) using isolated screw fixation. These patients were analyzed regarding demographical data (patient age, sex), type of accident, injury mechanism, Injury Severity Score (ISS), concomitant pelvic ring injuries, associated injuries, perioperative parameters and long-term results.

RESULTS: 44 patients were male and 13 were female with a mean age of 36.8 years. More than 90% sustained high-energy trauma with 34 sustaining multiple injuries or polytrauma. The mean Injury Severity Score was 11.5 points. All patients had fracture dislocations, which were reduced within 24 h after admission except one patient with secondary transfer. 11 patients presented with an initial sciatic nerve deficit. 29 patients had a single PW-fragment, 16 presented with two PW-fragments and in 12 patients marginal impactions were present. Surgery was performed in average after 7 days using the Kocher-Langenbeck appr [...]

[PubMed](#) · [DOI](#)

Research progress on debridement, antibiotics, and implant retention (DAIR) for the treatment of periprosthetic joint infection after artificial joint replacement.

Feng W, Zhang Z, Pu R, Wang H, Liu R, Sun Y, Zhang G

Periprosthetic joint infection (PJI) is one of the most serious complications following artificial joint replacement, posing significant treatment challenges and a heavy clinical burden. Debridement, antibiotics, and implant retention (DAIR) has become the preferred treatment strategy for acute PJI due to its advantages of minimal invasiveness, functional preservation, and cost-effectiveness. This article reviews the latest advances in optimizing the indications, timing of intervention, key technical details, and multimodal anti-infection strategies for DAIR. It focuses on discussing patient selection models (e.g., KLIC score), the impact of modular component exchange on biofilm eradication, the synergistic role of chemical debridement, and the central position of biofilm-active therapy in penetrating biofilms. Furthermore, this article proposes that future efforts should integrate molecular diagnostics, intelligent antimicrobial materials, and artificial intelligence prediction models to achieve individualized precision therapy for PJI, providing a theoretical basis for clinical decision-making.

[PubMed](#) · [DOI](#)

Tranexamic acid in high-risk shoulder arthroplasty patients: safety across thromboembolic, cardiac, renal, and neurologic risk profiles.

Parmar T, Adio A, Handy A, Daher M, Kellish A, Khan A, Abboud J

INTRODUCTION: Tranexamic acid (TXA) effectively reduces blood loss and transfusion requirements in shoulder arthroplasty. However, concerns regarding thromboembolic, neurologic, cardiac, and renal complications have limited its use in medically high-risk patients. This study evaluated temporal trends and the safety of TXA in high-risk patients undergoing total shoulder arthroplasty (TSA).

METHODS: A retrospective cohort study was conducted using TriNetX, identifying patients who underwent shoulder arthroplasty (2012-2025). Patients were stratified by preexisting high-risk conditions, including prior thromboembolism, renal failure, atrial fibrillation, seizure disorders, and visual disturbances. Within each subgroup, patients receiving TXA were propensity score-matched to those who did not. Perioperative TXA utilization trends were assessed. Ninety-day postoperative outcomes were compared, including transfusion requirements, thromboembolic events, cardiac complications, renal failure, neurologic events, infections, readmissions, emergency department visits, and mortality. Outcomes were reported as odds ratios (ORs) with 95% confidence intervals (CIs).

RESULTS: Reported perioperative TXA use began in 2012 and increased through 2025, with over 70% of patients in both standard- and high-risk cohorts receiving TXA. TXA use was associated with significantly lower transfusion rates [...]

[PubMed](#) · [DOI](#)

Osteoporosis treatment gap prior to femoral fracture and prevalence of pharmacological risk factors: a prospective observational study.

Böck L, Kerschan-Schindl K, Frossard M, Hajdu S, Crevenna R, Anditsch M, Stemer G, Antoni A

INTRODUCTION: Hip fractures are a major complication of osteoporosis and are associated with high morbidity, mortality, and costs. Despite clear guideline recommendations, pharmacological osteoporosis treatment remains underutilized. In addition, medications that increase fall and fracture risk further contribute to fracture occurrence. The aim of this study was to assess guideline concordance of osteoporosis therapy and to quantify the prevalence of medications associated with increased fall and fracture risk prior to the fracture in patients admitted with femoral fractures.

MATERIALS AND METHODS: In this prospective observational study, 145 patients aged ≥ 55 years admitted with femoral neck, per- and subtrochanteric, femoral shaft and distal femur fractures to a tertiary academic trauma center between April and June 2025 were included. Pre-admission medication was assessed using structured medication reconciliation. Guideline concordance of pre-fracture osteoporosis therapy was evaluated according to the current Austrian osteoporosis guideline (Dimai et al., 2024). Fall-risk-increasing drugs (FRIDs) were identified using the STOPPFall criteria, and fracture-associated medications were recorded based on Austrian osteoporosis guideline-defined drug classes.

RESULTS: Among 145 included patients (mean age 80.4 years; 71% female), 117 (81%) met criteria for pharmacological oste [...]

[PubMed](#) · [DOI](#)

Gait speed recovery after iliofemoral ligament-preserving hip arthroscopy for femoroacetabular impingement: associations with early postoperative gait kinematics.

Machida S, Tsutsumi M, Utsunomiya H, Kudo S

INTRODUCTION: Whether iliofemoral ligament-preserving hip arthroscopy facilitates recovery of gait speed-an indicator of functional recovery following hip arthroscopy in patients with femoroacetabular impingement-remains unclear. We investigated whether gait speed improved after iliofemoral ligament-preserving hip arthroscopy and examined whether pelvic jerk, defined as the time derivative of acceleration and assessed using inertial sensors at 1 month postoperatively (1M), was associated with gait speed at 6 months (6M).

MATERIALS AND METHODS: Eighteen patients who underwent primary hip arthroscopy for femoroacetabular impingement were included. Gait speed and pelvic jerk were assessed during a 10-m walk at a self-selected

speed, using an inertial sensor positioned at the third lumbar level. Peak pelvic jerks during the 1st and 2nd halves of the stance phase (1st - and 2nd -peak pelvic jerks) were calculated. Measurements were obtained preoperatively and at 1M and 6M. Gait parameters were compared across time points using a linear mixed model. Bayesian regression was used to compare the observed change in gait speed from preoperatively to 6M with previously reported values. Associations between gait variables at 1M and gait speed at 6M were also examined.

RESULTS: Gait speed was significantly higher at 6M than that pre surgery (preoperative, 1.15 [95% confidence interval, 1.0 [...]

[PubMed](#) · [DOI](#)

Computed tomography findings in 11,504 adult patients with traumatic brain injury: a large real-world cohort study with a S100B subgroup analysis.

Clar C, Puchwein P, Moshhammer M, Sadoghi P, Kramer D, Leithner A, Reinbacher P

OBJECTIVE: The aim of this study was to characterize acute cranial CT findings in a large cohort of adult patients with traumatic brain injury and to explore the diagnostic performance of S100B in a selected subgroup. The hypothesis was that most cranial CT scans in adult patients with traumatic brain injury would not demonstrate acute traumatic abnormalities, and that, within a selected subgroup, negative S100B levels would be associated with the absence of such findings on CT.

METHODS: This retrospective cohort study included 11,504 adult patients presenting with traumatic brain injury to a level I trauma center between 04/2016 and 05/2024. All patients underwent cranial CT, while serum S100B measurements were available in a selected subset. CT findings were classified as normal or showing acute traumatic abnormalities. In the S100B subgroup, biomarker levels were analyzed in relation to CT findings to assess their diagnostic performance for excluding acute pathology.

RESULTS: A total of 11,504 adult patients with traumatic brain injury were included; 27.4% were inpatients and 72.6% outpatients. Overall, 81.9% of CT scans showed no acute pathology, while 10.7% demonstrated abnormalities. 7.4% of all cases were excluded. Acute TBI-related interventions were rare (3.7% of inpatients vs. 0.3% of outpatients). In patients with dementia and those receiving anticoagulation, CT sc [...]

[PubMed](#) · [DOI](#)

Early post-operative CRP is a better predictor of DAIR failure than pre-operative CRP in total knee PJI.

Beadel H, Chaffey R, Kim K, Zhu M, Coleman B

INTRODUCTION: Debridement, antibiotics and implant retention (DAIR) surgery is often performed as a first-line treatment for periprosthetic joint infection (PJI), however, success rates are variable. This study aimed to determine whether post-operative C-reactive protein (CRP) levels predict DAIR failure.

MATERIALS AND METHODS: A multi-centre retrospective cohort study was performed over a 15-year period. All total knee arthroplasty (TKA) patients undergoing DAIR for first episode PJI were included. CRP was measured at admission and for 6 weeks post-operatively. Treatment success was defined as implant retention without the need for revision surgery or long-term suppressive antibiotics. Trends in CRP were compared between two groups: failed and successful DAIR. Receiver operating characteristic (ROC) curves were analysed.

RESULTS: 189 DAIR procedures were included, of which 49% (92) failed and 51% (97) were successful. Mean follow-up was 7.5 years. Overall, CRP trended down following surgery. Mean CRP was significantly higher in the failed DAIR group at all time points. The greatest difference was at week one post-operatively (mean 76 vs. 101, $P < 0.01$). This remained significant when patients experiencing DAIR failure within 90 days were excluded (mean 76 vs. 107, $P < 0.01$). CRP was most accurate as a predictor of failure at week one post-operatively, with an area under the [...]

[PubMed](#) · [DOI](#)

Tuberosity refixation in reverse proximal humerus fracture arthroplasty: a prospective cohort study with two-year follow-up.

Oberrauter CH, Hess F, Welter J, Pape HC, Mayer S, Dullenkopf A, Mazzucchelli R, Marintschev I

INTRODUCTION: To evaluate the outcomes of a simplified tuberosity refixation technique using a two-suture construct in patients with complex proximal humerus fractures treated with reverse shoulder arthroplasty (RSA). The technique reduces the number of fixation sutures rather than the traditional four to five, aiming to decrease operative complexity while promoting greater tuberosity (GT) healing and improving the healing rate. A secondary exploratory analysis compared clinical and radiological outcomes between cemented and uncemented humeral stem fixation.

MATERIALS AND METHODS: This prospective study included 73 patients who underwent RSA (Global Unite Fracture Reverse System, DePuy Synthes, Warsaw, Indiana, USA). Clinical and radiological outcomes were assessed at two-year follow-up. Bone quality was evaluated using the Deltoid Tuberosity Index (DTI). Functional outcomes and GT healing were analyzed, and stem fixation type was assessed in a secondary subgroup analysis.

RESULTS: The mean patient age was 79 ± 8 years (range 61-95), and 65 (89%) were female. Osteoporotic bone quality (DTI < 1.4) was observed in 51 (70%) of cases. At two years, the median outcome scores were: Subjective Shoulder Value, 90 (IQR 80-95); Constant-Murley score, 76 (IQR 73-81); active forward flexion, 140° (IQR 120-160); abduction, 140° (IQR 120-160); external rotation, 8 (IQR 8-8); and internal r [...]

[PubMed](#) · [DOI](#)

Is it time for a systematic and objective assessment of knee laxity? The impact of objective knee laxity after TKA on patient-reported outcomes: a comprehensive review.

Vermue H, Klincke V, Stroobant L, Tampere T, Victor J, Arnout N

PURPOSE: Over the past decade, robot assistance has emerged as an innovative and rapidly advancing tool in total knee arthroplasty (TKA). Despite technological advancements, the evaluation of the soft-tissue envelope during TKA often relies on subjective and rudimentary assessments by the surgeon. This literature review explores the relationship between objective intraoperative and postoperative laxity measurements in the coronal and sagittal planes and patient-reported outcome measures (PROMs) following TKA.

METHODS: A literature review was conducted using PubMed/MEDLINE and Embase to identify studies on intra- or postoperative laxity measurements in TKA and their correlation with PROMs. Inclusion criteria required studies on TKA patients with quantifiable coronal and/or sagittal laxity measurements and postoperative PROMs linked to these measurements.

RESULTS: Thirty-four studies met inclusion criteria: ten on sagittal laxity, twenty-seven on coronal laxity, and three on both. Follow-up ranged from 1 month to over 8 years. Sagittal laxity thresholds of 5-10 mm under applied loads of 89-133 N were associated with better functional outcomes, whereas excessive (> 10 mm) or overly tight (< 5 mm) laxity correlated with reduced satisfaction. Medial laxity in the coronal plane was linked to lower PROMs, while the impact of lateral laxity on PROMs was less conclusive.

CONCLUSIONS: [...]

[PubMed](#) · [DOI](#)

Silver-impregnated occlusive dressing is associated with a lower rate of acute surgical site infection after direct anterior total hip arthroplasty.

Tsai YY, Shen PH, Wu CC, Pan RY, Lin LC, Wang SH

INTRODUCTION: The direct anterior approach (DAA) for total hip arthroplasty (THA) has gained increasing adoption in recent years; however, it has been associated with a higher risk of wound-related complications compared with other surgical approaches. Silver-impregnated occlusive dressings have demonstrated antimicrobial potential, though their role in DAA THA remains underexplored.

MATERIALS AND METHODS: We conducted a retrospective cohort study with a historical control group, reviewing 321 consecutive primary DAA THAs performed by a single surgeon (September 2020-December 2025). Standard sterile gauze was used before August 2021 (n = 76) and a silver-impregnated occlusive dressing (AQUACEL® Ag SURGICAL; ConvaTec, Deeside, UK) thereafter (n = 245). Despite the chronological difference between cohorts, baseline demographic and clinical characteristics did not differ significantly between groups (Table 1). The primary endpoint was acute surgical site infection (SSI) within 30 days (CDC criteria); all patients had ≥ 3 months follow-up to capture early wound-related reoperations as secondary outcomes.

RESULTS: Four acute SSI events occurred (1.25%). SSI incidence was 3.94% (3/76) with gauze versus 0.41% (1/245) with the silver-impregnated dressing (Fisher's exact $p = 0.043$). This corresponds to an absolute risk reduction (ARR) of 3.54% (95% CI, 0.34-10.57) and a number needed to treat [...]

[PubMed](#) · [DOI](#)

Minimally invasive stabilisation of anterior pelvic ring injuries through anterior infix versus percutaneous anterior retrograde pubic screw: a prospective comparative cohort study.

Goda El-Hamalawy A, Abdelmoneim M, Al-Sayed M, Youness M, Sadek F, Kassem E

BACKGROUND: Management of unstable anterior pelvic ring fractures has shifted toward minimally invasive fixation techniques. This prospective randomized cohort study compares anterior subcutaneous internal fixator with percutaneous retrograde pubic ramus screw fixation, evaluating radiological outcomes, functional scores, complications, and peri-operative parameters.

METHODS: Fifty adult patients with unstable anterior pelvic ring injuries (Tile B & C; Young-Burgess lateral compression and vertical shear mechanisms) were enrolled between 2023 and 2025 and randomized into two equal groups: group 1 (internal fixator) and group 2 (retrograde pubic screw). Both groups underwent individualized posterior ring fixation as required. Outcomes assessed included operative time, blood loss, fluoroscopy time, union rate, Matta radiological score at 6 months, Majeed and Pelvic Outcome Scores at 12 months, and post-operative complications.

RESULTS: Management of anterior pelvic injury showed that retrograde screw fixation demonstrated significantly shorter operative time (32.0 ± 9.2 vs. 50.3 ± 8.4 min, $p < 0.001$) and less intraoperative blood loss (68.3 ± 16.2 vs. 122.3 ± 31.8 ml, $p < 0.001$). Radiological outcomes were comparable: excellent Matta score in 68% vs. 64% ($p = 0.834$). Union time showed no significant difference (14.1 ± 2.4 vs. 15.1 ± 1.5 weeks, $p = 0.076$). Functional outcomes at [...]

[PubMed](#) · [DOI](#)

Prioritisation of assessments, diagnostic classifications, and outcome measures in Perthes disease: a Delphi survey of international health professionals.

Ball S, Davies LM, Gray K, Pacey V

INTRODUCTION: The aim of this study is to reach consensus and prioritise the most clinically important assessments, diagnostic classifications, and outcome measures by an international panel of experts for Legg-Calve-Perthes disease.

MATERIALS AND METHODS: A three-round modified e-Delphi survey was conducted between April and May 2025. Participants rated the clinical importance of 14 clinical assessments, 26 radiological assessments, eight diagnostic classifications, and 26 outcome measures identified by a scoping review. Consensus was defined as $\geq 75\%$ agreement. In the final round, items in each category were prioritised for overall condition and by specific age groups (< 6 years, 6-8 years, > 8 years).

RESULTS: Thirty-eight health professionals, from nine countries participated in round 1. Thirty-four completed all three rounds, a retention rate of 89.5%. Consensus and prioritisation for clinical use were achieved for 10 clinical assessments, three radiological assessments, three diagnostic classifications, and three outcome measures.

CONCLUSION: Paediatric specialists have reached consensus and prioritised methods to evaluate and diagnose Legg-Calve-Perthes disease in clinical practice. These results represent an important step towards improving continuity of care and the foundation for future development of clinical practice guidelines.

[PubMed](#) · [DOI](#)

Compensatory activation of the deltoid and remaining rotator cuff muscles in patients with rotator cuff tears using real-time tissue elastography.

Hoshikawa K, Oishi R, Yuri T, Uno T, Nagai J, Giambini H, Mura N

INTRODUCTION: Rotator cuff (RC) tears are a prevalent condition leading to dysfunction and shoulder pain. Spatial changes in the behavior of the deltoid and remaining RC muscles during arm elevation in patients with RC tears have not been investigated. The purpose of this study was twofold: (1) to investigate factors related to the

changes in the deltoid muscle behavior during scapular plane abduction (scaption) in patients with RC tears, and (2) to examine the functional relationship between the deltoid and remaining RC muscles, particularly remaining infraspinatus (ISP) and teres minor (TMin) muscles, in these patients using ultrasound elastography.

METHODS: Twenty-four patients (mean age, 66 ± 6 years) with RC tears (mean antero-posterior diameter, 17.4 ± 8.3 mm) were evaluated. Antero-posterior and medio-lateral diameters of the posterosuperior RC and Subscapularis (SSC) tears were measured intra-operatively. Muscle elasticity was evaluated at passive and active states, as a proxy for muscle activation, at 30° , 60° , and 90° during scaption for the deltoid, ISP, and TMin muscles using real-time tissue elastography. Based on the deltoid muscle activity ratio at 30° and 90° scaption, patients were divided into normal and abnormal activation groups.

RESULTS: Activity of the deltoid muscle in the abnormal activation group exhibited consistently high activity across various ele [...]

[PubMed](#) · [DOI](#)

Pin orthosis extension block pinning versus conservative treatment for doyle type 4B mallet fractures.

Col KA, Demirsu O, Aydn M, Surucu S, Yilmaz M, Atlihan D

INTRODUCTION: Although numerous treatment options have been reported for mallet fractures, a universally accepted gold-standard approach has not yet been established. The purpose of this study was to compare the clinical and radiographic outcomes of the pin-orthosis extension-block pinning technique (PO-EBPT) with those of conventional conservative treatment in patients with Doyle type 4B mallet fractures.

MATERIALS AND METHODS: This study included 62 patients with Doyle type 4B mallet fractures involving 20-50% of the distal interphalangeal (DIP) joint articular surface, treated between March 2022 and April 2024. Patients were randomized into two groups: Group 1 (n = 33) underwent PO-EBPT, whereas Group 2 (n = 29) received conservative treatment with splint immobilization. Outcome measures included DIP joint extension lag, range of motion, fracture union, complication rates and functional outcomes according to the Crawford criteria. Follow-up evaluations were performed at 2, 4 and 6 weeks and at 3, 6 and 12 months.

RESULTS: A total of 62 patients were analyzed (33 PO-EBPT; 29 conservative). No statistically significant differences were observed between the groups with respect to sex, affected side, injured finger, or complication rates ($p = 0.461$, $p = 0.658$, $p = 0.763$ and $p = 0.165$, respectively). However, the PO-EBPT group demonstrated significantly improved DIP joint exten [...]

[PubMed](#) · [DOI](#)

Impact of cannabis use and tobacco smoking on outcomes of open carpal tunnel release surgery: a nationwide study in the United States.

Hoveidaei AH, Esmaeili S, Gilmor R, Chen Z, Conway JD, Ingari JV

PURPOSE: To evaluate the impact of cannabis use, tobacco smoking, and their concurrent use on postoperative complications following open carpal tunnel release (CTR) using a nationwide database.

METHODS: A retrospective cohort study was conducted using the PearlDiver database (2010-2022) to identify adult patients who underwent open CTR. Patients were categorized into four groups: cannabis-only users (n = 800), tobacco-only users (n = 167,803), concurrent users (n = 3529), and non-users (n = 167,803 matched controls). Postoperative complications, surgical site infection, wound disruption, joint stiffness, and complex regional pain syndrome, were assessed at 6 weeks and 3 months postoperatively.

RESULTS: At 6 weeks and 3 months, surgical site infection was more common among concurrent cannabis and tobacco users, cannabis-only users, and tobacco-only users. Wound disruption occurred more frequently in concurrent users and tobacco-only users, but not in cannabis-only users. Complex regional pain syndrome was associated only with tobacco use. No association was found between any substance use and joint stiffness.

CONCLUSION: Cannabis use, tobacco smoking, and especially concurrent use increase the risk of surgical site infection and wound disruption following open CTR. Tobacco use is independently associated with complex regional pain syndrome, while cannabis may have a protectiv [...]

[PubMed](#) · [DOI](#)

Timing and microbiological profile influence long-term outcomes after debridement, antibiotics, and implant retention (DAIR) in acute hip periprosthetic joint infection.

Muñoz-Mahamud E, Perdomo-Lizarraga JC, Combalia A, Alías A, Serra A, Fernández-Valencia JÁ, Verdejo MÁ, Soriano Á

INTRODUCTION: The strategy of debridement, antibiotics, and implant retention (DAIR) represents one of the main therapeutic modalities for acute periprosthetic joint infection (PJI). However, reported outcomes remain highly variable, with success rates ranging from 16 to 92%. This study aimed to evaluate long-term treatment failure-free survival and identify risk factors for treatment failure in patients with acute hip PJI treated with DAIR using time-to-event analytical methods.

MATERIAL AND METHODS: This retrospective study evaluated the treatment failure rate in 115 patients treated with the DAIR strategy for acute PJI following primary or aseptic revision hip arthroplasty between 1999 and 2018. Potential predictors of treatment failure-free survival, including patient characteristics, infection microbiology, and surgical timing, were analyzed using univariate tests, Cox proportional hazards regression (hazard ratio [HR]), and Kaplan-Meier survival estimates.

RESULTS: Among the 115 patients included, 56 were women and 59 were men; 48 were aged 75 years or younger and 67 were older than 75 years. In 88 cases, the infection occurred after primary arthroplasty, whereas 27 followed aseptic revision surgery. The mean follow-up was 7.1 years (SD 4.4). At five years, treatment failure occurred in 30.4% of cases. Procedures performed within seven days of diagnosis showed lower fai [...]

[PubMed](#) · [DOI](#)

Surgical treatment of varus unicompartmental knee osteoarthritis: indications, trends, and outcomes-a narrative review.

Burgio C, Bosco F, Cobisi C, Zepeda K, Giustra F, Capella M, Mayman D, Debbi E et al.

INTRODUCTION: Varus unicompartmental knee osteoarthritis (UC-KOA) is a frequent and heterogeneous clinical condition characterized by compartment-specific cartilage degeneration and frequently associated with lower-limb malalignment. The progressive understanding of knee biomechanics and alignment has led to the development of surgical strategies specifically tailored to address isolated compartment disease. The purpose of this narrative review is to evaluate indications, current trends, and clinical outcomes of the main surgical options for varus UC-KOA, with particular focus on realignment osteotomies and unicompartmental knee arthroplasty (UKA).

MATERIALS AND METHODS: A narrative review of the literature was conducted using major biomedical databases. Evidence related to biomechanics, alignment correction, osteotomy techniques, UKA indications, implant evolution, complication profiles, and comparative outcomes was analyzed and synthesized thematically.

RESULTS: Biomechanical and clinical evidence consistently identifies malalignment as a key driver of isolated compartment degeneration. Corrective osteotomies effectively restore physiological load distribution and provide durable outcomes in selected patients with extra-articular deformity. Advances in implant design and surgical techniques have substantially improved UKA survivorship, allowing broader patient selection wit [...]

[PubMed](#) · [DOI](#)

Preoperative NarxCare overdose risk scores greater than 100 are associated with worse PROMs improvements and dissatisfaction after primary THA.

Khan ST, Emara AK, Patel SV, Elmenawi KA, Ibaseta A, Pasqualini I, Zhang C, Cleveland Clinic Adult Reconstruction Research Group et al.

INTRODUCTION: The NarxCare Overdose Risk Score (ORS) is a weighted measure of patient-specific prescription drug use. In patients undergoing primary total hip arthroplasty (THA), this study aimed to evaluate (1) the trend of the ORS from preoperative to 1-year postoperatively, and (2) the association of preoperative ORS with achievement of clinically meaningful improvements in patient reported outcome measures (PROMs), and satisfaction at 1-year.

METHODS: All patients who underwent primary unilateral elective THA at a USA tertiary healthcare system from January 2016-December 2022 were eligible. Patients with incomplete PROMs or missing baseline NarxCare ORS were excluded, leading to the inclusion of 5,424 patients. Multivariable regression models were used to assess the association between ORS and 1-year PROMs, including achievement of the minimum clinically important

difference (MCID) and Patient Acceptable Symptom State (PASS) for the Hip Disability and Osteoarthritis Outcome Score (HOOS) subdomain scores, as well as satisfaction.

RESULTS: Preoperatively, 46.1% (n = 2501) had a NarxCare ORS of 0, 24.6% (n = 1336) had scores between 100 and 199, and 16.2% (n = 876) in the 200-299 range. After adjusting for confounding variables, a preoperative ORS of 100-199 was associated with failure to achieve MCID in HOOS Pain (OR 1.91;CI 1.32-2.77;p = 0.001), and HOOS JR (OR 1.54;1.10-2 [...])

[PubMed](#) · [DOI](#)

Only a partial shield: local vancomycin postpones but does not prevent hip and knee prosthetic infections.

Darup A, Ettinger M, Savov P, Brand S, Seeber GH, Stauss R

INTRODUCTION: Periprosthetic joint infection (PJI) represents a severe complication following total hip arthroplasty (THA) and total knee arthroplasty (TKA). The intraarticular application of vancomycin powder is increasingly integrated into perioperative standards as an innovative strategy for PJI prevention. Yet the efficacy of this method remains a topic of debate. The aim of this retrospective cohort study was to evaluate whether topical vancomycin application reduces the incidence of PJI after primary total joint arthroplasty (TJA) and whether it is associated with an increased risk of wound complications.

MATERIALS AND METHODS: In this retrospective monocentric cohort study, all primary THA and TKA procedures performed between January 1, 2022, and December 31, 2023, were analysed. From January 1, 2023, intraarticular vancomycin powder was implemented as part of the standard perioperative protocol for PJI prevention. The PJI rates, postoperative surgery-related complications, and time to infection were compared between the vancomycin and control groups.

RESULTS: A total of 1,499 patients were included in the study. Patients were divided into two groups: a vancomycin group (VG, n = 818) and a control group (CG, n = 681). No statistically significant group differences were observed for wound healing disorders (p = 0.775) or PJI rates (1.0% VG vs. 0.9% CG, p = 0.846). However [...]

[PubMed](#) · [DOI](#)

Validity and prognostic value of a novel intraoperative assessment tool for reduction in trimalleolar fractures: the SGSC checklist approach.

Gao H, Xu G, Ma J, Wang Y, Wang J, Zhou R, Liu Z, Wang B

BACKGROUND: Trimalleolar fractures present substantial surgical challenges, and the quality of fracture reduction significantly influences patient outcomes. Current intraoperative assessment methods, including fluoroscopy and arthroscopy, each possess inherent limitations when used independently. This study aimed to validate the effectiveness of a novel intraoperative evaluation tool, the SGSC (Step, Gap, Syndesmosis, Congruity) "traffic light" checklist, that integrates arthroscopic and fluoroscopic assessment, and to evaluate its predictive value for clinical prognosis.

METHODS: This single-center retrospective cohort study enrolled 36 patients who underwent surgical treatment for trimalleolar fractures. Validation proceeded in two stages: first, we calculated inter-rater reliability (Kappa statistic) and concordance with postoperative CT as the reference standard to establish the reliability and validity of the SGSC checklist (accuracy validation); second, patients were stratified into "All-Green" and "Non-All-Green" groups based on intraoperative SGSC assessment, and we compared AOFAS scores and early radiographic degenerative changes at final follow-up (prognostic validation).

RESULTS: For accuracy validation, inter-rater agreement between the two assessors demonstrated excellent reliability (Kappa = 0.827; 95% CI: 0.644-1.000). The overall concordance between SGSC check [...]

[PubMed](#) · [DOI](#)

Radiographic assessment of femoral, acetabular and global offset following hip spacer implantation in staged total hip arthroplasty".

Cavagnaro L, Ferrari E, Providenti V, Marciante V, Carrega G, Formica M

BACKGROUND: The number of total hip arthroplasties (THAs) performed worldwide is steadily increasing, and, consequently, the demand for revision procedures is expected to rise as well. two-stage revision procedure continues to be considered the gold standard in many clinical scenarios. During the first step, failure to restore femoral offset has been implicated as a contributor to instability and mechanical failure, although a clear consensus is still lacking.

OBJECTIVES: The primary aim of this study is to evaluate the restoration of biomechanical parameters of the hip-specifically leg length discrepancy (LLD), femoral offset (FO), acetabular offset (AO), and global offset (GO)-following implantation of a specific type of articulating hip spacer during the first stage of a two-stage revision for PJI. A secondary objective is to assess the correlation between variations in offset parameters and the rate of interstage mechanical complications.

MATERIALS AND METHODS: We retrospectively reviewed all patients undergoing staged revision with a specific articulating spacer between 2020 and 2022 at a single institution. Harris Hip Scores, Oxford Hip Scores, and Visual Analogue Scales for pain were obtained at different time points. Radiographic analysis included LLD, FO, AO and GO measurements on the affected (spacer and post-reimplantation) and the contralateral side. Complications [...]

[PubMed](#) · [DOI](#)

Simulated tears of antero-posterior rotator cuff force-couple and reduced glenoid concavity decrease anterior glenohumeral stability: a robot-assisted biomechanical analysis of a load and shift sequence.

Karlsfeld MJ, Wermers J, Tänzler S, Sußiek J, Herbst E, Katthagen JC

BACKGROUND: The glenoid concavity and the compression applied by the rotator cuff (RC) are essential for glenohumeral stability (GHS). This study aimed to determine how different simulated rotator cuff tears (RCT) and the glenoid depth influence GHS.

METHODS: A Load and Shift sequence was performed with eight fresh-frozen cadaveric shoulders in a robotic-assisted setup. Differently configured static loading of the reinforced RC and deltoid muscle (DLT) simulated intact RC and anterior, superior, anterosuperior, posterosuperior, mass, and complete RC plus DLT tears. Anterior dislocation forces and their changes were determined as indicators of stability. The glenoid depth and Bony shoulder stability ratio (BSSR) were defined as indicators of concavity. To assess GHS, the maximal force (Favg), the maximal force increase (dFavg), and their mean deviations (ΔF_{max} , ΔdF_{max}) to the intact configuration during the sequence were evaluated.

RESULTS: Simulated tears of the subscapularis tendon (SCP) ($\Delta F_{max} = 7.90$ N, $\Delta dF_{max} = 1.54$ N/mm) and simulated tears of the infraspinatus (ISP) + teres minor (TM) tendons ($\Delta F_{max} = 7.19$ N, $\Delta dF_{max} = 1.48$ N/mm) resulted in greater differences to the intact shoulder than simulated tears of the supraspinatus tendon ($\Delta F_{max} = 3.82$ N, $\Delta dF_{max} = 1.16$ N/mm). High correlations were observed between maximal force concerning glenoid depth ($r = 0.81$) and BSSR ($r = 0$ [...])

[PubMed](#) · [DOI](#)

Non-tobacco nicotine dependence is associated with increased complications following clavicle open reduction internal fixation.

Adio AA, Goyal RK, Skiena A, Daher M, Horneff JG, Kassam HF, Hill BW, Abboud JA

BACKGROUND: Tobacco dependence is well established to impair bone healing and increase postoperative complications. However, the independent effects of nicotine, separate from combustion-related factors, remain unclear. This study aimed to evaluate the impact of non-tobacco nicotine dependence (NTND) following open reduction and internal fixation (ORIF) of clavicle fractures, including midshaft and lateral fracture subtypes.

METHODS: A retrospective cohort study was performed using a large national database. Patients undergoing ORIF of clavicle fractures were identified using procedural and diagnostic codes. Patients with non-tobacco nicotine dependence (e.g., vaping, patches, or gum) were identified and propensity score matched to (1) patients without nicotine exposure and (2) patients with tobacco nicotine dependence. In a secondary analysis, patients were stratified by fracture type (midshaft and lateral), and outcomes were compared between nicotine exposure and no nicotine exposure within each subgroup. Outcomes were assessed at 90 days and 1 year and included medical complications, number of opioid prescriptions, and implant-related outcomes. Risk ratios (RRs) with 95%

confidence intervals (CIs) were calculated, with $p < 0.05$ considered significant.

RESULTS: After matching, 936 non-tobacco nicotine users were identified. Non-tobacco nicotine dependence was associated with [...]

[PubMed](#) · [DOI](#)

Association between timing of surgery and postoperative outcomes in older adults with distal femur fractures: a systematic review and meta-analysis.

Hoshika S, Yamamoto N, Saka N, Tsuge T, Sato T, Isaji Y, Nakao S

INTRODUCTION: The association between surgical timing and clinical outcomes in older adults with distal femur fractures remains unclear. We conducted a systematic review and meta-analysis to evaluate whether earlier surgery is associated with improved outcomes in this population.

MATERIALS AND METHODS: In accordance with the Preferred Reporting Items for Systematic Reviews and Meta-Analyses (PRISMA) guidelines, two independent reviewers searched MEDLINE, Embase, the Cochrane Central Register of Controlled Trials, ClinicalTrials.gov, and the World Health Organization International Clinical Trials Registry Platform from inception to October 8, 2024. The primary outcome was 30-day mortality, and secondary outcomes included longer-term mortality and postoperative complications. Risk of bias was assessed using the Quality in Prognosis Studies tool, and the certainty of evidence was evaluated using the Grading of Recommendations Assessment, Development and Evaluation approach.

RESULTS: Of 6,101 records screened, 16 retrospective cohort studies comprising 31,213 patients met the inclusion criteria. Early surgery was not associated with reduced 30-day mortality (adjusted odds ratio [OR]: 0.77, 95% confidence interval [CI]: 0.56-1.05; low-certainty evidence) or with longer-term mortality at 90 days (crude OR: 0.91, 95% CI: 0.51-1.60), 180 days (crude OR: 0.46, 95% CI: 0.12-1.77), or 1 [...]

[PubMed](#) · [DOI](#)

Is a central cavity necessary for bioactive glass-ceramic spacers in plated ACDF? A retrospective comparison of solid versus cavity designs.

Kang TH, Lee G, Kang B, Han J, Cho M, Lee JH

PURPOSE: Traditionally, interbody cages feature a central cavity for graft packing. However, for bioactive glass-ceramic (BGS-7) spacers achieving fusion via surface-mediated osseointegration, this cavity may be biologically redundant and structurally disadvantageous. We compared radiological and clinical outcomes of plated anterior cervical discectomy and fusion (ACDF) using BGS-7 spacers with or without a central cavity.

METHODS: We retrospectively reviewed patients undergoing plated ACDF using solid non-cavity (17 patients, 24 levels) or central-cavity (17 patients, 33 levels) BGS-7 spacers. Radiographic fusion was evaluated using dynamic interspinous distance (primary, ≤ 2 mm) and segmental angular motion (secondary, $\leq 4^\circ$), alongside CT-based osseointegration (direct contact $\geq 50\%$). Subsidence (≥ 2 mm) and clinical outcomes were evaluated at 12 months.

RESULTS: Both groups demonstrated comparable 12-month clinical improvements ($p > 0.05$). Radiographic fusion rates were equivalent between non-cavity and central-cavity groups based on the primary distance (91.7% vs. 84.8%, $p = 0.687$) and secondary angular criteria (79.2% vs. 63.6%, $p = 0.331$). The non-cavity group showed a trend toward superior segmental stability, with a smaller dynamic gap distance (0.82 ± 0.64 mm vs. 1.53 ± 1.99 mm, $p = 0.064$). CT-based fusion perfectly matched (90.5% vs. 90.9%, $p = 1.000$). Subsidence ra [...]

[PubMed](#) · [DOI](#)

Clinical outcomes of internal iliosciatic fixation using a conventional reconstruction plate via an anterior approach for acetabular fractures involving the posterior column.

Xiong H, Chen H, Cao S, Chen X, Wang F

BACKGROUND: Acetabular fractures involving the posterior column remain surgically challenging because of the complex three-dimensional anatomy and restricted operative corridor. As an alternative to traditional posterior approaches, internal iliosciatic fixation through an anterior approach has gained increasing acceptance. Reconstruction plates, which offer the advantages of low cost and convenient intraoperative bending, are extensively used in clinical settings.

METHODS: A retrospective case series was conducted on 36 consecutive patients with acetabular fractures involving the posterior column who underwent internal iliosciatic fixation using an intraoperatively contoured reconstruction plate between July 2019 and November 2022. The modified Stoppa approach (n = 26) or the pararectus approach (n = 10) was employed. Fracture reduction was assessed using the modified Matta criteria, and hip function was evaluated using the modified Merle d'Aubigné-Postel score at one-year follow-up. Operative time, blood loss, fracture union, and complications were recorded.

RESULTS: The mean age of the patients was 42.6 ± 14.5 years (range: 21-71 years), with 25 males and 11 females. Mechanisms of injury included traffic accidents (n = 21), falls from height (n = 10), skiing injuries (n = 3), and others (n = 2). According to the Letournel-Judet classification, there were 13 both-column fra [...]

[PubMed](#) · [DOI](#)

Novel UHMWPE sleeve for stabilizing loose suture anchors in rotator cuff repair.

Lin CK, Jaw FS, Sam SS, Siow TF, Yii SC, Chang YJ, Young TH

Suture anchors are frequently used for repairing torn rotator cuffs. However, post-operative anchor loosening is a common complication often caused by low bone density. The purpose of this study was to evaluate if a novel UHMWPE cortex locking sleeve could improve the anchor pullout strength after revision in synthetic bone. The initial failed anchor fixation was simulated by inserting a 4.5 mm anchor and pulling it out of the bone, which created an enlarged insertion hole for the revision anchor. After revision, two loosening conditions were simulated; (i) intra-operative anchor loosening where the revision anchor was directly pulled out from the block after insertion, and (ii) post-operative anchor loosening where the revision anchor was first subjected to cyclic loading and then pulled out. For both loosening conditions, the pullout strength and energy absorbed was recorded for the following configurations: (i) Group A, same-diameter anchor (SDA; 4.5 mm × 10 mm); (ii) Group B, larger-diameter anchor (LDA; 5.5 mm × 10 mm;); and (iii) Group C, same-diameter anchor combined with the novel sleeve (SDA + NS). For intra-operative anchor loosening, the highest pullout strength was recorded in the SDA + NS group, with a mean value of 370.077 ± 32.699 N, which was significantly greater than both SDA and LDA groups ($p < 0.05$). Similarly with post-operative anchor loosening, the SDAP [...]

[PubMed](#) · [DOI](#)

Impact of rheumatoid arthritis on complications and hospital outcomes following reverse shoulder arthroplasty: evidence from 389,135 hospitalizations in the national inpatient sample.

Mahamid A, Elgabsi M, Khatib M, Murad H, Qawasmi F, Lavon E, Zahalka S, Yassin A et al.

OBJECTIVE: To evaluate the impact of rheumatoid arthritis (RA) on perioperative complications and hospital outcomes following reverse total shoulder arthroplasty (RTSA) using the National Inpatient Sample (2016-2021).

METHODS: Adult patients undergoing RTSA were identified using ICD-10 procedure codes. Outcomes included prolonged length of stay (LOS), high hospital charges, inpatient complications, and mortality. Multivariable logistic regression adjusted for demographic, clinical, and hospital characteristics. Survey weights were applied to generate nationally representative estimates.

RESULTS: A total of 389,135 patients underwent RTSA, including 18,140 (4.66%) with RA. In unadjusted analyses, RA patients had higher rates of prolonged LOS (17.8% vs. 14.8%) and acute blood loss anemia (10.7% vs. 8.6%). After multivariable adjustment, RA remained an independent predictor of prolonged LOS (adjusted OR 1.18, 95% CI 1.14-1.23, $p < 0.001$) and acute blood loss anemia (adjusted OR 1.25, 95% CI 1.18-1.33, $p < 0.001$), but not urinary tract infection. In-hospital mortality was rare and could not be modeled due to zero events in the RA group. RA patients also had slightly longer hospitalizations and higher hospital charges.

CONCLUSION: RA is independently associated with prolonged hospitalization and increased risk of perioperative blood loss anemia following RTSA. These findings high [...]

[PubMed](#) · [DOI](#)

Long-term results of thumb carpometacarpal joint arthrodesis.

Vogler P, Benedikt S, Bode S, Keiler A, Stock K, Arora R

INTRODUCTION: Thumb carpometacarpal (CMC 1) arthrodesis remains an established treatment option for patients with high functional demands and strength requirements. Long-term data on mobility, complication rates and clinical outcomes remain limited. This study aimed to analyze long-term clinical and radiological results after CMC 1 arthrodesis.

MATERIALS AND METHODS: A single-center, retrospective study included patients who underwent CMC 1 arthrodesis between 2004 and 2024. Following medical record analysis, eligible patients were invited to a standardized clinical and radiological follow-up examination. Outcomes included DASH, PRWHE and MHQ scores, pain intensity, mobility, strength and patient satisfaction. Radiological assessment covered fusion signs, implant position and adjacent joint arthrosis. The median follow-up was 9 years.

RESULTS: 34 patients were included, 16 participated in follow-up. The median DASH score was 14 points, PRWHE 4 points and MHQ 80 (operated) vs. 84 (contralateral). Grip strength reached 85% of the contralateral side, with preserved pinch strength. Compensatory adaptation of adjacent joints was observed regarding range of motion. Patients reported no relevant resting pain and only minimal load-related discomfort. 15 of 16 patients were satisfied with the outcome. Non-union occurred in 4 of 34 cases (11.8%), associated with absent cancellous bone [...]

[PubMed](#) · [DOI](#)

Does the tendon retract in flexor hallucis longus muscle laceration? A cadaveric study.

Aydin M, Mahin Y, Yasar H, Çiçek F, Ceranoğlu FG, Mert A, Çınarolu S

INTRODUCTION: Flexor hallucis longus (FHL) tendon injuries are common, particularly in open injuries caused by sharp objects. However, data regarding FHL tendon retraction following hallux-level FHL tendon lacerations remain limited. This study investigated the retraction and shortening of the FHL tendon using cadaveric models.

MATERIALS AND METHODS: Six fresh-frozen below-knee amputated cadaveric feet (mean age: 65 years, range: 62-88) were utilized. FHL tendons were dissected, and lacerations were simulated at the hallux level. Tendons were subjected to various tensile forces (20-120 N) using a mechanical device. Force was measured in Newtons via a loadcell, and retraction distance was recorded in millimeters using an encoder. Data were summarized as mean ± standard deviation (SD), median, and quartiles.

RESULTS: The FHL tendon retracted gradually as applied force increased. An average retraction of 7 mm was measured at 20 N, reaching 21 mm at 60 N. Beyond 60 N (> 21 mm retraction), the other four toes also flexed. This occurs due to the anatomical connection between the FHL and flexor digitorum longus (FDL) tendons.

CONCLUSIONS: Only slight retraction was observed in hallux-level FHL tendon lacerations. At most, the findings suggest that distal FHL lacerations may exhibit limited proximal retraction under the tested conditions, which may have implications for surgical pla [...]

[PubMed](#) · [DOI](#)

Early-onset rice body synovitis of the knee in children under three years: a case series and review of the literature.

Zhu L, Liu C, Liu Y, Mao B, Huang Y, Wen J, Liu Y, Qian T

OBJECTIVE: Rice body synovitis reports have been rarely reported in children under three years of age. We aimed to explore the clinical characteristics and treatment experience of rice body-like synovitis in children's knees.

METHODS: We retrospectively analyzed the clinical and imaging data of 4 cases (5 knees) of rice body-like synovitis in children treated at our hospital from January 2014 to December 2021. Among the patients, 1 was male and 3 were female, with an average age of 22.5 months. The average length of symptoms before admission was 1.86 months. Clinical presentations included swelling and limping, with no fever or rash.

RESULTS: Three patients with unilateral onset and one patient with bilateral onset underwent unilateral arthroscopic excision of synovial lesions. All 4 patients had pathological findings consistent with chronic inflammation, leading to a postoperative diagnosis of juvenile idiopathic arthritis (JIA). Postoperatively, all patients received standardized pharmacological treatment and were followed for 24 to 48 months in collaboration with the rheumatology department. None of the 4 knees undergoing surgery experienced recurrence, and the symptoms of the knee treated conservatively resolved spontaneously. In the last follow-up session, MRI and ultrasound showed only mild synovial thickening, with effusion and loose bodies disappearing.

CONCLUSION: R [...]

[PubMed](#) · [DOI](#)

Extravertebral temperature exposure during spinal radiofrequency ablation: an experimental surrogate assessment of neural injury risk.

Weigert A, Hoffmann RT, Büttner A, Zaccaria R, Wegener V, Holzapfel BM, Wegener B

BACKGROUND: Radiofrequency ablation (RFA) has become an important minimally invasive option for the treatment of painful spinal tumors and benign bone lesions. While its effectiveness within the vertebral body is well established, there is ongoing concern about unintended heat exposure of nearby neural structures. Experimental data describing how heat propagates beyond the vertebra during spinal RFA are still limited. This experimental in vitro study aims to assess the extravertebral temperature exposure during spinal radiofrequency ablation as a surrogate parameter for potential neural injury risk under controlled conditions.

METHODS: This experimental in vitro study was performed on four fresh-frozen human lumbar spines. Radiofrequency ablation was carried out using a commercially available system and standard clinical protocols. Two anatomical configurations were examined: ablation within the vertebral body and ablation within the pedicle. Temperatures were measured at predefined locations outside the vertebra, including the spinal canal and cortical boundaries, using multiple thermocouples. To aid interpretation of radiofrequency-related measurement artifacts, an additional control setup using an electrothermal heating system without radiofrequency energy was included. Temperature data were evaluated descriptively.

RESULTS: In the control experiments, temperature profiles [...]

[PubMed](#) · [DOI](#)

Giant cell tumor of bone: exploring HtrA1 as a potential predictor of recurrence.

Mirioglu A, Dalkir KA, Gokmen MY, Bagir M, Ates KE, Deveci MA, Gönlüen G, Ozbarlas HS

INTRODUCTION: Giant cell tumor of bone (GCTB) is a locally aggressive tumor with a considerable risk of recurrence, but reliable predictive markers are limited. High-temperature requirement A serine peptidase 1 (HtrA1) has been implicated in the progression of several cancers and may also play a role in GCTB recurrence.

MATERIALS AND METHODS: This retrospective study included 34 patients with GCTB treated with curettage between 2002 and 2016. HtrA1 expression was evaluated separately in giant cells (GC) and mononuclear cells (MNC) using immunohistochemistry. Based on high or low expression in each cell type, patients were categorized into four expression patterns: High GC / High MNC, High GC / Low MNC, Low GC / High MNC, and Low GC / Low MNC. Clinical data, including recurrence status and time to recurrence, were analyzed.

RESULTS: The High GC / Low MNC group showed the highest recurrence rate at 71.4%, compared with 28.6% in the High GC / High MNC group and 31.6% in the Low GC / Low MNC group. No recurrence was observed in the Low GC / High MNC group, which included only one patient. The High GC / Low MNC group showed a non-significant trend toward higher recurrence compared with all other groups combined, with a similar trend toward shorter recurrence-free survival.

CONCLUSION: HtrA1 expression patterns in GCTB may provide preliminary insight into recurrence risk. Although [...]

[PubMed](#) · [DOI](#)

Coronal hamate body fractures with dorsal fourth and fifth carpometacarpal joint instability: fracture patterns, injury mechanisms, surgical strategies, and outcomes.

Lee SH, Chung HJ, Cha SM

INTRODUCTION: Coronal hamate body fractures are rare injuries, and studies regarding their morphology and management are limited. This study analyzed fracture patterns, injury mechanisms, associated injuries, surgical strategy, and clinical outcomes in patients with coronal hamate body fractures involving more than one-third of the articular surface and concomitant carpometacarpal (CMC) joint instability.

METHODS: Fifty-eight male patients were retrospectively reviewed. Fractures were classified as dorsal oblique or coronal splitting types based on computed tomography findings. Surgical treatment involved open reduction and

multiple screw fixation of the hamate using a K-wire-guided technique, with supplementary Kirschner wire stabilization for associated metacarpal base fractures or CMC joint instability. Radiographic and functional outcomes were evaluated after a minimum follow-up of 6 months.

RESULTS: Dorsal oblique fractures (63.8%) occurred more frequently than coronal splitting fractures (36.2%) and were most commonly associated with punching (OR, 4.9; $p = 0.01$), whereas falls from height were more common in coronal splitting types. Concomitant fractures occurred in 82.8% of cases, most frequently involving the fourth metacarpal base. Among 35 patients with a minimum 6-month follow-up, 34 achieved successful healing. In the 33 patients with functional outcome data, the [...]

[PubMed](#) · [DOI](#)

Effect of spinal fusion length on hip and knee osteoarthritis: a comparative cohort study of no fusion, short-segment fusion, and long-segment fusion.

Sülek Y, Demirkale ■, Erto■rul R, Özdemir HM

PURPOSE: Lumbar fusion alters spinal-pelvic biomechanics and may influence degenerative changes in the lower extremities. This study investigated whether fusion length is associated with radiographic progression of hip and knee osteoarthritis.

METHODS: In this retrospective, single-center cohort study, patients aged 40-65 years who underwent lumbar surgery between 2010 and 2021 were grouped as no fusion (N), short fusion (S; ≤ 3 levels), or long fusion (L; ≥ 4 levels). Hip joint space width was assessed using minimum joint space (MJS) and superointermediate joint space (SJS), including femoral head-standardized measures (sMJS and sSJS). Knee medial joint space width (KMJS) was measured when available. Spinopelvic parameters (LL, PI, PT, SS, PI-LL) and functional outcomes (ODI, HHS, WOMAC) were recorded. Multivariable linear and logistic regression analyses were performed to identify independent predictors of joint space narrowing and osteoarthritis risk.

RESULTS: Hip joint space narrowing differed significantly among groups, with the highest annual narrowing rates in the long-fusion group, followed by the short-fusion group, and the lowest in the no-fusion group ($p < 0.001$). In contrast, knee KL grade change and annual KMJS narrowing did not differ significantly between groups ($p > 0.05$). In logistic regression, each additional fused level increased the odds of postoperative MJ [...]

[PubMed](#) · [DOI](#)

Biomechanical comparison of inverted triangle and L-shaped screw configurations with medial buttress and anteromedial support plate in Pauwels type III femoral neck fractures.

Erem M, Demirta■ U, Y■ld■r■m S, Selçuk E, Çopuro■lu C

INTRODUCTION: Surgical stabilization of Pauwels Type III femoral neck fractures remains a significant challenge due to high vertical shear forces. While the medial buttress plate is a recognized solution, it requires extensive deep dissection. This study aims to compare the biomechanical performance of various screw configurations combined with either a medial buttress or anteromedial support plate.

MATERIALS AND METHODS: Twenty-five third-generation synthetic femurs were used to create a standardized 70-degree (Pauwels III) fracture model. Specimens were divided into five groups ($n = 5$): (A) Inverted triangle (IT) screws with a Pauwels screw, (B) Inverted triangle (IT) with medial buttress plate (MBP), (C) Inverted triangle with anteromedial support plate (ASP), (D) L-configuration with medial buttress plate (MBP), and (E) L-configuration with anteromedial support plate (ASP). Axial loading was applied at 2 mm/min until construct failure, defined objectively by the real-time force-distance curve.

RESULTS: Although no statistically significant difference was found between groups ($p = 0.102$), a large effect size was observed ($\eta^2 = 0.309$). Group C (IT + ASP) demonstrated the highest mean failure load (1695 ± 494.6 N). Conversely, Group E (L-configuration + ASP) exhibited the lowest stability (977.2 ± 195.4 N) with a remarkably narrow standard deviation. The majority of failures [...]

[PubMed](#) · [DOI](#)

Reverse total shoulder arthroplasty is a surgical option for patients over 60 years of age with locked shoulder dislocation.

Sebastia-Forcada E, Miralles-Muñoz FA, Martinez-Mendez D, Albero-Catala L, Climent-Peris V, Lizaur-Utrilla A

BACKGROUND: in locked dislocation of the shoulder, instead of reducing back to the glenoid, the humeral head remains incarcerated on the glenoid in a locked fashion. This clinical situation is fairly uncommon. It is essential to conduct an individual evaluation of each patient to determine the appropriate treatment.

OBJECTIVE: the aim of this study was to evaluate the functional outcomes of reverse total shoulder arthroplasty (rTSA) in the treatment of locked shoulder dislocation.

METHODS: patients with locked shoulder dislocation who underwent reverse shoulder prosthesis surgery and were admitted to our center between 2007 and 2023 were reviewed. The primary outcome was the Constant score. Secondary outcomes included the adjusted Constant, UCLA and DASH scores. Additionally, any signs of radiologic loosening were also documented.

RESULTS: the series consisted of 10 patients, six men and four women, with a mean age of 68.0 years. The average time from the traumatic injury to surgery was 7.5 months. All patients showed improved Constant, Adapted Constant, UCLA, and DASH scores compared to their preoperative values. When comparing the outcomes of chronic posterior and anterior dislocations, no differences in functional outcomes or shoulder motion were observed after rTSA implantation. There were no complications during or after surgery.

CONCLUSION: The results of the present [...]

[PubMed](#) · [DOI](#)

Does the order matter? Comparing the order of stem placement and fracture reduction in revision total hip arthroplasty for Vancouver B2 and B3 periprosthetic femur fractures.

Antonoli SS, Khury F, Kennedy MF, Haider MA, Duke A, Schwarzkopf R, Aggarwal VK

BACKGROUND: Vancouver B2 and B3 periprosthetic femur fractures (PPFF) have posed significant treatment challenges due to stem instability and lack of adequate femoral bone stock. This study investigated subsidence, survivorship, and outcomes of Vancouver B2 and B3 fractures, based on the order in which revision stem placement and fracture reduction occurred during revision total hip arthroplasty (rTHA).

METHODS: This retrospective, cohort study included 46 rTHAs between June 2011 and April 2023. Included patients underwent rTHA for Vancouver B2 or B3 PPFF with minimum one-year radiographic and two-year clinical follow-up. All patients were treated with diaphyseal-engaging tapered fluted titanium stems and stem modularity decisions were based on surgeon preference. Cohorts were separated based on if stem placement (SF, n = 19), or fracture reduction (RF, n = 27) occurred first. Patient demographics, intraoperative information, and clinical and radiographic outcomes were collected.

RESULTS: The SF and RF cohort showed no statistically significant differences in rate of subsidence ≥ 5 mm [26.3%[SF], 22.2%[RF], $P = 0.749$), rate of subsidence ≥ 10 mm (15.8%[SF], 14.8%[RF], $P = 0.928$), nor average subsidence (4.1 mm[SF], 4.4 mm[RF], $P = 0.861$). We found no statistically significant differences in surgery-related clinical outcomes or all-cause revision rates within a two-year follow- [...]

[PubMed](#) · [DOI](#)

Offset reconstruction in stemless and stemmed shoulder hemiarthroplasty: influence on long-term outcomes.

Trefzer R, Gurda A, Deisenhofer J, Weishorn J, Hariri M, Koch KA, Wolf M, Bühlhoff M

BACKGROUND: Shoulder hemiarthroplasty (HA) is a well-established treatment for various humeral-side pathologies. Overstuffing should be avoided because it is associated with glenoid wear; however, specific radiological reconstruction targets remain unclear. This study assessed the association between radiological joint restoration in stemmed and stemless HA and long-term functional outcomes.

METHODS: Patients who underwent stemmed or stemless HA between 2001 and 2011 were included in a long-term follow-up. Cases with incomplete or non-calibratable digital radiographs or concomitant glenoid procedures were excluded. Clinical outcomes were evaluated using the Constant-Murley Score (CMS) and satisfaction questionnaires. Radiographic parameters-lateral glenohumeral offset (LGHO), lateral humeral offset (LHO), and center of rotation (COR)-were measured on standardized anteroposterior radiographs pre- and postoperatively by two independent observers. Differences from baseline to final follow-up (Δ LHO, Δ LGHO, Δ COR) were calculated. Interobserver reliability was determined using the intraclass correlation coefficient (ICC). Associations between radiological measures and clinical outcomes were analyzed using linear regression;

intergroup comparisons used unpaired t-tests ($p < 0.05$).

RESULTS: Forty-eight patients (12 stemmed, 36 stemless HA) were examined after a mean follow-up of 16.6 [...]

[PubMed](#) · [DOI](#)

Emergency access to the subclavian vessels by non-thoracic surgeons: a cadaver-based learning model for orthopedic trauma surgery.

Grechenig P, Gänsslen A, Dauwe J, Wittig U, Sagmeister M, Koutp A, Puchwein P, Sadoghi P et al.

PURPOSE: The aim of this study was to evaluate the procedural time and safety of an infraclavicular approach to the subclavian vessels for damage control of major upper extremity vascular injuries. The study specifically focused on the performance of non-thoracic surgeons using a cadaveric model.

METHODS: A sample of 100 orthopedic trauma surgeons (residents, specialists, and attendings) was recruited from an AO cadaveric dissection course. Emergency infraclavicular access was performed on 50 cadavers over two consecutive days. The procedural time, successful vessel clamping, participant self-assessment, and the resulting learning curve were analyzed.

RESULTS: On day one, 27% (9/33) of participants who successfully achieved correct clamping of both subclavian vessels reported feeling confident. By day two, this proportion increased to 71% (55/77). Comparing day 1 to day 2 we found an improvement of 35% (subclavian artery) and 37% (subclavian vein) in the correct identification of subclavian vessels in our sample. The evaluations of this study show that there is no correlation between surgical experience and successful emergency access.

CONCLUSION: Anatomic dissection is of paramount importance for teaching rare and demanding surgical techniques. This study demonstrates that anatomical workshops significantly improve procedural safety and self-assessment when accessing subcla [...]

[PubMed](#) · [DOI](#)

Operative timing and surgical complexity in hand trauma: a multicenter analysis of replantation and non-replantation centers in Germany.

Nicklas A, Will Marks P, Dragu A, Schmidt H, Rotärmel A, HandTraumaRegister DGH

BACKGROUND: Hand injuries are common traumatic conditions that often require specialized surgical care. Differences in hospital structures and levels of specialization may influence the timeliness and complexity of treatment. This study evaluates quality of care across different hospital types, focusing on time-to-skin-incision and surgical management of finger and hand injuries.

MATERIALS AND METHODS: This retrospective multicenter study is based on data from the HandTraumaRegister of the German Society for Hand Surgery (HTR DGH). A total of 16,726 surgically treated cases with documented finger or hand injuries recorded between 2018 and 2023 were analyzed. Hospitals were categorized as non-replantation centers, replantation centers, or FESSH-accredited replantation centers. Demographic characteristics, injury-patterns, treatment modalities, and time-related parameters were assessed. Statistical analysis included descriptive statistics, group comparisons, and multivariate regression models to evaluate factors associated with time-to-skin-incision.

RESULTS: The study population consisted predominantly of male patients (75.79%), and 91.6% were right-handed. Soft tissue injuries without fractures or amputations accounted for 46.57% of cases. Non-replantation centers demonstrated the longest time-to-skin-incision (median 13 h), whereas replantation centers (median 3.75 h) and FE [...]

[PubMed](#) · [DOI](#)

Neglected acromioclavicular joint injury in coracoid process fractures: a retrospective cohort study.

Lee MS, Lee CH, Jo YH, Cho WT, Seo YW, Choi WS

INTRODUCTION: Coracoid process fractures are frequently associated with injuries to the superior suspensory shoulder complex, including the acromioclavicular (AC) joint. However, concomitant AC joint injuries may be overlooked in the acute phase, potentially limiting treatment options. This study aimed to investigate the

association between coracoid process fractures and SSSC injuries and to identify imaging-based predictors of concomitant AC joint injury.

METHODS: A retrospective single-center cohort study including 60 patients with coracoid process fractures was conducted between February 2016 and April 2023. Coracoid fractures were classified based on anatomical location, and associated SSSC injuries were categorized according to the involved structures. These injuries were evaluated using 3D shoulder CT. Morphological deformity of the SSSC was assessed by measuring the glenoclavicular distance (GCD)-a longitudinal parameter-on final follow-up clavicle AP radiographs. In base fractures, the medial displacement gap was measured on sagittal CT. Statistical analyses included logistic regression and ROC curve analysis to identify predictors of AC joint injury.

RESULTS: Among the 60 patients, fractures were located anteriorly in 27 cases, in the middle region in 5 cases, and at the base in 28 cases. AC joint injury was identified in 15 patients, all of whom had base-type fractu [...]

[PubMed](#) · [DOI](#)

Subclinical median nerve alterations after distal radius fractures: a comparative analysis of volar plate fixation and cast immobilization using dynamic ultrasonography and electrophysiology.

Karapinar SE, Arslan E, Oguzlar FC, Hatip AY, Akgun VO, Dincer R, Asci S, Kizilkaya V

BACKGROUND: Distal radius fractures may affect the median nerve through trauma-related and treatment-related mechanisms. While overt neurological deficits are uncommon, subclinical median nerve alterations may occur following both conservative and surgical treatment.

PURPOSE: To determine whether treatment modality influences subclinical median nerve behavior using combined dynamic ultrasonographic and electrophysiological assessment.

METHODS: This retrospective observational study included 42 patients with unilateral distal radius fractures. 18 patients were treated conservatively with circular casting, and 24 patients underwent volar plate fixation. Ultrasonographic assessment of the median nerve was performed at the distal radius level in neutral wrist position, as well as during wrist flexion and extension, using a dynamic ultrasonographic approach, and was combined with electrophysiological evaluation including sensory and motor nerve conduction studies. In the surgically treated group, only patients with Soong type 1 or type 2 plate positioning were included.

RESULTS: Clinical functional outcome scores were comparable between groups, despite a higher frequency of mild clinical signs such as positive Tinel test and night pain in the plate group. However, ultrasonographic evaluation demonstrated significantly greater position-dependent median nerve enlargement and flatte [...]

[PubMed](#) · [DOI](#)

Impulsivity and inattention in hand tendon injuries: a case-control study revealing distinct profiles for work-related accidents.

A■ta■ Ertan E, Demir Karak■/■ç G, Ertan MB

INTRODUCTION: The impact of attention deficit, impulsivity, and anger on various types of injuries is a subject of ongoing research. Hand tendon injuries are frequently encountered clinical conditions; however, their potential relationships with the aforementioned factors have not been previously investigated. This study aimed to examine the mechanisms of hand tendon injuries from certain psychiatric perspectives.

MATERIALS AND METHODS: 32 patients presenting to the physical therapy and rehabilitation outpatient clinic with hand tendon injuries and 32 healthy controls were evaluated. The assessment included sociodemographic data, causes of injury, Quick Disabilities of the Arm, Shoulder, and Hand (Q-DASH) and Visual Analog Scale (VAS) scores, as well as the physical and functional consequences of the injury. Patients and controls were assessed for attention deficit, impulsivity, and anger using the Adult ADHD Self-Report Scale (ASRS), Barratt Impulsivity Scale-11 (BIS-11), and State-Trait Anger Expression Inventory self-report scales. Statistical analyses of the results were performed.

RESULTS: Patients with tendon injuries were found to have significantly higher scores in ASRS attention and BIS-11 Motor impulsivity compared to the control group ($p = 0.048$, $p = 0.040$). Individuals with hand tendon injuries resulting from work accidents demonstrated significantly lower ASRS to [...]

Comparative analysis of the improvement in shoulder rotation in complex shoulder fractures treated by reverse shoulder arthroplasty compared to open reduction and internal fixation: a systematic review and meta-analysis.

Albadran A, Alsugair R, AlHarthi A, Alshangiti H, Alqahtani H, Alsugair G, Ababtain R, Alhajress D et al.

PURPOSE: To compare the clinical and functional outcomes of open reduction and internal fixation (ORIF) and reverse shoulder arthroplasty (RSA) in patients aged ≥ 65 years with complex proximal humeral fractures (PHFs).

METHODS: A systematic review and meta-analysis was conducted according to the PRISMA guidelines. PubMed, Web of Science, ScienceDirect, EBSCO, and the Cochrane Library were searched for randomized controlled trials (RCTs) and cohort studies published in English without date restrictions. Eligible studies compared RSA and ORIF in elderly patients with PHFs and reported functional, radiographic, or complication outcomes. Pooled data were analyzed using a random-effects model.

RESULTS: Twenty-two studies involving > 34,000 patients were included. RSA was associated with greater forward flexion and a trend toward improved abduction, whereas internal rotation favored ORIF without reaching significance. No significant differences were observed in external rotation. The functional scores (Constant-Murley, Oxford Shoulder Score) were similar, and the complication and reoperation rates did not differ significantly between the groups.

CONCLUSION: RSA offers modest advantages in forward flexion compared with ORIF but does not consistently improve overall functional scores or reduce complications. RSA may be preferred in elderly patients with poor bone stock or rotator c [...]

Changes in kinematic behavior and collateral ligament strain after medially stabilized TKA using a novel intraoperative navigation platform: a cadaveric study.

Van de Vyver A, Taylan O, Peersman G, Demeulenaere M, Bialy M, Lauwagie T, Scheys L

OBJECTIVES: This study aimed to analyze to what extent a novel intraoperative navigation platform (Next-AR, Medacta) for total knee arthroplasty (TKA) allows to restore the native knee joint kinematics and strains in the medial collateral ligament (MCL) and lateral collateral ligament (LCL) throughout a squatting motion.

MATERIALS AND METHODS: Computed tomography (CT) scans of 6 native cadaver legs were used to design patient-specific guides. Bony landmarks and virtual single-line collateral ligaments were identified to acquire real-time intraoperative feedback on bone resection, implant alignment, tibiofemoral kinematics, and collateral ligament elongations using the Next-AR system. The specimens were subjected to squatting (35°-100°) motion using a physiological ex vivo knee simulator while maintaining a constant vertical ankle load of 110 N through active quadriceps and bilateral hamstring controls. Subsequently, each knee underwent a medially-stabilized TKA (Medacta) with mechanical alignment technique and was retested under the same conditions as in their native state. The tibiofemoral and patellofemoral kinematics, along with collateral ligament strains, were computed from 3D marker trajectories using a six-camera optical system (Vicon). MCL and LCL insertions-ant, mid, and post bundles-were identified in relation to bone-pin markers using a wand.

RESULTS: Both native a [...]

Regional differences in the management of patients with mild traumatic brain injury and antithrombotic therapy-an Austrian survey.

Sebek B, Dammerer D, Putzer D, Moser L

BACKGROUND/PURPOSE: The number of patients with mild traumatic brain injury (TBI) receiving oral anticoagulation is increasing. Up-to-date diagnostic and treatment algorithms are missing in Austria. The aim of this survey is to collect and analyse Austria-wide data on the care of this patient group.

METHODS: The survey was carried out using "Jotform" between 27 November 2023 and 31 January 2024, asking questions relating to the diagnosis and treatment of patients with mild TBI receiving antithrombotic therapy. Patient

groups comprising individuals with alcohol addiction and hemophilia were incorporated as well with a view to the paucity of available literature on these groups. The survey was sent to all orthopedic-traumatological and neurosurgical departments in Austria.

RESULTS: 39 of 67 (58.2%) orthopedics & traumatology departments and 4 of 10 (40%) neurosurgery departments participated in the survey. The share of departments reporting routine submission of different patient groups to cranial computed tomography (CCT) was 38 (95%) for patients taking antiplatelet agents, 39 (97.5%) departments for patients taking vitamin K antagonists (VKAs), 39 (97.5%) for patients taking direct oral anticoagulants (DOACs), 11 (27.5%) for patients with alcohol addiction, and 28 (70%) for patients with known hemophilia. Considering the separate groups of patients not living in a nursing fa [...]

[PubMed](#) · [DOI](#)

Extracorporeal shockwave therapy in bone fractures: a systematic review.

Santos TIR, Souza E Silva LC, Bonifacio M, Campos Melo CM, Sena BG, Simões MC, Risso Parisi J, da Silva Fernandes AC et al.

INTRODUCTION: The increasing incidence of fractures and cases of delayed union or nonunion has encouraged the search for non-invasive therapies. Extracorporeal shockwave therapy (ESWT) has been proposed as a strategy to stimulate bone regeneration and improve fracture healing.

MATERIALS AND METHODS: A systematic review was conducted following PRISMA guidelines, with searches performed in PubMed, Scopus, Web of Science, and Embase (2005-2025). Clinical studies evaluating ESWT for fracture treatment with radiographic assessment of bone healing and the presence of a control group were included.

RESULTS: A total of 834 studies were initially identified, of which 7 were included in the final analysis. Most studies demonstrated positive effects of ESWT, including improved bone healing, pain reduction, and functional recovery. However, there was considerable heterogeneity in treatment protocols and relatively small sample sizes.

CONCLUSION: ESWT shows potential as a non-invasive therapy to stimulate fracture healing, particularly in cases of delayed union or nonunion. However, further controlled clinical trials and standardized protocols are needed to confirm its clinical effectiveness.

[PubMed](#) · [DOI](#)

Clinical outcomes and complication rates of minimally invasive plate osteosynthesis (MIPO) versus open reduction and internal fixation (ORIF) in midshaft clavicular fractures: a comparative study.

Lin G, Zheng S, Zhang Y, Liu Q, Liu H

BACKGROUND: Conventional open reduction and internal fixation (ORIF) remains the standard approach for midshaft clavicle fractures, yet its traumatic nature has driven the need for minimally invasive techniques. Locking minimally invasive plate osteosynthesis (MIPO) demonstrates clinical potential through biomechanical superiority and tissue-preserving characteristics.

OBJECTIVE: To compare the clinical efficacy and safety of MIPO versus ORIF for midshaft clavicular fractures.

METHODS: A retrospective cohort study included 203 patients (MIPO group:101 cases; ORIF group:102 cases) treated between 2022 and 2024. Baseline characteristics, functional outcomes (DASH and Constant-Murley scores), and complications were analyzed using Mann-Whitney U and chi-square tests.

RESULTS: No baseline differences were observed ($P > 0.05$). Functional outcomes were equivalent between groups (DASH score: $P = 0.906$; Constant-murley score: $P = 0.646$). The overall complication rate was significantly lower in the MIPO group than in the ORIF group (7.92% vs. 36.27%, $P < 0.001$), particularly in nonunion (0% vs. 5.88%, $P = 0.039$) and sensory disturbance (6.93% vs. 27.45%, $P < 0.001$). This advantage is also evident in the subgroup of patients with AO type 2 C fractures, where MIPO similarly demonstrates a significantly lower incidence of these two complications.

CONCLUSION: MIPO achieves comparable fun [...]

[PubMed](#) · [DOI](#)

Isolated scaphoidectomy for type II SLAC and SNAC wrists: retrospective case-series at long-term follow-up.

Altissimi M, Pataia E, Saracco M, Braghiroli L

BACKGROUNDS: Stage II scapholunate advanced collapse (SLAC) and Scaphoid non-union advanced collapse (SNAC) wrists are currently treated either by scaphoidectomy and Four Corner Fusion or by Proximal Row Carpectomy. In our retrospective study, isolated scaphoidectomy demonstrated clinical non-inferiority when compared to standard procedures and could be supplemented by a Four Corner Fusion only at a later time if needed.

METHODS: Between 2006 and 2011, 13 patients / 14 wrists (mostly males, manual workers) underwent the surgical procedure. 8 patients were affected by SNAC and 5 by SLAC. The average age was 56,5. All patients were manual workers. The mean follow-up was 15 years. These patients were reviewed a first time in 2012 and once again in 2024. Subjective outcome, DASH score, return to work, ROM and grip strength were evaluated. Radiographic assessment was done in all 14 wrists at the first follow-up and in 6 at the last long-time review. Statistical analysis was conducted using the SPSS software.

RESULTS: 12 patients had resumed the same work at a mean of 3 months after surgery. Only one patient required a Four Corner Fusion after 2 years. 11 of the 13 enrolled patients declared their satisfaction for the procedure at the longest follow-ups. Clinical results remained stable over time. Radiological deterioration was seen in 6/13 wrists at the first follow-up and in all [...]

[PubMed](#) · [DOI](#)

Loop versus anchor tenodesis of the long head of the biceps tendon: a prospective randomized controlled pilot study.

Bischofreiter M, Kölblinger C, Gattringer M, Azesberger A, Gruber MS, Rittenschober F, Kindermann H, Ortmaier R

BACKGROUND: The long head of the biceps tendon is a frequent source of anterior shoulder pain, often treated with tenodesis. While anchor tenodesis is well established, it carries risks of implant-related complications. This study evaluates an implant-free "loop tenodesis" technique as a potentially safer and cost-effective alternative.

METHODS: In this prospective, double-blinded, randomized exploratory study, 48 patients undergoing arthroscopic biceps tenodesis were randomized to either loop (n = 24) or anchor (n = 24) tenodesis. Clinical outcomes were assessed using ASES, Constant, and DASH scores, as well as range of motion, supination strength, satisfaction, aesthetic appearance, and return-to-sport. Statistical analysis was performed using the Mann-Whitney U and Wilcoxon signed-rank tests to compare non-parametric data.

RESULTS: Both groups showed significant improvements with similar gains in functional scores. Supination strength recovery was greater in the loop group (103% vs. 93%). There were no significant differences in range of motion or complications. Notably, only the anchor group showed cases of Popeye deformity (2 patients). Return to sport was achieved in 95.3% of patients, typically within 3 months. No revision or implant-related issues occurred.

CONCLUSION: Loop tenodesis offers equivalent clinical outcomes to anchor-based techniques, with fewer aesthetic [...]

[PubMed](#) · [DOI](#)

Split hamate fractures as part of complex ulnar carpometacarpal injuries: radiographic and functional outcomes.

Glaser J, Schmidt E, Aman M, Bickert B, Harhaus-Wähner L, Cordts T

BACKGROUND: Split hamate fractures, which most often are a complicating part of fracture dislocations of the 5th or 5th and 4th carpometacarpal joints, are rare and challenging injuries that can severely affect hand function. The injury usually requires surgical treatment. However, both short- and long-term data on the effectiveness of these interventions are limited. The objective of this study is to advance understanding of this rare hand surgical condition by investigating its epidemiology, etiopathogenesis, subsequent complications, and functional outcomes.

METHODS: Between 2010 and 2024, a total of 29 patients with a split hamate fracture were treated at a level-1 trauma center, of which 12 patients are included in this study. Using patient surveys (DASH, SF-36, PRWE), along with clinical and radiological examinations, the hand function of a total of 12 patients with split hamate fractures is

evaluated.

RESULTS: Twelve male patients with a split hamate fracture were included, with a mean age of 28.1 years at the time of surgery. In 10 patients (83%) the dominant right hand was affected, most involving, the injuries resulted from direct trauma or a fall; surgery was performed on average of 12 days after injury. Functional outcomes were favourable, with a mean DASH score of 14.1 and PRWE score of 14.8, indicating minimal residual reduction of function in most cases. Radiol [...]

[PubMed](#) · [DOI](#)

Outcomes and short-term survivorship of fixed-bearing all-poly unicompartmental knee arthroplasty. A retrospective single-center study of 42 prostheses with a main follow-up of 32.3 months.

Gaggiotti S, Gaggiotti G, Lustig S, Batailler C, Monllau JC, Gaggiotti S, Combalia A

PURPOSE: All Poly (AP) tibial unicompartmental knee arthroplasty (UKA) seems to have advantages in comparison with metal-based (MB) UKA, such as greater polyethylene thickness, wear resistance, bone stock preservation and lower implant costs. The objective of this retrospective study was to evaluate the short-term clinical-functional and radiographic results, complications and survivorship of AP UKA performed consecutively with a minimum follow-up of 2 years.

METHODS: Retrospective cohort analysis all consecutive knees operated with medial and lateral AP UKAs, including bilateral cases, with a minimum follow-up of 2 years. Demographic data were analyzed at the patient level and clinical-functional results were assessed per knee using the KSS, KOOS and VAS scales. Radiographically, mechanical axis (HKA) correction and the presence of radiolucency > 2mm were evaluated. Complications incidence and prosthetic survivorship was determined. Pre-postoperative comparisons were performed using linear mixed-effects models with random intercept per patient to account for the non-independence of bilateral cases. Subgroup analyses involving fewer than 10 observations were considered exploratory and no formal statistical inference is drawn. A value of $p < 0.05$ was considered statistically significant with a confidence interval (CI) 95%.

RESULTS: 42 AP UKAs were included, 5 lateral (11.9%) a [...]

[PubMed](#) · [DOI](#)

Mix and match in reverse total shoulder arthroplasty: overview of current prostheses systems and potential combinations.

Seiß D, Schlichter A, Graf JM, Stange R, Ohlmeier M

INTRODUCTION: Mix and Match is considered a viable option for minimizing morbidity in total hip arthroplasty (THA). However in reverse total shoulder arthroplasty (rTSA) this strategy remains comparatively new. This study provides an overview of currently available implant systems and the potential combinations of glenoid and humeral components.

MATERIALS AND METHODS: A review of commercially available shoulder arthroplasty systems was performed. Nineteen manufacturers were contacted and asked to provide detailed information on glenoid and humeral component sizes for their rTSA systems. The reported component sizes were compiled and analyzed for possible combinations.

RESULTS: Thirteen manufacturers (68%) consented to the publication of their component size data. Two manufacturers (11%) do not produce rTSA systems, and four (21%) did not respond to repeated inquiries. A total of 21 implant systems were reported, comprising 10 distinct implant sizes. Glenosphere and humeral inlay diameters ranged from 32 mm to 48 mm, with 36 mm being the most commonly available size (16 of 21 systems, 76%).

CONCLUSION: Mix-and-match implantation in rTSA is likely employed more frequently than documented, yet remains poorly investigated. This study demonstrates that numerous systems can theoretically be combined during revision surgery, although the biomechanical implications are not yet under [...]

[PubMed](#) · [DOI](#)

Radiolucent lines after robotic-assisted surface-matching kinematically aligned UKA: short-term incidence and lack of association with tibial alignment.

Itou J, Kuwasawa A, Nihei K, Okazaki K

PURPOSE: This study investigated the clinical outcomes of robotic-assisted unicompartmental knee arthroplasty (UKA) performed using a surface-matching kinematic technique that restores native joint morphology without alignment constraints. The primary aim was to determine the incidence and distribution of radiolucent lines (RLLs), and the secondary aim was to evaluate short-term revision rates and clinical outcomes.

METHODS: This retrospective cohort study reviewed 242 medial UKAs performed between November 2021 and August 2023 by the same high-volume surgeon using a surface-matching technique. Radiographic parameters and patient-reported outcome measures (PROMs) obtained before and 1 year after surgery were assessed, with radiographic evaluation performed independently of the clinical outcome assessment. A cemented JOURNEY II Uni implant was used in all cases, with intraoperative surface mapping used to guide kinematic placement of components without alignment constraints.

RESULTS: RLLs were seen in 21.9% of knees, exclusively on the tibial side and mainly in the most medial (zone 1: 14.8%, 95% confidence interval [CI] 10.9-19.9) and posterior regions (zone 9: 15.3%, 95% CI 11.3-20.3). A larger tibial component was the only independent predictor of RLLs (odds ratio 1.54, 95% CI 1.27-1.88; $p = 0.04$). All PROMs improved, with no differences between knees with and without RLLs. [...]

[PubMed](#) · [DOI](#)

Radiographic parameters and mechanical complications of articulating vs. non-articulating hip spacers.

Lenz J, Baumann F, Baertl S, Straub J, Rupp M, Alt V, Freigang V

BACKGROUND: Periprosthetic joint infection (PJI) is a serious complication of total hip arthroplasty. Standard treatment for PJI is a two-stage revision surgery with antibiotic-loaded cement spacers to control infection and provide temporary joint stability. This study compares the radiological outcomes and complication rates between articulating and non-articulating hip spacers in patients undergoing treatment for PJI and native joint infections.

METHODS: We retrospectively reviewed 71 hip spacers (34 articulating, 37 non-articulating) in 38 patients treated between 2004 and 2022. Data on leg length discrepancy, femoral offset, infection control, and mechanical complications were obtained. For infection analysis, only patients treated exclusively with one spacer type were included. After excluding eight patients with mixed spacer types (29 spacers), 30 patients with 42 spacers were included in this subgroup.

RESULTS: Articulating spacers were significantly better at preserving leg length, with a mean discrepancy of -3.7 mm compared to -16.9 mm for non-articulating spacers ($p = 0.025$). However, non-articulating spacers maintained femoral offset (1.1 vs. 0.7, $p < 0.001$) closer to physiological offset. The rate of mechanical complications was higher in the articulating spacer group, with spacer dislocations occurring in 45% of cases compared to 10% in the non-articulating group [...]

[PubMed](#) · [DOI](#)

The effect of Poller screw or monocortical plate augmentation on the stability of intramedullary nailing in proximal extra-articular tibial fractures: a biomechanical study.

■ahin MA, Durgut F, Yi■it ■, Akar MS, Özkul E, Atıç R, Çiftçi S

BACKGROUND: Proximal extra-articular tibial fractures present considerable biomechanical and technical challenges due to the short proximal fragment, the wide metaphyseal canal, and the deforming forces acting around the knee. These factors predispose isolated intramedullary (IM) nailing to malalignment, prompting the use of adjunct reduction aids such as Poller screws or supplemental plates. This study aimed to compare the mechanical behavior of IM nailing alone with that augmented by Poller screws or a monocortical plate in a standardized proximal tibia fracture model.

METHODS: Fifteen synthetic tibiae with an AO/OTA 41-A2.3 extra-articular proximal fracture model were randomized into three groups ($n = 5$ each): IM nailing alone, IM nailing plus two Poller screws, and IM nailing plus a four-hole medial monocortical plate. All specimens were instrumented with a 10 × 340-mm titanium nail and two

proximal and two distal locking screws. Mechanical testing included axial compression to 800 N, axial loading to failure, and torsion at a rate of 25°/min. Displacement, stiffness, maximum load, and torque parameters were compared across groups.

RESULTS: Under 800 N axial load, the plate-augmented group showed the lowest displacement and highest stiffness (2.40 ± 0.22 mm; 332.4 ± 21.7 N/mm), followed by the Poller screw group (2.48 ± 0.25 mm; 321.6 ± 19.4 N/mm) and the IM-only group (3 [...])

[PubMed](#) · [DOI](#)

Clinical cutoff value of lumbar bending range for predicting residual pelvic obliquity after total hip arthroplasty in dysplastic hip osteoarthritis.

Yokoi H, Osawa Y, Ozawa Y, Funahashi H, Takegami Y, Imagama S

INTRODUCTION: Residual pelvic obliquity (PO) after total hip arthroplasty (THA) for dysplastic hip osteoarthritis (DHOA) may adversely affect coronal balance and postoperative outcomes. Although lumbar bending range (LBR) is considered critical for PO improvement, the clinical cutoff value for predicting residual PO remains unclear. This study aimed to determine the cutoff value of LBR associated with residual PO after THA.

MATERIALS AND METHODS: Among 382 patients who underwent primary THA for unilateral DHOA between July 2019 and June 2024, 98 patients with preoperative downward PO of $\geq 2^\circ$ on the affected side were included. Patients were classified into a residual group (postoperative PO $\geq 2^\circ$ at 1 year) and an improvement group (postoperative PO $< 2^\circ$). Demographic data and radiographic parameters of the hip, lower limbs, and spine were compared. Multivariate logistic regression analysis was performed to identify factors associated with residual PO, and receiver operating characteristic (ROC) analysis was used to determine the optimal LBR cutoff value.

RESULTS: Residual PO was observed in 28 patients (29%). Compared with the improvement group, the residual group demonstrated significantly greater Crowe index values, preoperative radiographic leg length discrepancy, and preoperative PO. Affected-side LBR was significantly smaller in the residual group ($4.3^\circ \pm 3.0^\circ$ vs. $7.7^\circ \pm [...]$)

[PubMed](#) · [DOI](#)

Hidden blood loss after proximal femur fractures fixation: analysis of the first three postoperative days and influence of anticoagulants.

Mayor J, Winkelmann M, Schwinn LS, Özbek H, Aktas G, Omar-Pacha T, Oberthür S, Krammig RH et al.

PURPOSE: Hidden blood loss (HBL) is an underrecognized contributor to total perioperative blood loss after hip fracture surgery. This study aimed to quantify HBL during the first three postoperative days following proximal femur fracture fixation and to assess the impact of preoperative anticoagulant therapy on blood loss and transfusion requirements.

METHODS: A retrospective cohort study was conducted including patients aged ≥ 65 years who underwent surgical fixation of proximal femur fractures at a level I trauma center between January 2020 and December 2022. Total blood loss was estimated from perioperative hemoglobin changes and calculated blood volume. Patients were stratified according to anticoagulant use, and logistic regression analysis was performed to identify independent predictors of increased blood loss.

RESULTS: Among 260 patients analyzed, median total blood loss was 1184 ml, and 25% experienced losses exceeding 1675 ml. High blood loss correlated significantly with anticoagulant use, higher BMI, and longer operative time ($p < 0.001$). In multivariate analysis, anticoagulant therapy (particularly direct oral anticoagulants) remained an independent predictor of high blood loss (OR 3.06; 95% CI 1.47-6.37; $p = 0.003$). Patients in the highest quartile required significantly more transfusions, though short-term mortality did not differ.

CONCLUSION: Hidden blood loss [...]

[PubMed](#) · [DOI](#)

Visual processing and interference performance influences on knee angular impulse in ACLR individuals: a cognitive-biomechanical analysis of drop-jumps.

Santamaria-Guzman K, Holmes H, Kosek J, Peoples B, Harrison K, Campos-Vargas S, Weimar W, Neely K et al.

INTRODUCTION: Visual processing speed and cognitive interference control are critical for athletic performance and Anterior Cruciate Ligament (ACL) injury risk. However, how these cognitive functions relate to biomechanical performance after Anterior Cruciate Ligament Reconstruction (ACLR) remains unclear. This gap is clinically relevant given persistent re-injury rates of 20-25% in young athletes returning to sport, suggesting that rehabilitation may insufficiently address cognitive demands. This study examined cognitive differences between ACLR individuals and controls during drop-jump tasks, and their relationship with knee angular impulse.

MATERIALS AND METHODS: Thirty-two females (16 ACLR, 16 controls; 20 ± 1 years) completed cognitive assessments including the Stroop Color and Word Test, Trail Making Test, Digit Span Memory Test, and visual/auditory reaction time tests. Participants performed drop-jumps under four conditions: standard, choice, visual-cued, and audio-cued. Knee angular impulse was calculated for eccentric, concentric, and net phases. Binomial logistic regression identified cognitive predictors distinguishing groups, followed by factorial ANOVA to assess biomechanical differences. Spearman correlations examined relationships between cognition and knee angular impulse.

RESULTS: Three cognitive variables distinguished groups: cognitive interference score, v [...]

[PubMed](#) · [DOI](#)

Perceived leg length discrepancy after total hip arthroplasty is associated with global spinopelvic coronal flexibility.

Minoji S, Ishikura H, Kudo M, Hatano M, Nishiwaki T, Tanaka S

BACKGROUND: Perceived leg length discrepancy (pLLD) is a common source of dissatisfaction after total hip arthroplasty (THA), even in the absence of significant radiographic discrepancies. While spinopelvic factors have been increasingly recognized, most previous studies have focused on sagittal alignment or lumbar mobility, and the impact of preoperative global spinopelvic coronal flexibility on pLLD has remained unclear.

METHODS: We retrospectively reviewed 114 patients who underwent primary unilateral THA for osteoarthritis between January and December 2023. pLLD was assessed using a four-point scale at 6 months postoperatively. Patients who reported no perception of discrepancy were classified as the non-pLLD group, while those reporting mild, clear, or strong perception were grouped as the pLLD cohort. Preoperative spinal flexibility was measured on coronal radiographs during maximal lateral bending, calculating changes in spinopelvic angle (Δ SPA), reflecting thoracic-to-pelvic coronal flexibility, and lumbosacral angle (Δ LSA), reflecting lumbar-to-pelvic coronal flexibility. Secondary parameters included radiographic leg length discrepancy, leg lengthening, and sagittal spinopelvic alignment. Multivariable logistic regression analysis was used to identify independent predictors of pLLD.

RESULTS: pLLD was reported in 47 patients (41.2%). The pLLD group exhibited signific [...]

[PubMed](#) · [DOI](#)

Sex-related differences in final knee balance during robotic-assisted unicompartmental knee arthroplasty.

Stimolo D, Leggieri F, Secci G, Massenzi M, Civinini R, Innocenti M

INTRODUCTION: In robotic medial unicompartmental knee arthroplasty (mUKA), a preoperative alignment strategy can be defined and subsequently adjusted through stress testing to achieve optimal soft-tissue balance. However, the application of uniform planning principles does not necessarily yield equivalent balance in all patients. The aim of this study was to examine differences in intraoperative balance according to patient sex and the degree of preoperative varus deformity.

MATERIALS AND METHODS: A single-center retrospective cohort study including 254 MAKO® robotic-assisted medial UKAs (2018-2025) was performed using a uniform intraoperative planning protocol. Gap widths were measured in millimeters at full extension and 90° of flexion (F90) and classified as tight (< 0.5 mm and negative values), balanced (0.5-2 mm), or loose (> 2 mm). Patients were stratified by sex and preoperative varus severity, and gaps were analyzed quantitatively and categorically. The independent effects of sex, varus deformity, and operating surgeon on extension and flexion gaps has been assessed.

RESULTS: Mean preoperative alignment was $174.2 \pm 2.8^\circ$, with no sex-related difference in baseline varus ($p = 0.058$). Flexion gaps were significantly tighter in female patients ($p = 0.0045$). A tight F90 gap was observed in 17.2% more female than male patients ($p = 0.023$), with an absolute mean difference o [...]

A novel pinless augmented reality-based navigation system for total hip arthroplasty in the supine position: a comparative study of acetabular cup angle accuracy.

Kurishima H, Ogawa H, Yamada N, Noro A, Tomata Y, Tsukada S

INTRODUCTION: We developed a novel pinless augmented reality (AR)-based portable navigation system for total hip arthroplasty (THA) in the supine position to eliminate pin-site complications and intraoperative navigation abandonment, and to reduce registration time. This study compared the measurement accuracy of this new pinless portable navigation system with that of a conventional AR-based portable navigation system requiring pin insertion.

MATERIALS AND METHODS: This retrospective cohort study compared primary unilateral THAs performed using the novel pinless AR-based navigation system and a conventional AR-based navigation system. The primary outcome was the absolute difference between the acetabular cup angles displayed on the navigation screen and those measured on postoperative computed tomography. Secondary outcomes included target achievement rates within 5° and 10°. Pin-site complications, intraoperative navigation abandonment, and dislocations were also recorded.

RESULTS: A total of 228 hips were analyzed (Pinless: n = 115; Conventional: n = 113). While there was no significant difference in the median absolute difference for inclination (2.5° vs 2.4°; p = 0.366), the median absolute difference for anteversion was significantly larger in the pinless group than in the conventional group (2.9° vs 1.9°; p = 0.011). There were no significant differences in the percent [...]

PubMed · DOI

Impact of modified Henry approach with preservation of Flexor Carpi Radialis Tendon Sheath on wrist function in the treatment of distal radius fractures.

Ma D, Ji J, Zhu Y, Fang X, Wang Y, Fan J

OBJECTIVE: To compare the clinical efficacy of a modified Henry approach (preserving the FCR tendon sheath) versus the traditional approach (incising the sheath) for distal radius fractures (DRFs) treated with volar locking plates, and to evaluate the outcomes within 12 months postoperatively.

METHODS: A retrospective cohort study was conducted on 165 patients with unstable DRFs who underwent volar locking plate fixation from January 2023 to March 2025, who were retrospectively divided into the Preservation Group (n = 79, FCR tendon sheath preserved via the modified approach) and the Control Group (n = 86, FCR tendon sheath incised via the traditional approach). Perioperative indicators, radiological parameters, Visual Analog Scale (VAS) scores, Disabilities of the Arm, Shoulder and Hand (DASH) scores, grip strength, wrist range of motion (ROM) and complications were compared at the follow-up time points of 1 week, 1, 2, 3, 6 and 12 months postoperatively.

RESULTS: There were no significant differences in baseline data and postoperative radiological reduction quality between the two groups (P > 0.05). The Preservation Group had significantly lower VAS scores at all follow-up time points (P < 0.001), and lower DASH scores from 1 week to 6 months (P < 0.001), with no statistical difference at 12 months (P = 0.550). Grip strength of the Preservation Group was significantly super [...]

PubMed · DOI

Clinical Outcomes, Return to Sport and Psychological Readiness After Arthroscopic Bankart Repair Using Knotless All-Suture Anchors at a Minimum 2Year Follow-up.

Rab P, Rupp MC, Pfarrmaier A, Vieider RP, Shirinskiy IJ, Scheiderer B, Siebenlist S, Lacheta L

PURPOSE: To report the clinical outcomes, return to sport (RTS) and psychological readiness of patients who underwent arthroscopic Bankart repair with knotless all-suture anchors with a minimum follow-up of 2 years.

METHODS: In this retrospective case series, consecutive patients who underwent primary arthroscopic Bankart repair using knotless all-suture anchors between 08/2019 and 07/2022 were included. Patient-reported outcomes were assessed using the Western Ontario Shoulder Instability Index (WOSI), American Shoulder and Elbow Surgeons (ASES) score, Disabilities of the Arm, Shoulder and Hand questionnaire (DASH), Shoulder Instability-Return to Sport after Injury (SI-RSI) scale, subjective shoulder value (SSV), and the visual analogue scale (VAS) for pain. Patient satisfaction, RTS, return to preinjury level of sport, instability recurrence and revisions

were recorded. Receiver operating characteristic (ROC) curve was calculated to assess the discriminative performance of the SI-RSI scale, and the Youden's index was employed to determine the optimal cutoff for prediction of return to preoperative level of sports.

RESULTS: Of 57 patients eligible for inclusion, 46 patients (11.1% female, 28.7 ± 6.8 years at surgery) were available at a follow-up of 2.9 [2.3-3.4] years. Three patients (6.5%) reported a redislocation, one patient (2.2%) underwent a revision and was excluded f [...]

[PubMed](#) · [DOI](#)

Impact of frailty on 12-month patient-reported outcomes after posterior lumbar fusion surgery.

Haffer H, Muellner M, Chiapparelli E, Dodo Y, Camino-Willhuber G, Zhu J, Pumberger M, Shue J et al.

INTRODUCTION: Frailty might increase the risk for postoperative complications and death. It is not well understood how frailty influences patient-reported outcomes (PROs) in spine surgery. The objective is to investigate on the impact of frailty on PROs 1-year after lumbar fusion.

MATERIALS AND METHODS: This was a prospective observational study of patients undergoing open posterior lumbar fusion for degenerative conditions with 12-month follow-up (12 M-FU). The modified 5-item frailty index (mFI-5) score was determined preoperatively. PROs included the Oswestry-Disability-Index (ODI), 12-item Short-Form-Healthy-Survey with Physical (PCS-12) and Mental-Component-Score (MCS-12) and Numerical-Rating-Scale (back/leg pain) before surgery and 6 weeks, 3, 6 and 12 months (6 W, 3 M, 6 M, 12 M) postoperatively. PROs were compared between groups stratified as frail (mFI-5 ≥ 2), pre-frail (mFI-5 = 1) and non-frail (mFI-5 = 0). PRO changes at 12 M-FU were evaluated with substantial clinical benefit (SCB) metrics. Regression models were used to determine associations between mFI-5 and PROs.

RESULTS: 135 patients (52.6% female, 62.1 years, BMI 29.1 kg/m²) were analyzed. 31.3% and 51.8% of the frail and non-frail patients reached ODI-SCB, respectively. Frailty patients reached a significantly smaller ODI change at 12 M-FU than pre- and non-frail patients (15.2% vs. 19.3% and 19.0%, $p < 0.0$ [...])

[PubMed](#) · [DOI](#)

Rates of union and risk factors for continued nonunion following exchange nailing of tibial nonunion.

Mastracci JC, Averkamp B, Braswell M, Yu Z, Chen AT, Natoli RM, Farooq H, Mir H et al.

INTRODUCTION: The objective of this study was to evaluate the rate of nonunion repair success in tibia nonunions treated with exchange nailing. This retrospective cohort study was conducted across five academic Level 1 trauma centers and included 63 patients with tibia nonunions.

MATERIALS AND METHODS: Patients who sustained a tibia fracture (AO/OTA 42) treated with intramedullary fixation that developed a nonunion and were subsequently treated with exchange nailing were retrospectively reviewed. The primary outcome measure was nonunion repair success based on osseous union. Additional analyses included union rate by AO/OTA classification, nonunion type, implant(s) used, graft used, time from initial procedure, and infection status.

RESULTS: All patients sustained an AO/OTA type 42 fracture. Out of 63 patients, 47 tibias (75%) achieved osseous union after the index exchange nail procedure. The rates of nonunion revision success were similar across nonunion types and time from initial procedure until exchange nailing. There was no significant difference in union rate when infection was present. Complications included re-operation, readmission, infection, and implant failure.

CONCLUSIONS: This large, multicenter study with contemporary implants, instruments, and techniques for exchange nailing tibia nonunions demonstrates a nonunion repair success rate of 75%, consistent with [...]

[PubMed](#) · [DOI](#)

Robotic-assisted unicompartmental knee arthroplasty is associated with lower odds of prolonged hospitalization and no higher odds of high-charge admission during the index hospitalization.

Sun Q, Xu Y, Hong X, Dai M, Wang J, Xie H, Fu Z

BACKGROUND: Robotic-assisted unicompartmental knee arthroplasty (RA-UKA) may improve implant positioning, but its impact on index-hospitalization length of stay (LOS), billed charges, and early inpatient outcomes in routine practice remains unclear.

METHODS: Using the National Inpatient Sample (2016-2022), we identified primary medial UKA for osteoarthritis. RA-UKA was defined by ICD-10-PCS robotic-assisted lower-extremity procedure codes. Primary outcomes were prolonged hospitalization (LOS $\geq$ the 75th percentile, $\geq$ 2 days), high-charge admission (total hospital charges [TOTCHG] $\geq$ the 75th percentile, $\geq$ \$72,321), and in-hospital mortality. We performed univariable comparisons and multivariable logistic regression adjusting for demographics, payer, admission type, hospital teaching status, comorbidities, and calendar year fixed effects (YEAR 2016-2022) using the unweighted NIS discharge sample.

RESULTS: Among 7,154 UKAs, 1,297 (18.1%) were RA-UKA and 5,857 (81.9%) were C-UKA. Median LOS was 1 day (IQR 1-2) in both groups; however, prolonged hospitalization (LOS $\geq$ 2 days) occurred less frequently in RA-UKA (412/1,297 [30.7%] vs. 2,162/5,857 [37.2%]; $P < 0.001$). In adjusted analyses including YEAR fixed effects, RA-UKA was associated with lower odds of prolonged hospitalization (aOR 0.750, 95% CI 0.647-0.870; $P < 0.001$) and was not associated with high-charge admission (TOTCHG $\geq$ [...])

[PubMed](#) · [DOI](#)

Is biceps augmentation effective for scope-assisted lower trapezius tendon transfer in posterosuperior irreparable rotator cuff tears? A retrospective short-term clinical comparison.

Baek CH, Elhassan BT, Lim C, Kim JG, Kim BT, Kim SJ

INTRODUCTION: Although favorable clinical outcomes of scope-assisted lower trapezius tendon transfer (SALTT) have been reported in posterosuperior irreparable rotator cuff tears (PSIRCTs) patients, the limited restoring of static stability can be supported by biceps augmentation as a static stabilizer. This study aimed to compare the outcomes of scope-assisted lower trapezius tendon transfer (SALTT) with and without biceps augmentation in patients with posterosuperior irreparable rotator cuff tears (PSIRCTs).

MATERIAL AND METHODS: This retrospective clinical comparative study was performed with the inclusion criteria: patients who underwent SALTT for PSIRCT from January 2022 to April 2023; a follow-up period of more than 2 years; and availability for clinical assessment and MRI evaluation preoperatively and at 2 years postoperatively. The patients were grouped according to the management of biceps; Non-augmentation group ($n = 26$) and Augmentation group ($n = 27$). The clinical outcomes were evaluated using shoulder pain, patient-reported outcome measures (PROMs), and active range of motion (aROM). Radiological outcomes were performed to evaluate the progression of shoulder joint arthritis, biceps integrity, and graft integrity.

RESULTS: Both groups showed significant improvements in VAS score, PROMs, and aROM. Although postoperative PROMs of Non-augmentation group were significant [...]

[PubMed](#) · [DOI](#)

Robotic-arm assisted total hip arthroplasty using a short metaphyseal filling collared femoral implant through a direct anterior approach: minimum two-year outcomes.

Marchand R, Kaczynski E, Taylor K, Smitherberg C, Mont MA

INTRODUCTION: Short, collared, metaphyseal-filling femoral stems preserve proximal bone and achieve stable fixation while reducing stress shielding compared to diaphyseal-engaging stems. Early six-month data for one such stem demonstrated improved patient-reported outcomes with no implant-related complications. This study extends follow-up of the same cohort to a minimum of two years to evaluate (a) functional outcomes, (b) implant-related complications, and (c) survivorship.

MATERIALS AND METHODS: The prospective cohort of 120 patients underwent robotic-arm assisted total hip arthroplasty (THA) through a direct anterior approach (DAA) using a short, collared, metaphyseal-filling stem. Patients had a mean age of 69.2 years (range 37-90) and a body mass index of 28.2 (range 18.1-49.1), and they were 71% women and were followed for a mean of 2.6 years (range 2.0-3.4). The Hip Disability and Osteoarthritis Outcome Score for Joint Replacement (HOOS-JR) was compared with baseline using Student's t-tests. Complications included all-cause revision, periprosthetic joint infection, periprosthetic fracture, and dislocation at

final follow-up.

RESULTS: At mean follow-up, HOOS-JR scores improved from a mean of 54.5 ± 13.9 preoperatively to 94.4 ± 10.3 ($P < 0.001$). The overall survivorship was 98.1% following two septic revisions.

CONCLUSIONS: This short, collared, metaphyseal-filling st [...]

[PubMed](#) · [DOI](#)

Second contralateral hip fractures reduce survival, mobility and daily activity : a matched pair analysis.

Blattner A, Sabath F, Röttinger T, Lisitano L, Mayr E, Fenwick A

BACKGROUND: Hip fractures in older adults are associated with substantial morbidity, functional decline, and mortality. Patients who experience a first fragility fracture are at high risk of subsequent fractures, with repeated events further exacerbating functional impairment and survival outcomes. Second contralateral hip fractures, while clinically important, remain under-characterized in terms of timing, long-term functional impact, and mortality.

OBJECTIVES: To investigate the incidence, timing, survival, mobility, and daily activity outcomes of second contralateral hip fractures using a matched pair analysis, and to situate these findings within the broader context of repeated fragility fractures.

METHODS: A retrospective cohort study was conducted at a single Level I trauma center, including all patients treated operatively for hip fractures (AO 31A1-A3, 31B) between January 2016 and June 2020. Patients with a second contralateral hip fracture were matched 1:1 with patients with a single fracture based on age, sex, fracture type, and Charlson Comorbidity Index. Demographic data, functional status (Barthel Index, Parker Mobility Score), walking aid use, living situation, and mortality were assessed at admission, discharge, and follow-up (minimum 3 years, maximum 7 years). Statistical analysis included paired tests and survival analysis.

RESULTS: Of 1,933 patients, 148 ([...]

[PubMed](#) · [DOI](#)

Short-term clinical outcomes of robotic-assisted total knee arthroplasty at 12-month follow-up: a prospective, multicenter, concomitant comparison to conventional total knee arthroplasty.

Nöth U, Rivkin G, Caldora P, Heller KD, Thienpont E, Hannouche D, Perets I, Cholewa J

INTRODUCTION: There is limited data available on short-term outcomes on a cut-block positioning robotic system. The purpose of this study was to compare 12-month clinical outcomes between robotic-assisted (raTKA) and conventional total knee arthroplasty (cTKA) with multiple outcomes and surgical centers.

METHODS: This was a non-randomized controlled trial of patients who received either raTKA ($n = 120$) or cTKA ($n = 101$) at 6 different surgical centers. Variables of interest included occurrence of soft tissue release, complications and revisions at minimum one-year follow-up. Satisfaction, pain (numeric rating scale [NRS]), 5-dimensional European Quality of Life (EQ-5D-5 L) questionnaire (index and visual analog scale [VAS]), Oxford Knee Score (OKS), and the Forgotten Joint Score (FJS-12) were collected pre-operatively, and at six weeks, three months, and 12 months post-operative.

RESULTS: There were significantly less soft tissue releases with raTKA (28/120, 23.3%) vs. cTKA (51/99, 51.5%), ($p < 0.0001$). There were significantly fewer cases of medial/lateral instability in the raTKA group at six-weeks ($p = 0.038$) and three-months ($p = 0.007$) post-operative. At one-year follow-up, 96.3 and 92.5% of raTKA and cTKA patients were satisfied with the overall results of their surgery, respectively. Significantly more raTKA patients were very satisfied (32.1% vs. 14.6%) with their abil [...]

[PubMed](#) · [DOI](#)

Clinical and radiological outcomes of using locked intramedullary nails in the treatment of severe frontal plane lower limb deformity in adolescents with hypophosphatemic rickets (mid-term results).

Galal S, Al-Abri AM, Al-Habsi G, Al Ghammari M, Elissawi A, Abou Senna WG, Arafa AS

INTRODUCTION: Lower limb deformities in patients with hypophosphatemic rickets are multi-apical and require multiple osteotomies for correction. Intramedullary nails (IMNs) are used to fix multiple osteotomies. Authors of this study aimed to assess the clinical and radiographic outcomes of using IMNs for correcting lower limb deformity in adolescents with hypophosphatemic rickets.

METHODS: This prospective study included patients with hypophosphatemic rickets who underwent deformity correction between November 2020 and November 2023 using IMNs, with a minimum follow-up of 2 years. Clinical outcomes were assessed using the Lower Limb Deformity-Scoliosis Research Society (LD-SRS) score. Radiographic outcomes measured included the mechanical tibiofemoral angle (mTFA), mechanical axis deviation (MAD), mechanical lateral distal femoral angle (mLDFA), mechanical medial proximal tibial angle (mMPTA), and Stevens' knee joint zoning.

RESULTS: Twenty patients (25 limbs and 35 bones) were included (mean age: 16 years; range: 13-22 years). The LD-SRS score improved from 3.2 ± 0.4 preoperatively to 4.3 ± 0.7 postoperatively. Preoperative mechanical tibiofemoral angle was $27.8^\circ \pm 12.1^\circ$ and $17.33^\circ \pm 7.9^\circ$ in the varus and valgus groups, respectively, improving to $3.4^\circ \pm 6.4^\circ$ and $2^\circ \pm 5.2^\circ$, respectively. Preoperative mechanical axis deviation was 82.2 ± 35.6 mm and 41.8 ± 22.2 mm in the varus [...]

[PubMed](#) · [DOI](#)

Risk prediction and surgical decision-making for pathological fractures in phalangeal enchondroma: validation of the 0.51 threshold and outcome comparison of prophylactic intervention in 65 patients.

Zhao J, Mo Z, Lao Y, Chen J, Li Q, Lin J

BACKGROUND: Enchondroma is the most common benign bone tumor of the hand, predominantly affecting the phalanges and metacarpals, and is frequently complicated by pathological fractures. Although previous studies have explored risk factors based on tumor size, a precise quantitative cut-off value remains undefined. Furthermore, quantitative research regarding the impact of "delayed treatment" on long-term functional outcomes is lacking. This study aims to establish a risk threshold for pathological fractures in phalangeal enchondroma and to provide an evidence-based rationale for optimizing surgical decision-making by comparing the clinical outcomes of prophylactic surgery versus post-fracture management. **METHODS:** We retrospectively analyzed the clinical data of 65 patients with phalangeal enchondroma treated surgically at our institution between 2017 and 2025. Patients were categorized into a Fracture Group (n = 35) and a Non-fracture Group (n = 30) based on the presence of a complete preoperative fracture. The tumor-to-phalanx length ratio was measured using the Picture Archiving and Communication System (PACS), and the degree of cortical thinning was assessed according to the Takedani classification. Multivariate logistic regression analysis was employed to identify independent risk factors, and Receiver Operating Characteristic (ROC) curve analysis was used to determine the [...]

[PubMed](#) · [DOI](#)

Assessing PHILOS plate as an alternative fixation method for pediatric femoral neck fractures: a biomechanical comparison with cannulated screws.

Lim BS, Wang YP, Chung ST, Wang CH, Chang SW, Wang TM, Hong CK

BACKGROUND: Pediatric femoral neck fractures require stable fixation to avoid complications. It remains unclear whether fixation with the Proximal Humeral Internal Locking System (PHILOS) can serve as an alternative to cannulated screw fixation. The purpose of this study was to compare the biomechanical properties of PHILOS and cannulated screws for stabilizing unstable pediatric femoral neck fractures using a synthetic bone model.

MATERIALS AND METHODS: Twelve fourth-generation synthetic composite femurs were randomly assigned to screw fixation (Group S) or PHILOS fixation (Group P) (n = 6 each). A standardized vertically oriented Delbet type II osteotomy was created in all specimens. Group S was fixed with three 6.5-mm cannulated screws, whereas Group P received a PHILOS plate with 3.5-mm locking screws. Each specimen underwent a standardized loading protocol using a universal testing machine. Axial stiffness, cyclic displacement, ultimate failure load, and failure modes were recorded and statistically compared between groups.

RESULTS: No statistically significant difference was found in axial stiffness between Group P (746 ± 300 N/mm) and Group S (753 ± 256 N/mm) (p = 1.000). Displacement after cyclic loading was significantly greater in Group P (1.42 ± 0.3 mm) compared with Group S (0.57 ± 0.2 mm) (p = 0.004). The ultimate failure load was higher in Group

S (2378 ± 513 N) [...]

[PubMed](#) · [DOI](#)

Rerupture after flexor tendon repair of the hand and wrist: a retrospective risk factor analysis.

Marwieser SF, Vavricka S, Kaiser P, Schmidle G, Lindtner RA, Arora R, Konschake M

INTRODUCTION: Rerupture after primary flexor tendon repair of the hand and wrist is a serious complication that can substantially impair hand function and restrict activities of daily living, occupational performance, and recreation. Identifying factors associated with rerupture is essential to optimize surgical techniques and postoperative rehabilitation strategies and to improve patient outcomes. **METHODS:** A retrospective observational cohort study was conducted including patients who underwent surgical repair of flexor tendon injuries of the hand and wrist at a single institution between 2010 and 2022. The study included 292 patients with 429 complete tendon injuries. Collected data comprised demographic characteristics, injury mechanism and type, anatomical location, surgical timing and technique, and postoperative immobilization and rehabilitation protocols. Associations between potential risk factors and rerupture were explored using univariable analyses, followed by a multivariable logistic regression model to identify independent predictors. **RESULTS:** The overall rerupture rate was low but clinically relevant, occurring in a small proportion of patients and tendon lesions. Rerupture was more frequently observed in male patients, older individuals, and those with specific injury mechanisms. Surgical factors, particularly the need for pulley reconstruction, were associated [...]

[PubMed](#) · [DOI](#)

Defining orthoplastic limb salvage centers: a systematic review.

Moussa O, Pacheco FJ, Raasveld FV, Gonzalez MR, Oflazoglu K, Ritt MJPF, Valerio IL, Rakhorst H et al.

INTRODUCTION: Limb salvage centers have increased in number over time, but lack standardized defining criteria. This systematic review aimed to assess organizational features of limb salvage centers and determine whether orthoplastic centers, in comparison to vascular limb salvage centers, represent a distinct care model that may benefit from standardization.

METHODS: We conducted a systematic review of publications related to limb salvage centers by searching MEDLINE, Embase, Web of Science, and Cochrane databases from their inception through 2024. We quantified binary data extraction as a reporting score of 26 organizational features across six structural care domains for limb salvage centers, based on a validated quality measurement framework. Organizational features differentiating distinct center types were identified to establish a quality framework for orthoplastic centers. Statistical comparisons between center types were performed using appropriate tests ($p < 0.05$).

RESULTS: Of 118 included studies, orthoplastic ($n = 43$) and vascular ($n = 48$) centers represented 77% of all studies. Recent increases in orthoplastic publications show substantial variability in organizational features. Orthoplastic center literature more frequently reported plastic surgery consultation criteria ($p < 0.001$), surgical outcomes ($p < 0.001$), and centralized network integration ($p \leq 0.006$), [...]

[PubMed](#) · [DOI](#)

Male incidence rates outpacing females: results from a nationwide analysis on acetabular fractures in Germany.

Baumann F, Bärthel S, Szymanski D, Hierl K, Alt V, Freigang V

INTRODUCTION: Acetabular fractures are uncommon, but serious injuries. Demographic changes may have a significant impact on planning healthcare structures to improve treatment outcomes. Aim of this nationwide, registry-based retrospective controlled study was to identify incidence trends, demographic characteristics, and care structures of patients with acetabular fractures in Germany.

MATERIALS AND METHODS: We analyzed inpatient data from the Institute for the Hospital Remuneration System (InEK). Based on 52 095 patients with primary diagnosis of an acetabular fracture between 2019 and 2024, we calculated incidence rates for different age-groups and put a spotlight on geriatric acetabular fractures (> 65 years of age).

RESULTS: Incidence rates in patients under 65 years remained stable, whereas patients over 65 years showed a significant age-dependent increase with an exponential rise in men aged 80 + with the highest incidence being 122.4/100 000 inhabitants annually. We recorded high levels of co-morbidity and nursing care dependency for elderly patients after acetabular fracture. Although 43% of patients were treated in hospitals > 500 beds, acetabular fractures were managed across all hospital sizes.

CONCLUSIONS: There is a rapidly increasing incidence of geriatric acetabular fractures, predominantly driven by elderly male patients over 80 years. Patients over 65 years [...]

[PubMed](#) · [DOI](#)

Cemented total hip arthroplasty reduces early complications: a Japanese nationwide propensity-matched study.

Tanaka H, Tarasawa K, Mori Y, Baba K, Kurishima H, Fushimi K, Aizawa T, Fujimori K

INTRODUCTION: The optimal fixation method in total hip arthroplasty (THA) remains under debate. While cemented fixation has been associated with a lower risk of periprosthetic fracture, uncemented fixation predominates in Japan. This study aimed to compare early postoperative complications between cemented and uncemented fixation in elective THA using a nationwide inpatient database.

MATERIALS AND METHODS: We identified 198,102 patients aged ≥ 65 years who underwent primary THA for osteoarthritis, osteonecrosis, or rheumatoid arthritis between December 2011 and March 2023 from the Japanese Diagnosis Procedure Combination (DPC) database. After 1:1 propensity score matching for age, sex, body mass index (BMI), and Charlson Comorbidity Index, 36,859 patients were included in each fixation cohort. Surgical and medical complications, and in-hospital mortality were compared using multivariate logistic regression.

RESULTS: Cemented fixation was associated with a significantly lower risk of periprosthetic fracture (odds ratio [OR], 0.40; 95% confidence interval [CI], 0.30-0.53; $p < 0.001$), blood transfusion (OR, 0.76; 95% CI, 0.74-0.78; $p < 0.001$), and deep vein thrombosis (OR, 0.79; 95% CI, 0.74-0.84; $p < 0.001$). There were no statistically significant differences based on the predefined threshold ($p < 0.001$) in dislocation, infection, pulmonary embolism, cardiac or cerebrovascular [...]

[PubMed](#) · [DOI](#)

Healthcare utilization following hip fractures based on social vulnerability status in the US: an analysis of 2016-2020 nationwide readmissions data.

Tilve R, Zhou G, Khan ST, Koroukian SM, Deren M, Piuze NS

INTRODUCTION: Social vulnerability (SV) influences rehabilitation and postoperative care for patients with hip fracture. However, most previous work relies on area-level measures that overlook interindividual variation. The recent adoption of ICD-10 Z-codes allows clinical identification of patient-level SV and may offer a better understanding of its impact. This study aimed to evaluate healthcare utilization, including readmissions, discharge disposition, and length of stay (LOS) in surgically treated hip fracture patients with and without clinically acknowledged SV.

METHODS: Adults surgically treated for hip fracture between 2016 and 2020 were included from the Nationwide Readmissions Database. SV was defined as having at least one documented relevant ICD-10 Z-code. Primary outcome measures included complications, LOS, discharge disposition, and 30- and 90-day readmissions, stratified by SV and evaluated using chi-square analyses. Multivariable logistic regression assessed long LOS (≥ 5 days) and discharge to home, adjusting for age, insurance/income status, and substance use.

RESULTS: Patients with SV were younger (35.6% with SV vs. 50.1% without SV were 81+), had a lower median household income (38.8% with SV vs. 25.7% without SV were in the lowest quartile), and were more often insured by Medicaid (19.3% vs. 3.8%). Alcohol/drug use disorders were significantly more prevalent [...]

[PubMed](#) · [DOI](#)



Cumulative traumatic life events and increased risk for emergency department and inpatient utilization after physical injury hospitalization.

Huynh M, O'Connor S, McManamen M, Buggaveeti AE, Russo J, Wang J, Iles-Shih M, Shoyer J et al.

INTRODUCTION: Few large-scale investigations have examined the association between the cumulative burden of traumatic life events and population-based assessments of emergency department (ED) and inpatient utilization among physically injured trauma survivors.

METHODS: The investigation was a secondary analysis of a randomized clinical trial that examined the association between cumulative traumatic life events and ED/inpatient utilization after physical injury hospitalization. All 445 physically injured trauma survivors included in the investigation were aged ≥ 18 and underwent screening for elevated posttraumatic stress symptoms. Three types of patient characteristics were examined for potential significant associations with the longitudinal ED/inpatient utilization outcome: pre-injury, baseline at the time of index physical injury admission, and events that occurred in the year post-injury. Regression analyses were used to assess the association between these clinical and demographic characteristics and ED/inpatient utilization in the 12-36 months after the index injury admission.

RESULTS: Approximately one-third of patients reported ≥ 7 serious traumatic life events before the index injury admission. Approximately 62% of patients had one or more ED/inpatient visits over the course of the 12-36 months post-injury. Clinical and demographic characteristics that were significant [...]

[PubMed](#) · [DOI](#)

Agreement between ChatGPT and emergency physicians in laceration management: A prospective study.

Saiflam Gürmen E, Yorgancıoğlu M, Oral A

INTRODUCTION: Although large language models (LLMs) like ChatGPT are increasingly used in clinical reasoning, their reliability in procedural decision-making remains questionable. Wound management provides a standardized framework for evaluating their performance in practical scenarios. The primary goal of this study was to assess the level of agreement between ChatGPT's recommendations and emergency physicians' decisions in the management of traumatic lacerations.

METHODS: The Emergency Department of Manisa Celal Bayar University Hafsa Sultan Hospital served as the site of this prospective, comparative study, conducted from February to May 2025. A cohort of 792 patients with traumatic lacerations was included. Five structured questions were presented to ChatGPT (GPT-4 model) to obtain information on suture material, technique, count, wound classification, and the necessity of antibiotics. Physicians' documented decisions were compared with their recommendations. Concordance was measured using Cohen's kappa (κ), and subgroup analyses were conducted by wound size and location.

RESULTS: The median age of the cohort was 28 years (IQR 16-40, range, 2-80), the proportion of males was 79.9%. The leading cause of injury was metal-related trauma, accounting for 47.7%. Across all decision domains, there was considerable to almost perfect agreement between ChatGPT and physicians ($\kappa = 0$ [...])

[PubMed](#) · [DOI](#)

Video-assisted thoracoscopic surgery versus open thoracotomy for thoracic trauma in hemodynamically stable patients: An updated systematic review and meta-analysis.

Barrientos ML, Zapata CAL, López CAD

BACKGROUND: Video-assisted thoracoscopic surgery (VATS) is increasingly used in thoracic trauma, yet high-certainty comparative evidence against thoracotomy remains limited. We updated the only prior comprehensive meta-analysis (Wu et al., 2015) and re-derived all estimates from source-verified study-level data.

METHODS: Six databases were searched through December 2025. Studies comparing VATS with thoracotomy in hemodynamically stable adult patients with thoracic trauma were eligible. Random-effects models with restricted maximum likelihood estimation and Knapp-Hartung adjustment were used; heterogeneity was assessed with I^2 and 95% prediction intervals (PI). PROSPERO CRD420261322930.

RESULTS: Thirty studies were included. VATS was associated with fewer postoperative complications (RR 0.56, 95% CI 0.42-0.74; $I^2 = 0\%$), shorter hospital stay (MD -4.30 days, 95% CI -5.12 to -3.49), shorter operative time (MD -37.52 min, 95% CI -47.73 to -27.32), and less intraoperative blood loss (MD -119.09 mL, 95% CI -159.16 to -79.01). Perioperative mortality favoured VATS but was of borderline significance (RR 0.48, 95% CI 0.24-0.97; $I^2 = 0\%$) and was not significant when the single dominant study was removed (RR 0.56, 95% CI 0.27-1.19). The pooled VATS-to-thoracotomy conversion rate was 13.0% (95% CI 5.9-26.2%). Prediction intervals excluded the null for complications and hospital stay but [...]

[PubMed](#) · [DOI](#)

Immersive virtual reality simulation for technical skills acquisition in pneumothorax management and chest tube insertion: A randomized controlled trial among surgery interns.

Leon S, Rouhi AD, Baji D, Goldberg DW, Hwang J, Weber D, Fernandez FB, Cannon JW et al.

BACKGROUND: Immersive virtual reality (IVR) offers a cost-effective, high-fidelity approach to enhance technical skills training. This study assessed the efficacy of IVR simulation delivered on a head-mounted display (HMD) for technical skills acquisition in pneumothorax management and chest tube insertion (CTI) among first-year surgery residents.

METHODS: A novel IVR module on pneumothorax management and CTI was developed using Unity software by a multidisciplinary team of surgeon-educators and software engineers. Forty-five surgical residents were randomly assigned to either IVR instruction or traditional classroom didactics. Participants in both groups then performed CTI on a high-fidelity task trainer, and technical skills acquisition was measured using a validated ten-item Objective Structured Assessment of Technical Skills (OSATS) tool.

RESULTS: The IVR group ($n = 21$) outperformed the control group ($n = 24$) in mean OSATS score [39.8 (4.4) vs 36.7 (2.9), $p = 0.008$]. The IVR group also scored higher in rib blunt dissection [4.14 (0.48) vs 3.79 (0.59), $p = 0.035$], economy of time and motion [3.71 (0.72) vs 3.21 (0.51), $p = 0.009$], and required less tutor assistance [4.00 (0.89) vs 3.33 (0.70), $p = 0.008$]. Five HMDs were utilized at a total cost of \$1749.95.

CONCLUSION: In this randomized controlled trial, a novel, cost-effective IVR module demonstrated superior technical [...]

[PubMed](#) · [DOI](#)

Re-triage of severely injured patients in no-fault vs. at-fault states.

Visenio MR, Thivalapill N, Silver CM, Sullivan GA, Thomas AC, Reddy S, Ghneim M, Ryan JL et al.

IMPORTANCE: State no-fault laws enforce fewer administrative requirements for automotive insurance to pay hospitals for injury care costs than states without no-fault laws (at-fault states).

OBJECTIVE: Determine if no-fault states had lower re-triage (urgent inter-hospital transfer) of patients injured in motor-vehicle collision (MVC) than at-fault states.

DESIGN: This was an observational retrospective cohort study.

SETTING: FL, NY, MD and WI Healthcare Cost and Utilization Project State Emergency Department and Inpatient Databases from 2016 to 2021.

PARTICIPANTS: Severely injured [injury severity score (RISS) > 15] Medicare patients.

EXPOSURE: FL and NY had no-fault laws, while MD and WI were at-fault states.

MAIN OUTCOME: Re-triage was defined as same-day inter-hospital transfer to a high-level trauma center. Mixed effects multivariable logistic regression estimated the association between no-fault states and re-triage controlling for age, sex, race, comorbidity, RISS, diagnosis, trauma designation, hospital for-profit, adjusting for clustering within hospitals and states. A moderation analysis with an interaction term for mechanism * fault laws was conducted to estimate whether no-fault laws moderated adjusted odds of re-triage differently among patients

severely injured in MVC versus falls.

RESULTS: In total 6447 Medicare patients in 345 hospitals were severely inju [...]

[PubMed](#) · [DOI](#)

Comparable outcomes for non-operative intracranial hemorrhage at Level III vs Level I/II trauma centers.

Lord S, Arda Y, Karikis I, DeWane MP, Paranjape CN, Ng-Kamstra JS, Maurer L, Kaafarani HMA et al.

INTRODUCTION: Most patients with isolated, closed blunt intracranial hemorrhage (ICH) do not require neurosurgical intervention. However, outcome data for those managed entirely at Level III trauma centers remain incompletely characterized.

METHODS: Using the 2017-2020 Trauma Quality Improvement Program (TQIP), we compared clinical outcomes for non-operative ICH patients treated at Level I/II versus Level III trauma centers. The primary endpoint was composite in-hospital adverse event counts per patient, modeled using negative binomial regression and reported as adjusted incidence rate ratios (aIRRs). Secondary analyses included patients with ISS > 15, ICU admission by trauma center level, predictors of ICU admission, and unplanned ICU transfers. To address triage selection bias, we also compared time to ED discharge in a cohort of patients who met the same inclusion and exclusion criteria and were transferred from a Level III trauma center.

RESULTS: A total of 24,132 patients were included after complete-case exclusion (Level I/II: 22,846; Level III: 1286). Composite adverse event counts were 0.067 events per patient at Level I/II centers and 0.042 at Level III centers. After adjustment, there was no statistically significant difference in composite adverse event counts between groups (aIRR 0.82, 95% CI 0.56-1.18). Mortality was lower at Level III centers (aOR 0.12, 95% CI 0 [...])

[PubMed](#) · [DOI](#)

Drug screening in patients with depressed GCS: An evaluation of current practice and potential bias.

Serio F, Jenkins PD, Barton C, Loss L, Tinoco-Garcia L, Kiraly L, Brasel KJ

BACKGROUND: Substance use contributes to trauma morbidity, mortality, and recidivism, yet screening practices among trauma patients are inconsistent. We examined urine drug screening (UDS) utilization among trauma patients presenting with depressed Glasgow Coma Scale (GCS).

METHODS: We retrospectively reviewed adult trauma activations (age ≥18) with admission GCS ≤ 12 at a Level I trauma center (2020-2022). The primary outcome was whether UDS was obtained. Secondary measures included demographic and clinical predictors, UDS results, and referral for screening, brief intervention, and referral to treatment (SBIRT). Logistic regression assessed associations with testing.

RESULTS: Of 661 patients, 39% underwent UDS; 67% of these were positive. Patients aged 31-50 were more likely to be screened, while those ≥ 81 and those discharged from the emergency department were less likely. Cannabis was the most commonly detected drug. Nearly all patients with positive UDS were referred for SBIRT.

CONCLUSIONS: Trauma patients with depressed GCS are under-screened despite high positivity. Standardized screening may improve detection and intervention.

[PubMed](#) · [DOI](#)

Effectiveness of magnets in removing metallic foreign bodies.

Aslantürk O, Ergen E, Karakaplan M, Özdeğir H, Al F, Ertem K

BACKGROUND: Metallic foreign bodies (MFB) in the extremities are a common clinical problem that requires removal to prevent complications and to avoid contraindications to magnetic resonance imaging. While magnetic extraction represents an alternative to conventional surgical techniques, systematic data regarding its efficacy and clinical utility remain limited. This study aimed to evaluate the clinical efficacy of magnetic extraction for MFBs in extremities, its impact on radiation exposure, and success rates across different anatomical locations and foreign body types.

METHODS: We retrospectively reviewed data on patients who underwent MFB removal surgery between 2017 and 2025 at a tertiary trauma center. Of the 364 patients reviewed, 51 met the inclusion criteria after excluding non-metallic foreign bodies, cases in which magnets were not used, and incomplete records. Demographic data, anatomical location, foreign body characteristics, use of image intensifier, and magnet efficiency were recorded.

RESULTS: A magnet was used for 52 MFBs in 51 patients. Magnetic extraction of MFBs was successful in 87% (45/52) of cases. The success rate was 100% for metal splinters but decreased to 63% for the needle-type foreign bodies ($p < 0.001$). Magnetic extraction succeeded in all upper-extremity cases (28/28, 100%) and in 68.2% (15/22) of lower-extremity cases; the single patient with [...]

[PubMed](#) · [DOI](#)

The economic value and financial burden of informal caregiving during 3 months after geriatric fracture admission: A longitudinal cohort study.

Raats JH, Sluijter AJ, Lintmeijer L, van der Velde D, Schuijt HJ

BACKGROUND: While fragility fractures result in significant healthcare expenditure, the (in)direct costs of fracture-related informal care are rarely acknowledged. This study quantifies the economic value of informal care and the financial burden among working-age caregivers of older adult patients following a fracture-related hospital admission.

METHODS: This longitudinal cohort study included working-age (<67 years) informal caregivers of patients aged ≥ 65 years who were admitted for any fracture. Caregivers reported pre-fracture caregiving details and at 2 weeks, 4 weeks and 3 months post-discharge. The primary outcome measure was the economic value of informal care in Euro (€) per week, estimated using the opportunity cost method. Secondary outcomes included a composite financial burden score ranging from 0 (low) to 6 (high) and dichotomized (low vs. high) financial worry score. Linear mixed models were constructed to assess associations of time and caregiver/patient characteristics with informal caregiving costs.

RESULTS: The 67 included caregivers were on average 54 years old, mostly female (66%), and patients' children (79%). Mean weekly informal care costs were €91.66 (95%CI 54.62-128.71) before admission, €244.80 (95%CI 164.62-342.99) at 2 weeks, €214.57 (95%CI 149.46-279.68) at 4 weeks, and €133.58 (95%CI 92.01-175.16) at 3 months post-discharge. In linear mixed mo [...]

[PubMed](#) · [DOI](#)

Epidemiology and outcomes of geriatric patients with severe injury in Korea: Nationwide community-based study, 2016-2023.

Lee SY, Song KJ, Hong KJ, Park JH, Kim TH, Jeong J, Choi SK, Choi YH

INTRODUCTION: As the number of geriatric populations increases in the aging era, the injury of the older adults is also increasing. This study aimed to investigate the epidemiologic trends and injury characteristics of geriatric patients with severe injuries by age groups and mechanism, using a nationwide database in Korea.

MATERIALS AND METHODS: A cross-sectional observational study was conducted for all emergency medical service (EMS) treated geriatric (≥ 65 years) patients with severe injuries in Korea between 2016 and 2023. Patients who met the criteria for a field-based injury triage scheme or had a prehospital physiologic abnormality were defined as severe injury. Exposure was age group: young old (65-74), middle old (75-84), and oldest old (≥ 85 years) groups. The distribution of age groups was investigated by year. Injury characteristics and outcomes were compared by age group and mechanism. A Multivariable logistic regression analysis was used to investigate the predictors of mortality of geriatric patients with severe injuries.

RESULTS: Among 319,567 patients with severe injuries transferred by EMS over 8 years, 92,085 (28.8%) were over 65 years of age. The proportion of older adults increased from 25.7% in 2016 to 32.0% in 2023. Among older adults, young old was 46.4% (42,742), middle old was 39.3% (36,182), and oldest old was 14.3% (13,161). The proportion of oldest [...]

[PubMed](#) · [DOI](#)

Emergency department burden of e-scooter injuries in Victoria, Australia: A statewide analysis of injury patterns and prevention implications, 2022-2025.

BACKGROUND: E-scooter use has expanded rapidly in Australia alongside evolving legislation and changes in e-scooter availability. However, the population-level impact on injury burden and emergency care services remains incompletely characterised, particularly in the absence of standardised external cause coding for e-scooter injuries.

METHODS: We conducted a retrospective statewide analysis of emergency department (ED) presentations for e-scooter-related injuries in Victoria, Australia, using the Victorian Emergency Minimum Dataset (VEMD). Cases from January 2022 to December 2025 were identified using predefined text-search terms applied to narrative fields, with manual validation. Descriptive analyses examined temporal trends, demographic characteristics, injury patterns, health service use, and seasonal variation. Temporal patterns were interpreted in the context of evolving e-scooter legislation and shared-hire programme changes.

RESULTS: A total of 4694 e-scooter-related ED presentations were identified, including 79 pedestrians struck by an e-scooter. Annual presentations increased from 825 in 2022 to 1,373 in 2024 and remained high in 2025 (1358). Riders were predominantly male (71.4%) with a mean age of 31.1 years; however, 15.4% of presentations involved children aged 2-14 years. Falls accounted for 74.1% of injuries, and 67.2% occurred on roads, streets, or highways [...]

[PubMed](#) · [DOI](#)

Dual free flap reconstruction: A configuration-based algorithm using parallel and serial designs.

Ha Y, Kim JT, Hong SE, Kim YH

BACKGROUND: In severe trauma, extensive or geometrically complex soft-tissue defects often exceed the capacity of a single free flap. Dual free flaps offer a versatile solution, yet guidance on flap choice and configuration remains limited. We assessed outcomes using a configuration-based algorithm distinguishing parallel and serial designs.

METHODS: From January 2015 to July 2025, 32 patients at two centers underwent dual free-flap reconstruction for wide or elongated defects, predominantly in the setting of trauma. Flap selection was guided by defect geometry and recipient-vessel accessibility: parallel configurations were used for wide defects (first flap to the main recipient vessel, second flap to the serratus anterior branch of the thoracodorsal system), whereas serial configurations were applied for elongated defects (first flap to proximal inflow, second flap to the distal descending branch of the lateral circumflex femoral artery). Patient characteristics, flap types and dimensions, complications, and clinical outcomes were analyzed.

RESULTS: The mean age was 51.1 years, and trauma was the most common etiology (43.8%). Parallel configurations were used in 27 cases (84.4%) and serial configurations in 5 (15.6%). The mean surface areas of the first and second flaps were 281.7 ± 134.9 cm² and 247.9 ± 179.0 cm², respectively, with a total reconstructed area of 529.6 ± 24 [...]

[PubMed](#) · [DOI](#)

More than meets the eye: the continuing importance of tertiary trauma survey even in moderate (ISS 9-15) trauma.

Knight C, Burke JR, Hind S, Smith SR

INTRODUCTION: Tertiary Trauma Surveys (TTS) are designed to identify injuries which are not recognised during the primary and secondary surveys. These injuries occur in up to 40% of patients and are commonly referred to in the literature as "missed", we prefer the term "unrecognised" as this term does not infer the failure of primary and secondary survey, rapid assessments designed to identify life and limb threatening injuries. The incidence and definition of unrecognised injuries varies across studies, and several risk factors have been identified internationally. The aim of our study was to investigate the sensitivity and specificity of TTS and the incidence and risk factors for unrecognised injuries in a large UK Major Trauma Centre (MTC).

METHODS: Data was prospectively collected for all major trauma patients admitted to our MTC over six months. All patients over 16 years with an ISS ≥ 9 were included. Patients with a length of stay of less than 24 h or palliated were excluded. All patients were examined using a specific TTS protocol. Patient records were reviewed one year post admission to identify any injuries diagnosed after discharge.

RESULTS: A total of 293 patients were included. Unrecognised injuries were identified at TTS in 18.8% (55/293) patients, most commonly affecting the face or head and eyes. Patients with unrecognised injuries had a higher admission ISS (p [...])

[PubMed](#) · [DOI](#)

CTA-detected incidence and descriptive fracture patterns of arterial injury in 539 surgically treated tibial plateau fractures.

Valderrama J, Narváez F, Cancino J, Raffo JF, Bosselin R, Zamora F, Siegel S, Scheu M et al.

BACKGROUND: Tibial plateau fractures (TPF) are complex injuries associated with significant soft tissue and periarticular complications. Although vascular injuries are well described in lower limb trauma, their incidence in tibial plateau fractures has not been clearly established.

METHODS: Retrospective cohort study including patients with surgically treated TPF at a level I trauma center between 2016 and 2024. CTA was selectively performed based on clinical judgment. Post hoc descriptive risk scenarios were constructed to organize observed fracture patterns. Analysis was primarily descriptive.

RESULTS: 539 surgically treated TPF were included. Mean age was 43.7 years (18-75), 73.84% were male. Left knee was affected in 54.92%. High-energy trauma accounted for 70.87%. The CTA yield was 3.66% (6/164), corresponding to an overall observed incidence of 1.11% (6/539). Certain fracture patterns and clinical scenarios were more frequently observed among cases with arterial injury; however, these observations are based on very few events and should be interpreted cautiously as descriptive findings only.

CONCLUSION: The CTA-detected incidence of arterial injury in this series of surgically treated tibial plateau fractures was 1.11%. Certain fracture patterns and clinical contexts were observed among the small number of cases with arterial injury; however, these findings are descriptive [...]

[PubMed](#) · [DOI](#)

Diagnostic value of serum troponin in stable patients at risk for blunt cardiac injury.

Cleary HL, Bernard MD, Davenport DL, Bernard AC

INTRODUCTION: There are no gold standard criteria for diagnosing blunt cardiac injury (BCI). While the combination of electrocardiogram (ECG) and serum troponin is considered the most sensitive screening method for those being considered for discharge, little evidence exists regarding the diagnostic value and clinical utility of initial and serial serum troponin levels in stable patients who are being admitted with a possible BCI. Therefore, this study seeks to determine if troponin level is an independent predictor of adverse cardiac events in stable, admitted patients at risk for BCI.

METHODS: This was a five-year retrospective study using the trauma database at a University Level I Trauma Center. The study population included adult trauma patients presenting with a physician diagnosis of BCI or a sternal fracture who met prespecified stability criteria (SBP $\geq$ 90 mmHg, HR < 110bpm, shock index < 1, GCS $\geq$ 14). The data collected included all troponin values and ECG interpretations as well as echocardiogram interpretations, when performed. A patient was classified as having an adverse cardiac event if they were diagnosed with a new arrhythmia requiring treatment, had cardiac surgery, or suffered cardiac-related mortality. Sensitivity, specificity, positive and negative likelihood ratios, and diagnostic odds ratios were calculated to analyze the diagnostic performance of the di [...]

[PubMed](#) · [DOI](#)

Prevalence and presenting symptoms of intracranial hemorrhage in older adults presenting with subacute head injuries in a level-2 trauma center: A retrospective cohort study.

Raats JH, Schuijt HJ, Hoogervorst ELJ, Velde DV

INTRODUCTION: Due to our ageing population, the incidence of subacute head injuries such as chronic subdural hematoma is increasing. As current clinical decision rules regarding CT imaging are in context of acute head injuries, it remains unclear what presenting symptoms of subacute (>48 h after initial trauma) head injuries in older adults are relevant for clinical practice. To aim of this study is to describe the prevalence of traumatic intracranial hemorrhage in older adults presenting to the emergency department with subacute head injuries and identify symptoms associated with traumatic ICH.

METHODS: This retrospective cohort included patients > 65 years who underwent CT imaging for subacute head injuries in the emergency department of a single level-2 trauma center between 2020-2024. The primary outcome was traumatic ICH. Secondary outcomes included hospital length of stay, anticoagulation reversal, neurosurgical intervention, and 90-day mortality.

RESULTS: 374 patients were included, with a mean age of 79.1 (SD 6.7) and 170 (54.5%) were male. 62 (16.6%) patients had ICH, of which 40 (68%) were large lesions with mass effect or epidural hematoma. In patients with ICH, median hospital length of stay was 7 (IQR 4- 11) days, 8/18 (44%) anticoagulated patients received pharmacological reversal, 25 (40%) received neurosurgical intervention, and five (8%) died within 90 days. [...]

[PubMed](#) · [DOI](#)

The use of temporary intravascular shunting to improve resource utilization is safe and effective for penetrating peripheral vascular trauma.

Strobl HV, Coopwood CK, Bolaji T, McKinley WI, Hoefer LE, Polcari AM, Dowzicky PM, Zakrison TL et al.

BACKGROUND: Temporary Intravascular Shunting (TIVS) is well described in peripheral vascular trauma, typically for orthopedic fixation or damage control. Due to a high volume of penetrating trauma, our urban Level 1 trauma center began utilizing TIVS frequently to minimize resource constraints. Our hypothesis is that TIVS is not inferior to immediate repair.

METHODS: Penetrating peripheral arterial injuries were identified from the institutional trauma registry from 5/2018-8/2024. Patients were categorized as having undergone immediate repair (ImR) vs delayed repair (DR), following TIVS placement. The DR group was further divided into resource utilization (DR-RU) and damage control (DR-DC) subgroups. DR-RU included patients receiving < 6 units of packed red blood cells intraoperatively, with a final intraoperative lactate < 5.0 mmol/L, and no other body cavity explored. DR-DC included shunted patients not meeting these criteria. The DR-RU subgroup was compared to both ImR and DR-DC using univariable tests.

RESULTS: Of 228 patients included, 118 were in the ImR group, 47 in the DR-RU subgroup, and 63 in the DR-DC subgroup. Trauma surgeons performed 76% of the ImR, 99% of the TIVS and 97% of the definitive repairs in the DR group. Median shunt dwell time was 12.9 h (IQR 9.7-17.7) in the DR-RU group. When comparing ImR to DR-RU respectively, there was no statistically significant [...]

[PubMed](#) · [DOI](#)

Cranial CT scans rarely result in neurosurgical transfers in older adults over 80 years with acute head injuries: A retrospective cohort study.

Raats JH, Schuijt HJ, Kroes T, Hoogervorst ELJ, van der Velde D

PURPOSE: Due to an ageing population and low threshold for imaging in older adults with acute head injuries, the resulting number of cranial CT scans continues to increase. While current clinical decision rules aim to identify all intracranial traumatic injuries, the clinical significance of these findings in older adults remains disputed. This study describes traumatic findings on CT imaging and neurosurgical care transfers of older adults with acute head injuries.

METHODS: This retrospective cohort included patients ≥ 65 years who received cranial CT imaging for an acute head injury in a single level-2 trauma center, without neurosurgical care available, between 2020-2024. Intracranial hemorrhage was classified as a minor or major lesion (i.e., with or without mass effect). The primary outcome measure, transfer rate to neurosurgical care, was compared between age groups 65-80 and > 80. Secondary outcomes included hospital length of stay, anticoagulation reversal, 30-day readmission, and in-hospital and 90-day mortality.

RESULTS: In 4581 included CT reports, minor and major traumatic cranial lesions were found in 315 (6.9%) and 88 (1.9%) patients, respectively. The prevalence of traumatic findings was comparable between age groups. Five out of 45 patients aged > 80 with major lesions were transferred to neurosurgical care, compared to 15 out of 43 patients aged 65-80 (11.1% [...])

[PubMed](#) · [DOI](#)

Impact of "Kids' Court" on traffic infraction recidivism in Dar es Salaam: A novel approach to a troublesome problem.

Hopkins M, Reich J, Kalolo S, Swai N, Rweyendela AG, Stoffan A, Guerrero A

Road Traffic Injuries are a major problem in Sub-Saharan Africa. Recidivism after road traffic infraction has not previously been studied in this setting. This study evaluated a novel elementary school based program to reduce repeat minor traffic offenses near schools by having children serve as "judges" to rhetorically interrogate offenders. Offenders were randomly allocated into standard penalties or participation in Kids' Court (412 total) and followed via quarterly interrogation of a traffic police database for 1 year. While the number of repeat offenders was similar in both groups (88 vs. 85, $p = 0.84$), the total number of offenses was improved in the Kids' Court group and approached significance but did not achieve it (344 vs. 404, $p = 0.065$). The average time to first repeat offense was significantly decreased in the Kids' Court group (127 vs. 154 days, $p = 0.0019$). This program reduces the number of repeat offenses and lengthens the time to repeat offense. It likely has a yet unquantifiable long-term effect on the children that participate as "judges" in the program.

[PubMed](#) · [DOI](#)

Development and measurement properties of the Safety Threats and Adverse Events in Trauma (STAT) taxonomy: A critical review.

Nazir A, Shore E, Keown-Stoneman C, Grantcharov T, Nolan B, FIRST60 Investigators

BACKGROUND: Adverse events are common in trauma care, with most occurring during the initial resuscitation phase. The Safety Threats and Adverse Events in Trauma (STAT) taxonomy is a comprehensive tool that was created to evaluate both technical and non-technical errors in trauma resuscitations. The objective of this study is to critically appraise the development and measurement properties, including sensibility, reliability, and validity of the STAT taxonomy.

METHODS: A literature search was conducted in MEDLINE, Embase, and CINAHL, supplemented by citation searching. Studies describing the development, evaluation, or adaptation of the STAT taxonomy were included. Evidence was synthesized narratively across tool development, sensibility, reliability, and validity.

RESULTS: Seven studies met inclusion criteria, encompassing tool development, pilot testing, reliability assessment, and adaptation. The STAT taxonomy was developed using a systematic review and a RAND-modified Delphi process involving 22 trauma experts, resulting in 67 adverse events across nine domains. Early evidence supports strong content validity, with comprehensive coverage of both technical and non-technical domains. Feasibility was demonstrated across simulation and clinical trauma video review settings. Inter-rater reliability was consistently high, with agreement rates of 90.1% in simulation studies and [...]

[PubMed](#) · [DOI](#)

ED discharge after trauma transfer has increased over time: A single tertiary care center analysis.

Etra JW, Pfaff AC, Kaul S, Seshadri A, Berrigan M, Dombek M, Morris RE, Fleishman A et al.

OBJECTIVES: Regionalization of care and changes in hospital and provider networks have led to increased transfer of patients to an accredited trauma center. We aimed to identify trends in trauma transfers and injury patterns that are likely to result in tertiary center admission versus discharge from the emergency department (ED).

METHODS: All patients transferred from 2014 to 2021 to a single level I tertiary trauma center were identified. Using demographic, encounter, and diagnosis coding data from our trauma registry, specific injuries were stratified based on body system or type of injury. Numbers of transfers by year were analyzed using Wilcoxon rank sum test and chi-squared or Fisher's exact test. Univariable and multivariable logistic regression was used to identify injury patterns that correlated with tertiary hospital admission/intervention or discharge from the ED.

RESULTS: 9307 trauma transfer patients were included. On multivariable regression, thoracic or orthopedic injuries were less likely to result in discharge from the ED (OR 0.34, 95% CI 0.27-0.42; OR 0.33, 95%CI 0.28-0.39). GCS of < 14 was also associated with a decreased likelihood of discharge (OR 0.13, 95% CI 0.08-0.22). Soft tissue injuries, spinal injuries, and head injuries were more likely to be discharged from the ED without requiring admission (OR1.24, 95%CI 1.09-1.40; OR 1.68, 95%CI 1.43-2.00; OR [...])

[PubMed](#) · [DOI](#)

Implementation of a smartphone-based ultrasound program to improve timely diagnosis of life-threatening injuries in Cameroon: Results from a prospective, multisite feasibility study.

Rosenbloom L, Emeh AN, Tambe J, Haddad K, Marpha M, Maffo CJ, Mbebi EF, Driban M et al.

BACKGROUND: Undiagnosed hemorrhage is the leading cause of preventable trauma death in Cameroon, yet few injured patients receive imaging to diagnose hemorrhage. A smartphone-based ultrasonography (SBU) curriculum was developed to rapidly train frontline healthcare providers in Cameroon to perform and interpret eFAST. SBU has demonstrated promising educational efficacy, but its clinical feasibility in this technology-constrained setting is unknown. We evaluate the feasibility and acceptability of a SBU pilot at three trauma centers in Cameroon.

METHODS: We implemented a six-month feasibility pilot at three Cameroonian hospitals participating in the Cameroon Trauma Registry (CTR). Trauma providers were trained to perform SBU eFAST using a novel 5-hour curriculum, then asked to perform SBU on all injured patients as part of the trauma evaluation. Feasibility was assessed as the proportion of CTR patients with completed eFAST. Acceptability was assessed as the proportion of users rating SBU ≥ 68 on the previously validated System Usability Scale (SUS).

RESULTS: Trauma care providers completed SBU eFAST on 87% (608 of 701) of eligible patients, compared to a pre-study diagnostic imaging rate of 5.5% (38 of 648). Completion was highest at the rural and regional low-volume centers (100%, $n = 42$ and $n = 75$, respectively) and lower at the high-volume urban tertiary referral center (8 [...])

[PubMed](#) · [DOI](#)

Insurance-based disparities in traumatic brain injury length of stay and discharge dispositions: A retrospective cohort analysis of the national trauma data bank.

Strelko O, Spiro E, Lui A, Tarapore PE, Huang MC, Manley GT, Yue JK, DiGiorgio AM

BACKGROUND: Insurance-based disparities associated with outcomes in traumatic brain injury (TBI), including acute care discharge dispositions, remain understudied. This study aims to evaluate the associations between Medicaid versus private insurance (PI) on hospital length of stay (LOS) and discharge disposition in patients with TBI across severities.

METHODS: This is a retrospective multicenter cohort study utilizing data from United States trauma centers from the National Trauma Data Bank (NTDB) years 2014-2016. Inclusion criteria were: identified to have TBI based on ICD-9 codes. Exclusion criteria were: missing gender, race, Glasgow Coma Scale (GCS) score, or LOS; not admitted as inpatient to hospital; did not have Medicaid or PI; age < 14 or > 89 years; extracranial AIS= 6. Patients were grouped into those with Medicaid and those with PI.

RESULTS: Of 808,120 NTDB patients with TBI, 188,756 met inclusion criteria (Medicaid: $N = 58,867$, PI: $N = 129,889$). Overall, PI patients were older and more frequently female, with differences in racial distribution and TBI severity compared to Medicaid patients. On multivariable analysis, Medicaid patients demonstrated longer inpatient LOS across all TBI severities, corresponding to a 7.3% increase after mild TBI (95% CI: 6.4%-8.2%, $p < 0.001$), a 7.4% increase after moderate TBI (95% CI: 2.8%-12.2%, $p = 0.001$), and an 8.7% increase af [...]

[PubMed](#) · [DOI](#)

Early operative management in trauma: A nationwide comparison of time to surgery and survival at trauma centers and non-trauma centers.

Eriksson J, Mattsson A, Larsson E, Oldner A, Oelreich EV

BACKGROUND: Trauma is a leading cause of death and disability, both in Sweden and globally. Timely surgical intervention and level of care have been identified as important determinants of outcome, yet it is not known whether time to surgery differs between trauma centers and non-trauma centers in Sweden, or whether previously observed survival differences also apply to patients undergoing early operative management.

METHODS: This retrospective national cohort study included adult trauma patients (≥ 18 years) who underwent surgery within six hours of hospital arrival between 1 January 2019 and 21 March 2023. Data were obtained from Swedish national quality registries and administrative registers held by the National Board of Health and Welfare (Socialstyrelsen). The primary outcome was time from hospital arrival to initiation of surgery. Secondary outcomes were time to urgent surgery and 30-day mortality. Associations between trauma center status and outcomes were

examined using regression models adjusted for demographic and injury-related factors.

RESULTS: A total of 1129 adult trauma patients who underwent surgery within six hours of hospital arrival were included, the majority presenting with moderate to severe injuries; of whom 659 (58.4%) were treated at trauma centers and 470 (41.6%) at non-trauma centers. Patients treated at trauma centers had higher injury severity and [...]

[PubMed](#) · [DOI](#)

Turret truck-related head and neck injuries in a Japanese wholesale market: Clinical characteristics, risk factors, and the need for preventive measures.

Kushi K, Hifumi T, Kamijo E, Ito K, Shirasaki K, Ro B, Mochizuki T, Nonaka T et al.

BACKGROUND: Turret trucks are widely used to handle cargo in Japanese wholesale markets. Under Japanese law, helmet use is not mandatory for turret truck operators, which potentially increases the risk of head and neck injuries. This study aimed to describe the clinical characteristics, risk factors, and outcomes of turret truck-related head and neck injuries as the first reported analysis of this injury mechanism.

METHODS: We conducted a retrospective observational study of patients with turret truck-related head and neck injuries who were transported to St. Luke's International Hospital between January 2011 and June 2024. Demographic data, injury characteristics, and outcomes were collected. The primary outcome was the necessity for hospital admission, and the secondary outcome was functional status at discharge, measured using the modified Rankin Scale (mRS).

RESULTS: Of the 67 patients analyzed, 48 (72%) were turret drivers and 19 (28%) were non-drivers. The cohort was predominantly male (97%) with a median age of 64 years. Twenty-four patients (36%) required hospital admission, with a higher proportion among turret drivers (44%) than among non-drivers (16%). Among turret drivers, all patients with severe injuries (Injury Severity Score ≥ 16) and those with significant head and neck injuries (Abbreviated Injury Scale ≥ 3) required hospitalization. Despite injury severity, m [...]

[PubMed](#) · [DOI](#)

Emergency medical services response times to motor vehicle crashes increased in the USA over the period 1987-2020.

Dubois S, Gravelle P, Savage DW, Maxwell H, Bédard M

INTRODUCTION: The timing of Emergency Medical Services (EMS) notification, crash scene arrival, and hospital arrival may impact motor vehicle fatalities. We examined EMS response time intervals over the past three decades, considering the effects of weather, vehicles involved, time of day, and location.

METHODS: We used the Fatal Accident Reporting System to compute and describe annual (1987-2020) EMS response time intervals. This included total time (i.e., crash-to-hospital), as well as the intervals between four key timepoints: crash, crash notification, crash scene arrival, and hospital arrival. We examined the proportion of fatal crashes with total intervals under 60 min (i.e., the "golden hour"), and where the crash arrival-to-hospital interval was under 30 min (the "beneficial timeframe"). Additionally, analyses were stratified by crash factors including weather (poor/clear) number of vehicles involved (single/multiple), time of day (early morning/rest of the day), and location (urban/rural).

RESULTS: A total of 310,001 fatal crashes were analyzed. Between 1987-1994 total median response times ranged between 40 and 42 min. By 1999, intervals had increased to 45 min; elevated intervals were evident through 2009. By 2020, observed intervals had returned to 41 min. Paralleling this pattern, crashes with "golden hour" intervals decreased from 77.0% in 1987 to 72.4% in 2009 [...]

[PubMed](#) · [DOI](#)

All-terrain vehicle related urethral injuries: An evaluation of the National Trauma Database.

Hertz A, VanDyke M, Hudak S

BACKGROUND: U.S. Consumer Product Safety Commission reported that from 2018 to 2020 there were 94,700 off-highway vehicle accidents with 92% of those injuries related to all-terrain vehicles (ATV). An estimated 300 deaths were in children less than age 16. There has not been any previous evaluation of urethral injuries resulting

from ATV accidents. We sought to define the volume and demographics of ATV-related urethral injuries.

METHODS: Data was obtained from the National Trauma Database (NTDB) Trauma Quality Program over a 7-year period (2016-2022). ICD-10 codes were used to identify patients who sustained urethral injury in ATV-related accidents. Injury and patient related demographic data was reviewed. R: A Language and Environment for Statistical Computing was used for data review and statistical analysis. Data were presented and analyzed using standard statistical methods **RESULTS:** During this time frame, there was a total of 7679,101 injuries recorded in the NTDB. There was a total of 7398 urethral injuries (0.1%), with 181 (2.5%) of those being related to ATV-accidents. 90.6% of patients were male, and there was a median age of 30 (range 5-76, IQR 29). The median injury severity score (ISS) was 17 (range 4-50, IQR 15). There was no association between age and higher ISS ($r = 0.07$, $p = 0.36$). Passengers were more likely to be less than age 16 ($p = 0.001$). Interestingly, [...]

[PubMed](#) · [DOI](#)

Challenges and learning curve in adopting resuscitative endovascular balloon occlusion of the aorta for trauma patients: A retrospective multicenter study.

Kim K, Kang BH, Kim DH, Yu B, Chang SW, Jung PY, Heo Y, Kang WS

BACKGROUND: This study aimed to assess the learning curve of REBOA from the first case in trauma centers using cumulative sum (CUSUM) analysis.

MATERIALS AND METHODS: This study enrolled consecutive trauma patients who visited five trauma centers from December 2015 to December 2021. To monitor the effectiveness of REBOA, we performed risk-adjusted cumulative sum (RA-CUSUM) analysis for mortality due to exsanguination. For individual risk adjustment, we implemented the least absolute shrinkage and selection operator (LASSO) logistic regression model. We then calculated the CUSUM for: (1) the time from common femoral artery (CFA) access to confirmation of REBOA placement (RP-CUSUM), (2) the total occlusion time (OT-CUSUM), (3) door-to-balloon time (DB-CUSUM), and (4) the time from injury to admission (IA-CUSUM). To determine whether observed deviations were statistically significant, a V-mask was superimposed on the CUSUM curve.

RESULTS: A total of 251 patients were enrolled. The overall mortality rate was 67.7% (170/251), and the mortality rate due to exsanguination was 49.0% (123/251). The RA-CUSUM model was developed using a LASSO logistic regression approach. In three hospitals, the RA-CUSUM showed an improvement after 5-34 procedures. However, the RA-CUSUM showed fluctuations with deterioration in the other two hospitals. The RA-CUSUM fluctuations exceeded the V-mask contr [...]

[PubMed](#) · [DOI](#)

Venous thromboembolism burden in IVC injuries: Is there an optimal strategy in prevention?

Himmler A, Preston JD, Guindi A, Wagner V, De Leon Castro L, Fransman RB, Nguyen J, Smith RN et al.

INTRODUCTION: Venous thromboembolism (VTE) is a common complication in trauma patients; however, there have been limited studies on VTEs and the prevention thereof in patients with inferior vena cava (IVC) injuries. This study aimed to examine VTE incidence and the preventative efficacy of VTE prophylaxis regimens after operative IVC repair.

METHODS: A 12-year retrospective review was performed of all patients who presented with IVC injuries to an urban level 1 trauma center. A subgroup analysis of VTE incidence, prophylaxis regimen (i.e., prophylactic anticoagulation [PAC] and/or antiplatelet [AP]) efficacy, and other clinical variables was performed on patients who underwent a primary or patch IVC repair and survived > 72 h.

RESULTS: A total of 132 patients presented with IVC injuries requiring operative management, with 56% overall survival. Among the 66 patients who received primary or patch repair and survived > 72 h, 27% had a VTE during index hospitalization and/or readmission. VTEs occurred a median of 9 days post-injury, and the most common type of VTE was deep vein thrombosis, followed by IVC thrombosis and pulmonary embolism. There were no significant differences in any clinical variables compared between patients who developed VTEs during index hospitalization and those who did not, including VTE prophylaxis timing, hospital or ICU length-of-stay, and injury sever [...]

[PubMed](#) · [DOI](#)

Association between social vulnerability, urbanicity, and post-injury outcomes following nonfatal motor vehicle crashes.

Grossman MK, Rink AJ, Foley LD, Ott RL

BACKGROUND: Motor vehicle crashes (MVCs) are a leading cause of injury and death in the United States. Community-level factors, such as social vulnerability and urbanicity, have been associated with risk of death; less is known about how these factors impact nonfatal, post-injury outcomes. This study examined the association between social vulnerability and urbanicity with hospital length of stay (LOS) and hospital discharge disposition among MVC patients.

METHODS: Patients aged 18 years and older who were admitted to a Montana regional trauma center with a non-fatal injury following an MVC from 2016 to 2024 were included in the study. The CDC Social Vulnerability Index (SVI) was used to quantify social vulnerability at the census tract level and scores were divided into tertiles representing low, medium, and high vulnerability. Urbanicity was defined using RUCA codes based on patient residence. Generalized estimating equations with a binomial distribution were used to estimate the joint association between SVI and urbanicity with discharge disposition (home vs. facility) and prolonged LOS (≥ 7 days), controlling for injury severity, patient demographics, and comorbidities.

RESULTS: Of the 668 patients, 529 (79%) were discharged home and 179 (27%) had a prolonged LOS. Among metropolitan patients, higher SVI rankings were associated with increased odds of discharge home; patien [...]

[PubMed](#) · [DOI](#)

Integrated surgical management of forequarter lateral implosion injury: Technical considerations and early outcomes.

Chua SKK, Balasubramaniam S, Cole PA, Tan BY

Concomitant ipsilateral fractures of the clavicle, scapula, and ribs, termed forequarter lateral implosion injury, represent a severe but underrecognized injury pattern resulting from high-energy lateral shoulder trauma. While the surgical indications for isolated chest wall and shoulder girdle injuries are well described, guidance on the integrated management of this combined injury complex remains limited. We describe a reproducible multidisciplinary approach for the concurrent surgical fixation of clavicle, scapula, and rib fractures, illustrated through two cases of forequarter lateral implosion injury resulting from high-energy road traffic accidents. Preoperative planning incorporated CT three-dimensional (3D) reconstructions and patient-specific 3D printed models to facilitate pre-operative planning, with a multidisciplinary team involved for incision planning and fixation sequencing. Surgery was performed in a single setting with the patient in the lateral position, utilizing muscle-sparing approaches and a staged fixation strategy to address the clavicle, scapula, and ribs through coordinated exposures. Simultaneous osseous stabilization allowed restoration of the superior shoulder suspensory complex integrity and chest wall mechanics, enabling immediate postoperative shoulder mobilization and aggressive pulmonary rehabilitation. Both patients demonstrated early pain r [...]

[PubMed](#) · [DOI](#)

Palliative and end of life care in older major trauma - A point prevalence evaluation in England, Wales and Scotland.

Cole E, Jarman H, PECOT collaborators

INTRODUCTION: Traumatic injury in older people is a significant health burden with higher mortality rates than younger cohorts. Survival following older trauma may be complicated by the patients pre-injury state and clinical uncertainty. Timely identification of palliative and end-of-life care needs may be challenging for acute clinical teams, and treatment escalation planning is not routinely embedded in trauma care. This point prevalence snap-shot aimed to evaluate treatment escalation discussions and palliative/end of life care (EoLC) practice in older major trauma patients at a national level.

METHODS: A one-day point prevalence "flash-mob" audit was conducted across Major Trauma Centres (MTCs) and Trauma Units (TUs) in England, Wales and Scotland. All trauma patients aged ≥ 65 years in hospital were eligible for inclusion. Patients with and without treatment escalation plans (TEPs) and those on care pathways were analysed.

RESULTS: Data from 957 patients in 49 hospitals were included and median time from injury was 11 days (interquartile range 4-24). A TEP or equivalent was documented in 393 patients (41.0%). Among patients with a TEP, there were more aged > 85 years (165/393 (41.9%)), than in those without a TEP (167/564 (29.6%)), $p < 0.001$). Clinical frailty scoring was performed in 657 patients (68.6%), and where recorded, TEPs were associated with increased frailty (CF [...])

[PubMed](#) · [DOI](#)

Interhospital variation in highest-level trauma activation and its association with mortality: A 37-center cohort study of level I and II trauma centers in the US.

Orlando A, Mazumder H, Leichtle S, West MA, Joseph B, Kaafarani H, Hunt JP, Shen Y et al.

BACKGROUND: Trauma team activation protocols are critical for mobilizing resources in the care of severely injured patients. In the US, the American College of Surgeons (ACS) specifies minimum criteria for the highest-level (full) trauma activation (fTA), but hospitals retain discretion to add criteria, potentially leading to variability in activation practices and resource utilization. The extent of this variation and its impact on patient outcomes is unknown. The aim of this study was to quantify inter-hospital variability in fTA use and its relationship to mortality.

METHODS: We conducted a multicenter, retrospective cohort study of adult trauma patients treated at 37 Level I and II trauma centers across the United States from 2017 to 2019; transfers were excluded. Mixed-effects logistic regression models were used to quantify inter-hospital variability in fTA utilization and total mortality (death+hospice), adjusting for 12 patient and hospital-level characteristics. Correlation analyses assessed the relationship between adjusted hospital-specific fTA rates and adjusted total mortality.

RESULTS: Overall, 158,696 patients were included, with 34,374 (21.7%) receiving a fTA. The median age was 53 yrs, with 59% male, 71% White, 88% blunt, and a median Injury Severity Score of 9. Use of fTA varied widely (3.3% to 54.1%, median [IQR]=19.3% [13.6-27.3%]) and the adjusted odds of [...]

[PubMed](#) · [DOI](#)

Alternative trials of recipient site vessel in maxillomandibular reconstruction with multisegment fibular flaps including only one pedicle anastomosis.

Akbay E, Gülmez M■

BACKGROUNDS: The objective of this study is to closely examine the preferred anastomoses of the superior thyroid artery, facial artery, lingual artery, maxillary artery, and superficial temporal artery anastomoses for the reconstruction of maxillomandibular defects caused by firearm injuries. The study will also examine the reasons for choosing these arteries, their advantages and disadvantages, surgical techniques, complications, and their postoperatively clinically and scintigraphically assessable viability.

METHODS: The present study encompasses a cohort of ten patients who sustained maxillomandibular injuries and underwent surgical intervention employing a multisegment fibular flap. In all cases, following bone fixation, microvascular anastomoses were sutured under a microscope using 9-0 nylon sutures. Flap viability was initially assessed clinically; in addition, all patients underwent a bone scan on the fifth day after surgery using an intravenous infusion of Tc-99m methylene diphosphonate (MDP).

RESULTS: In the study, 10 patients who underwent maxillomandibular multisegment flap surgery received the most appropriate microvascular anastomosis for their pathology. Of these patients, the six most demonstrative cases, which best represent the relevant artery used for anastomosis, have been detailed.

CONCLUSIONS: Although the aim of this study was not to establish a defini [...]

[PubMed](#) · [DOI](#)

An epidemiological analysis of extracorporeal membrane oxygenation use in trauma.

Wallis MC, Rizzo JA, O'Neil ER, Sams VG, Schauer SG

BACKGROUND: Trauma is the leading cause of death in younger adults and children. Severe polytrauma predisposes patients to the failure of multiple organ systems, including the cardiovascular and respiratory systems, at times necessitating extracorporeal membrane oxygenation (ECMO). ECMO use in the setting of trauma is

increasing, yet little data exists describing current practice patterns. We performed an epidemiological analysis of ECMO use and outcomes in trauma patients.

STUDY DESIGN AND METHODS: We analyzed data from the Trauma Quality Improvement Program Registry from 2017 to 2023. Procedure codes were used to identify the application of ECMO. Pediatric patients were those < 18 years of age. We analyzed two groups from this data set: one including all patients who required ECMO and a separate group including only pediatric cases. We used descriptive and inferential statistical methods.

RESULTS: There were 8,014,737 encounters of which 1919 had documented ECMO use. Within that group, 224 were < 18 years of age. The median time from hospital arrival to the first initiation of ECMO was 44 h (interquartile range [IQR] 5-147). The incidence per year ranged from 1.9 to 2.9 events per 10,000 encounters. Survival ranged from 59% to 68% per year. The number of facilities with documented ECMO use annually ranged from 103 to 158 and overall increased during the time of the study. | [...]

[PubMed](#) · [DOI](#)

The role of palliative care in older persons trauma.

Baxter M

[PubMed](#) · [DOI](#)

Shock and Trauma Risk Index Nomogram (STRIN): A bedside physiology-based tool for early-high-acuity care triage in major trauma.

Saronni S, Mecca G, Rampin C, Agarossi A, Ripoll-Gallardo A, Colzani G, Cimbanassi S, Chiara O

BACKGROUND: Major trauma is a leading cause of mortality and permanent disability, especially in younger adults. Early identification of high-risk patients is critical for appropriate care. Tools like the Shock Index (SI) or New Trauma Score (NTS) enable rapid bedside assessment but show variable performance across populations.

OBJECTIVE: To develop and internally test the Shock Trauma Risk Index Nomogram (STRIN), integrating Shock Index (SI), Glasgow Coma Scale (GCS), oxygen saturation (SpO₂), and mean arterial pressure (MAP) for early risk stratification of time-critical resource use in major trauma.

METHODS: Data of a retrospective cohort (March 2022-December 2024) from a single level-1 centre retrieved from Lombardy registry was considered. Adult patients (≥16 years) with major trauma and complete vital signs were included. Composite outcome was emergency surgery (ES) within 6 h, massive transfusion (MT) (≥10 units of packed red blood cells within 24 h or ≥4 units within first hour), Intensive Care Unit (ICU) admission and interventional radiology procedures (IR). The model was developed using multivariable logistic regression with Restricted Cubic Splines. Performance was assessed through discrimination (AUROC), calibration (Hosmer-Lemeshow test, Brier score), and internal testing via bootstrap resampling. Comparative performance against SI and NTS included AUROC, Net Re [...]

[PubMed](#) · [DOI](#)

Predictors of traumatic brain injury amongst secondary school students in England: A retrospective cohort study using electronic health records.

Benavente MT, Wilson R, Jackson J, Birnie K, Mytton J, Ijaz S, Booker M, Burrell A et al.

BACKGROUND: Traumatic brain injury (TBI) in children and adolescents is a leading cause of disability and mortality, with long-term health-related consequences. There is little evidence describing the children and adolescents most at risk of TBI.

OBJECTIVE: To identify the demographic and clinical predictors of TBI in secondary school-aged children in England.

PARTICIPANTS AND SETTING: Linked Clinical Practice Research Datalink and Hospital Episode Statistics Admitted Patient Care data were used to identify patients aged 11-18 years registered with a GP surgery in England between 2013 and 2021.

METHODS: Multivariable Cox regression was used to assess the association between demographic and clinical risk factors and time to medically attended TBI.

RESULTS: The analytical sample included 402,249 children, 2.3% of which had a TBI presenting to primary or hospital care when aged 11-18 years. In the fully adjusted model, increased risk of TBI was associated with being male; less socioeconomically deprived; having a history of fracture, abuse, depression, or previous TBI; having two or more previous GP visits, having more previous Emergency Department visits; and having fewer hospital admissions.

CONCLUSION: Using a nationally representative dataset we were able to identify which children were most at risk of TBI in their secondary school years. TBI is often preventable and targ [...]

[PubMed](#) · [DOI](#)

Delivering psychosocial care and support in traumatic injury follow-up: A qualitative study of clinician's experiences.

Wake E, Ranse J, Marshall AP

INTRODUCTION: The purpose of trauma follow-up care after hospital discharge is to support patients with serious injuries recover to their fullest extent. Whilst the physical health of patients is a central focus of follow-up care, psychosocial support is described as challenging and inadequate, with few trauma follow-up services able to provide integrated psychosocial support inclusive of routine access to psychosocial experts. As trauma follow-up care within Australia and New Zealand (Australia and New Zealand) is predominantly provided by nursing and medical trauma clinicians, exploring how psychosocial care and support is provided by these clinicians is warranted. The broad aim of this qualitative study was to explore trauma clinicians' experiences in providing trauma follow-up care with this paper focusing on the provision of psychosocial care and support.

METHODS: Semi-structured interviews were used to explore the experiences of trauma clinicians in providing follow-up care after hospital discharge to patients who have experienced serious injury. Twenty participants were purposively selected from public hospitals throughout Australia and New Zealand who provided trauma follow-up care after hospital discharge. Qualitative data were transcribed verbatim and analysed using inductive content analysis.

RESULTS: Psychosocial care and support provided by trauma service clinici [...]

[PubMed](#) · [DOI](#)

Diagnostic criteria and clinical course of inappropriate antidiuresis and cerebral salt wasting syndrome following traumatic brain injury: A retrospective cohort of 351 severe trauma patients.

Laurent C, Deffontis L, Bouchdoug K, Dagod G, Girard M, Festoc M, Capdevila X, Charbit J

BACKGROUND: Hyponatremia is common in traumatic brain injury (TBI) population, and is associated with poor outcomes. The main mechanisms are the syndrome of inappropriate antidiuresis (SIAD) and cerebral salt wasting syndrome (CSW). This study aimed to assess hyponatremia prevalence and time of onset in patients with TBI and assess differences between SIAD and CSW.

METHOD: This retrospective cohort study was conducted between 2015 and 2018 in our level 1 trauma center. Patients admitted to the intensive care unit with TBI were included. Three subgroups were determined using urinary clinical and biological criteria: SIAD, CSW, and Undetermined. Predictive factors were assessed for each subgroup, especially the influence of free water, sodium, or fluid intake.

RESULTS: Among 351 trauma patients with TBI, 57 (16 % [95 %CI 12 %20 %]) developed hyponatremia within 30 days. 30 (9 % [95 %CI 6 %12 %]) developed a SIAD, 13 (4 % [95 %CI 2 %6 %]) a CSW, 14 (4 % [95 %CI 2 %6 %]) an undetermined form. CSW subgroup had higher Simplified Acute Physiology Score II (SAPS II), Injury severity score ISS, more cerebral injuries, intracranial pressure monitoring, mechanical ventilation than the SIAD subgroup. Fluid, sodium and free water intake in the 48 h preceding hyponatremia did not influence CSW or Undetermined hyponatremia development, whereas excessive free water intake in the 48 h precedi [...]

[PubMed](#) · [DOI](#)

Video review analysis of early common femoral arterial access in trauma: Can we identify occult shock?

Murali S, Cannon J, Joergensen S, Zone AI, Forsythe L, Brinson M, Chreiman K, Abella BS et al.

BACKGROUND: Early common femoral artery (CFA) access in trauma resuscitation has the potential to improve hemodynamic monitoring and facilitate interventions. However, data on its utilization and impact are limited. This study aimed to collect objective data on early CFA access.

METHODS: We conducted a prospective observational study using trauma video review at a Level 1 trauma center. Critically injured trauma patients were included based on predefined criteria. Video and chart review were used to collect data on patient demographics, injury characteristics, procedural details of CFA access (including time of procedural milestones), and hemodynamic measurements. Trauma team leaders were also surveyed to determine their impressions on early arterial access.

RESULTS: Among 72 patients, 34.7% underwent early CFA access. Early CFA access was associated with blunt mechanism ($p < 0.001$), lower presenting GCS ($p = 0.006$), and worse functional status at discharge ($p = 0.028$). The median time from arrival to visualizing an arterial waveform was 16.5 min (IQR 13.2, 26.2). Ultrasound improved first-pass success rates (median attempt number: 1 vs 2, $p = 0.030$). Noninvasive systolic blood pressure measurements were, on average, 13.5 mmHg higher than invasive measurements; the discrepancy was more pronounced in patients with arterial SBP < 90 mmHg, and 9 paired measurements revealed occu [...]

[PubMed](#) · [DOI](#)

Outcomes and safety of selective non-operative management of penetrating liver injuries: A prospective study.

Kapnias K, Kivilo M, Saar S, Talving P, Navsaria P

BACKGROUND: Highly selective nonoperative management (SNOM) of penetrating liver injuries is increasingly implemented, however, still underutilized. This ambivalence is derived from the anxiety that hollow viscus injury may be missed, especially after gunshot wounds (GSWs). As a result, exploratory laparotomy is routinely performed in patients with penetrating abdominal trauma (PAT), frequently resulting in both negative and non-therapeutic laparotomies. The aim of this study was to assess the safety of nonoperative management (NOM) of penetrating liver injuries.

PATIENTS AND METHODS: A prospective study including patients with penetrating liver injuries admitted to a high-volume level I trauma centre was conducted over a 52 - month period (April 2015 - July 2019). Patients suitable for SNOM included: 1) penetrating injury to the right upper quadrant or right thoracoabdominal area 2) absence of peritonism 3) hemodynamic stability 4) GCS > 13 and no spinal cord injury 5) evidence of liver injury and absence of hollow viscus organ injury on CT scanning. Patients presenting with peritonism, hemodynamic instability, evisceration, or unreliable physical examination underwent immediate laparotomy. Primary outcome was failure of NOM and mortality.

RESULTS: Overall, 243 patients had liver injury and met inclusion criteria. Immediate laparotomy (IL) and SNOM groups included 168 (68 %) [...]

[PubMed](#) · [DOI](#)

Epidemiology of ranger injuries among members of the Game Rangers Association of Africa: Service-related risk and the call for targeted prevention strategies.

Viljoen C, van Mechelen W, Sykes L, Clark J, de Bruin L, Mushonga T, Rensburg DCJV

OBJECTIVES: To determine the epidemiology of ranger injuries, risk factors for injury, and health support needs among members of the Game Rangers Association of Africa (GRAA).

METHODS: We employed a retrospective cross-sectional design to collect data online using a questionnaire developed specifically for this study. We used convenience sampling to include 120 rangers aged 18 years or older who were actively working in Africa.

RESULTS: Our findings showed a career injury prevalence of 52 % and period prevalence (past 12 months) of 33 %. The lower limb was the most frequently injured anatomical region (38 %), followed by the upper limb (31 %) and trunk (19 %), specifically affecting the hand (19 %), ankle (14 %), and foot (13 %). The most common injury types included skin lacerations (29 %), followed by muscle injuries (23 %), and fractures (10 %). Working for more than 15 years as a ranger is associated with an increased risk for career injury (OR=2.63) compared to working for 10 years or less. Injuries resulted in a median of 7 days (IQR=0-20 days) lost from work, with most injuries requiring either emergency medical care (31 %) or consultation with a medical doctor (26 %). Rangers need better

equipment, cardiovascular fitness training programs, safety training, and mental health support.

CONCLUSION: One in every two rangers associated with the GRAA sustains an injury acro [...]

[PubMed](#) · [DOI](#)

The effectiveness of an early management guideline for older patients following blunt thoracic injury.

O'Donohoe RB, Mathew J, Winter N, Fitzgerald M, Hendel S, Dockrell L, Lodge M, Hooper A et al.

BACKGROUND: Thoracic injury is a leading cause of trauma related morbidity and mortality worldwide. Older patients (aged ≥ 65 years) who have sustained rib fractures following blunt chest wall trauma are at greater risk of serious pulmonary complications, prolonged hospital stay, and death when compared with the general population. The Blunt Chest-Wall Injury and Rib Fracture Pathway (BCWP) was introduced at our institution in July 2021 and advocates for early, aggressive analgesia, specialist support from the Acute Pain Service as well as consideration of regional techniques and Intensive Care Unit admission.

METHODS: A retrospective review of the prospectively maintained Alfred Hospital Trauma Registry identified all patients aged ≥ 65 years admitted to our Level 1 Adult Trauma Centre following blunt chest wall injury between the 1st of January 2019-1st of January 2024. The primary outcome measure was proportion of patients receiving an adequate analgesia (AA) regimen. Secondary outcomes included in-hospital mortality and hospital LOS.

RESULTS: 2488 patients were admitted during the 5-year study period, with 1168 admitted prior and 1320 post BCWP introduction. Mean age was 78.3 (SD 8.4) and 78.8 (SD 8.3) while most patients [57.0 % (n = 666) & 56.3 % (n = 743)] were male. There was a significant difference in the proportion of patients who received AA, with 87.7 % (n = 1158) [...]

[PubMed](#) · [DOI](#)

Posterior interosseous artery flap for severe hand injuries: Outcomes of reconstruction combined with local and regional flaps.

Ertem H, Adıgüzel F, Özde H, Aslantürk O, Ertem K

BACKGROUND: Severe hand injuries with extensive soft-tissue loss present a significant reconstructive challenge. Achieving stable coverage while preserving hand function often necessitates the combined use of regional and local flaps. The posterior interosseous artery (PIA) flap, in combination with regional flaps harvested from non-salvageable digits, may offer an effective solution.

METHODS: A retrospective analysis was performed on patients with severe hand injuries who underwent reconstruction using a PIA flap between 2022 and 2025. Patients were treated with either an isolated PIA flap or a PIA flap combined with local flaps, including fillet flaps harvested from non-salvageable digits or rotational flaps, depending on the extent and location of the defect. Demographic characteristics, injury mechanisms, defect locations, and surgical details were recorded. Postoperative complications and functional outcomes were assessed using fingertip-to-palm distance, Quick DASH score, and VAS for pain.

RESULTS: Fourteen patients with severe hand injuries were included in the study. The mean age was 39.4 years, and the mean follow-up period was 19.7 months. Five patients with complex, multi-site defects underwent combined reconstruction using a PIA flap with local flaps (four with fillet flaps, one with a rotational flap), while nine patients were treated with an isolated PIA flap. S [...]

[PubMed](#) · [DOI](#)

Trends and outcomes following diagnostic laparoscopy for blunt abdominal trauma in the United States.

Atkins KK, Gallaher JR, Schneider AB, Charles AG

BACKGROUND: The role of diagnostic laparoscopy in adults with blunt abdominal trauma and the effect of negative laparoscopy on mortality is not well delineated.

METHODS: We reviewed the National Trauma Data Bank (2007-2019) for adults sustaining blunt abdominal trauma who underwent operative intervention. We performed a doubly robust, augmented inverse propensity weighted multivariable logistic regression to estimate the effect of a negative diagnostic laparoscopy on mortality in

adults with operative blunt abdominal trauma.

RESULTS: 87,864 patients met the inclusion criteria. Diagnostic laparoscopy occurred in 6.6% (n = 5816) of patients, with a 21.1% (n = 1226) conversion to laparotomy rate. The rate of negative diagnostic laparoscopy was 28.6% (n = 1665). Negative laparoscopy patients had a 49% reduction in odds of mortality (OR 0.51, 95%CI 0.47 - 0.56, $p < 0.001$) compared to negative laparotomy patients. Patient's that underwent laparoscopy, found to have intra-abdominal injury, had a similar reduction in odds of mortality compared to negative laparotomy patients (OR 0.54, 95% CI 0.51 - 0.57, $p < 0.001$).

CONCLUSION: Diagnostic laparoscopy may be safe for adults with blunt abdominal trauma and prevent significant morbidity and mortality from a negative laparotomy.

[PubMed](#) · [DOI](#)



European Journal of Trauma & Emergency Surgery

Volume 52, Issue 1 · April 2026 · 120 new articles

Impairment in activities of daily living and textbook outcome after major emergency abdominal surgery.

Mima K, Nitta H, Shikiji Y, Nakashima R, Uemura S, Morinaga T, Itoyama R, Umezaki N et al.

PURPOSE: Textbook Outcome (TO) has been a useful tool to assess benchmark patient-centred surgical outcomes in elective abdominal surgery. This study aimed to evaluate the association of preoperative activities of daily living (ADL) status with TOs following major emergency abdominal surgery.

METHODS: This retrospective study included 301 consecutive patients who underwent major emergency abdominal surgery for high-risk acute abdominal conditions, including perforated peptic ulcer, intestinal ischemia, bowel obstruction, bowel perforation, and severe intraabdominal bleeding or infection, at Kumamoto Regional Medical Center between April 2019 and March 2025. Preoperative ADL was evaluated using the Barthel Index (range, 0 to 100), with higher scores indicating greater independence. Barthel Index ≤ 85 was defined as ADL impairment. According to a recent international consensus, TO was defined as discharge to "home" without major postoperative complications (Clavien-Dindo grade $\geq$ III). Multivariable logistic regression analysis was conducted to assess the association of preoperative ADL with TO achievement, adjusting for clinical features and surgery-related factors.

RESULTS: Among the 301 patients, 90 (30%) had preoperative ADL impairment (Barthel Index ≤ 85) and 211 (70%) achieved TO. Multivariable analysis revealed that preoperative ADL impairment was independently associated [...]

[PubMed](#) · [DOI](#)

Early calcium replacement after minimal blood transfusion is associated with adverse outcomes.

Hewgley WP, Griffin RL, Baird E, Hashmi ZG, Baker SJ, Jansen JO, Kerby J, Holcomb JB et al.

PURPOSE: Calcium disturbances are common in patients with traumatic injuries, exacerbated by shock and blood product resuscitation. Recently, empiric calcium supplementation has been recommended with the first unit of blood transfusion, despite a lack of data supporting this practice. We hypothesize that there are risks associated with calcium administration in patients receiving small volume transfusion.

METHODS: This is a retrospective review of patients admitted to our Level 1 trauma center from 2018 to 2024 who received 1-3 units of red blood cells (RBCs). Patients were stratified by calcium levels and whether they received early calcium supplementation. A logistic regression was used to estimate odds ratios for the association between calcium levels, calcium supplementation, and outcomes. The primary outcome was in-hospital mortality. Secondary outcomes included acute kidney injury, myocardial infarction, acute respiratory distress syndrome, and venous thromboembolism. The model was adjusted for age, sex, ISS and shock index.

RESULTS: A total of 2,680 patients were included. When stratified by calcium levels and supplementation status, patients were similar in terms of age, sex, mechanism and injury severity score, but shock index was higher among

patients receiving calcium supplementation. After adjustment, hypocalcemic patients who received calcium supplementation had [...]

[PubMed](#) · [DOI](#)

Peripheral perfusion index as a predictor of severe shock in trauma patients: a prospective observational study.

Sundar H, Purushothaman V, Mathuram SR, Chandy GM, Sharma SL, Nayak S, James JD

BACKGROUND: Early recognition of haemorrhagic shock is critical in trauma care, particularly in low- and middle-income countries where delayed referral is common. Conventional vital signs may remain normal despite ongoing hypoperfusion. The Perfusion Index (PI), derived from pulse oximetry, is a non-invasive and continuously available marker of peripheral perfusion. This study evaluated the ability of PI to identify severe haemorrhagic shock in trauma patients and compared its performance with the Shock Index (SI) and Modified Shock Index (mSI).

METHODS: This prospective, single-centre study was conducted at a Level 1 trauma centre between January and May 2024. Adult Priority 1 trauma patients were enrolled and classified as having no/mild shock (ATLS grades 1-2) or severe shock (grades 3-4). PI was recorded during initial assessment. SI, mSI, lactate, injury severity, transfusion requirement, ICU admission, and 24-hour mortality were analysed. Receiver operating characteristic (ROC) analysis and multivariable logistic regression were performed.

RESULTS: A total of 120 patients were included (mean age 40.4 ± 17.2 years; 90% male), predominantly with blunt trauma (85% road traffic injuries). Severe shock was present in 35 patients (29.2%). Median PI was significantly lower in severe shock compared with no/mild shock (0.54 [IQR 0.28-1.31] vs. 1.33 [0.71-2.16], $p = 0.008$). Mean [...]

[PubMed](#) · [DOI](#)

A prospective study on abdominal trauma: patterns of injury, clinical presentation, organ involvement, associated injuries, and outcomes at a tertiary hospital in Mogadishu, Somalia.

Mohamud AA, Hassan AI, Ali AK, Jubur AA, Ahmed MR, Mohamed SS, Mohamed MM, Bashir AM

BACKGROUND: Trauma is a major global public health problem and one of the leading causes of death and disability worldwide, particularly among individuals younger than 44 years of age. Approximately 6 million people die each year as a result of traumatic injuries, with nearly 90% of these deaths occurring in low- and middle-income countries. Previous epidemiological studies have reported postoperative complication rates ranging from 18% to 42% and mortality rates between 8.5% and 13%.

OBJECTIVE: To evaluate the patterns of injury, clinical presentation, organ involvement, associated injuries, management strategies, and outcomes among adult patients with abdominal trauma in Mogadishu, Somalia.

METHODS: A prospective observational study was conducted from August 2023 to July 2024 at Mogadishu Somali-Türkiye Recep Tayyip Erdoğan Training and Research Hospital. A total of 146 adult patients with abdominal trauma were enrolled. Data were collected using a pretested structured questionnaire and analyzed using Stata version 16. Univariable and multivariable logistic regression analyses were performed to identify factors independently associated with in-hospital mortality.

RESULTS: The study population consisted predominantly of young adult males, with a mean age of 30.18 ± 9.2 years, and males accounted for 97.3% of all patients. Penetrating trauma was the predominant mechanism of [...]

[PubMed](#) · [DOI](#)

Comparison of fluoroscopic versus electromagnetic distal locking in humeral shaft osteosynthesis using intramedullary nail: a retrospective study.

Madeja R, Szeliga J, Voves J, Pometlova J, Konde A, Janosek J, Dousa P

PURPOSE: Intramedullary nailing is a standard method for osteosynthesis of diaphyseal humeral fractures. Distal locking, however, typically requires fluoroscopic guidance, which increases radiation exposure. This study aimed to compare standard fluoroscopy-guided distal locking with the Ezy-Aim system (Austofix, Australia) based on electromagnetic pulse detection from the perspectives of surgical times, radiation exposure, and complications.

METHODS: A retrospective study included 58 patients treated for diaphyseal humeral fractures (AO/OTA 12A, 12B) between 2019 and 2023. Group A (n = 28) received osteosynthesis using the Ezy-Aim system; Group B (n = 30) underwent fluoroscopy-guided distal locking. Data on operative time, fluoroscopy time, radiation dose, and complications were analysed.

RESULTS: In all, distal locking was not successful at the first attempt in 14.3% (4/28) of screws placed with Ezy-Aim and 20.0% (6/30) when fluoroscopy was used. Time and dose analyses were performed on the remaining 48 patients. The total surgical time was significantly longer with the Ezy-Aim system (median 58 min) compared to fluoroscopy (median 52 min, $p = 0.034$), primarily due to the time required for the device assembly. However, distal screw placement was faster with Ezy-Aim (median 12 min vs. 14 min, $p = 0.019$). Radiation exposure was significantly lower in the Ezy-Aim group (median [...])

[PubMed](#) · [DOI](#)

Patterns of patient worry following major emergency abdominal surgery: a 180-day follow-up study.

Parner MK, Jensen LR, Gamst-Jensen H, Burcharth J, Kokotovic D

PURPOSE: Major emergency abdominal surgery is associated with high postoperative morbidity and mortality. Beyond physical complications, patients experience psychological distress. This study aimed to describe the distribution of patient-reported worry across independent cross-sectional samples and its association with Days Alive and out of Hospital (DAOH) over 180 days postoperatively.

METHODS: Prospective cohort at Copenhagen University Hospital Herlev (March 30, 2023-March 31, 2024). Degree of Worry (DOW) was surveyed at discharge and postoperative day (POD) 30, 90, and 180, categorized as physical, mental, and physical-generic domains of worry. DAOH was computed cumulatively and incrementally for 0-30, 30-90, and 90-180 days. Associations between high DOW (hDOW) and subsequent DAOH were tested with Mann-Whitney U.

RESULTS: hDOW was reported by 36.1% at discharge, 22.4% at POD 30, 45.0% at POD 90, and 32.0% at POD 180. Across 180 days, the most frequent worries were relapse (21.3%), disease (14.1%), functional level (12.0%) and daily living (9.1%). The proportion of physical worries was significantly higher at POD 30 than at discharge and significantly lower at POD 180; mental worries varied between time points without statistically significant differences. hDOW at POD 30 was associated with fewer cumulative DAOH by POD 90 ($p = 0.008$).

CONCLUSION: Patient-reported worry f [...]

[PubMed](#) · [DOI](#)

Outcomes and complications of metallic vs. bioabsorbable screw fixation for displaced medial epicondylar fractures in pediatric and adolescent patients.

Jhala T, Scherer S, Esser M, Lieber J, Fuchs J, Dietzel M

PURPOSE: The use of bioabsorbable screws in orthopedics and trauma surgery is increasing. The major advantage of not having to remove the implant is of particular interest in pediatric surgery. Due to drawbacks, most notably implant fracture, bioabsorbable screw osteosynthesis has not yet surpassed the gold standard of metallic screw osteosynthesis. This single-centre study compares metallic screw and bioabsorbable screw osteosynthesis of medial humeral epicondyle fractures in children and adolescents over a two-year period.

MATERIALS AND METHODS: Retrospective analysis of surgically treated (screw fixation with either bioabsorbable magnesium-based or metallic screw) patients with a medial humeral epicondyle fracture between 2020 and 2022 at the author's institution.

RESULTS: Seventeen patients with medial epicondyle fractures underwent surgical treatment using either bioabsorbable screws (n = 8) or metallic screws (n = 9). Baseline characteristics, surgical approach, and postoperative care were similar between groups. Two complications (Clavien-Dindo I) occurred in the bioabsorbable group due to implant failure (screw breakage), though without clinical sequelae. These implant failures led to a significantly increased number of postoperative radiographs in the bioabsorbable group (median 4.8 vs. 2.5; $p = 0.014$). Functional outcomes (LOM, MEPI) and follow-up durations were com [...]

[PubMed](#) · [DOI](#)

Characteristics and temporal trends of mobility scooter accidents in the Netherlands: a multicenter cohort study.

Stensen Z, Peters A, Keereweer D, Krul I, van Osch F, de Groot B, Barten D

PURPOSE: Mobility scooters are increasingly used by older individuals. The aim of this study was to assess the characteristics and temporal trends of mobility scooter accidents (MSAs) in the Netherlands over a 10-year period using a representative dataset derived from fourteen Emergency Departments (EDs).

METHODS: In this retrospective multicenter cohort study, we analyzed MSA-related ED visits between January 2014 and December 2023, using the Dutch Injury Surveillance System (DISS). Information about age, sex, hospital admission, type and location of injury, mechanism of injury, and operational errors was analyzed. Injury severity was classified using the maximum abbreviated injury score (MAIS). Trend analyses were only performed for ED visits of patients with MAIS2+ injuries.

RESULTS: In total, 2,702 MSA-related ED visits were identified. Forty-nine percent of patients was male and the median age was 75 years (IQR 64.0-83.0). The most common mechanism of injury was a mobility scooter rollover (34.2%). From 2014 to 2023, the number of MSA-related ED visits of patients with MAIS2+ injuries increased by 38.0%. In the 70-79 age group, this rise was 91%. The total number of traumatic brain injuries (TBI) rose steeply by 103.0%, which includes a modest (+34.0%) increase in moderate-severe TBI. The hospital admission rate was 30.1%, with a median length of stay of 7 days (IQR 3.0 [...])

[PubMed](#) · [DOI](#)

"Is the effect of intraoperative 3D imaging beneficial in plate fixation of distal radius fractures: a matched-pair analysis".

Zumsteg JS, Pieringer A, Hinkelmann MA, Isenmann K, Pape HC, Allemann F, Barthel C

BACKGROUND: Distal radius fractures (DRFs) are among the most common adult fractures and are frequently treated with volar plate fixation. While intraoperative 3D imaging has been proposed to improve detection of malreduction and implant malposition, its impact on clinical and radiological outcomes remains unclear. This study aimed to evaluate whether intraoperative 3D imaging provides advantages over conventional 2D fluoroscopy. Outcome criteria were accuracy of fracture reduction, plate positioning relative to the watershed line, complication rates, operative time, and functional outcomes following volar plate fixation of distal radius fractures.

METHODS: This retrospective single-centre study included patients aged ≥ 16 years who underwent volar plate fixation for distal radius fractures between 2016 and 2024. Patients were assigned to a 3D imaging group or a conventional 2D fluoroscopy group. To ensure comparability, patients were matched 1:1 using exact AO/OTA subgroup matching and propensity score matching for age and sex. Outcomes included operative duration, radiographic reduction quality, plate positioning, complications, and wrist range of motion. Matched-pair comparisons were performed using appropriate paired statistical tests.

RESULTS: Of 1,259 patients screened, 418 patients (209 matched pairs) were included after exact AO/OTA subgroup matching and propensity score matching [...]

[PubMed](#) · [DOI](#)

High-velocity gunshot chest trauma: tailoring patient care through insights from a mass casualty experience.

Azulay AA, Stein M, Tabo LY, Lebenthal A, Ruderman L, Refaely Y

OBJECTIVES: This study describes the management strategies and clinical outcomes of a cohort of patients sustaining high-velocity military rifle gunshot wounds to the thorax during a mass casualty incident (MCI). The analysis focuses on damage control surgical techniques, the incidence and timeline of venous thromboembolism (VTE), regional pain management, and early psychiatric protocols.

METHODS: A descriptive retrospective analysis was conducted on patients presenting with a single penetrating high-velocity ballistic wound to the thorax on October 7, 2023. Inclusion criteria were limited to patients with a projectile trajectory traversing the lung parenchyma, managed by the cardiothoracic surgery department. Injuries to immediately adjacent structures (subclavian vessels, brachial plexus, clavicle, and scapula) were recorded as associated injuries. Patients presenting with multi-system trauma involving entirely separate, distant anatomical

regions were excluded.

RESULTS: The cohort comprised 11 male patients (age range: 19-49 years; 9 soldiers, 2 civilians) injured by 7.62 mm assault rifle rounds. Six patients (54.5%) required emergency thoracotomy, sternotomy, or VATS within the first 24 h due to ongoing hemorrhage; primary hemorrhage control consisted of deep parenchymal lung suturing (n = 5) and temporary chest packing (n = 1). Four patients (36.4%) underwent delayed, el [...]

[PubMed](#) · [DOI](#)

Elevated risk of wound and systemic complications in MRSA-colonized patients undergoing lower extremity orthopedic trauma surgery.

Areti AS, Hafeez H, Sangineni PV, Argao L, Deveza L, Dawson J

PURPOSE: Preoperative colonization with methicillin-resistant *Staphylococcus aureus* (MRSA) is a known risk factor for adverse outcomes in arthroplasty but remains under-investigated in orthopedic trauma. This study evaluates the impact of preoperative MRSA colonization on postoperative complication rates in patients undergoing lower extremity orthopedic trauma surgery.

METHODS: We conducted a retrospective cohort study using the TriNetX Research Network, identifying adult patients who underwent lower extremity orthopedic trauma procedures over a 20-year period. Patients with a documented MRSA diagnosis within three months before surgery were matched 1:1 to MRSA-negative controls based on demographics and comorbidities. Postoperative outcomes were assessed within three months of surgery, including surgical site infections (SSIs), wound dehiscence, organ dysfunction, thromboembolic events, and healthcare utilization. Odds ratios (ORs) and p-values were calculated, with statistical significance set at $p < 0.05$.

RESULTS: A total of 1,868 MRSA+ and 629,342 control patients were identified. After matching, MRSA+ patients remained at significantly increased risk for wound dehiscence (OR = 1.80, $p = 0.005$), pneumonia (OR = 2.33, $p < 0.001$), sepsis (OR = 3.71, $p < 0.001$), acute renal failure (OR = 1.45, $p < 0.001$), blood transfusion (OR = 1.74, $p < 0.001$), deep vein thrombosis (OR = 2 [...])

[PubMed](#) · [DOI](#)

A global contrast shortage: a descriptive study exploring IV contrast utilization and outcomes in trauma patients at a level I trauma center.

Zakaria H, Johansen C, Waldrop C, Wan B, Graham C, Fetzer D, Minei J, Bruns B et al.

PURPOSE: Computed Tomography (CT) with intravenous (IV) contrast is critical to the evaluation of the trauma patient. A pandemic-related lockdown occurred in Spring 2022 in the city where iohexol iodinated contrast media is manufactured, resulting in a global shortage. We explored the potential impact of this shortage on contrast utilization and patient outcomes.

METHODS: A retrospective study was performed of all trauma patients who underwent CT between January and December 2022 at a level I trauma center. Pre-contrast conservation (PRE) period was defined as January to April 2022, and post-conservation (POST) as May to December 2022. Demographics, utilization rates (bolus/patient) and outcomes were evaluated. Chi-square tests were performed for categorical variables and Mann-Whitney U tests for continuous variables. Statistical significance was set at $p < 0.05$.

RESULTS: 1,097 patients were included; 509 in the pre-shortage period, 588 post-shortage. 857 CTs with contrast (1.68 contrast bolus/patient) were performed pre-conservation and 979 post-conservation (1.66 contrast bolus/patient) ($p = 0.04$). Maximum Abbreviated Injury Scale (AIS) head, head/neck, and extremity were higher post-shortage ($p < 0.05$); however, mortality was higher in the pre-conservation group ($p < 0.0012$). There were 9 cases of Acute Kidney Injury (AKI, 8 PRE, 1 POST), 22 delayed diagnoses (14 PRE, 8 P [...])

[PubMed](#) · [DOI](#)

Trauma stabilization points in armed conflict: operational evolution and strategic reflections from mosul to Gaza.

Marta C, Luca P, Kai-Hsun H, Areej AF, Athanasios G, Wessam EG, Jonas Z, Johan VS et al.

BACKGROUND: Forward trauma care remains a critical gap in emergency responses to armed conflict, particularly where conventional medical evacuation is disrupted. Trauma Stabilization Points (TSPs) were introduced in previous conflicts to bring life-saving care closer to the point of injury. To improve standards and adaptability, the World Health Organization convened a Technical Working Group to develop operational guidance for TSPs, drawing on diverse field experiences, including the ongoing Gaza deployments.

CASE PRESENTATION: Between February and July 2024, three TSPs were established in Gaza, managing over 4,000 consultations. Initially focused on trauma stabilization and referral, these sites quickly adapted to minor injuries and non-traumatic conditions, reflecting population needs and access challenges. Referral rates varied across sites due to hospital proximity, ambulance availability, and shifting frontlines. Security threats limited forward deployment and safe patient access, requiring high mobility and rapid relocation. Experience from Gaza highlighted key operational principles: locating TSPs near the point of injury; integrating within a functioning trauma referral pathway supported by evacuation capacity and hospital readiness; maintaining clear clinical functions and staffing standards; and using standardized documentation for quality assurance and continuity. [...]

[PubMed](#) · [DOI](#)

Acute-on-chronic versus isolated acute subdural hematoma in complicated mild traumatic brain injury: association with radiological mass effect markers in a multicenter cohort.

Crombé A, Benhamed A, Seux M, Frassin L, Matichard R, L'Huillier R, Millon D, Gorincour G

PURPOSE: The clinical impact of acute-on-chronic subdural hematoma (acSDH) in mild traumatic brain injury (mTBI) remains incompletely characterized. We assessed whether acSDH, compared with isolated acute SDH (aSDH), is associated with worse neurological presentation and increased radiological markers of mass effect.

METHODS: We conducted a retrospective multicenter cohort study using the IMADIS teleradiology head trauma workflow (103 emergency departments in France, January 2020 to December 2022). Adult patients with complicated mTBI (i.e., GCS 13-15 plus intracranial hemorrhage or skull fracture) and acute SDH on non-contrast head CT were eligible. Exposure was acSDH versus aSDH based on structured radiology reports. Outcomes were (1) GCS < 15 at presentation, (2) SDH maximal thickness ≥ 7 mm, (3) radiological brain herniation, and (4) intermediate or high QueBIC risk category. Univariable and multivariable logistic regression models were used to estimate associations.

RESULTS: Among 935 included patients, 107 (11.4%) had acSDH. Patients with acSDH were older (≥ 75 years: 57.0% vs. 39.7%, $p < .0001$) and more frequently had dementia (8.4% vs. 3.7%, $p = .0465$). Compared with aSDH, acSDH was associated with GCS < 15 (37.4% vs. 21.7%, $P = .0010$) and confusion (41.0% vs. 25.8%, $p = .0016$). On CT, acSDH was associated with SDH thickness ≥ 7 mm (54.2% vs. 20.4%, OR 4.50, 95%CI 2.9 [...])

[PubMed](#) · [DOI](#)

Persistent neuropathic pain symptoms after FPV drone-related thoracic blast injury without rib fractures: a pilot observational study from the Ukraine War.

Battle C, Baker E, Giamello JD, Melchio R, Dmytriiev D

PURPOSE: The aim of this exploratory pilot study was to describe First Person View drone (FPV) related, blast-mechanism thoracic trauma with no rib fractures in Ukrainian military patients sustained during the Russia-Ukraine Conflict, and report the prevalence of persistent neuropathic pain symptoms at three months among patients admitted to a rehabilitation and pain medicine service.

METHODS: A retrospective, single-centre observational, exploratory study design was used. Military patients (≥ 16 years old) with blast-mechanism thoracic injuries with no rib fractures, admitted to a hospital's rehabilitation and pain medicine service in Ukraine for ≥ 24 h were included. The PainDETECT survey was completed at three months post-injury. Participant characteristics were summarised using descriptive statistics. Exploratory univariate and multivariate analyses were performed to explore possible associations between variables and the development of persistent neuropathic pain symptoms.

RESULTS: 217 patients were included in the final analysis. Median age was 39 years (IQR: 32-47.5). 131 (60.4%) patients were admitted to critical care with 91 (41.9%) patients requiring mechanical ventilation. The median hospital length of stay was 22 days (IQR: 16-25) and critical care length of stay was 4 days (IQR: 2-5). Persistent neuropathic pain symptoms were reported in 173 out of 217 patients ([...])

The association between pancreatic enzyme serum concentrations and blunt pancreatic trauma: a systematic review.

Sykes B, Gracias D, Bendinelli C, Balogh ZJ

PURPOSE: Blunt pancreatic trauma is uncommon. When suspected, serum amylase and lipase are measured, but their diagnostic value remains uncertain. This systematic review evaluated the diagnostic performance of serum amylase and serum lipase in detecting blunt pancreatic trauma in adults.

METHODS: MEDLINE, Embase, Scopus and Cochrane Central were searched for studies evaluating serum amylase and/or lipase in adults with blunt pancreatic trauma. Observational studies reporting diagnostic accuracy or clinical outcomes were included. Two reviewers independently screened studies and assessed risk of bias using the Newcastle-Ottawa Scale.

RESULTS: Four observational studies met inclusion criteria. A total of 4937 patients with blunt abdominal trauma, including 150 with pancreatic injury were investigated. Two studies evaluated serum lipase alone and two compared amylase and lipase. Pancreatic injuries were graded according to the American or Japanese Association for the Surgery of Trauma. Early serum amylase measurement (< 90 min post-injury) demonstrated low sensitivity. In contrast, testing performed more than 3 h after injury has shown improved diagnostic accuracy with one study suggesting combined serum amylase and lipase testing can achieve a sensitivity of 85% and specificity of 100%.

CONCLUSION: Early serum pancreatic enzymes measurement does not appear to aid initial management [...]

[PubMed](#) · [DOI](#)

Extension of the fluidics NTF-Biobank of polytraumatized patients with extracellular vesicles: improved vesicle preservation through local processing.

Weber B, Leppik L, Han J, Becker N, Wang T, Groven RVM, Balmayor ER, Zhao Q et al.

PURPOSE: Extracellular vesicles (EVs) show diagnostic relevance in trauma, prompting the development of the NTF■EV■Biobank as an extension of the established Network Trauma Research (NTF) plasma/serum biobank. After optimizing the EV isolation and characterization standard operating procedure (SOP) at a single center, this multicenter study evaluates its transferability and examines how sample transport affects EV quality before large■scale implementation in a nationwide biobank setting.

METHODS: Plasma and serum from polytrauma patients and healthy controls (n=10) were processed under the standardized NTF■Biobank protocol and distributed to three centers. EVs were isolated using the unified NTF■EV■Biobank workflow, exchanged between centers, and analyzed for protein markers, particle size and concentration (NTA), total protein content, and morphology (TEM) to compare EV quality across sites.

RESULTS: Implementation of the EV biobank protocol across participating centers, as well as inter-center transport on dry ice, was successfully performed without complications. All samples arrived at the analyzing centers in a frozen state. EVs exhibited typical vesicular morphology as confirmed by transmission electron microscopy and remained positive for CD9 and CD81, indicating that transportation did not affect morphology or surface marker expression. However, transportation did reduce [...]

[PubMed](#) · [DOI](#)

Whole blood first resuscitation and association of blood product utilization based on mechanism of injury.

Hadley J, Conner J, Thomas C, Llorente K, Gilani A, White B, Miller Iii P, Nunn A et al.

INTRODUCTION: Studies evaluating the effect of whole blood (WB) vs. component-based resuscitation accounting for mechanism of injury are lacking. We sought to evaluate the effects of a WB first resuscitation strategy regarding mechanism of injury.

METHODS: This is a retrospective cohort study of trauma patients who received any blood product during resuscitation from January 2016 to November 2021. The primary outcomes were total blood volume utilized within 24 h and 30-day mortality. Multivariable gamma regression with a log link was used for blood volume and logistic

regression for mortality. A likelihood ratio test evaluated whether injury mechanism modified the treatment effect.

RESULTS: Between January 2016 and November 2021, 1,013 injured patients received blood products (785 WB-first). We did not detect statistically significant effect modification by injury mechanism for either outcome (interaction $p = 0.44$ for blood volume, $p = 0.25$ for mortality). In multivariable gamma regression, WB-first resuscitation was associated with a ratio of mean total blood volume of 0.71 (95% CI 0.55 to 0.90, $p = 0.005$) relative to components, representing approximately 29% lower utilization. WB-first was not associated with 30-day mortality (OR 0.90, 95% CI 0.48 to 1.71, $p = 0.73$).

CONCLUSIONS: A WB-first resuscitation strategy is associated with lower blood volume utilized within the f [...]

[PubMed](#) · [DOI](#)

Reported complications after spinopelvic fixation for U-type sacral fractures in older adults: a systematic review with narrative synthesis.

Giordano V, Pinheiro HRS, Pires RE, Varga P, Freitas A, Giannoudis PV

BACKGROUND: Spinopelvic fixation (SPF), alone or combined with iliosacral or transiliac-transsacral fixation, is often used for highly unstable posterior pelvic ring injuries and some fragility fracture patterns in older adults. However, the evidence specifically describing complications after SPF in older patients with U-type sacral fractures is limited.

METHODS: This systematic review was conducted according to the Cochrane Handbook and PRISMA recommendations. PubMed, Embase, and grey-literature sources were searched from January 1, 2000 to July 31, 2025. Eligible studies included adults aged 60 years or older with U-type sacral fractures or closely related spinopelvic dissociation/FFP IVb patterns treated with SPF, and reporting at least one complication. Because of the small number of eligible studies, probable cohort overlap, heterogeneity in fracture definitions, treatment indications, fixation constructs, and outcome reporting, findings were synthesized narratively rather than pooled quantitatively.

RESULTS: Three studies met the eligibility criteria, comprising 157 patients overall; however, only a subset underwent SPF, and two studies were produced by the same group with probable cohort overlap. The available cohorts mixed bilateral sacral fragility fractures, FFP IVb patterns, and comparative SPF-versus-trans-sacral constructs. Reported adverse events included wound [...]

[PubMed](#) · [DOI](#)

Influence of allograft augmentation on surgical outcomes and complications in proximal humerus fractures: a matched-pair analysis.

Barthel C, Rudolph F, Kalbas Y, Russek R, Pape HC, Allemann F

BACKGROUND: Proximal humerus fractures predominantly affect elderly patients and pose a growing clinical burden. This study evaluates the impact of allograft augmentation on operative time, fracture reduction quality, and complication rates compared with standard fixation across different fracture morphologies.

METHODS: This retrospective matched-pair study included patients surgically treated for proximal humerus fractures with locking plate fixation at a single tertiary trauma center between 01.01.2014 and 31.12.2024. Patients treated with allograft augmentation (Group A) were matched 1:1 to patients without allograft (Group NA) using exact matching for Mayo FJD fracture morphology and sex, and propensity score matching for age. Radiological outcomes were assessed using pre- and postoperative neck-shaft angles, with normalization defined as correction into the physiological range (120°-150°). Complications and operative parameters were recorded. Paired statistical analyses were applied.

RESULTS: A total of 1,012 patients were treated during the study period; after selection and matching, 102 matched patient pairs (204 patients) were analyzed. Operative time was longer with allograft augmentation (141.3 ± 40.3 vs. 126.9 ± 50.0 min; $p = 0.08$), reaching statistical significance only in varus posteromedial fractures (144.4 ± 37.8 vs. 122.2 ± 39.6 min; $p = 0.007$). Although overa [...]

[PubMed](#) · [DOI](#)

Impact of a comprehensive trauma pathway on workflow and clinical processes in a level III trauma center.

Neitzert L, Druetto C, Forno D, Molino P, Vitale D, Grio M

PURPOSE: Organized trauma systems improve coordination of care and may enhance the efficiency of trauma management, potentially improving outcomes and reducing hospital length of stay and costs. The aim of the study was to evaluate key process-of-care indicators and clinical outcomes before and after the introduction of a comprehensive trauma pathway.

METHODS: We conducted a single-center retrospective-prospective cohort study at Rivoli Hospital, Italy. Trauma patients admitted to the Emergency Department (ED) between 2024 and 2025 who met the criteria for trauma team activation were prospectively included. Outcomes were compared with a retrospective cohort from the period prior to the implementation of the dedicated trauma pathway (2023-2024).

RESULTS: A total of 465 patients were included. Implementation of a comprehensive trauma pathway was associated with significant improvements in several time-dependent quality indicators, including ED boarding time (888 vs. 1021 min; $p = 0.02$), time to imaging (48 vs. 78 min; $p < 0.01$), time to surgery (183 vs. 237 min; $p = 0.03$), time to ICU admission (226 vs. 343 min; $p = 0.02$), and time to interhospital transfer (238 vs. 305 min; $p = 0.04$). We observed a trend toward a reduction in early mortality (2% vs. 3.6%; $p = 0.41$) and 30-day mortality (3.7% vs. 5.9%; $p = 0.62$).

CONCLUSIONS: The adoption of a structured multidisciplinary trauma pathway [...]

[PubMed](#) · [DOI](#)

The DRoPH score: development of a tool for predicting reduction difficulty in proximal humerus fracture osteosynthesis.

Barthel C, Rudolph F, Kraus M, Pape HC, Allemann F

BACKGROUND: Proximal humerus fractures (PHFs) are among the most common and challenging fractures due to complex anatomy and variable fracture patterns. While anatomical reduction is critical for successful outcomes, no tool currently predicts the intraoperative difficulty of achieving it. Despite advances in surgical techniques, complications persist. Existing classification systems, such as Codman's, Neer's, AO/OTA, and Mayo-FJD, focus on fracture morphology and displacement but do not address intraoperative complexity. This study aims to develop a new score for the early prediction of intraoperative reduction difficulty in proximal humerus fracture surgery, the DRoPH score (Difficulty of Reduction of Proximal Humerus Fractures).

METHODS: An online survey was conducted among experienced upper extremity surgeons in Switzerland to identify factors influencing the difficulty of reducing and retaining proximal humerus fractures. Inclusion criteria for surgeon grading comprised experience with more than 100 surgically treated proximal humerus fractures. Eligible surgeons rated the relevance of factors influencing reduction difficulty using a five-point Likert scale. A rank-based weighting system derived from expert ratings was applied, assigning two points to the three highest-ranked factors. The resulting DRoPH-Score ranged from 0 to 10, with higher values indicating greater surgical difficulty [...]

[PubMed](#) · [DOI](#)

Interface stability of intramedullary nail locking configurations under combined axial and torsional loading.

Marianne H, Martin W, Claus G, Anja B, Peter A

The biomechanical performance of intramedullary nails in metaphyseal regions strongly depends on distal locking configuration. This study aimed to evaluate screw-bone interface stability under combined axial and torsional cyclic loading for different distal locking configurations of intramedullary nails, focusing on the effects of screw number, spatial distribution, interscrew distance, and multiplanar versus coplanar fixation using a standardized synthetic bone model. Six distal locking configurations ($n = 5$ per group) were tested under stepwise cyclic axial compression, superimposed with a constant torsional moment (± 2 Nm, 2 Hz). Primary endpoints were cycles to failure, cycles to 1 mm axial displacement, and cycles to 0.5° rotation. All constructs failed by progressive screw cut-through within the synthetic bone. The three-screw multiplanar configuration exhibited the highest endurance ($\approx 36,000$ cycles to failure) and lowest micromotion ($\approx 34,000$ cycles to 1 mm displacement), significantly outperforming the two-screw configurations overall ($p = 0.002$). Increasing interscrew distance from 10 mm to 34 mm improved migration resistance by approximately 25%, whereas the short 10 mm spacing led to premature failure. Blocked and free-hand configurations showed comparable behavior to the standard setup, indicating no added benefit from toggle elimination alone. Locking configuration [...]

Three-dimensional finite element analysis for internal fixation of mandibular condylar fractures: a scoping review.

Kakei Y, Sader R, Ebener P, Tanaka A, Akashi M, Kuroda R, Thoenissen P

PURPOSE: Three-dimensional finite element analysis (3D-FEA) has emerged as an essential computational tool for evaluating the biomechanical behaviors of internal fixation systems for mandibular condylar fractures. This scoping review maps and synthesizes the existing literature on 3D-FEA studies investigating the internal fixation of mandibular condylar fractures and explores, in a descriptive manner, differences in methodological focus between European and non-European research centers.

METHODS: PubMed and Cochrane Library databases were systematically searched following PRISMA-ScR guidelines. Original English-language research articles using 3D-FEA to evaluate internal fixation of mandibular condylar fractures were included. Data were extracted using a standardized charting form.

RESULTS: Thirty-six studies met the inclusion criteria. European studies ($n = 12$) predominantly focused on development of novel plate designs and mechanical testing protocols, whereas Asian and other nationality studies ($n = 24$) emphasized material comparisons between titanium and bioabsorbable systems, patient-specific modeling, and loading condition analysis. Consensus findings include: (1) double-plate fixation provides superior stability; (2) among 3D plate designs, trapezoid plates demonstrate the best performance for subcondylar and condylar base fractures, whereas alpha and lambda plates may [...]

[PubMed](#) · [DOI](#)

Initial surgical management of injuries to the upper extremities for patients with multiple and/or severe injuries - a systematic review and clinical practice guideline update.

Berk T, Könsgen N, Lechler P, Imach S, Schär M, Bieler D, Horst K, Hildebrand F et al.

PURPOSE: Our objective was to update the evidence-based and consensus-based recommendations for the initial surgical management of upper extremity injuries in patients with suspected multiple and/or severe injuries based on current evidence. This guideline topic is part of the 2025 update of the German Guideline on the Treatment of Patients with Multiple and/or Severe Injuries.

METHODS: MEDLINE and Embase were systematically searched to September 2024. Further literature reports were obtained from clinical experts. Randomised controlled trials (RCTs) or observational studies reporting risk-adjusted outcomes were included if they compared early versus delayed surgical treatment for fractures, vascular injuries, or nerve injuries affecting the upper extremities in patients with multiple and/or severe injuries. Studies comparing limb salvage versus amputation or comparing amputation criteria for the upper extremities were also included. We considered patient-relevant outcomes such as mortality and limb salvage, as well as the sensitivity and specificity of scores for predicting upper extremity amputation. Risk of bias was assessed at the outcome level using ROBINS-I for observational studies and AMSTAR-2 for systematic reviews. We used available meta-analyses if possible; alternatively, we synthesised the evidence narratively. We used GRADE to rate the certainty of evidence. Expe [...]

[PubMed](#) · [DOI](#)

Transosseous sutures versus vertical cerclage wire for distal pole patellar fractures: a retrospective comparative cohort study.

Laso JI, Muñoz JT, Bianchi S, Martínez D, Rojas T, Franulic N, Olivieri R

PURPOSE: To compare transosseous sutures (TOS) and vertical wire cerclage (VCW) for treating distal pole patellar fractures, evaluating clinical outcomes, patient-reported outcome measures (PROMs), radiographic outcomes, and complications.

METHODS: We conducted a retrospective comparative cohort study of patients surgically treated for patellar distal pole fractures (AO/OTA 34A1b) with either TOS or VCW between 2015 and 2023. Patient characteristics, surgical technique, postoperative data, patellar height, and complications were recorded. Functional outcomes were evaluated at a minimum of 12 months postoperatively using PROMs (Kujala, Lysholm, and Bostman scores). The primary outcome of this study was functional recovery, assessed via the Bostman score at 12 months.

RESULTS: A total of 46 patellar distal pole fractures were included (30 TOS and 16 VCW). Baseline characteristics showed no statistically significant differences between groups. No significant differences were found regarding clinical or functional outcomes. Regarding radiographic outcomes, patellar height remained stable; however, both groups showed significant patellar length variation, with elongation being significantly more pronounced in the TOS group (mean 3.77 mm) compared to the VCW group (mean 0.44 mm; $p = 0.001$). While overall complication rates showed no statistically significant difference, all five ca [...]

[PubMed](#) · [DOI](#)

Refracture risk following implant removal in consolidated hip fractures: a multicenter retrospective cohort study of 575 patients.

Mohammed J, Mili-Schmidt V, Wadsten M, Juto H, Wolf O, Crnalic S, Sköldenberg O, Mukka S et al.

PURPOSE: The risk of re-fracture after implant removal in healed hip fractures remains a clinical concern. This study aimed to determine the incidence of re-fracture following implant removal after osteosynthesis of a hip fracture.

METHODS: We conducted a retrospective multicenter cohort study including patients aged ≥ 50 years who underwent implant removal between 2003 and 2023 after radiologically confirmed consolidation of a hip fracture. Patients were identified using procedural codes. Baseline variables included age, sex, ASA classification, fracture type (femoral neck or trochanteric), and implant type (pins/screws, sliding hip device [SHD], or short/long cephalomedullary nail [CMN]). Patients were followed from implant removal until re-fracture, conversion to arthroplasty, death, or end of follow-up, with a minimum follow-up of 1 year. Cox proportional hazards regression was used to assess associations between implant type and re-fracture risk, adjusting for age and sex. Because the proportional hazards assumption was violated, a time-stratified Cox regression and restricted mean survival time analyses were applied.

RESULTS: A total of 575 patients (median age 73 years, IQR 65-81) were included, with a median follow-up of 53 months (IQR 18-100). Lateral hip pain was the most common indication for implant removal (72.5%). The overall re-fracture incidence was 10.4%. Ris [...]

[PubMed](#) · [DOI](#)

Physical functionality two years after major trauma: predictors from a single center cohort study.

Fetz K, Koetsenruijter J, Rodrigues Recchia D, Rutetzki J, Lefering R, Kaske S

PURPOSE: Traumatic injuries are a major global public health concern, often leading to impaired physical function and disability. Long-term physical function impairments can greatly affect a patient's health related quality of life, influencing their emotional state and social interactions. This study examines patient-reported physical functionality before and after major trauma, identifying key predictors of poor functional recovery.

METHODS: This retrospective single-center cohort study investigates physical functionality 23 months after a traumatic injury using standardized questionnaires combined with clinical data. Inclusion criteria were Injury Severity Score (ISS) ≥ 9 , requiring ICU treatment, and age ≥ 18 resulting in an eligible sample of 515 patients. Health outcomes were assessed using the Trauma Outcome Profile (TOP), the Abbreviated Injury Scale (AIS), and the ISS.

RESULTS: Physical functionality significantly decreased from 94.8 (SD 11.5) to 72.9 (SD 26.7) 23 months post-trauma with almost half of participants still scoring below the critical cutoff of 80 points. The strongest predictor for functional physical impairment was pre-trauma physical functionality (OR 8.21, $p < 0.001$). Other determinants included age (age 50-59 versus 18-29 OR 2.75, $p = 0.001$), as well as an impacted lower extremity including pelvic (OR 1.60, $p = 0.018$).

CONCLUSION: The most importan [...]

[PubMed](#) · [DOI](#)

Association between fresh frozen plasma transfusion and mortality stratified by Glasgow Coma Scale scores in isolated traumatic brain injury: a nationwide cohort study in Japan.

Nagamura T, Aoki M, Yamada K, Terayama T, Seno S, Tomura S, Kiyozumi T

PURPOSE: Fresh frozen plasma (FFP) is widely used to manage trauma-induced coagulopathy. However, its effectiveness in isolated traumatic brain injury (TBI) management remains controversial, particularly when stratified by severity based on the Glasgow Coma Scale (GCS) scores. This study aimed to examine the association between FFP administration within 24 h and mortality in patients with isolated TBI, overall and stratified by GCS on arrival.

METHODS: We conducted a multicenter retrospective cohort study using the Japan Trauma Data Bank (2019-2023). Adults with isolated TBI (intracranial AIS ≥ 3 and extracranial AIS < 3) were included. Patients were classified into FFP and non-FFP groups based on transfusion within 24 h of admission. Subgroup analyses were performed by TBI severity (severe: GCS ≤ 8 ; mild-to-moderate: GCS > 8), craniotomy status, and FFP dose. The primary outcome was 28-day mortality. Propensity score matching was applied to adjust for confounding.

RESULTS: Among 12,480 patients (median age 72 years [IQR: 55, 82]; 66.6% male), 513 received FFP. After matching, FFP was not significantly associated with 28-day mortality (OR 0.80; 95% CI 0.59-1.08). In subgroup analyses, FFP was associated with higher mortality in mild-to-moderate TBI (OR 2.38; 95% CI 1.07-5.31) and lower mortality in severe TBI (OR 0.61; 95% CI 0.42-0.88). No significant differences were observed [...]

[PubMed](#) · [DOI](#)

Percutaneous fixation of posterior pelvic ring lesions in the elderly: current evidence and recent advances.

Liodakis E, Schreiber S, Giannoudis VP, Fritz T, Giannoudis PV

Transsacral percutaneous fixation is increasingly seen as a safe, effective minimally invasive treatment for geriatric posterior ring fractures. The use of Navigation appears a safe and efficient method for percutaneous screw placement. Multiple new implants have been described for management of these injuries. Level I evidence is still lacking for treatment options in the management of posterior pelvic ring lesions with percutaneous fixation.

[PubMed](#) · [DOI](#)

Infrared thermography and dual-layer spectral CT for the detection of femoral vascular compromise: an experimental large animal study.

Hübner CT, Ricklin J, Müller AL, Klingebiel FK, Stoeck CT, Weisskopf M, Carneiro Gomes A, Cinelli P et al.

BACKGROUND: Rapid identification of vascular compromise remains a clinical challenge in the acute trauma setting. Conventional diagnostic tools such as CT angiography (CTA) and Doppler-ultrasound might provide decent anatomical and functional assessment. However, they are limited by ionizing radiation, contrast exposure or time requirements. Although not validated, infrared thermography (IRT) might offer a non-invasive alternative capable of detecting surface temperature asymmetries that may reflect underlying perfusion deficits. This study aimed to evaluate the feasibility of thermographic imaging in detecting vascular lesions.

METHODS: As part of a porcine polytrauma model (n = 28), vascular lesion of the right femoral artery was defined as a $\geq 50\%$ diameter reduction on CTA. Temperature differences between both extremities were measured using infrared thermography. SDCT based iodine concentrations distal to the stenosis were quantified to assess perfusion changes. Correlation and regression analyses were performed between stenosis degree, temperature difference, and tissue iodine uptake.

RESULTS: Six animals fulfilled the criteria for relevant vascular compromise. This resulted in a mean vessel diameter reduction behind the lesion of $68\% \pm 18\%$. The vascular lesion group demonstrated a mean inter-limb temperature difference of $6.93^\circ\text{C} \pm 1.78^\circ\text{C}$ compared to $0.71^\circ\text{C} \pm 1.27^\circ\text{C}$ in [...]

[PubMed](#) · [DOI](#)

Fixation strategies for periprosthetic femur fractures around knee replacements: a comprehensive single-center analysis of 102 consecutive cases.

Nomdedéu J, González-Morgado D, Minguell-Monyart J, Joshi-Jubert N, Teixidor-Serra J, Tomàs-Hernández J, Andrés-Peiró JV

PURPOSE: To evaluate 1 year mortality, complication rates, and functional outcomes after internal fixation of femoral periprosthetic knee fractures following total knee arthroplasty, and to identify predictors of adverse outcomes and increased healthcare resource utilization.

METHODS: This retrospective cohort study included 102 consecutive patients with femoral periprosthetic knee fractures classified as UCPF VB3 and VC3 treated with internal fixation at a single tertiary center between 2010 and 2023. Interprosthetic fractures and combined fixation techniques were excluded. Primary outcomes were 1 year mortality and postoperative complications. Secondary outcomes included length of stay, discharge destination, and ambulatory status at follow up.

RESULTS: Patients were markedly frail, with a mean age of 82.4 ± 10.8 years and mean Charlson Comorbidity Index of 5.6. One year mortality was 16.7% and was associated with older age, higher ASA class, higher Charlson index, impaired pre fracture ambulation, and fracture related infection. Overall complication rate was 29.4%, including 15.7% infections and 13.7% mechanical failures. Open reduction was more frequent in plate fixation and resulted in longer operative times, without increasing complication rates. Median hospital stay was 18 days, 66.3% required discharge to nursing facilities, and only 7.8% regained independent ambulation [...]

[PubMed](#) · [DOI](#)

Epidemiology and management of pediatric femoral fractures at a level I trauma center: a 10-year retrospective monocentric study.

Dietz SO, Traub F, Schmidt E, Schippers P, Wegner E, Schierjott L, Gercek E, Nowak TE

PURPOSE: Pediatric femoral fractures are relatively uncommon but represent severe injuries frequently requiring inpatient trauma care. Epidemiological characteristics, fracture patterns, and treatment strategies vary between institutions and regions. This study aimed to analyze the epidemiology, fracture distribution, treatment modalities, complications, and fracture-predisposing comorbidities of pediatric femoral fractures treated at a German level I pediatric trauma center over a 10-year period.

METHODS: All patients aged 0-14 years treated for femoral fractures between January 2012 and December 2021 were retrospectively reviewed. Fractures were categorized using the AO Pediatric Comprehensive Classification of Long-Bone Fractures (PCCF). Demographic variables, fracture characteristics, management approaches, time to consolidation, complications, and relevant comorbidities predisposing to fracture were analysed descriptively in accordance with STROBE recommendations.

RESULTS: A total of 183 patients met the inclusion criteria (67.2% male; mean age 5.2 years). Children aged 2-5 years constituted the largest group (42.1%). Femoral shaft fractures were the most frequent femoral fracture type (70%). Surgical management, most frequently elastic stable intramedullary nailing (ESIN), was increasingly utilized with advancing age. The overall complication rate was 14.2%, with the ma [...]

[PubMed](#) · [DOI](#)

Blood-based biomarkers GFAP/UCH-L1 for the diagnosis of mild traumatic brain injury (mTBI): a single-center implementation experience.

Maegle M, Breidenbach L, Kaufmann J, Keles Y, Uhl E, Schroeder P

INTRODUCTION: Mild traumatic brain injuries (mTBI) affect millions of people worldwide every year as one of the most common clinical presentations in the emergency department. Diagnosis is mainly based on clinical criteria and computed tomography scans. The use of computed tomography causes high costs, long waiting times in daily clinical practice and radiation exposure. GFAP (glial fibrillary acidic protein) and UCH-L1 (ubiquitin carboxyl-terminal hydrolase-L1) turned out to be potential biomarkers for the diagnosis of mTBI. This study retrospectively evaluates the possible use of these biomarkers combined as negative predictors for excluding brain injuries in patients with suspected mTBI in the emergency department.

METHODS: Adult patients (n = 320) registered in the emergency department at a level 1 trauma emergency center in Germany (Cologne Merheim Medical Center/CMMC) between 11/2023 and 04/2024, with suspected mTBI, Glasgow Coma Scale (GCS) score 13-15 and within 12 h after trauma were considered. All evaluable patients underwent cranial CT (cCT) scans and blood tests for GFAP and UCH-L1 serum concentrations.

RESULTS: Biomarkers GFAP and UCH-L1 were tested positive in 261 patients (82%) while CT detected intracranial injuries in only 29 patients (9%). Biomarkers combined had a sensitivity of 97% and a negative predictive value (NPV) of 98% in mTBI diagnosis with a nega [...]

[PubMed](#) · [DOI](#)

Resuscitative endovascular occlusion of the aorta restores cerebral metabolic markers of ischaemia induced by haemorrhagic shock.

Bader SE, Magnuson A, Brorsson C, Wallin G, Löfgren N, Löfgren F, Blind PJ, Öman M et al.

BACKGROUND: Resuscitative endovascular balloon occlusion of the aorta (REBOA) is used as an adjunct in haemorrhagic shock (HS) to restore proximal perfusion. Its cerebral metabolic effects during haemorrhagic shock, particularly in the presence of elevated intracranial pressure (ICP), remain incompletely characterised.

METHODS: In a controlled porcine model, HS was induced by controlled bleeding to a mean arterial pressure of approximately 40 mmHg. Animals were allocated to a normal ICP group or an experimentally elevated ICP group. Total REBOA (zone 1) was applied for 90 min. Cerebral metabolism was assessed using intracerebral microdialysis with measurements of lactate, pyruvate, and lactate-pyruvate ratio (LPR). Cerebral haemodynamics and ICP were continuously monitored.

RESULTS: HS was associated with a statistically significant increase in LPR in both groups, indicating cerebral metabolic disturbance. Following aortic occlusion, LPR gradually decreased toward baseline in both groups. Animals with elevated ICP demonstrated a transient delay in metabolic normalisation during the early post-occlusion phase. Statistically significant differences between groups were limited to the first 10 min following occlusion. Overall metabolic trajectories were similar thereafter.

CONCLUSIONS: Total REBOA restored cerebral metabolic markers of ischaemia during haemorrhagic shock, as ref [...]

[PubMed](#) · [DOI](#)

Delayed single-stage surgical management using a peroneus brevis muscle flap for fracture-related infection after lateral malleolus fixation: a retrospective case series.

Dyreborg K, Jørgensen AI, Peterlin AAN, Gottlieb H

BACKGROUND: Fracture-related infection (FRI) following lateral malleolus fracture fixation complicated by soft tissue breakdown represents a challenging clinical problem with no established standard of care. This study evaluates a treatment strategy consisting of prolonged antibiotic suppression until fracture union, followed by delayed single-stage surgical management.

METHODS: This retrospective consecutive case series included 11 patients treated between 2020 and 2023 at a multidisciplinary orthopaedic infection unit. All patients were managed according to a standardized protocol involving oral antibiotic suppression until radiologically confirmed fracture consolidation. This was followed by a single-stage surgical procedure including implant removal, radical debridement, deep tissue sampling, local antibiotic therapy, and soft tissue reconstruction using a distally based peroneus brevis muscle flap with split-thickness skin grafting.

RESULTS: Eleven consecutive patients (median age 58 years) were treated according to the protocol. Six patients underwent antibiotic suppression for a median duration of 147 days prior to surgery. Among patients available for outcome assessment, infection control was achieved in 10 of 10 surviving patients. One patient died of unrelated causes within 14 days postoperatively and was not classified as treatment failure. Soft tissue complication [...]

[PubMed](#) · [DOI](#)

Mixed reality improves agreement on surgical approach selection and patient positioning in tibial plateau fracture planning compared to CT, 3DCT and 3D printing.

Dust T, Henneberg JE, Hartel M, Reiter A, Kylies J, Korthaus A, Streckenbach A, Keller J et al.

PURPOSE: Tibial plateau fractures require precise preoperative planning, particularly regarding treatment concept, patient positioning, and surgical approach. Mixed reality (MR) may enhance three-dimensional understanding of fracture morphology beyond conventional imaging and physical 3D models. This study investigated whether MR improves agreement on key preoperative planning decisions compared to computed tomography (CT), 3D computed tomography (3DCT), and 3D-printed fracture models.

METHODS: Twelve orthopaedic trauma surgeons (6 junior, 6 senior) evaluated 22 surgically treated tibial plateau fractures (AO/OTA type B or C). Each case was assessed in four steps using (1) CT, (2) 3DCT reconstructions, (3) physical 3D-printed models, and (4) MR visualization using Microsoft HoloLens 2. Raters selected (i) treatment

concept, (ii) patient positioning, and (iii) surgical approach. Interobserver agreement was analysed using Fleiss' kappa (κ) and percentage match (PM).

RESULTS: Mixed Reality (MR) yielded the highest interobserver agreement for both surgical approach (PM 32%, $\kappa = 0.30$) and patient positioning (PM 57%, $\kappa = 0.35$), particularly among junior surgeons (approach PM 39%, $\kappa = 0.28$; positioning PM 57%, $\kappa = 0.39$). Compared to CT, 3DCT, and 3D printing, MR demonstrated consistent improvements, while agreement on treatment concepts remained high across all modalities (PM > 82% [...])

[PubMed](#) · [DOI](#)

Timing of fracture fixation for femur and pelvis fractures in patients with severe traumatic brain injury - an analysis of the TraumaRegister DGU®.

Bayer TP, Teuben MPJ, Halvachizadeh S, Shehu A, Lefering R, Pfeifer R, Pape HC, Jensen KO

PURPOSE: Fracture fixation timing and strategy in polytrauma patients with traumatic brain injury (TBI) remain controversial. This study investigates treatment patterns and outcomes for femoral and/or pelvic fractures stratified by TBI severity.

METHODS: Patients in the TraumaRegister DGU® (2016-2022) with pelvic and/or femoral fractures (AIS ≥ 3) and TBI (head AIS ≥ 3) were included. Strategies were non-operative management (NOM), early total care (ETC), and damage-control orthopedics (DCO). Outcomes included treatment allocation, fixation timing, and in-hospital mortality.

RESULTS: 985 patients were included (mean age 52.5, SD 26.3 years; ISS 27.8, SD 8.1). Allocation was NOM in 320 (32.5%), ETC in 336 (34.1%), and DCO in 329 (33.4%) patients. Head AIS was 3 in 48.5%, 4 in 31.1%, and 5 in 20.3%. NOM patients were older, had the highest ISS and estimated mortality, and showed the largest proportion of critical TBI (AIS 5: NOM 30.9%, ETC 14.3%, DCO 16.1%). Femoral ETC was mainly performed within the first day (median 0, IQR 0-1 days), whereas pelvic ETC was delayed with increasing TBI severity (median 3, IQR 0-5 days for head AIS 3; 5, IQR 0-7 days for AIS 4). Observed mortality was 37.2% after NOM, 9.2% after ETC, and 10.3% after DCO.

CONCLUSION: ETC in patients with moderate TBI (AIS 3) was associated with reduced observed mortality relative to NOM and matching DCO. Increase [...]

[PubMed](#) · [DOI](#)

From single to dual platelet inhibition FIBTEM assay: fibrinogen replacement thresholds for critical bleeding after trauma.

Latona A, Winearls J, Ho A, Mitra B

PURPOSE: The FIBTEM assay in Rotational Thromboelastometry (ROTEM®) measures fibrin-based clot formation. The reagent transitioned from single to dual-platelet inhibition with addition of tirofiban to cytochalasin D, offering more complete platelet suppression. The aim of this study was to assess the impact of this change on transfusion thresholds in trauma.

METHODS: In this retrospective cohort study, we included adult patients with major trauma across Queensland who presented between 2022 and 2025 and had results available for paired ROTEM and conventional coagulation tests on arrival. Patients were sub-grouped into single and dual-platelet inhibition assay groups. Correlations between FIBTEM-A5 and Clauss fibrinogen were analysed, comparing correlation strength and diagnostic accuracy for fibrinogen levels of < 2.0 g/L.

RESULTS: The relationship between FIBTEM-A5 and Clauss fibrinogen was similar using single platelet inhibition assays (R^2 0.61; 95%CI: 0.51-0.70) and dual-platelet inhibition assays (R^2 0.70; 95%CI: 0.62-0.77; $p = 0.94$). Predicted FIBTEM-A5 amplitudes at 2 g/L fibrinogen were 7.83 mm (95%CI: 7.42-8.23) and 7.54 mm (95%CI: 7.22-7.87). Diagnostic performance for detecting fibrinogen < 2.0 g/L was comparable between assays, for both FIBTEM-A5 ≤ 10 mm and FIBTEM-A5 ≤ 8 mm (all $p > 0.05$). For the dual-platelet inhibition assay, sensitivity and specificity were 97. [...]

[PubMed](#) · [DOI](#)

Pattern of blast injuries. A systematic review: Part 2 - Landmines, unexploded ordnance and terrorism.

Neubert A, Gaeth C, Seidelmann M, Bieler D

PURPOSE: This systematic review analyzes the patterns and distribution of blast injuries from landmine explosions, UXOs and terror attacks including suicide bombing, in civilians and military personnel.

METHODS: A search was conducted in two databases on 26. March 2025 for observational studies reporting blast injuries in civilian and military populations of all ages. Studies were clustered by explosion typ. A narrative data synthesis was performed. The JBI tool for case series was used.

PROSPERO: CRD42023391736.

RESULTS: 126 studies were included in the overall systematic review. 45 studies on terror & suicide bombings and 12 studies on landmines & UXOs are analyzed in the present paper. Studies on landmines from 1980 to 2006 included 13,270 individuals (age: 24.0 years (Range 8.2 years - 32.2 years); mainly male (88.9%)) in post-conflict low- and middle-income countries. Lower extremity injuries including amputations were most frequent, followed by eye injuries. Studies on terrorist attacks with a total of 12,518 mainly civilian victims (average age: 33,1 years; 61.6% males) reported on incidents in post-conflict regions and high-income cities mainly from the early 2000s. The injury pattern was multidimensional, with extremity injuries most frequent (28%-30%), including fractures, lacerations, and traumatic amputations (18%-20%).

CONCLUSION: This Systematic Review Part 2 [...]

[PubMed](#) · [DOI](#)

Is nail plate fixation the way forward for distal femur fractures with long segmental metaphyseal comminution? - A study of 56 patients.

Mahesan H, Sundaram VP, Devendra A, Ramesh P, Dheenadhyalan J, Rajasekaran S

BACKGROUND: Mechanical instability due to long metaphyseal comminution and medial bone gap after lateral locking plate results in varus collapse and implant failure. We believe that fixation in the form of Nail and plate combination provides better stability and prevents complications.

MATERIALS: We performed a retrospective analysis of 56 distal femur fractures with long metaphyseal comminution treated in our institution with nail and plate construct from 2017 to 2023. The mean age was 43.1 years and the mean follow up was 24 Months (4-36 months). Of these 56 patients, 16 were closed injuries underwent primary fixation, and 40 were open injuries requiring staged reconstruction. At the final follow up, patients were assessed for shortening, knee range of movements, Sander et al. functional score, radiological union and loss in alignment.

RESULTS: The mean length of metaphyseal comminution was 10.15 cm(5-18 cm). Primary Iliac crest bone grafting was done in 25 patients. Of the 56 patients, 51 achieved primary union and the mean time to union was 6 months. Of the remaining, 3 patients united after secondary bone grafting and 2 patients had nonunion. The mean loss in aLDFA at the final follow up was 1.75° in 7 patients. Shortening of the limb was noted in 16 patients, and 4 patients had significant shortening of 2 cm. Three patients developed deep SSI, of which one patient requi [...]

[PubMed](#) · [DOI](#)

Impact of modified ERAS protocol implementation on length of stay and complications in elderly hip fracture patients: a before-and-after study.

Liu Y, Ou G, Xie M, Wu F, Wang X, Yang B

AIM: Hip fractures in elderly patients pose significant healthcare challenges with high complication rates and prolonged hospital stays. Enhanced Recovery After Surgery (ERAS) protocols have shown promise in various surgical specialties, but their application in elderly hip fracture patients, particularly in Traditional Chinese Medicine (TCM) hospital settings, remains limited.

METHODS: This retrospective before-and-after study was conducted at a TCM hospital from July 2022 to July 2025. A total of 300 elderly hip fracture patients (≥ 65 years) were included: 146 in the control group (July 2022 - October 2023) receiving conventional care and 154 in the ERAS group (February 2024 - July 2025) receiving modified ERAS protocol incorporating TCM elements. Primary outcomes were length of hospital stay and perioperative complications within 30 days. Secondary outcomes included functional recovery [Harris Hip Score (HHS), Modified Barthel Index (MBI)], time to first mobilization, pain scores, and healthcare costs.

RESULTS: The ERAS group demonstrated significantly shorter length of stay compared with controls (12.38 ± 3.75 vs. 16.24 ± 5.10 days; mean difference 3.86 days, 95% CI: 2.84-4.88; $P < 0.001$). Total perioperative complications were reduced (7.79% vs. 15.07%; risk difference: -7.28%, 95% CI: -14.5% to -0.1%; $P = 0.047$). The ERAS group showed superior functional outcomes with [...]

[PubMed](#) · [DOI](#)

Prehospital endotracheal intubation and 30-day survival in severe trauma: a matched cohort study from Navarra, Spain.

Zulet Murillo D, Ferraz Torres M, Fortún Moral M, Belzunegui Otano T, Tamayo Rodríguez I

PURPOSE: Prehospital endotracheal intubation (ETI) is frequently used in major trauma, but its association with 30-day survival remains debated. We evaluated the association between prehospital ETI and 30-day survival and assessed whether sex modifies the age-related increase in mortality risk.

METHODS: We conducted a retrospective cohort study using the Navarra Major Trauma Registry (2010-2019). Patients aged ≥ 15 years with New Injury Severity Score (NISS) ≥ 15 were included; patients who died before hospital admission or had incomplete records were excluded. To mitigate confounding by indication, intubated and non-intubated patients were matched 1:1 using prehospital Glasgow Coma Scale (GCS), respiratory distress, and prehospital cardiac arrest as matching variables. Because the proportional hazards assumption for Cox regression was not met, parametric Weibull survival models were fitted, including a sex-by-age interaction term.

RESULTS: Overall, 1,909 patients were included; 212 (11.1%) underwent prehospital ETI. Matching yielded 190 pairs ($n=380$). In the matched cohort, prehospital ETI was not independently associated with 30-day survival (adjusted HR 0.91, 95% CI 0.65-1.27). Mortality risk was independently associated with lower GCS (per 1-point increase: HR 0.86, 95% CI 0.82-0.91) and respiratory distress (HR 7.63, 95% CI 4.34-13.4). A significant sex-by-age interaction [...]

[PubMed](#) · [DOI](#)

Mechanical and manual cardiopulmonary resuscitation in traumatic cardiac arrest: a retrospective observational study.

El-Menyar A, Ahmed K, Howland IR, Mollazehi M, Mekkodathil A, Rizoli S, Cameron P, de Oliveira Manoel AL et al.

BACKGROUND: Prehospital Traumatic cardiac arrest has a high fatality rate despite advances in trauma systems and resuscitation strategies. Mechanical chest compression devices have been increasingly adopted to optimize cardiopulmonary resuscitation (CPR) during Emergency Medical Services (EMS) transportation. We aimed to evaluate the impact of CPR (mechanical during transportation versus manual CPR only at the scene) on survival among trauma patients.

METHODS: A retrospective analysis was conducted for patients who received CPR at the scene and during transportation to the hospital between 2016 and 2024.

RESULTS: A total of 610 patients sustaining blunt traumatic cardiac arrest were included. The mean age was 34 ± 12.6 years; 94.6% were male, and 5.4% were female patients. Two hundred twenty (36.1%) received mechanical CPR during transportation to the hospital, and 390 (63.9%) received manual CPR only at the scene. Compared with the manual CPR group, the mechanical CPR group had higher rates of primary chest injury (30.5% vs. 20.5%; $p = 0.006$), bystander CPR (15.9% vs. 6.7%; $p = 0.001$), adrenaline administrations (95% vs. 26.4%; $p = 0.001$), and initial non-shockable rhythm (80% vs. 27.2%; $p = 0.001$). The two groups were comparable in median Injury Severity Score (ISS), head Abbreviated Injury Score (AIS), and total transport time. Mechanical CPR was associated with a markedly [...]

[PubMed](#) · [DOI](#)

Cervical spine clearance in pediatric trauma patients - Consensus algorithm of the Pediatric Spinal Trauma Group of the Spine Section of the German Society for Orthopedic and Trauma Surgery (DGOU).

Bolte J, R  ther H, Brecht P, Wiersbicki DW, Youssef Y, Jarvers JS, Disch AC

PURPOSE: Although cervical spine injuries in pediatric patients are rare, clinical and radiological examinations are clinical routine. Currently, no standardized decision algorithm does exist for detection of these injuries in the

European region. The proposed algorithm is aiming to safely identify significant injuries through consistent diagnostics while minimizing immobilization and radiation exposure at the same time.

METHODS: The Pediatric Spinal Trauma Group of the Spine Section of the German Society for Orthopedic and Trauma Surgery (DGOU) developed an algorithm based on own scientific investigations and considering primary and secondary literature. The algorithm was developed through a formal consensus process and thereby aligns with international recommendations.

RESULTS: The algorithm uses the initial pediatric Glasgow Coma Scale (pGCS) to group patients to three treatment pathways. Resulting recommendations for imaging, clinical re-assessment, and further management include the child's age, trauma mechanism, and clinical evaluations.

CONCLUSION: Despite their rarity, potential cervical spine injury in children requires accurate diagnosis. Clinical examinations can be challenging, especially in children with compromised consciousness. Increased radiation exposure from X-rays and computer tomography (CT) scans must be considered, particularly as magnetic resonance imaging [...]

[PubMed](#) · [DOI](#)

Development of a low-cost external fixation system: an off-label solution for resource-limited settings.

Seiser M, Deininger C, Fierlbeck J, Hollensteiner M, Coles P, Reuter C, Tempfer H, Traweger A et al.

PURPOSE: External fixators are essential for the surgical management of long-bone fractures, particularly in resource-limited settings where access to commercial systems is often restricted by cost and logistics. This study aimed to develop a low-cost external fixation concept using widely available construction materials and evaluate its biomechanical performance under axial loading.

METHODS: An external fixator composed of M10 threaded rods, washers, and nuts was developed. Five configurations were tested: SP (Steinmann pins), KT (threaded K-wires), KS (smooth K-wires), and BKS (bilaterally applied, smooth K-wires). A commercial external fixator served as control (CG). Juvenile bovine ulnae with a standardized fracture gap were subjected to cyclic axial loading. Construct stiffness, elastic deformation, and plastic deformation were assessed.

RESULTS: CG showed the highest stiffness (107.80 ± 14.12 N/mm), followed by SP (87.73 ± 17.92 N/mm), BKS (83.81 ± 10.01 N/mm), KS (49.84 ± 3.80 N/mm), and KT (46.66 ± 2.39 N/mm). Post-hoc analyses showed no significant differences between CG vs. SP and CG vs. BKS, whereas KT and KS were significantly less stiff. Elastic deformation followed a similar pattern. Plastic deformation was greatest in CG, while SP and BKS showed significantly lower values.

CONCLUSIONS: The threaded-rod fixator provides relevant mechanical stability at extreme [...]

[PubMed](#) · [DOI](#)

Supplementary fixation in multifragmentary midshaft clavicle fractures: do interfragmentary screws or suture cerclage improve outcomes compared with plate-only fixation?

Barthel C, Hinkelmann MA, Schibler D, Frei F, Pape HC, Allemann F

BACKGROUND: Clavicle fractures are common injuries. As displaced or multifragmentary midshaft patterns carry a substantial risk of malunion and nonunion with nonoperative treatment, they have contributed to the increasing use of surgical fixation. Modern operative techniques include plate fixation, intramedullary devices, and hybrid strategies to address comminution and improve fragment stability. Supplementary interfragmentary screws and cerclage constructs have been proposed to enhance fragment control and optimize load transfer in multifragmentary fracture patterns. This study aimed to compare three fixation strategies for operatively treated midshaft clavicle fractures managed with plate osteosynthesis: Group IS (interfragmentary screw), Group CA (cerclage augmentation), and Group POF (plate-only fixation), with respect to operative time, complication rates, and potential biological and mechanical differences.

METHODS: This retrospective study included patients aged ≥ 16 years who underwent plate fixation for multifragmentary clavicle fractures between 2016 and 2024. Patients were categorized into three groups: Group IS (supplementary interfragmentary screw), Group CA (suture cerclage augmentation), and Group POF (plate-only fixation). Demographic, surgical, and radiographic data were collected. Callus formation was assessed by three

independent observers. Statistical anal [...]

[PubMed](#) · [DOI](#)

Cerebral glycerol during haemorrhagic shock in normal and raised intracranial pressure resuscitated with total REBOA: an experimental porcine study.

Bader S, Magnuson A, Brorsson C, Wallin G, Löfgren N, Löfgren F, Blind PJ, Öman M et al.

BACKGROUND: Haemorrhagic shock (HS) is frequently associated with secondary cerebral injury due to compromised cerebral perfusion. Resuscitative endovascular balloon occlusion of the aorta (REBOA) is an effective haemorrhage control strategy; however, concerns persist regarding its cerebral effects, particularly in the presence of elevated intracranial pressure (ICP). Cerebral microdialysis (CMD) derived glycerol (CGly) is a recognised marker of cellular membrane stress and injury.

OBJECTIVE: To investigate CGly dynamics during prolonged total REBOA (tREBOA) in HS and to compare cerebral cellular responses between animals with normal and elevated ICP.

METHODS: In this experimental porcine study, eighteen pigs were subjected to controlled HS and resuscitated with tREBOA for 90 min. Animals were allocated to either a normal ICP group (NICPG, n = 9) or an elevated ICP group (EICPG, n = 9). Continuous monitoring of proximal arterial pressure, ICP, and cerebral perfusion pressure was performed. CGly concentrations were measured using CMD throughout the experimental protocol.

RESULTS: tREBOA effectively restored proximal arterial pressure and cerebral perfusion pressure in both groups. CGly concentrations remained relatively stable during the haemorrhage phase and increased progressively during prolonged aortic occlusion. Importantly, CGly responses did not differ significantly be [...]

[PubMed](#) · [DOI](#)

Transport and destination hospital for patients with suspected multiple and/or severe injuries - a systematic review and clinical practice guideline update.

Pavlu F, Könsgen N, Bieler D, Goossen K, Gliwitzky B, Häske D, Faul P, Wölfl C et al.

PURPOSE: Our aim was to update the evidence-based and consensus-based recommendations on criteria for transport and destination hospital for patients with suspected multiple and/or severe injuries based on available evidence. This guideline topic is part of the 2025 update of the German Guideline on the Treatment of Patients with Multiple and/or Severe Injuries.

METHODS: MEDLINE and Embase were systematically searched to January/February 2024. Further literature reports were obtained from clinical experts. Randomised controlled trials (RCTs) or observational studies reporting risk-adjusted outcomes were included if they compared different transport strategies or destination hospitals for patients with (suspected) multiple and/or severe injuries. We considered patient-relevant outcomes such as mortality and neurological outcomes. Risk of bias was assessed at outcome level using ROBINS-I for observational studies and RoB 2 for RCTs. We conducted meta-analyses if possible; alternatively, we synthesised the evidence narratively. We used GRADE to rate the certainty of evidence. Expert consensus was used to develop recommendations and determine their strength.

RESULTS: A total of 127 studies were identified. They addressed helicopter vs. ground emergency medical services (EMS) transport, ground EMS with vs. without a physician, on-scene time, and handover on arrival at the hospital [...]

[PubMed](#) · [DOI](#)

Does the addition of peripheral nerve blocks improve analgesia during distal radius fracture reduction? A prospective, randomized, multicenter clinical trial.

Guerrero-Serrano JA, Santamaría-López A, García-Lamas L, Jiménez-Díaz V, González-Fernández Á, Cecilia D

PURPOSE: This study aimed to evaluate whether adding a median nerve and a superficial branch of the radial nerve block to hematoma block improves analgesia during distal radius fracture reduction and to identify factors associated with increased pain perception.

METHODS: A prospective, randomized, controlled study was conducted in two tertiary hospitals, including 180 adult patients with displaced distal radius fractures requiring closed reduction. Patients were allocated either to an

isolated hematoma block group or to a combined block group receiving hematoma block with median nerve and superficial branch of the radial nerve blocks. Within the combined group, half of the median nerve blocks were performed under ultrasound guidance and half using anatomical landmarks. The primary endpoint was digital pain during traction and fracture reduction, assessed using a visual analog scale (VAS, 0-10). Secondary outcomes included wrist pain at different procedural stages and the influence of demographic and clinical variables including age, sex, laterality, participating hospital, body mass index (BMI), fracture type (AO/OTA), and baseline psychiatric or chronic analgesic medication use.

RESULTS: Digital pain was significantly lower in combined group compared with isolated hematoma block during traction (4.54 ± 2.53 vs. 7.39 ± 1.76 ; $p < 0.001$) and reduction (5.36 ± 2.56 vs. $8.33 \pm 1. [...]$

[PubMed](#) · [DOI](#)

Under pressure - a sensor-based analysis of simulated prehospital pressure dressing application for hemorrhage management.

Brauckmann V, Menzel J, Welke B, Decker S, Macke C

BACKGROUND: The concept of "x-ABCDE", with "X" representing exsanguinating hemorrhage, highlights the priority of immediate bleeding control in trauma care. Despite being a fundamental intervention, pressure dressing techniques lack standardization. Studies show high variability in achieved pressures, often below therapeutic targets or exceeding harmful levels, with no correlation between subjective estimates and objective measurements.

OBJECTIVES: To objectively assess current practices of pressure bandage application and identify influencing factors including material type, technique, and provider experience.

METHODS: This prospective, single-center, simulation-based observational study was conducted from August 2024 to January 2025 at a Level I Trauma Center. Emergency medical providers applied pressure dressings according to personal preference using available materials. Pressure distribution was measured via two calibrated capacitive pressure sensors (61 measurement fields). Analysis included correlation tests, group comparisons, and multiple regression.

RESULTS: A total of 124 emergency medical providers (75% male; 36% paramedics, 44% emergency medical technicians, 13% trainees, 7% emergency physicians) completed pressure dressing applications, with 116 datasets included in final analysis. Mean maximum pressure was 169.35 ± 84.33 mmHg (range 50-412 mmHg) with applicati [...]

[PubMed](#) · [DOI](#)

Clinical impact of early postoperative routine radiology in isolated pelvic injuries: a retrospective study.

Bolierakis E, Ioannou P, Schubert SL, Groven RVM, Bläsius FM, Bode M, Alabdulrahman H, Hildebrand F et al.

PURPOSE: The clinical value of routine immediate postoperative conventional radiography following surgical fixation of pelvic fractures remains debated. This study assessed the impact of early postoperative X-rays on management decisions and revision surgery in patients with isolated pelvic ring or acetabular fractures.

METHODS: We conducted a retrospective cohort study at a Level 1 trauma center, including adults undergoing surgical treatment for isolated pelvic ring or acetabular fractures between 2020 and 2023. All patients received high-quality intraoperative fluoroscopy and postoperative conventional radiographs. The primary outcome was any change in standard postoperative care based on radiographic findings. The secondary outcome was revision surgery prompted by conventional postoperative imaging.

RESULTS: A total of 216 patients were included (median age 63 years), with 156 pelvic ring injuries and 60 acetabular fractures. Postoperative radiographs were obtained at a median of 3 days post-surgery. Changes in postoperative care based on radiographs occurred in 2 patients (3.3%) within the acetabular fracture group. Revision surgery triggered by conventional imaging was required in 1 patient (0.6%) among pelvic ring injuries. In all cases, additional CT imaging preceded definitive management changes.

CONCLUSION: Routine immediate postoperative conventional radiography a [...]

[PubMed](#) · [DOI](#)

Older age and post-traumatic organ failure a TraumaRegister DGU® analysis of 34,469 elderly major trauma patients.

Bewersdorf TN, Lefering R, Stein S, Streblow J, Findeisen S, Schmidmaier G, Grossner T

PURPOSE: As the number of elderly major trauma patients (EMTP) increases continuously, this study sought to determine the incidence of organ failure (OF) and multiple organ failure (MOF) and their impact on mortality and recovery status in this cohort.

METHODS: Based on the German TraumaRegister DGU® database this study included EMTP aged between 65 and 99 years with an Injury Severity Score (ISS) of ≥ 16 between 2014 and 2023. OF was defined as Sequential Organ Failure Assessment (SOFA) score of ≥ 3 and MOF as failure of ≥ 2 organs.

RESULTS: The database revealed 34,469 EMTP, here 37% suffered from OF (16.0%) or MOF (21.0%). Mortality was significantly higher in OF and MOF patients compared to non-OF patients (non-OF: 7.4%; OF: 45.0%; MOF: 59.0%; $p < 0.001$). 80.9% of all deaths were associated with failure of at least one organ and 51.1% of non-survivors suffered from MOF. Highest odds ratios (OR) for mortality were found for liver (OR: 5.13, $p < 0.001$), central nervous system (OR: 3.63, $p < 0.001$) and renal failure (OR: 3.25, $p < 0.001$), followed by cardiovascular (OR: 1.95, $p < 0.001$) and lung failure (OR: 1.16, $p = 0.010$). Haemostatic failure had no impact on mortality.

CONCLUSION: OR and mortality differ significantly between distinct OFs due to varying degrees of success in treatment options and systemic impact of these OFs. Clinicians should be aware that OF and MOF o [...]

[PubMed](#) · [DOI](#)

Denis pattern-dependent biomechanical behavior of posterior trans-ilial plate fixation compared with bilateral triangular osteosynthesis in unilateral sacral fractures: a finite element study.

Yang J, Böker KO, Li C, Schilling AF, Ritter Z, Acharya M, Lehmann W

OBJECTIVE: Posterior trans-ilial plate osteosynthesis (TPO) is commonly used for the stabilization of unilateral vertically unstable sacral fractures. However, its biomechanical performance across different Denis fracture patterns remains unclear. This study aimed to investigate the fracture pattern-dependent biomechanical behavior of posterior trans-ilial plate osteosynthesis (TPO) in unilateral vertically unstable sacral fractures, using bilateral triangular osteosynthesis (BTO) as a high-stability reference construct.

METHODS: A validated three-dimensional finite element model of the lumbopelvic complex was developed to simulate unilateral vertically unstable sacral fractures corresponding to Denis type I, II, and III patterns. Two posterior fixation strategies, TPO and BTO, were constructed for each fracture type. Seven physiological loading conditions were applied, including standing, flexion, extension, axial rotation, and lateral bending. Von Mises stress distribution and fracture displacement were evaluated. Fracture micromotion was quantified by calculating relative displacement changes between paired points along the fracture gap.

RESULTS: When interpreted relative to the high-stability BTO reference construct, TPO showed fracture pattern-dependent variation in fracture displacement. The difference was most pronounced in Denis type I fractures and progressively decre [...]

[PubMed](#) · [DOI](#)

Factors associated with in-hospital mortality in necrotising soft tissue infections. a multicentre retrospective cohort study.

Podda M, Ceresoli M, Viridis F, Rosa F, Pilia T, Murzi V, Dessì A, Tedesco S et al.

PURPOSE: Necrotising soft tissue infections (NSTIs) are rare but life-threatening conditions associated with high mortality rates. This multicentre study aimed to identify admission variables associated with in-hospital mortality.

METHODS: This retrospective study included adult patients with surgically confirmed NSTIs treated at four high-volume academic referral centres in Italy between 2010 and 2024. Demographic, clinical, physiological, and laboratory variables available at hospital admission were analysed. Categorical variables were compared between survivors and non-survivors using the chi-square test or Fisher's exact test, while quantitative variables were compared using the Student's t test or Mann-Whitney U test. Multivariable logistic regression analyses were performed to identify independent predictors of in-hospital mortality. Results were reported as ORs with 95%CI. A

prognostic nomogram was developed from the final multivariable model, including variables independently associated with mortality.

RESULTS: A total of 379 patients were included. In-hospital mortality was 16.7%. In subgroup comparisons, mortality was 17.0% in necrotising fasciitis and 22.4% in Fournier's gangrene ($P = 0.275$). In the overall NSTI population, age (aOR 1.061, 95%CI 1.037-1.087) and serum creatinine at admission (aOR 1.301, 95%CI 1.040-1.629) were independently associated with mortality [...]

[PubMed](#) · [DOI](#)

Influence of frailty on short-term mortality in older patients with multiple trauma in the emergency department.

Çelik ■, Toker A, Biçer S, Cebecio■lu ■K

PURPOSE: To investigate the association between frailty and 28-day mortality in older patients presenting to the emergency department with multiple trauma and to assess the discriminative performance of PRISMA-7 and the Trauma-Specific Frailty Index (TSFI) in frailty assessment in relation to commonly used trauma severity scores.

METHODS: This prospective observational study included patients aged ≥ 65 years with multiple trauma admitted to the emergency department between July 1 and August 31, 2024. Frailty was assessed using PRISMA-7 and TSFI. Patients were stratified into frail and non-frail groups and further categorized according to age groups (65-74, 75-84, and ≥ 85 years). Clinical characteristics, trauma severity scores, and outcomes, including 28-day mortality, were compared. Receiver operating characteristic (ROC) analysis was used to evaluate discriminative performance, and logistic regression analysis was performed to assess the association between scores and mortality.

RESULTS: A total of 144 patients were included. Frailty was significantly associated with increased 28-day mortality, higher ICU admission rates, and longer hospital stays ($p < 0.001$). Mortality increased with age, reaching 30.0% in patients aged ≥ 85 years ($p = 0.006$). Frailty scores (PRISMA-7 and TSFI) increased significantly with age, whereas ISS did not differ significantly across age groups. R [...]

[PubMed](#) · [DOI](#)

Mortality prediction in older adults with earthquake-related trauma: a model combining MEWS and laboratory biomarkers.

Ozen A, Acehan S, Satar S, Bulut EA, Gulen M, N■g■z K, Sevd■mbas S, Sonmez GO et al.

PURPOSE: This study aimed to identify laboratory parameters and clinical scores that can predict in-hospital mortality among patients aged 65 years and older who presented to the emergency department (ED) with earthquake-related trauma following the February 6 and 20, 2023 earthquakes.

METHODS: In this retrospective observational study, 362 patients with earthquake-related trauma between February 6-21, 2023, were included. Demographic characteristics, laboratory results, and clinical scores such as the Modified Early Warning Score (MEWS), Injury Severity Score (ISS), and Shock Index (SI) were analyzed.

RESULTS: Of the included patients, 53.9% ($n = 195$) were female, and the median age was 73 years (IQR: 69-80). The overall mortality rate was 9.4%. Multivariable logistic regression analysis identified MEWS (OR: 3.535; 95% CI: 2.397-5.212; $p < 0.001$), urea (OR: 1.009; 95% CI: 1.002-1.015; $p = 0.008$), and hs-Troponin I (OR: 1.001; 95% CI: 1.000-1.002; $p = 0.016$) as independent predictors of mortality. Receiver operating characteristic (ROC) analysis demonstrated an AUC of 0.936 for the combined model including MEWS, urea, and hs-Troponin I (95% CI: 0.875-0.996; $p < 0.001$). Among the variables, MEWS showed the highest AUC (0.903; 95% CI: 0.829-0.976). The optimal cutoff value for MEWS was determined as 1.5, with a sensitivity of 70.6% and a specificity of 97.6%.

CONCLUSIONS: In o [...]

[PubMed](#) · [DOI](#)

The economic burden in terms of cost of illness and generic health-related quality of life of posttraumatic long bone non-unions among the adult population of the Netherlands from a societal perspective.

van der Broeck LCA, Shchurov DG, Gabrio A, Lodewijks AJL, Poeze M, Blokhuis TJ, Evers S

PURPOSE: This study assessed the societal economic burden in terms of cost of illness and health-related quality of life (HRQoL) of posttraumatic long bone non-unions in the Netherlands.

METHODS: An incidence-based bottom-up approach was used, focusing on adult patients with posttraumatic long bone non-unions. The analysis included healthcare costs, patient and family costs, and productivity losses, measured using the iMCQ and iPCQ questionnaires. Cost evaluations followed Dutch costing guidelines, and productivity losses were calculated using the friction cost method. HRQoL was assessed with the EQ-5D-5L. A deterministic one-way sensitivity analysis varied baseline characteristics by $\pm 10\%$. Scenario analyses were conducted from healthcare and patient perspectives, as well as for patient subgroups.

RESULTS: Average costs per patient during the three months before the initial visit at a non-union clinic were €8,928 (healthcare, N = 78), €1,360 (patient and family, N = 58), and €4,313 (productivity losses, N = 59), the average of the observed data calculates to 10,831 (N = 44). The mean EQ-5D-5L utility score was 0.390 (± 0.29 SD). Subgroup analysis showed no significant cost increase for patients with infections or open fractures. However, a higher Non-Union Severity Score and a lower HRQoL were significantly associated with higher total costs.

CONCLUSION: Posttraumatic long [...]

[PubMed](#) · [DOI](#)

Nonoperative treatment of Maisonneuve fractures.

Frühauf J, Bartonířek J, Fojtík P, Horáková A, Tušek M

PURPOSE: The aim of this study is to describe the basic pathoanatomical characteristics of a stable Maisonneuve fracture and mid-term results of its nonoperative treatment.

METHODS: The study included 17 prospectively collected patients with a mean age of 59 years. The postinjury ankle CT had to meet the following criteria: nondisplaced or minimally displaced (up to 1 mm) fracture of medial malleolus, medial clear space less than 3 mm, nondisplaced or minimally displaced (up to 2 mm) fracture of posterior malleolus, anatomical position or minimal malposition of the distal fibula in the fibular notch (widening of the tibiofibular space up to 2 mm or external rotation of the distal fibula up to 10°). The average follow-up period was 34 months, the final follow-up included CT examination and functional evaluation based on AOFAS and FAOS scores.

RESULTS: A medial malleolus fracture was recorded in 12% cases, a posterior malleolus fracture in 29% patients and a Tillaux-Chaput tubercle fracture in 18% cases. All fractures of the proximal fibula, medial and posterior malleolus healed within 3 months. The position of the distal fibula in the fibular notch did not change in 11 cases compared to the post-injury CT scan, improved slightly in 5 cases, and worsened slightly in 1 case. The average final AOFAS hindfoot score was 96.9 points and the average final FAOS score was 98.7%.

CONCL [...]

[PubMed](#) · [DOI](#)

Age-related differences in prehospital triage and critical interventions: a multicenter TraumaRegister DGU® analysis.

Koch DA, Hagebusch P, Hackenberg L, Pavlu F, Jakobi T, Münzberg M, Lefering R, Schweigkofler U et al.

PURPOSE: Older trauma patients are known to be undertriaged in the prehospital setting. However, it remains unclear whether these disparities translate into differences in the delivery of critical prehospital interventions. This study aimed to evaluate age-related differences in prehospital triage, allocation, and treatment within a physician-staffed emergency medical service system.

METHODS: A retrospective multicenter analysis of the TraumaRegister DGU® was conducted, including trauma patients aged ≥ 20 years between 2020 and 2023 with trauma team activation and subsequent intensive care unit admission. Patients were stratified into two age groups, 20-59 vs. ≥ 60 years, with additional subgroups up to 90-99 years. Prehospital interventions, transport modality, and trauma center allocation were analyzed. Furthermore, predefined high-risk scenarios (severe motor vehicle trauma, traumatic brain injury with GCS ≤ 8 , and pneumothorax) were evaluated.

RESULTS: A total of 67,069 patients were included, with comparable injury severity across age groups (mean ISS 20.2). Older patients were less frequently transported by air and less often allocated to supra-regional trauma centers. While overall intervention rates were similar, differences emerged in high-risk scenarios. Older patients received endotracheal intubation and tranexamic acid less frequently despite comparable injury severity [...]

[PubMed](#) · [DOI](#)

Prediction of risk of death in severely injured patients: the revised injury severity classification score, version 3 (RISC III).

Lefering R, Imach S, Bieler D

PURPOSE: Trauma Registries with a focus on severely injured patients use survival as their primary outcome. In order to compare hospitals, interventions, and changes over time, a precise risk of death prediction is mandatory. The German TraumaRegister DGU® uses the RISC II model (Revised Injury Severity Classification, version II) since 2013. The aging trauma population and an improved handling of missing data required the present revision.

METHODS: A total of 53,738 seriously injured trauma patients documented in 2022-2023 served as basis for development and validation (3:1 ratio). Missing values should now be considered to be within normal physiological range except in patients with specific findings indicative for an altered physiology. These findings were suggested by clinical experts and validated by registry data. The increasing age of trauma patients was addressed by additional categories and higher point weights. A logistic regression analysis provided new point weights for all predictors. Precision (observed versus predicted mortality) and discrimination (area under the receiver operating characteristic curve, AUROC) were calculated in the development and validation dataset.

RESULTS: Patients in both datasets were well comparable, with a mean age of 55 years, 69% males, and an average Injury Severity Score (ISS) of 18 points. The rate of missing values ranged from 0% [...]

[PubMed](#) · [DOI](#)

Dislodgement forces and cost effectiveness of prehospital securement of peripheral intravenous catheters on moist skin: a randomized controlled clinical trial in healthy volunteers.

Vogeler J, Schumann S, Heinrich S, Hell J, Schmutz A

OBJECTIVES: Peripheral intravenous catheters (PIVC) are widely used clinical devices for administering fluids or essential drugs to patients. The failure rate of PIVC remains high and Emergency Medicine has been shown to be a risk factor for dislodgement. When the skin is moist from sweat or fluids, standard intra-hospital dressings and securements fail. In emergency situations, a failed catheter can then critically delay intravenous therapies. The most effective dressing to prevent accidental removal of a prehospital PIVC remains unclear. It was the aim of this study to compare the force required to dislodge a PIVC with four different methods of securing PIVCs used in emergency medicine. In addition, the costs were calculated.

METHODS: Artificial sweat was applied to the skin of 180 volunteers. PIVCs were attached onto the forearm using four different securements (elastic gauze, cohesive gauze, clingfilm and a velcro securing device). Continuously increasing traction force was applied until dislodgement of the respective securement. For statistical tests, either Friedman's test or repeated measures ANOVA was used.

RESULTS: Clingfilm showed the greatest resistance to increasing pulling force with cohesive bandages as close second. The velcro securing device was strongest at resisting low level of forces but fell off sharply at higher force. Elastic bandages were the weakest [...]

[PubMed](#) · [DOI](#)

Injury to the thorax is a dominant contributor to post-traumatic neutrophil mediated inflammatory response.

van Eerten FJC, Duebel LV, de Fraiture EJ, Hanlo MF, Breuking EA, Brands SR, van den Bosch TD, van Wessem KJP et al.

PURPOSE: Traumatic injury induces a neutrophil associated inflammatory response mediated by the release of damage-associated molecular patterns (DAMPs). This response can be maladaptive, increasing risk of infection. Since blunt thoracic injuries are associated with high infection rates, we hypothesized that these injuries elicit a

stronger neutrophil associated inflammatory response than injuries to other body regions.

METHODS: A cohort study was conducted in adult patients presenting at the trauma bay of a level 1 trauma center between March 2020 and November 2022. These patients underwent routine analysis of their neutrophil associated inflammatory response by phenotyping blood neutrophils using automated flow cytometry. Injury per body region was counted when the Abbreviated Injury Severity (AIS) scores were ≥ 3 to focus on substantial injuries. The primary outcome was an extensive post-traumatic inflammatory neutrophil response.

RESULTS: 861 patients were included. Blunt internal thoracic injuries showed the strongest association with severe neutrophil associated inflammation (OR 5.7), followed by abdominal (OR 2.6), pelvis (OR 2.5) and lower extremities (OR 2.3), even when corrected for ISS, age and hemodynamic instability. In multivariable analysis, only thoracic internal injuries remained a statistically significant contributor to excessive neutrophil associated inflam [...]

[PubMed](#) · [DOI](#)

Major improvement using peroperative 3D compared to 2D imaging for SI screw fixation in traumatic pelvic fractures.

van Eerten FJC, Benders KEM, van Baal MCPM, Hietbrink F

INTRODUCTION: Pelvic fractures, though accounting for only 3% of skeletal injuries, carry a high mortality rate, primarily due to the associated risk of hemodynamic instability. Sacroiliac (SI) screw fixation is a common surgical intervention, which can be technically challenging due to the complex pelvic anatomy. Imaging techniques have evolved from 2D to 3D imaging for its improved accuracy in screw placement. The impact of this transition for SI screw placement was evaluated.

METHOD: A single-center observational retrospective study was performed between 2013 and 2023. Patients ≥ 18 years who underwent SI screw placement following pelvic injuries were analyzed. 2D imaging was used until 2021, after which 3D imaging was implemented. Outcomes included the need for revision surgery during initial admission and < 18 months after trauma.

RESULTS: 2D imaging was used in 42 patients and 3D imaging in 44 patients. Baseline characteristics and injury severity of both groups were comparable. In the 2D group, 18 patients (42.9%) had suboptimal placement of the SI screw. Of these, 3 (7.1%) required revision surgery due to malpositioned screws. In contrast, no patients (0.0%) in the 3D group had suboptimally positioned screws or underwent revision surgery, ($p < 0.001$, $p = 0.22$).

DISCUSSION: Intraoperative 3D imaging enhanced the accuracy of SI screw placement, leading to an encouraging [...]

[PubMed](#) · [DOI](#)

Minimally invasive versus open fixation for flail chest in elderly patients: a prospective cohort study on perioperative safety and long-term outcomes.

Tian Y, Zheng Z, Ji Z, Lu W, Song N, Wang J

OBJECTIVE: To systematically evaluate the perioperative safety and short- and long-term efficacy of minimally invasive fixation using a reversed shape-memory alloy plate versus conventional open fixation for the treatment of flail chest in elderly patients.

METHODS: This prospective cohort study enrolled 130 elderly patients (≥ 60 years) with flail chest between January 2022 and January 2026. After 1:1 propensity score matching, 100 patients were included (minimally invasive group, $n = 50$; open group, $n = 50$). Perioperative parameters (operative time, blood loss, C-reactive protein(CRP) at 24 h/48 h/72 h, complications), short-term outcomes (resting/cough-evoked Numerical Rating Scale(NRS) pain scores, non-steroidal analgesic drug duration, rescue analgesic frequency, Patient-Controlled Analgesia Pump(PCA) consumption, time to ambulation, hospital stay), and long-term outcomes (implant dislodgement, 6-month pulmonary function, Douleur Neuropathique 4 Questions Naïve(DN4), Medical Outcomes Study 36- Item Short Form Health Survey(SF-36)) were compared.

RESULTS: The minimally invasive group had significantly fewer incision-related complications ($P = 0.046$), lower resting NRS scores on postoperative days 1 and 3 ($P < 0.01$; $P = 0.041$), lower cough-evoked NRS score on day 1 ($P = 0.011$), shorter non-steroidal analgesic duration ($P < 0.01$), and lower PCA consumption ($P = 0.027$). At 6 [...]

Epidemiology and outcomes of traumatic sternal fractures and associated blunt cardiac injury: a nationwide cohort study in the Netherlands.

Klei DS, Benders KEM, Leenen LPH, van Wessem KJP

PURPOSE: Comprehensive data on epidemiology, trauma mechanisms, associated injuries, and outcomes of traumatic sternal fractures are scarce. This study analysed nationwide data to improve diagnosis and management within the Dutch healthcare system.

METHODS: This nationwide retrospective cohort study using the Dutch National Trauma Registry included adult patients admitted with traumatic sternal fractures between 2015 and 2023. Patients with prehospital cardiopulmonary resuscitation or penetrating trauma were excluded. Incidence, patient characteristics, trauma mechanisms, associated injuries, and in-hospital outcomes were analysed. Subgroup analyses evaluated patients with concomitant blunt cardiac injury (BCI).

RESULTS: Of 568,399 adult trauma admissions, 4,765 patients (0.84%) sustained traumatic sternal fractures. Median age was 62 years; 60% were male. Motor vehicle accidents (48%) and falls (28%) were the leading mechanisms. 35% were severely injured (ISS ≥ 16). Associated injuries included rib fractures (51%), spinal fractures (36%), and lung contusions (18%). Critical care unit admission was 40%, with median mechanical ventilation duration of 4 days; median hospital stay was 5 days. In-hospital mortality was 5.7%, and 30-day mortality 6.0%. BCI occurred in 9.5% of patients and was associated with a higher number of injuries and increased injury severity, emergency inte [...]

Trauma patients ≥ 65 years old constitute 40% of severely injured adult trauma patients in Sweden - a 10-year analysis of geriatric trauma prevalence.

Bredberg S, Modalen E, Jöneby C, Nilsson D, Lapidus O

BACKGROUND: The elderly population in Sweden is steadily increasing. Age is an independent predictor of mortality in trauma patients. There is growing evidence that geriatric patients constitute an increasing proportion of the Swedish trauma population. No studies have analyzed temporal trends in the proportion of geriatric trauma patients in the context of the national trauma population in Sweden.

AIM: To determine the proportion of geriatric patients in the Swedish trauma population and analyze temporal trends in geriatric trauma prevalence.

METHODS: The study cohort was 11,969 severely injured trauma patients (NISS > 15) treated in Sweden during 2013-2022, obtained from the Swedish Trauma Registry. The proportion of geriatric patients was compiled annually. Temporal trends were analyzed using weighted linear regression models. Two subgroups were defined based on injury severity (NISS ≥ 25 and NISS ≥ 40).

RESULTS: The proportion of patients ≥ 65 years in the adult trauma population increased from 30% to 40% during 2013-2022 ($p = 0.003$), while the proportion of patient ≥ 80 years also increased, from 8.6% to 17% ($p = 0.009$). Similar increases were also seen in the NISS ≥ 25 and NISS ≥ 40 subgroups.

CONCLUSION: There is likely an increase in geriatric trauma prevalence in Sweden, but the full extent of this temporal trend remains uncertain. Trauma patients ≥ 65 years old co [...]

A totally laparoscopic strategy combining microwave ablation and autologous blood transfusion for patients with initial WSES grade IV traumatic splenic rupture in selected transient responders: a proof-of-concept study.

Feng Z, Xie D, Hong W, Zheng C

PURPOSE: Splenic artery embolization (SAE) is the standard intervention for transient responders with high-grade splenic trauma. When SAE is unavailable, contraindicated, or declined, alternative minimally invasive approaches may be warranted. This study evaluates a totally laparoscopic strategy combining microwave ablation (MWA) for hemostasis with intraoperative autologous blood transfusion (ABT) in selected transient responders with initial

WSES Grade IV splenic rupture.

METHODS: This retrospective study included 10 consecutive transient responders (October 2021-June 2024) with initial WSES Grade IV splenic rupture who underwent totally laparoscopic MWA combined with intraoperative ABT. All patients exhibited transient hemodynamic response to limited resuscitation (≤ 1500 mL fluids) and preoperative shock index > 0.9 . Intraoperative injury was graded by the AAST system. Outcomes included technical success, transfusion requirements, complications, and follow-up.

RESULTS: All 10 patients underwent successful spleen-preserving laparoscopy without conversion. AAST grades: II (n = 2), III (n = 4), IV (n = 4). Mean operative time was 191.3 ± 101.7 min. Mean blood salvage was 1587.5 ± 663.9 mL, yielding 490.8 ± 80.2 mL autologous red cells. No patient required postoperative allogeneic transfusion. Hemoglobin, platelets, and lactate improved significantly by postoperative day 3 (P [...])

[PubMed](#) · [DOI](#)

Pediatric pancreatic trauma: a 10-year retrospective analysis from two high-complexity centers in Medellín, Colombia.

Barrientos ML, Calderon SP, Zapata CAL, Lopez CAD

PURPOSE: Pediatric pancreatic trauma is uncommon, but it is associated with significant morbidity and an ongoing debate regarding its management. Hemodynamic stability and pancreatic duct involvement are key factors in treatment decisions.

METHODS: A retrospective study analyzed 25 pediatric patients with pancreatic trauma treated between 2010 and 2020 at two high-complexity hospitals in Medellín. Clinical variables, trauma mechanism, imaging findings, management, and complications were analyzed.

RESULTS: Blunt trauma was the predominant mechanism (64%). Preoperative computed tomography was performed in 12 patients (48%); radiological AAST grading was not retrievable in 13 (52%). Immediate surgery was performed in 13 patients (52%), predominantly driven by associated intra-abdominal injuries rather than the pancreatic injury itself. Non-operative management was initially attempted in 9 patients (36%), with a 56% success rate; failures (44%) were related to clinical deterioration and progression of peripancreatic collections. Clinically relevant postoperative pancreatic fistula (ISGPS Grade B) occurred in 3 of 13 operated patients (23%), with a mean closure time of 55.6 days (range 18-110); no Grade A or Grade C fistulas were recorded. Other complications included post-traumatic pancreatitis (24%), pancreatic pseudocyst (8%), and organ/space surgical site infection (8%); most [...]

[PubMed](#) · [DOI](#)

Influence of Antithrombotics on Severely Injured Patients Over 50 With Blunt Abdominal Trauma.

Eibinger N, Limberger B, Hörlesberger N, Puchwein P, Lefering R, Bayer TP, Zumsteg JS, Pape HC et al.

INTRODUCTION: Severe abdominal trauma represents a critical subset of injuries, particularly in the aging European population, where the prevalence of preexisting anticoagulant or antiplatelet therapy is increasing. While the impact of such medications on outcomes in traumatic brain injury is well studied, limited data exist regarding their influence in patients with abdominal trauma.

MATERIALS AND METHODS: This retrospective cohort study used data from the TraumaRegister DGU® between January 2015 and December 2023. Inclusion criteria were patients aged ≥ 50 years with severe blunt abdominal trauma (AIS Abdomen ≥ 3) without relevant head injury (AIS Head ≤ 3) from Austria, Germany, or Switzerland. Patients were grouped according to pre-injury antithrombotic medication (no medication [NM] vs. antithrombotic medication [AM], with subgroups). Statistical analysis included descriptive comparisons, standardized mortality ratios (SMR) using RISC III, and multivariate logistic regression to identify independent predictors of hospital mortality.

RESULTS: Of 328,281 patients, 4,069 met inclusion criteria (2,831 NM; 1,238 ATM). Patients in the AM group were older (74.1 vs. 62.3 years) and had higher mortality (25.0% vs. 10.4%), particularly in the DOAC subgroup (30.3%). SMR analysis demonstrated increased mortality in the AM group compared to expected rates. In multivariate logistic re [...]

[PubMed](#) · [DOI](#)

Predicting limb amputation in trauma-related necrotizing fasciitis of the extremities: insights from a 10-year cohort.

Yang S, Ren Z, Li Y, Zhao Z, Sun C, Liu L, Jin L, Hou Z

BACKGROUND: Necrotizing fasciitis (NF) is a rapidly progressive, life-threatening soft tissue infection that in most cases requires aggressive surgical management. Major limb amputation is a devastating limb-related outcome. We retrospectively analyzed NF cases to quantify the rate of amputation and identify associated risk factors.

METHODS: We retrospectively reviewed the medical records of all patients with trauma-related extremity NF treated at Hebei Medical University Third Hospital from Jan 2014 to Dec 2024. Demographics, comorbidities, admission laboratories and key time intervals were compared between patients who did and did not undergo major limb amputation. Multivariable logistic regression identified independent predictors, and receiver operating characteristic (ROC) analysis assessed the discriminative power of continuous variables.

RESULTS: Of 134 patients, 22 (16.4%) required amputation. Compared with the non-amputation group, amputees had a markedly longer pre-hospital delay (median 242 h vs. 65 h), more multiple injuries (63.6% vs. 13.4%) and a higher prevalence of diabetes (59.1% vs. 21.4%). Admission lactate dehydrogenase (LDH) was substantially higher (median 589 vs. 182 U/L). In multivariable analysis, elevated LDH, prolonged time from traumatic injury to hospital admission, multiple injuries and diabetes were independently associated with amputation. LDH [...]

[PubMed](#) · [DOI](#)

Is 18 F-FDG PET-CT-guided geographic debridement superior to conventional debridement in appendicular chronic osteomyelitis? a retrospective comparative cohort study.

Elsheikh A, Elseidy H, Saad H

BACKGROUND: Determining the surgical extent of chronic osteomyelitis and fracture-related infection (FRI) remains challenging. Adequate resection margins prevent recurrence; however, no standardised preoperative tool objectively defines resection borders.

OBJECTIVE: To compare PET-CT-guided surgical planning versus the local standard-of-care imaging pathway (predominantly plain radiographs, with selective CT or MRI in a minority of patients) for chronic limb osteomyelitis and FRI outcomes.

METHODS: Retrospective comparative cohort study of 126 patients treated 2019-2024. Group 1: 84 patients underwent 18 F-FDG PET-CT imaging and map-based geographic debridement. Group 2: 42 patients received surgery after local standard-of-care imaging pathway (predominantly plain radiographs, with selective CT or MRI).

PRIMARY OUTCOME: infection recurrence at 12 months. Multivariable logistic regression controlled for confounders.

RESULTS: PET-CT group achieved 3.6% recurrence versus 16.7% in the conventional group ($p = 0.011$), representing 78.5% relative risk reduction and a number-needed-to-treat of 8. Critically, the PET-CT cohort had significantly worse baseline complexity: 3.3-fold longer infection duration (43.1 vs. 13.1 months, $p < 0.001$) and 3.6-fold more prior surgeries (2.48 vs. 0.69, $p < 0.001$). Multivariable analysis confirmed PET-CT-guided surgery as an independent protective [...]

[PubMed](#) · [DOI](#)

Digital coordination in mass casualty incidents: a retrospective analysis of prehospital distribution times before and after IVENA-MANV implementation.

Brauckmann V, Ringe B, Caviglia M

BACKGROUND: Digital coordination tools have been proposed to improve management during mass casualty incidents (MCIs). In 2021, the IVENA-MANV system was introduced in the Hannover region as an extension of the established IVENA eHealth platform to enable digital patient tracking and hospital capacity management. However, no real-world evaluations have yet quantified their operational impact.

OBJECTIVES: This study aimed to provide the first real-world evaluation of the IVENA-MANV digital coordination system and to determine how its implementation affected prehospital distribution processes and time intervals during mass casualty incidents.

METHODS: A retrospective observational study to assess the impact of the IVENA-MANV digital coordination platform on prehospital times during MCIs in Germany was conducted using dispatch records and IVENA-MANV logs from all MCIs between January 2018 and July 2025 in the Hanover-Region in Germany. Primary outcomes were the triage-to-evacuation (TtE) and total prehospital time (TPHT) per patient. Mann-Whitney U tests compared pre- and post-implementation groups, and multiple linear regression models examined associations between IVENA-MANV use, MCI-level, and time intervals.

RESULTS: Seventy-three MCIs met inclusion criteria, including 188 documented individual casualty characteristics. 41.1% of incidents used IVENA-MANV. Most MCIs were tra [...]

[PubMed](#) · [DOI](#)

The real-world management of acute mesenteric ischemia in Spain: results from a multicenter national survey.

González-Castillo AM, Calsina Juscafresa L, Gelabert A, Velescu A, Marrano E, Vikatmaa P, Tolonen M

BACKGROUND: Acute mesenteric ischemia (AMI) remains one of the most lethal vascular emergencies, with in-hospital mortality rates frequently exceeding 50%. Although early diagnosis and timely revascularization are critical, clinical practice remains highly variable. This study aimed to evaluate the real-world management of AMI in Spain, identifying gaps in resources, protocols, and interhospital coordination.

METHODS: A national cross-sectional survey was conducted between March and August 2024 using the Survio® platform. The questionnaire was distributed to general surgeons through national surgical societies and included 27 items covering hospital infrastructure, clinical protocols, diagnostic and therapeutic availability, personal experience, and perceived system-level barriers.

RESULTS: A total of 291 surgeons responded. The median age was 40 years (IQR: 17). Most were consultants (76.3%) working in tertiary (42.6%), secondary (33%), or community hospitals (24.4%). While 97.6% reported access to 24/7 multiphasic CT and 90.4% to round-the-clock radiology, only 51.9% had 24/7 interventional radiology and 60.1% vascular surgery. Just 26.8% had institutional AMI protocols. The median distance to a referral center was 25 km (range: 2-250 km), and 68.4% reported difficulty in patient transfers. While 96.9% felt competent managing AMI, only 36.4% were familiar with the term "int [...]

[PubMed](#) · [DOI](#)

Changes in operative and in-hospital outcome of proximal femoral fractures before and after the implementation of work hour restrictions in Switzerland.

Tewes A, Canal C, Schöb O, Neuhaus V

PURPOSE: Surgeon fatigue is a recognized risk factor for patient safety; however, reduced working hours may limit surgical training exposure. In 2005, Switzerland introduced a maximum 50-hour workweek for resident surgeons. In view of current discussions on further reductions (42 + 4 h), concerns persist regarding training quality and workforce capacity. This study evaluates the impact of the 2005 Swiss labor law on patient outcomes, operative characteristics, and intraoperative teaching utilizing a standardized trauma procedure.

METHODS: A retrospective analysis of anonymized data from a national database was performed. All patients undergoing operative fixation of trochanteric femur fractures (ICD-10 S72.1) were included. Two four-year periods were compared: pre-regulation (2001-2004) and post-regulation (2016-2019). Primary endpoints were in-hospital complications, mortality, and length of stay. Secondary endpoints included patient characteristics, surgeon seniority, teaching status, operative duration, and time to surgery.

RESULTS: Post-regulation patients were older and exhibited higher American Society of Anesthesiologists classifications despite fewer comorbidities. Complications increased from 9.1% to 17.7%, and mortality from 1.6% to 3.5%. Length of stay decreased from 14 to 9 days. Operative duration decreased by 10 min across all surgeon levels. Resident surgeons i [...]

[PubMed](#) · [DOI](#)

Early discharges in severely injured patients by ISS score: exploring injury patterns and coding practices.

Parouei F, Krüsemann EJZ, Poeze M, de Jongh MAC, Hietbrink F, Landelijke Beraadsgroep Traumachirurgen (LBTC)

PURPOSE: This study examines the prevalence of early discharge (ED) among patients classified as severely injured in the Dutch National Trauma Registry (DNTR) and evaluates whether identifiable patient-, injury-, and system-related factors are associated with the occurrence of ED in this population.

METHODS: This retrospective cohort study analyzed DNTR data from 2015 to 2022, focusing on severely injured patients (Injury Severity Score [ISS] ≥ 16). Patients were grouped as early discharge (ED; discharged within 48 h without in-hospital mortality) or non-ED (NED). Patients' characteristics were compared using descriptive statistics, and a mixed-effect model identified factors associated with ED.

RESULTS: From 2015 to 2022, a total of 37,626 severely injured patients were registered including 1,454 (3.9%) ED patients. This proportion increased from 3.2% in 2015 to 4.8% in 2022. ED was more common in younger patients, males, general practitioner (GP) or self-referred cases, those with severe head injury (Abbreviated Injury Scale [AIS] ≥ 3) and fewer registered injury codes. In the mixed-effects model, younger age (OR up to 2.0), increasing head injury severity (OR: 1.16 per AIS point), less than 3 registered AIS codes (OR: 0.60), referral by a general practitioner (OR 1.68) or self-referral (OR 1.92), and admission year (OR: 1.08 per year) independently predicted early discharge [...]

[PubMed](#) · [DOI](#)

Trauma team activation for pediatric patients in Denmark: a multicenter study of criteria, organization, and injury severity.

Nordestgaard CH, Gude MF, Leth SV, Raaber N

BACKGROUND: Pediatric trauma team activation (TTA) is intended to ensure timely management of severely injured children, yet both prehospital visitation practices and in-hospital TTA criteria vary across trauma systems. The aim of this study was to describe and compare pediatric TTA criteria, organizational models, and injury severity across all Danish level I trauma centers.

METHODS: We conducted a retrospective multicenter cohort study including all pediatric trauma patients (< 18 years) admitted via TTA at the four Danish level I trauma centers between 2014 and 2024. Center-specific prehospital and in-hospital TTA protocols were obtained through structured inquiries. Injury severity was assessed using Injury Severity Score (ISS) and Abbreviated Injury Scale (AIS). Overtriage was defined as TTA in patients with ISS < 15.

RESULTS: A total of 3,452 pediatric trauma patients were included. Two distinct TTA models (single-criterion and point-based) and four organizational structures were identified. Overall, 14% of patients had ISS ≥ 15 , corresponding to high overtriage across all centers. Overtriage rates ranged from 81% to 95% and were highest at centers using point-based TTA models. Mortality was low and did not differ significantly between centers. Among the most severely injured patients (ISS ≥ 25), head and thoracic injuries predominated.

CONCLUSION: Pediatric trauma triage [...]

[PubMed](#) · [DOI](#)

Does the operative technique (MIPO vs. ORIF) influence surgical outcomes and complications in proximal humerus fractures? A matched-pair analysis.

Barthel C, Rudolph F, Kraus M, Kalbas Y, Russek R, Pape HC, Allemann F

INTRODUCTION: Proximal humerus fractures account for approximately 4-6% of all fractures and are among the most common fractures in the elderly. Surgical treatment is increasingly applied for displaced and unstable fracture patterns. The two most commonly used techniques are open reduction and internal fixation (ORIF) via a deltopectoral approach and minimally invasive plate osteosynthesis (MIPO) using a deltoid-split or anterolateral approach. ORIF allows direct fracture visualization and anatomical reduction but is associated with more extensive surgical exposure. MIPO aims to preserve soft tissues and vascular supply through indirect reduction techniques. This study compares ORIF and MIPO in proximal humerus fractures with regard to surgical parameters, radiographic outcomes, and complications.

METHODS: A retrospective single-center cohort of 392 proximal humerus fractures was analyzed. Seventy-nine patients treated with MIPO were matched to 79 patients treated with ORIF using exact matching for fracture type (Mayo FJD classification) and propensity score matching for age and sex. Primary outcomes included operative time, reduction quality, and complication rates.

RESULTS: The MIPO group showed significantly shorter operative times (97.3 ± 30.8 min vs. 112.2 ± 34.0 min, $p = 0.004$). No significant differences were observed in reduction quality or overall complication rates [...]

[PubMed](#) · [DOI](#)

Risk factors and predictors of in-hospital mortality in necrotizing fasciitis: a five-year retrospective cohort study from a tertiary care center in Colombia.

Barrientos ML, Cordoba ASR, Valencia MDML, Valencia MMM, Toro DAM, Zapata CAL, Delgado CA

PURPOSE: Necrotizing fasciitis (NF) is a life-threatening soft tissue infection with limited epidemiological data from Latin America. This study aimed to characterize clinical outcomes, identify factors associated with in-hospital mortality, and develop a clinical prediction model in patients with NF at a tertiary care center in Medellín, Colombia.

METHODS: A retrospective cohort study was conducted including all adult patients (≥ 18 years) with surgically or histopathologically confirmed NF between January 2019 and December 2024. Demographic, clinical, microbiological, and therapeutic variables were collected. Bivariable analysis and multivariable logistic regression were performed to identify independent mortality predictors. A clinical risk score was developed and internally validated using bootstrap resampling and 10-fold cross-validation.

RESULTS: Of 112 identified patients, 111 met inclusion criteria. Overall in-hospital mortality was 23.4% (26/111). Deceased patients were significantly older (median 69.5 vs. 52 years; $p < 0.001$). Serum lactate demonstrated strong prognostic value (AUC 0.719; 95% CI 0.596-0.842; 2.9 vs. 1.8 mmol/L; $p = 0.003$). Septic shock was present in 34.2% and doubled mortality risk (34.2% vs. 17.8%; $p = 0.060$). Polymicrobial infections predominated (74.8%), with a predominantly gram-negative profile and high antimicrobial resistance rates (carbapen [...]

[PubMed](#) · [DOI](#)

ESTES recommendations regarding distal radius fractures.

Lameijer CM, Teuben MPJ, van Delft EAK, Tomazevic M, Kastelec M, Nau C

Distal radius fractures (DRFs) account for 15% of all fractures in adults. Optimal treatment of DRFs depend on patient and fracture characteristics. Non- or minimally displaced DRFs can be treated nonoperatively with a cast. Although surgical treatment is becoming more popular, the majority of patients with a DRF are still treated nonoperatively. In addition, these considerations may be different for elderly patients. Conflicting evidence exists on the different aspects of treatment of DRFs. These recommendations regarding distal radius fractures describe considerations regarding diagnostic modalities, associated soft tissue injuries, different treatment options, rehabilitation protocol, considerations for treatment of elderly patients, follow-up decisions and possible complications. The aim is to aid the orthopedic traumatologist in decision making and treatment of distal radius fractures.

[PubMed](#) · [DOI](#)

Small traumatic intracranial hemorrhages identified in routine radiology reports are associated with a low risk of adverse events: a retrospective cohort study.

Gerdås K, Wigermo A, Offenbartl M, Vestlund S, Rezk F, Vedin T

PURPOSE: To determine whether routinely available radiology reports, together with basic clinical data, can identify patients with traumatic intracranial hemorrhage (TICH) who are at low risk of adverse events.

METHODS: This retrospective cohort study of adults with TICH in Region Jönköping, Sweden (2019-2021). Clinical data, findings from radiology reports and outcomes were extracted from medical records. Hemorrhage size was classified as small (≤ 4 mm or described as minimal/very small/discrete) or larger. Adverse events were defined as neurosurgical intervention or death directly attributable to the TICH. Risk difference (RD), relative risk (RR), and Firth's penalized logistic regression were used to assess associations with adverse events.

RESULTS: Among 527 included patients, 195 (37%) had small TICHs. None of these patients experienced adverse events, compared with 13.6% neurosurgical interventions and 13.0% trauma-related deaths in the group with larger TICHs (RD 24.5% points, 95% CI 18.4-29.6; RR 97.1, 95% CI 6.1-1557.1; $p < 0.001$). Small TICH size had the strongest association with absence of adverse events. Normal neurological status and GCS 14-15 were also associated with a low risk of adverse events. Anticoagulant or antiplatelet therapy showed no significant association with adverse events.

CONCLUSION: Routinely available radiology reports, combined with basic c [...]

[PubMed](#) · [DOI](#)

From fracture to joint injury: association between CT-based fracture characteristics and surgically treated non-bony injuries in tibial plateau fractures.

Neidlein C, Colcuc S, Ullerich F, Schöllmann H, Steinhausen ES, Kröpil P, Brinkmann N, Dudda M et al.

PURPOSE: Tibial plateau fractures (TPF) are often associated with non-bony injuries that may impair knee stability and long-term outcomes. However, it remains unclear in which patients preoperative MRI is indicated to detect these injuries, and clear recommendations for surgical decision making are lacking. This study aimed to develop a CT-based predictor of surgically relevant non-bony injuries.

METHODS: This retrospective single-center study included all intra-articular TPF treated between January 2022 and May 2025 at a Level I trauma center. Fractures were classified according to the 10-segment and three-column classifications. Operative reports were reviewed to identify non-bony injuries requiring surgical treatment. Descriptive statistics and multivariate logistic regression were performed to determine predictors of surgically relevant non-bony injuries.

RESULTS: Among 243 patients (mean age $49,3 \pm 14,9$ years), 34,6% had at least one surgically treated non-bony injury. Posterior column involvement was significantly associated with a higher injury rate (41,4%), particularly affecting meniscus (20,7%), Anterior cruciate ligament (ACL) (17,2%), and medial collateral ligament (MCL) (12,3%). The posterolateral-central (PLC) segment showed the highest segment-specific injury rate (37,6%) and was an independent predictor of ACL injury. Increasing fracture complexity was associa [...]

[PubMed](#) · [DOI](#)

Transferrals and clinical pathways of unstable pelvic fractures over the last 10 years - a retrospective analysis of the Trauma Register DGU®.

Weuster M, Pfeifer R, Seekamp A, Lippross S, Lefering R, Dgu T, Menzendorf L

PURPOSE: Pelvic fractures are classified as stable or unstable. They correlate with a severity of trauma and the initial medical treatment is decisive. This study evaluated the transferrals of such fractures and described the initial treatment as well as the clinical course.

METHODS: We analysed retrospective data from a large cohort of the TraumaRegister DGU® (TR-DGU), covering the period from 2014 to 2023, comprising a total of $n = 397,910$ patients. All patients aged ≥ 16 years were included. Injury patterns were described according to the Abbreviated Injury Scale (AIS), the mechanically unstable fractures were classified with an $AIS \geq 3$. We considered all participating hospitals within Germany. The patients were subdivided in three groups: Group 1 = primary admitted patients with outcome, group 2 = pre-treated patients transferred in from other hospitals, and group 3 = primary admitted and early (< 48 h) transferred out.

RESULTS: The majority of the patients was male and about 53 years old. Blunt trauma was the leading trauma mechanism. Concomitant injuries ($AIS \geq 2$) affected thorax (56%), spinal cord (41%), lower extremities (38%), head (31%) and abdomen (24%). Among primary admitted cases with pelvic fractures ($n = 36,398$), 21,091 cases (57.9%) had an unstable pelvic fracture (AIS pelvis 3-5). Level 1 trauma centers not only treated 12,836 primary admitted cases with unst [...]

[PubMed](#) · [DOI](#)

Surgical treatment of coronal shear fractures: short- to mid-term results and risk factor analysis.

Bauer A, Cornel L, Sauter M, Jakobi T, Münzberg M, Klug A

BACKGROUND: Coronal shear fractures of the distal humerus are rare but severe injuries. Reconstruction is often challenging, especially in comminuted cases, which is why many surgeons opt for an elbow arthroplasty in those cases. However, total elbow arthroplasty is associated with a variety of potential problems itself. Therefore, the aim of this study was to present the functional and clinical outcome of coronal shear fractures treated by osteosynthetic reconstruction in a short- to mid-term follow-up, and to identify possible risk factors for an inferior outcome.

METHODS: We performed a retrospective follow-up assessment of 51 consecutive patients (30 women; median age 56 years, (IQR 39-62)) who underwent osteosynthetic reconstruction for coronal shear fractures between 2012 and 2022 after a minimum follow-up period of two years. The Mayo Elbow Performance Score, Oxford Elbow Score, and Disabilities of the Arm, Shoulder and Hand score were evaluated, and all available radiographs were analyzed. All complications and revision procedures were assessed. Univariable and multivariate regression analyses were performed to identify potential risk factors for a poor outcome following osteosynthetic reconstruction.

RESULTS: After a median follow-up period of 43 (IQR 28-78) months, the median Mayo Elbow Performance Score was 100 (IQR 85-100), the median Oxford Elbow Score was 42 (IQ [...])

[PubMed](#) · [DOI](#)

Publisher Correction: Experience with patient-specific guides, instead of general surgical experience, improves the accuracy of 3D-guided corrective osteotomies.

Buist ML, Meesters AML, Colaris JW, Doornberg JN, Ten Duis K, Tabernée Heijtmeyer SJC, Jutte PC, Pijpker PAJ et al.

[PubMed](#) · [DOI](#)

Assessment of a new technique for biological augmentation in sternoclavicular joint dislocations: a cadaveric feasibility and biomechanical evaluation.

Somberg O, Förster E, Rosteijs T, Bernstorff M, Schildhauer T, Königshausen M

AIMS: Biological augmentation is necessary in chronic sternoclavicular dislocations. However, the placement of drill holes for a graft can be difficult at that region. The aim of this study was to evaluate, as an early proof-of-concept, the feasibility of using a locally available clavicular periosteal flap (CPF) or sternal periosteal flap (SPF) for sternoclavicular joint (SCJ) stabilization and to biomechanically test the periosteal flaps.

METHODS: A cadaveric anatomic feasibility and biomechanical study (level of evidence V) was performed on nine SCJs. The CPF was mobilized from three sides by lifting the flap off the bone, preserving its attachment near the SCJ. Two 1.6 mm drill holes were created in the sternum, and the CPF was folded over the SCJ and fixed with a non-resorbable wire. Additionally, the non-resorbable wire was used in terms of a figure-of-8-bracing through a 1.6 mm clavicular drill hole to secure the joint position. Flap dimensions and sternal overlap were measured. Vice versa SPFs were prepared in selected specimens and fixed in a similar fashion. The CPFs were biomechanically tested using a uniaxial tensile testing machine to obtain values for stress at 50% strain (σ (50%)), maximum force (Fmax), maximum stress (σ_{max}), strain at maximum force (ϵ (Fmax)), and strain at failure (ϵ at failure).

RESULTS: CPF mobilization and fixation were successfully achieve [...]

[PubMed](#) · [DOI](#)

Differentiating injury patterns and outcomes in accidental, suicidal and occupational falls from heights.

Beyersdorf C, Hussmann B, Wergen N, Lefering R, Schiffner E, Maus U, Jaekel C, Dgu T

BACKGROUND: Falls from significant heights are a leading cause of severe injuries and can result from accidents, suicide attempts, or occupational incidents. While previous studies have explored general injury patterns associated with falls, little is known about the specific differences between accidental, suicidal, and occupational falls. This study aims to identify distinct injury patterns and outcomes among these fall types, thereby enhancing primary care and decision-making algorithms.

METHODS: We conducted a retrospective analysis using data from the German TraumaRegister DGU® (2014-2023), focusing on patients who experienced falls from heights. Only patients admitted to trauma centers with trauma team activation and a Maximum Abbreviated Injury Scale (MAIS) severity ≥ 3 were included. All falls were categorized in accidental, suicidal, or occupational and distinguished between high (≥ 3 m) and low (< 3 m) falls. Statistical analyses included chi-squared tests for categorical variables and Mann-Whitney U tests for continuous variables, with significance set at $p < 0.05$.

RESULTS: Among 188,708 patients, 29,991 (15.9%) experienced high falls, while 50,674 (26.9%) fell from low height. High falls were predominantly accidental (82.3%), with suicidal falls accounting for 17.7%. Accidental high falls mainly caused thoracic (58.9%) and head injuries (46.2%), whereas suicidal f [...]

Predictive performance of ISS and NISS for clinical outcomes in severely injured trauma patients: a retrospective registry study.

Säkkinen A, Partio N, Mattila V, Mäenpää H, Riuttanen A

BACKGROUND: The Injury Severity Score (ISS) and New Injury Severity Score (NISS) are widely used in evaluation of injury severity in trauma patients. The aim of this study is to assess the predictive accuracy of these scoring systems on clinical outcomes.

METHODS: We conducted a retrospective registry study of 1,112 severely injured trauma patients (NISS ≥ 16) treated at Tampere University Hospital (TAUH) between 2015 and 2024. Outcomes included in-hospital mortality, prolonged hospital and ICU stay (≥ 75 th percentile), in-hospital intubation, and blood transfusions. Predictive performance was assessed using the area under the receiver operating characteristic curve (AUC) and the Hosmer-Lemeshow (H-L) test for calibration. Subgroup analyses were performed for patients with significant (AIS ≥ 3) head, thorax, and extremity injuries.

RESULTS: A total of 848 (76%) patients had a higher NISS than ISS with a median increase of 9 points. Both scoring systems generally demonstrated fair to considerable ($0.7 < \text{AUC} < 0.9$) discrimination for mortality and intubation, and poor to fair ($0.6 < \text{AUC} < 0.8$) discrimination for blood transfusions and prolonged hospital/ICU stay. ISS had superior discrimination for blood transfusions in the total cohort (AUC: 0.669 vs. 0.630, $p = 0.019$), head (AUC: 0.737 vs. 0.672, $p = 0.004$), and thorax (AUC: 0.656 vs. 0.613, $p = 0.006$) subgroups, and for intu [...]

PubMed · DOI

Surgical and anesthetic activity in a french emergency field hospital responding to devastating floods.

Gaudray E, Brac L, Agopian P, Arnaud I, Ribelles P, Buland S, Couret A, Simonet J et al.

PURPOSE: Floods are the most frequent natural disasters, causing widespread mortality and devastation. In 2023, Storm Daniel triggered catastrophic flooding in Derna, Libya, leading to over 11,300 deaths and leaving more than 10,000 people missing. In response, the French Emergency Medical Team (EMT-FRA) ESCRIM, deployed via the European Union Civil Protection Mechanism, arrived to provide critical medical aid. **METHODS:** The team set up a field hospital equipped with adult and pediatric care, ICU, radiology, and laboratory services, adhering to strict aseptic conditions. Data collection was performed daily by designated physicians, with records transcribed for analysis. Over the course of the 22-day mission, the EMT-FRA provided care to over 1,600 patients, with 7.6% undergoing surgery. **RESULTS:** The anesthesia team performed 64 procedures, including 11 critical care assessments and 65 pre-anesthetic consultations. Surgical cases were primarily related to infections (49%), visceral issues (27%), and orthopaedic injuries (25%), including wound debridement, VAC therapy, amputations, and laparotomy. The surgical team carried out 64 surgeries and 126 consultations, with 83% of patients receiving emergency care for the first time. A significant number of procedures (78%) were dirty or infected, reflecting the local health challenges. **CONCLUSIONS:** The deployment of EMT-FRA demonstrated [...]

PubMed · DOI

Hook plate versus locking plate fixation in lateral clavicle fractures: operative efficiency and cumulative surgical burden in a propensity score-matched cohort study.

Schibler D, Hinkelmann MA, Frei F, Pape HC, Allemann F, Barthel C

INTRODUCTION: Hook plates remain controversial in the management of lateral clavicle fractures despite their biomechanical advantages. Concerns include implant-related complications, subacromial irritation, and the need for routine implant removal. The aim of this study was to compare operative efficiency and surgical burden of hook plate fixation with locking plate fixation in lateral clavicle fractures. **METHODS:** A retrospective cohort analysis was conducted at a single tertiary trauma center between January 1, 2016, and December 31, 2024. Surgically treated lateral clavicle fractures were identified. Propensity score matching was performed based on age, sex, and AO fracture classification to reduce confounding, and matched pairs were compared between hook plate fixation (Group HP) and locking plate fixation (Group LCP). Outcome measures included operative time, complications, implant removal, radiographic callus formation, and refracture rates. **RESULTS:** A total of 101 lateral clavicle fractures were identified. After propensity score matching, 44 matched pairs (Group HP vs. Group LCP) were

analyzed, with comparable baseline characteristics between groups. Operative time was significantly shorter in Group HP (77.8 ± 28.4 vs. 93.5 ± 30.2 min, $p = 0.018$). Complication rates were higher in Group HP but did not reach statistical significance (18.2% vs. 4.5%, $p = 0.093$). Implant re [...]

[PubMed](#) · [DOI](#)

Diagnostic value of hypoxia-inducible factor 1 alpha and adrenomedullin in acute mesenteric ischemia.

Bayindir E, Küçükceran K, Özer MR, Ayrancı MK, Kiliç ■, Kökbudak N, Bayindir GK

BACKGROUND: This study aims to evaluate the diagnostic significance of adrenomedullin and hypoxia-inducible factor 1 alpha (HIF1-alpha) levels in an animal model of acute mesenteric ischemia.

MATERIALS AND METHODS: In our study, a total of 21 male New Zealand rabbits were used, and the animals were divided into three groups for the study. Blood samples were taken for adrenomedullin and HIF1-alpha levels at 0, 1, 3, and 6 h from the control, sham, ischemia groups.

RESULTS: HIF levels of the ischemia group at the 3rd and 6th hours were statistically significantly higher than those in the sham ($p = 0.002$, $p < 0.001$, respectively) and control groups ($p = 0.002$, $p < 0.001$, respectively). Adrenomedullin levels of the ischemia group at the 1st, 3rd, and 6th hours were statistically significantly higher than those in the sham ($p = 0.004$, $p < 0.001$, $p < 0.001$, respectively) and control groups ($p < 0.001$, $p < 0.001$, $p < 0.001$ respectively).

CONCLUSION: In this experimental model of acute mesenteric ischemia, hypoxia-inducible factor-1 alpha and adrenomedullin levels were significantly higher in the ischemia group compared with the control and sham groups. Adrenomedullin demonstrated an earlier increase, while hypoxia-inducible factor-1 alpha increased significantly at later time points. These biomarkers may provide supportive diagnostic value when interpreted alongside clinical and im [...]

[PubMed](#) · [DOI](#)

Videolaryngoscope-assisted fiberoptic intubation in anticipated difficult airways of maxillofacial trauma: a prospective randomised comparison with videolaryngoscope-guided nasotracheal intubation.

Jain K, Nair R, Goyal A, Makkar JK, Singla K, Arora S, Bhatia N

BACKGROUND: Maxillofacial trauma poses airway management challenges due to edema, distorted facial anatomy and associated bleeding. Further, its fixation requires unhindered access to facial fractures, often necessitating securing the airway naso-tracheally during the intraoperative period. Traditional technique of nasotracheal intubation is associated with 17%-77% incidence of epistaxis. Fiberoptic bronchoscope helps guide nasotracheal tube under vision, decreasing incidence and severity of epistaxis. Moreover, as patients with maxillofacial trauma have difficult airways, every effort should be made to secure these challenging airways with best possible technique that increases first-attempt success rate. Use of fiberoptic bronchoscope as a part of hybrid videolaryngoscope-assisted fiberoptic intubation(VAFI) technique may provide these benefits. Hence, we conceptualized this study comparing the VAFI technique with traditional videolaryngoscope-assisted technique for securing nasotracheal tube in patients with anticipated difficult airways following maxillofacial trauma. **METHODS:** Following a written, informed consent, we included adult patients with maxillofacial trauma, scheduled for fracture fixation under general anaesthesia, requiring nasotracheal intubation. All patients were randomised into two groups of 30 patients each, with patients in group-VLS undergoing videolaryng [...]

[PubMed](#) · [DOI](#)

Malignancy risk and management outcomes in adult intussusception: a single-institution retrospective study.

Chang HC, Kang JC, Pu TW, Chen CY

PURPOSE: Adult intussusception is rare and frequently caused by a pathological lead point, including malignancy. We assessed malignancy rates by CT-based anatomical subtype/location, identified predictors of malignancy, and summarized management outcomes to support risk-stratified decision-making in the emergency setting.

METHODS: We conducted a single-institution retrospective study at Tri-Service General Hospital (Taipei, Taiwan),

including consecutive adults with CT-diagnosed intussusception from January 2015 to August 2024. We collected demographics, symptoms, hemoglobin, CT-based type/location, management, and histopathology. Malignancy was defined by index-admission pathology (malignant vs. benign). Anemia was defined as hemoglobin < 13 g/dL in men or < 12 g/dL in women. Predictors of malignancy were evaluated using univariable and multivariable logistic regression. Primary analyses included pathology-confirmed cases; sensitivity analyses additionally included presumed benign cases. RESULTS: Forty-seven patients were identified; 38 had pathology-confirmed outcomes. In the primary cohort (n = 38), 18 (47.4%) were malignant. Lower hemoglobin was associated with malignancy (OR per 1 g/dL increase, 0.709; 95% CI, 0.529–0.950; p = 0.021), and anemia strongly predicted malignancy (OR, 11.667; 95% CI, 2.438–55.830; p = 0.002). In multivariable analysis adjusting for age and colo [...]

[PubMed](#) · [DOI](#)

Operative burden and resuscitation resource utilisation in patients with traumatic shock requiring urgent surgical or endovascular intervention.

Noonan M, O'Loan M, Hendel S, Fitzgerald M, Ban EJ, Huynh F, McKay S, Scorer A et al.

BACKGROUND: Patients with traumatic shock represent a high-risk subgroup within major trauma populations, yet the operative burden and resuscitation resource utilisation associated with urgent surgical and endovascular intervention in trauma systems managing predominantly blunt injury remain incompletely described. AIMS: To describe the epidemiology, resuscitation resource utilisation, and operative burden of patients presenting with traumatic shock who require urgent surgical or endovascular intervention. METHODS: We conducted a retrospective observational study at an Australian level 1 trauma centre using prospectively maintained trauma registries. Adult major trauma patients (≥ 16 years, Injury Severity Score ≥ 13) meeting institutional shocked trauma activation criteria between December 2022 and December 2024 were included. Resource utilisation, blood product transfusion, operative and endovascular interventions, procedural timing, and critical care outcomes were described. Comparisons were performed between shocked patients requiring urgent surgical or endovascular intervention and those managed without urgent procedures. RESULTS: Of 3667 major trauma patients, 324 (8.8%) met shocked trauma criteria, and 138 (42.6% of shocked patients) underwent urgent surgical or endovascular intervention. These patients demonstrated substantial operative complexity; 37.7% required combin [...]

[PubMed](#) · [DOI](#)

Incidence of ROTEM®-defined coagulopathy at hospital arrival among polytrauma patients receiving fibrinogen concentrate in prehospital setting: a prospective observational study.

Vrbica K, Keller F, Stingl J, Vanickova K, Stepanova R, Dolecek M, Hrdy O, Gal R

PURPOSE: Trauma-induced coagulopathy affects approximately one-quarter of polytrauma patients and is associated with increased mortality. Given the central role of fibrinogen and its early depletion after injury, supplementation is recommended after hospital admission in the presence of hypofibrinogenaemia or suspected major haemorrhage. As coagulopathy develops within minutes of trauma, prehospital fibrinogen administration may help mitigate its progression before hospital arrival. This study evaluated the incidence of coagulopathy at hospital admission, assessed using ROTEM®, among polytrauma patients receiving fibrinogen concentrate in the prehospital setting. METHODS: In this single-centre, prospective observational study, adult patients with polytrauma (Injury Severity Score ≥ 16) were enrolled after prehospital administration of fibrinogen concentrate. The primary outcome was the presence of coagulopathy at hospital arrival, defined according to predefined ROTEM® Delta thresholds (FIBTEM MCF ≤ 8 mm and/or EXTEM CT ≥ 80 s and/or EXTEM MCF ≤ 49 mm). The secondary outcomes were laboratory coagulation status, 28-day all-cause mortality, and thromboembolic complications. RESULTS: Thirty patients were enrolled, of whom 28 were included in the final analysis. Twenty-seven patients received 4 g of fibrinogen concentrate and one received 2 g due to short transport time. ROTEM®-def [...]

[PubMed](#) · [DOI](#)

Timing of laparoscopic cholecystectomy after percutaneous transhepatic gallbladder drainage and the risk of bailout surgery: a restricted cubic spline analysis.

Yoshitake H, Uchida K, Kataoka N, Mizobata Y

BACKGROUND: The optimal timing of laparoscopic cholecystectomy (LC) after percutaneous transhepatic gallbladder drainage (PTGBD) for acute cholecystitis remains controversial. Previous studies have often categorized surgical timing using arbitrary cutoffs and have not adequately evaluated potential non-linear associations. In particular, the interval from PTGBD to LC has rarely been analyzed as a continuous variable when assessing operative difficulty. This study aimed to assess the relationship between the interval from PTGBD to LC and intraoperative technical difficulty using restricted cubic spline (RCS) analysis. **METHODS:** This retrospective single-center study included patients with acute cholecystitis who underwent LC after PTGBD between November 2014 and November 2024. The primary outcome was bailout surgery (conversion to open surgery, subtotal cholecystectomy, or fundus-first approach). Secondary outcomes included operative time, intraoperative blood loss, postoperative length of stay, and major postoperative complications (Clavien–Dindo $\geq$ III). The interval from PTGBD to LC was analyzed as a continuous variable using multivariable regression models incorporating RCS. Models were adjusted for age, C-reactive protein (CRP) at diagnosis, Tokyo Guidelines 2018 severity grade, previous abdominal surgery, anticoagulant therapy, and Charlson Comorbidity Index. **RESULTS:** A total [...]

[PubMed](#) · [DOI](#)

Early orthoplastic coverage reduces fracture-related infections in Gustilo III B/C tibial fractures - a systematic review and meta-analysis.

Ruprecht N, Klingebiel FK, Hübner CT, Hax J, Teuber H, Jensen KO, Hierholzer C, Pfeifer R et al.

BACKGROUND: Gustilo-Anderson type III B/C open tibial fractures are among the most complex and high-risk injuries in trauma care. Their management demands a multidisciplinary approach, and early intervention is critical. While early debridement has been established as essential, the optimal timing for definitive orthoplastic coverage remains less clearly defined. **OBJECTIVE:** To evaluate whether early orthoplastic coverage reduces the risk of fracture-related infections (FRIs) and complication rates compared to late coverage in type III B/C open tibial fractures. **METHODS:** We conducted a PRISMA-compliant systematic review and meta-analysis of studies comparing early versus late orthoplastic treatment in patients with Gustilo-Anderson III B/C open tibial fractures. Databases included PubMed and EMBASE. Primary outcome was FRI; secondary outcomes included amputation, non-union, and revision surgery rates. Risk of bias was assessed using the Newcastle-Ottawa Scale. **RESULTS:** Of 3746 studies screened, five studies with a total of 454 patients met the inclusion criteria. Cutoff for early vs. late treatment was defined variably across studies (either within 72–168 h). Meta-analysis showed that early treatment significantly reduced the odds of infection (OR: 0.34; 95% CI: 0.23–0.52; $p < 0.001$) compared to late treatment. Subgroup analyses confirmed consistent benefits regardless of timing [...]

[PubMed](#) · [DOI](#)

Long-term renal function after high-grade renal trauma: a nuclear imaging based cohort study.

Saxena M, Dheer Y, Roy S, Singh U, Deswal S, Kumar N, Singh A, Jaiswal V et al.

PURPOSE: Renal trauma constitutes approximately 1% of all trauma cases, yet data on long-term renal function following severe injury remain scarce. This study aimed to evaluate long-term functional and morphological outcomes in patients sustaining Grade 3 or higher renal trauma, and to identify predictors of residual renal function using nuclear imaging. **METHODS:** This retrospective study included patients with Grade ≥ 3 renal injuries who presented for follow-up ≥ 6 months after trauma at a tertiary care centre between January 2018 and June 2023. Patients with nephrectomy or comorbidities affecting renal function were excluded. Serum renal function tests, DTPA scans (differential GFR), and DMSA scans (cortical morphology and dye uptake) were performed. Multivariable linear regression was used to determine predictors of residual function based on injury grade, gender, and duration of follow-up. **RESULTS:** Thirty-six patients (mean age 20.50 ± 12.08 years; 75% male) were included with equal representation across Grades 3–5. Mean follow-up was 35.92 ± 18.09 months. The affected kidney demonstrated reduced function (mean GFR 24.96 ± 12.50 mL/min; split function $31.83 \pm 12.69\%$; dye uptake $35.71 \pm 13.40\%$). Grade 5 injuries were significantly associated with lower differential GFR and dye uptake ($p < 0.01$), while longer follow-up duration and female sex predicted better renal recovery ([...])

[PubMed](#) · [DOI](#)

Experience with patient-specific guides, instead of general surgical experience, improves the accuracy of 3D-guided corrective osteotomies.

Buist ML, Meesters AML, Colaris JW, Doornberg JN, Ten Duis K, Tabernée Heijtmeyer SJC, Jutte PC, Pijpker PAJ et al.

PURPOSE: This study investigated whether the accuracy of 3D-guided osteotomies is influenced by surgeon experience. Two types of experience were assessed: (1) general surgical experience, defined as self-reported years in practice as an orthopaedic trauma surgeon, and (2) prior use of patient-specific guides (PSGs). Secondary aim was to evaluate whether accuracy varies by anatomical location. **METHODS:** 24 orthopaedic-trauma surgeons performed 75 corrective osteotomies on cadaveric long-bones (43 single-cut, 32 double-cut; 107 cuts in total). Each osteotomy was preoperatively planned on CT-derived 3D-reconstructions, and PSGs were manufactured to guide the cuts. Postoperative CT-scans were used to compare planned and executed osteotomy planes. Surgeons completed a questionnaire reporting years of practice and annual PSG usage. A Spearman's rank-order correlation was used to test differences in accuracy versus experience, a Kruskal-Wallis test was performed to explore differences across anatomical locations. **RESULTS:** General surgical experience ranged from 0 to 25 years (median 9), and PSG-usage from 0 to 20 cases annually (median 2). No association was found between general surgical experience and osteotomy accuracy ($p = 0.406$ and $p = 0.548$). In contrast, PSG-experience correlated with improved accuracy for angular and translational deviation of the osteotomy plane ($p = 0.027$ and [...])

[PubMed](#) · [DOI](#)

Analysis of DRG remuneration of acute fracture procedures versus elective total arthroplasty for the German health care system - Is elective total arthroplasty really more cost-effective?

Hierl K, Kerschbaum M, Baumann F, Alt V

PURPOSE: Elective arthroplasty is considered a cost-effective procedure compared to trauma-related surgical interventions. This study aims to compare remuneration, certain ratios regarding implant costs, operation (OR) time and length of stay (LOS) as well as contribution margins including personnel costs for acute fracture interventions and elective arthroplasty procedures at the hip and shoulder joint in the aG-DRG system. **METHODS:** Based on billing data of 212 inpatient treatment cases from a University Orthopaedic Trauma Department reimbursements of nail fixation and hemi-arthroplasty (HA) for proximal femoral fractures versus elective total hip arthroplasty (THA) for osteoarthritis were analysed. Locking plate fixation and reverse total shoulder arthroplasty (rTSA) for proximal humeral fractures were compared to elective rTSA for osteoarthritis. Different ratios of the DRG remuneration in relation to LOS, OR time and implant costs but also personnel costs and contribution margins were calculated. **RESULTS:** DRG reimbursement was highest for fracture arthroplasty of the shoulder (9.050 EUR, rTSA) and the hip (6.930 EUR, HA). Regarding LOS, DRG reimbursement was highest for elective rTSA (1.641 EUR/day). Nail fixation of proximal femoral fractures showed the highest DRG reimbursement regarding OR time (111 EUR/min) and implant costs (15 EUR/EUR). The contribution margin was hig [...]

[PubMed](#) · [DOI](#)

Percutaneous screw osteosynthesis for the treatment of intra-articular displaced calcaneus fractures.

Wilmsen L, Neubert A, Deichsel A, Schulz D, Hoelscher-Doht S, Windolf J, Icks A, Thelen S

PURPOSE: This systematic review investigates how percutaneous screw osteosynthesis for the treatment of intra-articular displaced calcaneus fractures differs from open reduction and internal fixation (ORIF). **METHODS:** Randomized controlled trials and controlled clinical trials investigating adults with closed intra-articular displaced calcaneus fracture (Sanders type II - IV) treated with percutaneous screw osteosynthesis or ORIF were included. Severe vascular and neurological diseases or polytrauma patients were excluded. On January 29, 2024, five databases and two trial registries were searched. The Risk of Bias Tool of Cochrane was used. The protocol was registered on PROSPERO (CRD42021244695). **RESULTS:** Five studies with 654 participants (682 calcaneus fractures) were included. The risk of bias was moderate to high. None of the single studies found a significant difference regarding severe complications. However, meta-analysis revealed a lower risk for the percutaneous screw osteosynthesis group compared to ORIF (relative risk (RR) 0.40 (95% Confidence Interval [0.17; 0.92]; $I^2 = 0\%$; $p = 0.03$). One study showed less pain in the percutaneous screw osteosynthesis group up to 4 weeks postoperatively. No study assessed health-related quality of life. **CONCLUSION:** Percutaneous screw osteosynthesis in intra-articular displaced calcaneus fractures might be beneficial to ORIF in terms [...]

[PubMed](#) · [DOI](#)

Conservative versus operative treatment of LC-1 pelvic ring fractures: radiographic and long-term functional outcomes.

Metzger F, Buschbeck S, Swartman B, Schweigkofler U

PURPOSE: To compare radiographic and long-term functional outcomes of conservatively and operatively treated traumatic lateral compression type 1 (LC-1) pelvic ring fractures, considering fracture morphology and injury severity. **METHODS:** This single-center observational cohort study included adult patients with traumatic LC-1 pelvic ring injuries treated between 2012 and 2020. Radiographic outcomes were assessed using the Matta–Tornetta system, and functional outcomes were evaluated at long-term follow-up using the normalized Majeed Pelvic Score and SF-12. Baseline characteristics and complications were analyzed descriptively. **RESULTS:** Seventy-seven patients were included (38 conservative, 39 operative). Operatively treated patients had higher injury severity and more complex fracture morphology. Despite worse initial alignment, radiographic outcomes at follow-up were similar between groups, with most patients achieving excellent or good results. Radiographic consolidation was observed in nearly all patients. Long-term functional outcomes did not differ significantly between groups. **CONCLUSION:** In this observational cohort with substantial baseline differences, no significant differences in long-term functional outcomes were observed between treatment strategies. These findings support an individualized, morphology-based treatment approach. However, results should be interpreted [...]

[PubMed](#) · [DOI](#)

Correction: Clinically relevant bleeding risk in low-energy fragility fractures of the pelvis in elderly patients.

de Herdt CL, Loggers SAI, van Embden D, Bijlsma T, Joosse P, Ponsen KJ

[PubMed](#) · [DOI](#)

A novel concept for splint fixation of midshaft clavicle fractures: a biomechanical feasibility study.

Ziegenhain F, Zderic I, Richards RG, Gueorguiev B, Groven RVM, Varga P, Hildebrand F, Pape HC et al.

BACKGROUND: Minimally invasive treatment of clavicle shaft fractures has become increasingly popular due to reduced soft tissue damage, smaller incisions, lower rates of infections, and minimal scarring. Most commonly described options are titanium elastic nailing and plate osteosynthesis. Clinical studies report promising results with good healing tendencies, particularly in children and adolescents. However, there are still some downsides to consider—titanium elastic nails have been associated with soft-tissue irritation, implant associated discomfort, and notable complications rates including lateral perforation of the cortex and medial protrusion of the implant. Implant removal after a few months is standardly carried out, leading to a second surgery with corresponding risks and costs. Furthermore, biomechanical studies report a low rotational stability for these implants. Therefore, the aim of this biomechanical feasibility study was to explore a new concept for minimally invasive splint fixation of clavicle shaft fractures in comparison to the titanium elastic nailing. **METHODS:** Twelve anatomical composite clavicles with standardised oblique midshaft fractures AO/OTA 15.2 A were randomized to two groups for fixation with either a 2.5 mm titanium elastic nail (Group 1) or a 5.0 mm rib splint (Group 2). Each specimen underwent (1) non-destructive quasi-static pure-bending te [...]

[PubMed](#) · [DOI](#)

Development and external validation of a novel prediction model for the TraumaTriage App.

Gulickx M, Lokerman RD, van der Sluijs R, Tuinema RM, van Vliet R, Hietbrink F, Groenwold RHH, van Heijl M

PURPOSE: Accurate pre-hospital trauma triage is essential for optimizing survival and functional outcomes within inclusive trauma systems. A prediction model incorporated into the TraumaTriage App (TTApp) has been shown to improve patient allocation in the Netherlands, but was developed based on a relatively small cohort. This study aims to redevelop and improve the performance and applicability of the TTApp's prediction model before nationwide implementation. **METHODS:** In this prospective multicenter cohort study, data from all trauma patients transported within two emergency medical service (EMS) regions (i.e., Brabant and Utrecht), between February 2015 and October 2019, were used to develop and externally validate a prediction model to identify severely injured adults (Injury Severity Score [ISS] ≥ 16) using routinely available pre-hospital data. A Gradient Boosting

Decision Tree (GBDT) algorithm was applied, and model performance was evaluated in terms of discrimination and calibration. RESULTS: The development cohort included 51,001 patients (median age 63.8 years, median ISS 9), and the external validation cohort included 29,737 patients (median age 62.1 years, median ISS 9). In external validation, the GBDT model showed excellent discrimination (c-statistic 0.850; 95% CI, 0.837–0.863) and good calibration (calibration-in-the-large 0.009; slope 0.952). Sensitivity was 91. [...]

[PubMed](#) · [DOI](#)

Association between high ratio of platelet concentrate to red blood cell and survival among massively transfused patients.

Aoki M, Okada Y, Fujiwara G, Katsura M, Matsumoto S, Kiyozumi T, Tomura S, Matsushima K

PURPOSE: Although balanced transfusion protocols (platelet concentrate (PC): red blood cell (RBC) = 1:1) are standard for severe trauma, the impact of a higher PC: RBC ratio (> 1:1) on survival remains unclear. This study aimed to investigate the effectiveness of a high PC to RBC among severely injured patients requiring massive transfusion. METHODS: We performed a retrospective cohort study using the Japan Trauma Data Bank (2019–2023). Adult patients receiving massive transfusion (> = 10 units of RBC) within the first 24 h of injury were included. Patients were classified into high PC (> 1) and low PC groups (< = 1). The primary outcome was 24-hour mortality, analyzed using inverse probability of treatment weighting and modified Poisson regression. Additionally, we performed spline curve analysis to investigate the nonlinear relationship between the PC: RBC ratio and outcome. RESULTS: Among 3,067 patients (mean age 57.1 years, 67.2% male), 934 were in the high PC group and 2,133 in the low PC group. The high PC group had lower 24-hour mortality (8.9% vs. 15.7%). Adjusted analysis showed a significantly lower risk of mortality in the high PC group (risk ratio 0.54; 95% CI 0.43–0.69). A nonlinear analysis suggested that increasing PC: RBC ratio was associated with lower mortality. CONCLUSION: A high PC: RBC ratio was associated with lower 24-hour mortality in severely injured pa [...]

[PubMed](#) · [DOI](#)

Treatment and outcomes of periprosthetic femoral fractures in patients aged 90 years or older - a retrospective bicentre cohort study with 107 patients.

Müller F, Wulbrand C, Probst A, Langenhan R

PURPOSE: The surgical treatment of periprosthetic femoral fractures (PFFs) is highly demanding, especially in very old patients. For patients aged 90 years or older, data regarding internal fixation (IF) and revision arthroplasty (RA) are lacking. METHODS: This retrospective case series was conducted in two different regional centres in one country. We enrolled patients with an age of 90 years or older, and who had PFFs with cemented or cementless standard stems. The treatment period was from 01/2010 to 12/2024. The exclusion criteria were interprosthetic femoral fractures or periprosthetic acetabular fractures. Treatment was performed with IF (locking plates) or RA (noncemented revision stems). The primary end point was any revision within the follow-up period. The secondary end points were death, serious adverse events, and function with Parker score. The minimum follow-up for living patients was one year after surgery. Living patients received a final contact by phone. RESULTS: A total of 107 patients were included in this study. The mean age of the cohort was 92.0 years (range 90 to 100), and most patients were female (n = 85). Following the Vancouver classification and its recommendations, 17 out of 22 patients with type B1 fractures were treated with IF, 44 out of 48 patients with type B2/3 fractures were treated with RA, and 34 out of 37 patients with type C fractures we [...]

[PubMed](#) · [DOI](#)

The graft-reposition-on-flap technique for fingertip amputations: a patient-reported outcomes study of 111 patients.

Thushara KR, Sankaran A, Koller MN

PURPOSE: Graft-reposition-on-flap (GRF) is a reconstruction method described for non-replantable fingertip amputations. We hypothesised that this technique provides acceptable outcomes for such an injury, as measured by a patient-reported survey.

METHODS: This retrospective study evaluated 111 patients [118 digits], operated between January 2019 and March 2023 for Allen 3 or 4 fingertip amputations. All patients had crush avulsion amputations not fit for replantation and had undergone flap cover with grafting of the amputated nail bed and bone. 80 digits had

undergone a flap with bone and nail bed graft, while 38 digits received only a nail bed graft with the flap. The flaps used were based on the defect, including local and pedicle flaps. Outcomes were assessed using an original patient-reported questionnaire administered telephonically at least 1 year after injury. The survey contained six items rated on a 1-10 Likert scale, with higher scores indicating better outcomes.

RESULTS: Nail growth was present in 105 of the 118 digits (89%). 107 patients of 111 (96%) were willing to undergo a similar procedure again if indicated. The median rated score for the Length of the finger was 9 [Interquartile range (IQR) 8-10], for Fingertip shape 8 [IQR 7-9], for Donor appearance 9 [IQR 8-10], for Nail appearance 8 [IQR 5-10], and for Sensibility 9 [IQR 7.5-10]. The median Overall Satisfisf [...]

[PubMed](#) · [DOI](#)

Mechanical comparison of different plate-and-screw configurations for repairing forearm diaphysis fractures: a systematic review.

Zdero R, Brzozowski P, Schemitsch EH

PURPOSE: This is the first systematic review of all studies that adjusted plate and/or screw variables to determine the most mechanically stable implant configuration for forearm fracture repair. METHODS: PubMed and Scopus archives were searched using the phrase “biomechanics” AND “fracture” AND “plate” AND (“forearm” OR “radius” OR “ulna”). Eligibility criteria were applied: (i) studies that mechanically analyzed implant performance by modulating plate and/or screw variables, such as plate geometry, plate material, screw number, etc.; (ii) forearm diaphysis “shaft” fractures of the radius and/or ulna; (iii) studies published at any time; (iv) English language studies. RESULTS: There were 34 eligible papers. There was greater mechanical stability for these configurations: (i) plates used alone or double-stacked; (ii) plates that were longer, wider, thicker, straight or curved at the bottom, non-ribbed at the bottom, and/or non-grooved at the sides; (iii) plates made of material with higher elastic modulus; (iv) plates that were locking; (v) plates not elevated above bone; (vi) plates affixed to the anterior side of bone; (vii) screws that were longer, thicker, and/or numerous. There was a range of numerical outcomes: (i) interfragmentary motion; (ii) peak stress in cortical bone or plates; (iii) cortical bone stress under the plate to assess bone “stress shielding”; (iv) fatigu [...]

[PubMed](#) · [DOI](#)

Postoperative complications of xenogenic dural substitutes in hemicraniectomy.

Tanrikulu L, Alhasan AA, Dömer P, Woitzik J

PURPOSE: Decompressive hemicraniectomy (DHC) is a treatment option for refractory intracranial pressure (ICP) elevation in cases of cerebral infarction, hemorrhage, and traumatic brain injury. Several surgical techniques are used for musculocutaneous, bony, and dural exposure. The method of dural closure, specifically whether to use expanding duraplasty with or without xenogenic dural substitutes, remains a topic of debate. This study focuses on the occurrence of postoperative surgical side effects associated with these approaches. METHODS: A total of 258 patients (mean age 52.8 ± 16.7 years; 140 male, 118 female, 79 infarctions, 57 traumatic brain injuries, 55 subarachnoid hemorrhages, 32 intracranial hemorrhages, 30 subdural hematomas, 5 other indications) who underwent DHC between 2015 and 2024 at our institution were reviewed. Complete clinical and radiological data were available for all patients. The patient population was divided into two groups: • Group A: 201 patients who received the xenogenic dural substitute. • Group B: 57 patients who did not receive a dural substitute. RESULTS: Baseline demographics (age, gender, underlying pathology) were comparable between both groups. No significant differences were found in the following parameters: • Duration of DHC surgery: 102.0 min (Group A) vs. 108.0 min (Group B); $p = 0.24$. • Postoperative CSF leakage: 3.5% (7/201) in Gr [...]

[PubMed](#) · [DOI](#)

Rethinking routine chest radiography after central venous catheterisation: a rapid review of current evidence and alternatives.

Orso D, Guglielmo N, Pangallo R, Montanar V, Favaro S, Bove T

BACKGROUND: Despite growing evidence supporting ultrasound (US)- and ECG-guided protocols, chest radiography (CXR) remains the most widely used method for post-procedural confirmation of central venous catheter (CVC) position in clinical practice. METHODS: We conducted a rapid review of observational studies, randomised controlled trials, systematic reviews, and guidelines comparing CXR with US- and ECG-based

confirmation modalities in adult patients across ICU, emergency, and perioperative settings. Systematic searches were performed in PubMed, Scopus, and Web of Science through December 2025. Outcomes of interest included diagnostic accuracy for tip position and complications, procedural time, and clinical and economic impact. FINDINGS: Ninety-four studies were included, comprising more than 20,000 CVC placements. Evidence from meta-analyses, multicentre cohorts, and comparative studies suggests that US-based protocols – particularly those incorporating agitated saline (“bubble test”) and right-atrial visualisation – and intracavitary ECG (IC-ECG) can achieve high diagnostic performance for tip confirmation in appropriately selected patients, with several reports describing sensitivity and specificity of approximately 95% or higher. Across studies, bedside strategies enabled faster confirmation than CXR (often 15–40 min), avoided radiation exposure, and facilitated earlier de [...]

[PubMed](#) · [DOI](#)

Comparison of CATCH, PECARN and CHALICE rules in children with minor head trauma.

Ferhatlar ME, Cumhuri BA, Koca Y, Bozan O, Kalkan A

PURPOSE: The aim of this study is to prospectively compare the diagnostic performance of the PECARN, CATCH, and CHALICE clinical decision rules in pediatric patients presenting with minor head trauma, within a rural secondary-level emergency department characterized by limited resources, high patient volume, and restricted access to specialist physicians.

METHODS: The study was designed as a prospective observational study. The PECARN, CATCH, and CHALICE clinical decision rules were applied to the patients included in the study, and the data were recorded. The data included in the study were statistically analyzed using the SPSS (Statistical Package for Social Sciences; SPSS Inc., Chicago, IL) 27 program. A value of $p < 0.05$ was considered statistically significant.

RESULTS: A total of 1,032 patients were included in the study. The PECARN did not miss any of the 15 patients with pathological brain CT and had 100% sensitivity and 52.1% specificity. CATCH failed to detect pathological brain CT in 1 patient and had 93.3% sensitivity and 53.3% specificity. CHALICE failed to detect pathological brain CT in 2 patients, with 86.7% sensitivity and 87.8% specificity. Overall, PECARN was more successful in detecting pathological brain CT, but it also had a high false-positive brain CT rate.

CONCLUSION: PECARN criteria are more sensitive than other criteria in predicting brain CT patho [...]

[PubMed](#) · [DOI](#)

Effect of head injury on blood transfusion requirements in severe trauma: a post-hoc analysis of the RESTRIC trial.

Nagasawa H, Omori K, Tanaka N, Maekawa C, Hamada M, Ota S, Ohsaka H, Yanagawa Y et al.

PURPOSE: To evaluate whether head-region injury, as defined by Abbreviated Injury Scale (AIS) coding, is associated with transfusion requirements in patients with severe trauma. **METHODS:** This study is a post-hoc analysis of the RESTRIC trial, a multicentre, cluster-randomised, crossover, non-inferiority trial comparing restrictive (7–9 g/dL) and liberal (10–12 g/dL) blood transfusion strategies for patients with severe haemorrhagic trauma. Adult patients with severe trauma anticipated to require transfusion were included. Head injury was defined as an AIS score ≥ 2 in the head region. The primary outcome was cumulative red blood cell (RBC) transfusion volume within 28 days. Key secondary outcomes included RBC transfusion volume within 24 and 48 h. Propensity score matching (PSM) was used to balance baseline injury severity using Injury Severity Score (ISS) and, separately, AIS scores for body regions excluding the head. Sensitivity analyses were conducted using generalised linear mixed models (GLMM) with random intercepts for study centre. **RESULTS:** Among the 411 patients, 157 (38.2%) had head AIS ≥ 2 . In unadjusted analyses, 28-day RBC transfusion volumes were similar between patients with and without head injury (median 10 vs. 10 units). After adjustment using PSM and GLMM, no evidence of an independent association between head-region injury and RBC transfusion volume was obse [...]

[PubMed](#) · [DOI](#)

Emergency laparotomies in trauma management: a six-year case series.

Ahrens G, Hoffmann M, Proksch N

INTRODUCTION: Emergency laparotomies, defined as procedures performed within 24 hours of hospital admission following trauma, are associated with considerable mortality. Notably, blunt abdominal injuries carry

higher mortality rates compared to penetrating injuries. The primary aim of this study was to evaluate whether emergency trauma laparotomies performed at our Level I trauma center—serving a population with a comparatively higher proportion of penetrating injuries—achieve outcomes comparable to contemporary benchmark data. **METHODS:** We conducted a retrospective analysis of patients treated between 2019 and 2024 in the surgical trauma bay at Asklepios Klinik St. Georg, Hamburg, who underwent laparotomy within 24 hours of hospital admission. Mortality was defined as survival until hospital discharge. Functional outcomes were assessed using the Glasgow Outcome Scale (GOS). **RESULTS:** A total of 64 patients were included. Of these, 39.0% sustained penetrating trauma and 61.0% underwent laparotomy following blunt trauma. Among penetrating injuries, stab wounds predominated (92.0%), while 8.0% resulted from gunshot wounds. The most frequent mechanisms of blunt trauma were traffic accidents (30.8%) and falls from height (25.6%). Overall survival was 71.9% (mortality 20.0% for penetrating trauma; 33.3% for blunt trauma). Prehospital cardiopulmonary resuscitation (CPR) had been performed [...]

[PubMed](#) · [DOI](#)

Correction: Comparison of total intravenous anesthesia vs. inhalational anesthesia on brain relaxation, intracranial pressure, and hemodynamics in patients with acute subdural hematoma undergoing emergency craniotomy: a randomized control trial.

Preethi J, Bidkar PU, Cherian A, Dey A, Srinivasan S, Adinarayanan S, Ramesh AS

[PubMed](#) · [DOI](#)

Non-scaphoid carpal fractures: lessons learned from a series of 86 cases.

Herrero Alonso S, Fernández González-Cuevas J

PURPOSE: To describe the epidemiology, injury mechanisms, diagnostic approach, management, and clinical outcomes of non-scaphoid carpal fractures, which are frequently overlooked or diagnosed late. **METHODS:** A retrospective review was conducted of patients with non-scaphoid carpal fractures treated at our institution between 2013 and 2020. Demographic data, fracture characteristics, time to diagnosis, treatment strategy, rehabilitation requirements, and clinical outcomes, including complications, were analyzed. **RESULTS:** 76 patients with 86 fractures were included; 54 (71%) were male. The most commonly affected bones were the triquetrum (37%) and hamate (33%). Isolated fractures accounted for 89% of cases. Advanced imaging was frequently required, with CT performed in 53% and MRI in 7% of patients. The median diagnostic delay was 6 days (range: 1–40). Conservative treatment was used in 84% of cases. Surgical management was significantly more frequent in lunate (75%) and hamate fractures (25%) ($p = 0.008$). Surgically treated patients required rehabilitation more often than those managed conservatively (58% vs. 31%, $p = 0.012$). Functional range of motion ($> 40^\circ$ flexion/extension) was achieved in 97% of patients. However, 16% developed chronic wrist pain, 8% reported prolonged soreness, and one patient developed reflex sympathetic dystrophy. **CONCLUSIONS:** Although functional outcomes [...]

[PubMed](#) · [DOI](#)

A machine-learning-based tool for stroke risk prediction in blunt cerebrovascular injury: development and preliminary evaluation.

Wagner V, Preston JD, De Leon Castro A, Mueller WF, Nguyen J, Garcia-Toca M, Benjamin ER, Todd SR et al.

PURPOSE: Stroke risk correlates with the Biffi grading system in blunt cerebrovascular injury (BCVI). Although anti-thrombotic therapy is the mainstay of stroke prevention, no point-of-care clinical decision-support tool exists to guide timing for therapy. We sought to develop an interactive online calculator that incorporates patient-specific demographic and injury characteristics to estimate stroke risk and risk reduction with anti-thrombotic (AT) administration.

METHODS: Data from BCVI patients ($n = 1,197$) at a Level I Trauma Center were retrospectively collected. Six machine learning methods were employed to predict stroke risk with and without AT therapy. Class imbalance was addressed using downsampling and/or class weighting. Model performance was assessed using 10-fold cross-validation. The model was implemented as an R-based Shiny online application.

RESULTS: Stroke rate among the population was 4%, and the strongest predictors for stroke were the greatest Biffi grade of carotid (aOR [95%CI] = 2.02 [1.62-2.53]) and vertebral injuries (1.44 [1.18-1.77]). The least absolute shrinkage and selection operator (LASSO) model outperformed all others, achieving 66% [33%-100%] sensitivity

and 74% [62%-82%] specificity for stroke prediction, with an area under the receiver operating characteristic curve of 0.79 [0.57-0.95]. This model was integrated into an interactive online to [...]

[PubMed](#) · [DOI](#)

Association between screw angulation and union after inverted-triangle cannulated screw fixation of femoral neck fractures.

Çatal T, Bayrak A, Tüngen M, Sapmaz E, Kızılkaya ME, Mercan N, Kural C, Duramaz A

PURPOSE: The aim of this study was to evaluate whether screw angulation relative to the inter-ischial(horizontal) axis and femoral anatomical axis was associated with union after inverted-triangle cannulated screw fixation of femoral neck fractures and to explore angle intervals associated with higher union rates. **METHODS:** Between January 2017 and December 2022, 83 patients operated on with closed reduction-percutaneous cannulated screws by the same surgical team were retrospectively analyzed. The angles of the calcar, anterosuperior, and posterosuperior screws were measured according to both the femoral anatomical axis and the horizontal axis. Multivariable logistic regression analyses adjusted for lateral angulation and Garden or Pauwels classification were additionally performed for the horizontal-axis screw-angle parameters. Exploratory angle intervals were assessed using quartile and sliding-window analyses. **RESULTS:** A total of 64 patients achieved union, with a mean union time of 8.69 ± 3.03 weeks. The nonunion group had higher lateral angulation and greater postoperative neck length($p = 0.006$, $p = 0.042$, respectively). In angle values relative to the horizontal axis, calcar and anterosuperior screws exhibited higher measurements in the union group($p = 0.027$, $p = 0.013$, respectively). In exploratory sliding-window analyses, high union intervals were identified as follows: [...]

[PubMed](#) · [DOI](#)

Response to the letter to the editor to discuss "Factors to consider when selecting an injured control group in case-control studies on skiing injuries".

Rutsch N, Gewiess J, Albers CE, Tinner C

[PubMed](#) · [DOI](#)

A study on prolonged use of resuscitative endovascular balloon of the aorta (REBOA) with shunt flow in a porcine hemorrhagic shock model.

Kiriü N, Saitoh D, Yamamura K, Sasa R, Kiyozumi T, Tomura S

PURPOSE: To evaluate the future efficacy and prolonged availability of double-balloon resuscitative endovascular balloon occlusion of the aorta (DB-REBOA) to determine whether bypassing upstream blood at the occlusion site improves survival in a porcine hemorrhagic shock model. **METHODS:** Twenty-five pigs (approximately 40 kg, estimated circulating blood volume 25%) were subjected to hemorrhagic shock by controlled phlebotomy. Animals were treated with standard REBOA (control, $n = 12$), internal shunt DB-REBOA (internal shunt, $n = 4$; an intraluminal conduit was set parallel to the DBREBOA in the aorta), or external shunt DB-REBOA (external shunt, $n = 9$; upstream blood from the occlusion site was shunted by connecting the internal carotid and femoral arteries with a pressure-resistant tube). Zone 1 aortic occlusion was maintained for 90 min, followed by balloon deflation. Survival, systolic blood pressure, and serum lactate levels were analyzed. **RESULTS:** Ten of 12 control animals developed cardiac arrest shortly after balloon deflation. All animals in the internal shunt group survived, whereas 5 of 9 animals in the external shunt group experienced cardiac arrest. Internal shunt DB-REBOA maintained stable systolic BP during prolonged occlusion, resulting in significantly lower serum lactate levels at 90 min compared to controls. External shunt DB-REBOA resulted in unstable hemodynam [...]

[PubMed](#) · [DOI](#)