

Electrochemical TEMPO-Catalyzed Oxidative Ugi-Type Reaction

Na Pan, Maegan Xinen Lee, Louis Bunel, Laurence Grimaud,* and Maxime R. Vitale*

Cite This: *ACS Org. Inorg. Au* 2021, 1, 18–22

Read Online

ACCESS |

Metrics & More

Article Recommendations

Supporting Information

ABSTRACT: Oxidative isocyanide-based multicomponent reactions (oxidative IMCRs) are very useful tools for the rapid construction of molecular diversity starting from readily available and stable substrates. Despite all their benefits, such multicomponent reactions are underdeveloped and strictly limited to 3-component processes. Indeed, in the presence of several reaction partners, the oxidation event needs to be rigorously chemoselective, which becomes incredibly more intricate as the number of reactive components increases. Nonetheless, we could overcome this significant pitfall and reach the first oxidative Ugi-type 4-IMCR by capitalizing on a very mild and green TEMPO-catalyzed electro-oxidation process. Employing alcohols as aldehyde surrogates and in the notable absence of any supporting electrolyte, this transformation proved to be extremely chemoselective in the presence of an amine and was compatible with a wide range of alcohols, amines, isocyanides, and carboxylic acids.


KEYWORDS: Oxidative Ugi reaction, Multicomponent reaction, Electrosynthesis, Indirect anodic oxidation, TEMPO

INTRODUCTION

Multicomponent reactions (MCRs) are now established as very efficient tools for the rapid and efficient preparation of elaborated chemicals, starting from relatively simple building blocks.¹ Inasmuch as they meet many of the criteria dictated by green chemistry,^{2–5} this sustainable strategy has found many applications ranging from the production of fine chemicals to various uses in material sciences.^{1,6–11} Isocyanide-based processes (IMCRs) have partaken substantially to the blossoming of this field. Since Passerini and Ugi demonstrated that the ambivalent reactivity of isocyanides could be successfully harnessed in multicomponent processes,^{12,13} IMCRs have not ceased to bring stimulating developments.^{1,6–11,14–21} Nevertheless, one of the major impediments of IMCRs is that, because often relying on the use of $\text{Csp}^2=\text{O}$ or $\text{Csp}^2=\text{N}$ partners, such reactions largely rely on the use of carbonyl/imine derivatives, which may reveal themselves to be unstable enough to impede their preparation and, accordingly, their use.

To circumvent this problem, the concept of generating the adequate carbonyl/imine electrophiles by means of the in situ oxidation of the corresponding alcohol/amine emerged.^{22,23} As the first milestone in oxidative IMCRs, Zhu et al. succeeded in performing an alcohol-based 3-component Passerini reaction by employing 2-iodoxybenzoic acid (IBX) as the stoichiometric oxidant (Scheme 1, eq (a)).²⁴ Whereas a catalytic version of this multicomponent process was later reported,^{25,26} more attention was subsequently paid to oxidative 3-component Ugi–Joullé-type IMCRs, in which an amine or an aniline derivative serves as the imine/iminium precursor (Scheme 1, eq (b)).^{27–41}


Scheme 1. Oxidative IMCRs—Aim of This Work



Despite all of these significant achievements, neither catalytic nor stoichiometric methods could be applied to related 4-component processes so far. This significant pitfall probably results from the problematic, yet compulsory, chemoselective oxidation required by the presence of an additional reactant. In particular, the 4-component Ugi coupling based on the use of alcohols remains an unsolved challenge, with the amine partner being also prone to

Received: March 13, 2021

Published: April 27, 2021




oxidation. In fact, alcohol-based Ugi-type processes could only be performed via a one-pot two-step sequence, wherein the stoichiometric or catalytic oxidation of an alcohol to its aldehyde form is achieved first, followed by the subsequent addition of the other three components.^{42–45}

In this specific context, we recently began to study the synergistic merger of electrosynthesis with IMCRs. Performing an anodic oxidation, direct or indirect, is particularly appealing as it avoids the use of stoichiometric oxidants by employing perfectly sustainable reagents: electrons and protons.^{46–51} Moreover, such oxidation methods are typically very mild, giving better chances to reach better chemoselectivity. After establishing the proof of concept via the development of an electrochemical oxidative Ugi–Joullie reaction,⁴¹ we now report the first alcohol-based oxidative Ugi reaction, thanks to an indirect anodic oxidation process (Scheme 1, eq (c)).

RESULTS AND DISCUSSION

At the onset of this work, we decided to study the model oxidative Ugi reaction between *tert*-butylisocyanide **1a** (1.5 mmol, 1 equiv), benzylamine **2a** (1.1 equiv), acetic acid **3a** (2.5 equiv), and methanol **4a** (3 equiv). TEMPO (2,2,6,6-tetramethyl-1-piperidinyloxy) was chosen as the redox mediator to promote the oxidation of methanol into formaldehyde (Table 1).^{52,53}

Table 1. Optimization of the Reaction Conditions



entry	variation from above	yield (%) ^a
1	none	68 (70) ^b
2	<i>t</i> -BuNC (0.5 mmol), [0.17 M]	39
3	DCM ^c as solvent	75
5	Ni foam as cathode	69
6	stainless steel as cathode	68
7	Ni as anode	54
8	stainless steel as anode	53
9	MeCO ₂ H (1.5 equiv)	57
10	MeCO ₂ H (3.5 equiv)	69
11	MeOH (1.5 equiv)	59
12	MeOH (6 equiv)	74
13	TEMPO (5 mol %)	70 (72) ^b
14	TEMPO (20 mol %)	67
15	no TEMPO	40
16	no electrolyte	78 (74) ^b
17	no electrolyte, TEMPO (5 mol %)	76 (74) ^b
18	no electricity	NR ^d

^aNMR yields using 1,3,5-trimethoxybenzene as an internal standard.

^bIsolated yield. ^cDichloromethane. ^dNo reaction.

In initial reaction conditions, employing an undivided cell, TEMPO (10 mol %), Et₄NBF₄ (1 equiv) as supporting electrolyte in acetonitrile (0.5 M), with a nickel plate cathode and a carbon graphite anode, we were pleased to observe the desired 4-component IMCR adduct **5a** in 70% isolated yield after 2.7 F/mol of electricity (Table 1, entry 1). This result was very encouraging as, despite the presence of benzylamine **2a**, this oxidation method allowed the desired selective oxidation


of methanol **4a**, without interfering with the concomitant multicomponent coupling. Moreover, it showcased the synthetic utility of this strategy, with which it became possible to avoid the use of toxic formaldehyde in Ugi processes.

Therefore, we embarked on the optimization of the reaction conditions. Whereas lowering the reaction concentration was detrimental (Table 1, entry 2), using dichloromethane as solvent or changing the cathode material to nickel foam or stainless steel had little influence over the reaction efficiency (Table 1, entries 3–6). On the other hand, substituting the graphite anode by other electrode materials gave lower yields (Table 1, entries 7 and 8). We then evaluated the influence of the starting quantities of acetic acid and methanol. Lowering them up to 1.5 equiv resulted in a noticeable diminution of yields, whereas increasing them did not significantly improve the formation of **5** (Table 1, entries 9–12). Pleasingly, this reaction was equally efficient when employing only 5 mol % of TEMPO (Table 1, entry 13), whereas the use of 20 mol % had no positive impact (Table 1, entry 14). The fact that some product was obtained without TEMPO indicated that direct anodic oxidation of methanol may happen, although with lesser efficiency (Table 1, entry 15). Most interestingly, likely because of the intrinsic conductivity of the reaction mixture, this electro-oxidative Ugi reaction could be achieved in the absence of any supporting electrolyte, using either 10 or 5 mol % of TEMPO (Table 1, entries 16 and 17). In the latter reaction conditions, which we kept for the rest of our study, the desired 4-IMCR adduct **5** was isolated in satisfying 74% yield, equaling a fair 55% Faraday yield.

We then investigated the scope of this new oxidative IMCR (Scheme 2). The tolerance to various isocyanides was surveyed first (Scheme 2, red box). Employing benzylamine, acetic acid, and methanol, this reaction was amenable for the use of diverse alkyl- or benzyl-substituted isocyanides **1b–e**. The corresponding multicomponent adducts **5–8** were obtained in good yields ranging from 74 to 83%. Using less nucleophilic and/or more sterically demanding partners such as methyl isocyanoacetate **1f** or 2,6-xylyl isocyanide **1g** resulted in a small decrease of reaction efficiency. Nevertheless, the desired Ugi products **9** and **10** were isolated in acceptable 55 and 52% yields, respectively. Of note, this electrochemical multicomponent reaction could be transposed to gram-scale, demonstrating its genuine synthetic utility and versatility. Indeed, while this scope evaluation was realized on a typical 1 mmol scale, starting from 5.0 mmol of isocyanide **1b** straightforwardly yielded 1.08 g of adduct **6** (75% yield).


Next, different amine partners were evaluated (Scheme 2, orange box). Replacing benzylamine **2a** with primary amines such as methylamine **2b** or *n*-butylamine **2c** had no detrimental impact over the reaction efficiency. A similar trend was also observed when employing bulkier α,α -di- or α,α,α -trisubstituted primary amines, such as cyclohexylamine **2d** and *tert*-butylamine **2e**, provided that 5% (v/v) of water was added to ensure the homogeneity of the reaction mixture. In these cases, **13** and **14** were obtained in respectable 78 and 88% yields. More elaborated amine partners could also be used successfully; allylamine **2f**, methoxyethylamine **2g**, and 3,4-dimethoxyphenyl ethylamine **2h** all afforded the desired Ugi adducts **15–17** in good yields, up to 87%.

Several carboxylic acids **3b–k** were also tested (Scheme 2, green box). Aliphatic acids were well-tolerated in this reaction, as witnessed by the good yields which were obtained when butyric acid **3b** and pivalic acid **3c** were employed (83 and

Scheme 2. Scope of the Electrochemical Oxidative Ugi Reaction^a

^aReaction conditions: isocyanide **1** (1.5 mmol, 1 equiv), amine **2** (1.65 mmol, 1.1 equiv), carboxylic acid **3** (3.75 mmol, 2.5 equiv), and alcohol **4** (4.5 mmol, 3 equiv) in acetonitrile (3 mL, [0.5 M]) at room temperature; graphite as working electrode, Ni as counter electrode, constant current electrolysis ($i = 14.5 \text{ mA}$, $j = 5 \text{ mA cm}^{-2}$) for 7.5 h, 2.7 F mol^{-1} . *Gram-scale reaction starting from 5.0 mmol isocyanide **1b** (1.08 g of multicomponent adduct **6** was obtained). **Water 5% (v/v) was added.

Scheme 3. Mechanistic Proposal



71%). This oxidative multicomponent process also appeared to be compatible with various aromatic and heteroaromatic acids **3d–i**, as well as acrylic acids **3j,k**, with which the corresponding coupling products **20–27** were isolated likewise with satisfactory yields (68–81%).

We next questioned if this unconventional oxidative IMCR was specific for the use of methanol (Scheme 2, blue box). A wide range of alcohol partners was also compatible with this electro-oxidative process. Different aliphatic alcohols **4b–h** led straightforwardly to the corresponding Ugi adducts **28–34** in

reasonable yields (53–69%). On the other hand, 2-phenylethanol **4i** and benzyl alcohol **4j**, possibly less nucleophilic due to the presence of the aromatic ring, afforded only little of the desired products **35** and **36**. Halogen-, alkene-, and alkyne-substituted alcohols **4k–o** also proved to be suitable reaction partners. The corresponding multicomponent coupling products **37–41** were isolated in 40–64% yields.

Based on our previous work and literature,^{14–21,41,52–54} we propose that TEMPO is oxidized at the graphite anode to generate the corresponding oxoammonium species TEMPO⁺.

The latter would, in turn, promote the selective oxidation of the alcohol partner into the corresponding aldehyde, leaving the amine partner untouched. At this stage, we surmise that the presence of the carboxylic acid is critical for chemoselectivity. Indeed, it probably ensures the protection of the amine via an acid/base equilibrium in favor of the non-nucleophilic ammonium carboxylate form, thereby preventing it to interact with TEMPO⁺. Then, the newly formed aldehyde, as in a classical 4-component Ugi reaction, would generate the corresponding iminium, electrophilic enough to react with the isocyanide. The resulting nitrilium cation would be trapped by the remaining carboxylate anion, and after 1,5-acyl transfer, the multicomponent product would be formed. To complete the catalytic cycle, either through comproportionation or direct anodic oxidation,⁵⁴ TEMPO⁺ would be regenerated and, to keep the electroneutrality of the global system, proton reduction would occur at the nickel cathode.

CONCLUSION

In conclusion, we have developed the first oxidative 4-IMCR by implementing a very chemoselective TEMPO-catalyzed electrochemical process. This alcohol-based Ugi-type reaction allows the use of unstable/toxic aldehydes to be avoided (such as formaldehyde) and is compatible with a wide range of alcohols, amines, isocyanides, and carboxylic acids. A small library of the corresponding Ugi adducts (37 examples) was rapidly obtained, showcasing the synthetic utility of this approach for the rapid construction of high molecular diversity in a sustainable manner. Further studies concerning this promising synergistic merger of electrosynthesis with IMCRs is currently underway and will be reported in due course.

ASSOCIATED CONTENT

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acsorginorgau.1c00003>.

All experimental details including further optimization studies, detailed procedures, and proper characterization of all products; copies of ¹H and ¹³C NMR (PDF)

AUTHOR INFORMATION

Corresponding Authors

Laurence Grimaud – Laboratoire des biomolécules, LBM, Département de chimie, École Normale Supérieure, PSL University, Sorbonne Université, CNRS, 75005 Paris, France; orcid.org/0000-0003-3904-1822; Email: laurence.grimaud@ens.psl.eu

Maxime R. Vitale – Laboratoire des biomolécules, LBM, Département de chimie, École Normale Supérieure, PSL University, Sorbonne Université, CNRS, 75005 Paris, France; orcid.org/0000-0002-6740-2472; Email: maxime.vitale@ens.psl.eu

Authors

Na Pan – Laboratoire des biomolécules, LBM, Département de chimie, École Normale Supérieure, PSL University, Sorbonne Université, CNRS, 75005 Paris, France; Shanghai Engineering Research Center of Molecular Therapeutics and New Drug Development, SCME, East China Normal University, Shanghai 200062, China

Maegan Xinen Lee – Laboratoire des biomolécules, LBM, Département de chimie, École Normale Supérieure, PSL University, Sorbonne Université, CNRS, 75005 Paris, France
Louis Bunel – Laboratoire des biomolécules, LBM, Département de chimie, École Normale Supérieure, PSL University, Sorbonne Université, CNRS, 75005 Paris, France

Complete contact information is available at:
<https://pubs.acs.org/doi/10.1021/acsorginorgau.1c00003>

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

The authors gratefully thank the ANR (ANR-20-CE07-0020, ELMER), the CNRS, the ENS and Sorbonne Université for financial support. N.P. thanks the China Scholarship Council (CSC) for funding her Ph.D. studies.

REFERENCES

- (1) *Multicomponent Reactions*; Zhu, J., Bienaymé, H., Eds.; Wiley-VCH: Weinheim, Germany, 2005.
- (2) Anastas, P.; Eghbali, N. Green Chemistry: Principles and Practice. *Chem. Soc. Rev.* **2010**, *39*, 301–312.
- (3) Cioc, R. C.; Ruijter, E.; Orru, R. V. A. Multicomponent Reactions: Advanced Tools for Sustainable Organic Synthesis. *Green Chem.* **2014**, *16*, 2958–2975.
- (4) *Green Chemistry: Theory and Practice*, Anastas, P. T.; Warner, J. C., Eds.; Oxford University Press: Oxford, 1998.
- (5) Zimmerman, J. B.; Anastas, P. T.; Erythropel, H. C.; Leitner, W. Designing for a Green Chemistry Future. *Science* **2020**, *367*, 397–400.
- (6) Dömling, A.; Wang, W.; Wang, K. Chemistry and Biology Of Multicomponent Reactions. *Chem. Rev.* **2012**, *112*, 3083–3135.
- (7) Biggs-Houck, J. E.; Younai, A.; Shaw, J. T. Recent Advances in Multicomponent Reactions for Diversity-Oriented Synthesis. *Curr. Opin. Chem. Biol.* **2010**, *14*, 371–382.
- (8) Garbarino, S.; Ravelli, D.; Protti, S.; Basso, A. Photoinduced Multicomponent Reactions. *Angew. Chem., Int. Ed.* **2016**, *55*, 15476–15484.
- (9) Ruijter, E.; Orru, R. V. A. Multicomponent Reactions - Opportunities for the Pharmaceutical Industry. *Drug Discovery Today: Technol.* **2013**, *10*, e15–e20.
- (10) Touré, B. B.; Hall, D. G. Natural Product Synthesis Using Multicomponent Reaction Strategies. *Chem. Rev.* **2009**, *109*, 4439–4486.
- (11) *Multi-Component and Sequential Reactions in Polymer Synthesis - Advances in Polymer Science*; Theato, P., Ed.; Springer International Publishing: Cham, Switzerland, 2015; Vol. 269.
- (12) Passerini, M. Sopra gli isonitrili (I). Composto del *p*-isonitril-azobenzolo con acetone ed acido acetico. *Gazz. Chim. Ital.* **1921**, *51*, 126–129.
- (13) Ugi, I.; Meyr, R.; Fetzer, U.; Steinbrückner, C. Versuche mit Isonitrilen. *Angew. Chem.* **1959**, *71*, 386.
- (14) Dömling, A.; Ugi, I. Multicomponent Reactions with Isocyanides. *Angew. Chem., Int. Ed.* **2000**, *39*, 3168–3210.
- (15) Dömling, A. Recent Developments in Isocyanide Based Multicomponent Reactions in Applied Chemistry. *Chem. Rev.* **2006**, *106*, 17–89.
- (16) Akritopoulou-Zanze, I. Isocyanide-Based Multicomponent Reactions in Drug Discovery. *Curr. Opin. Chem. Biol.* **2008**, *12*, 324–331.
- (17) El Kaïm, L.; Grimaud, L. The Ugi-Smiles and Passerini-Smiles Couplings: A Story About Phenols in Isocyanide-Based Multicomponent Reactions. *Eur. J. Org. Chem.* **2014**, 7749–7762.
- (18) Várad, A.; Palmer, T.; Notis Dardashti, R.; Majumdar, S. Isocyanide-Based Multicomponent Reactions for the Synthesis of Heterocycles. *Molecules* **2016**, *21*, 19.

- (19) Sadjadi, S.; Heravi, M. M.; Nazari, N. Isocyanide-Based Multicomponent Reactions in the Synthesis of Heterocycles. *RSC Adv.* **2016**, *6*, 53203–53272.
- (20) Giustiniano, M.; Moni, L.; Sangaletti, L.; Pelliccia, S.; Basso, A.; Novellino, E.; Tron, G. Interrupted Ugi and Passerini Reactions: An Underexplored Treasure Island. *Synthesis* **2018**, *50*, 3549–3570.
- (21) Wang, Q.; Wang, D.-X.; Wang, M.-X.; Zhu, J. Still Unconquered: Enantioselective Passerini and Ugi Multicomponent Reactions. *Acc. Chem. Res.* **2018**, *51*, 1290–1300.
- (22) Moni, L.; Banfi, L.; Riva, R.; Basso, A. External-Oxidant-Based Multicomponent Reactions. *Synthesis* **2016**, *48*, 4050–4059.
- (23) Basso, A.; Moni, L.; Riva, R. Multicomponent Reactions under Oxidative Conditions. *Multicomponent Reactions in Organic Synthesis*; Wiley-VCH Verlag GmbH & Co. KGaA: Weinheim, Germany, 2014; Vol. 8, pp 265–300.
- (24) Ngouansavanh, T.; Zhu, J. Alcohols in Isonitrile-Based Multicomponent Reaction: Passerini Reaction of Alcohols in the Presence Of *o*-Iodoxybenzoic Acid. *Angew. Chem., Int. Ed.* **2006**, *45*, 3495–3497.
- (25) Brioché, J.; Masson, G.; Zhu, J. Passerini Three-Component Reaction of Alcohols under Catalytic Aerobic Oxidative Conditions. *Org. Lett.* **2010**, *12*, 1432–1435.
- (26) Karimi, B.; Farhangi, E. One-Pot Oxidative Passerini Reaction of Alcohols Using a Magnetically Recyclable TEMPO under Metal- and Halogen-Free Conditions. *Adv. Synth. Catal.* **2013**, *355*, 508–516.
- (27) Ngouansavanh, T.; Zhu, J. IBX-Mediated Oxidative Ugi-Type Multicomponent Reactions: Application to the N and C1 Functionalization of Tetrahydroisoquinoline. *Angew. Chem., Int. Ed.* **2007**, *46*, 5775–5778.
- (28) Islas-Jácome, A.; Gutiérrez-Carrillo, A.; García-Garibay, M.; González-Zamora, E. One-Pot Synthesis of Nuevamine Aza-Analogues by Combined Use of an Oxidative Ugi Type Reaction and Aza-Diels-Alder Cycloaddition. *Synlett* **2014**, *25*, 403–406.
- (29) Wang, J.; Sun, Y.; Jiang, M.-H.; Hu, T.-Y.; Zhao, Y.-J.; Li, X.; Wang, G.; Hao, K.; Zhen, L. Iminium Ion and *N*-Hydroxyimide as the Surrogate Components in DEAD-Promoted Oxidative Ugi Variant. *J. Org. Chem.* **2018**, *83*, 13121–13131.
- (30) Ye, X.; Xie, C.; Pan, Y.; Han, L.; Xie, T. Copper-Catalyzed Synthesis of α -Amino Imides from Tertiary Amines: Ugi-Type Three-Component Assemblies Involving Direct Functionalization of Sp^3 C-Hs Adjacent to Nitrogen Atoms. *Org. Lett.* **2010**, *12*, 4240–4243.
- (31) Ye, X.; Xie, C.; Huang, R.; Liu, J. Direct Synthesis of α -Amino Amides from *N*-Alkyl Amines by the Copper-Catalyzed Oxidative Ugi-Type Reaction. *Synlett* **2012**, *23*, 409–412.
- (32) Rueping, M.; Vila, C. Visible Light Photoredox-Catalyzed Multicomponent Reactions. *Org. Lett.* **2013**, *15*, 2092–2095.
- (33) Vila, C.; Rueping, M. Visible-Light Mediated Heterogeneous C-H Functionalization: Oxidative Multi-Component Reactions Using a Recyclable Titanium Dioxide (TiO_2) Catalyst. *Green Chem.* **2013**, *15*, 2056–2059.
- (34) Rueping, M.; Vila, C.; Bootwicha, T. Continuous Flow Organocatalytic C-H Functionalization and Cross-Dehydrogenative Coupling Reactions: Visible Light Organophotocatalysis for Multicomponent Reactions and C-C, C-P Bond Formations. *ACS Catal.* **2013**, *3*, 1676–1680.
- (35) Chen, Y.; Feng, G. Visible Light Mediated Sp^3 C-H Bond Functionalization of *N*-Aryl-1,2,3,4-Tetrahydroisoquinolines via Ugi-Type Three-Component Reaction. *Org. Biomol. Chem.* **2015**, *13*, 4260–4265.
- (36) Dighe, S. U.; Kolle, S.; Batra, S. Iron-Catalysed Oxidative Ugi-Type Multicomponent Reaction Using (Arylmethyl)Amines as Imine Precursors. *Eur. J. Org. Chem.* **2015**, 4238–4245.
- (37) Guerrero, I.; San Segundo, M.; Correa, A. Iron-Catalyzed C(Sp^3)-H Functionalization of *N,N*-Dimethylanilines with Isocyanides. *Chem. Commun.* **2018**, *54*, 1627–1630.
- (38) Gini, A.; Uygur, M.; Rigotti, T.; Alemán, J.; García Mancheño, O. Novel Oxidative Ugi Reaction for the Synthesis of Highly Active, Visible-Light, Imide-Acrinium Organophotocatalysts. *Chem. - Eur. J.* **2018**, *24*, 12509–12514.
- (39) Brandhofer, T.; Gini, A.; Stockerl, S.; Piekarski, D. G.; García Mancheño, O. Direct C-H Bond Imidation with Benzoyl Peroxide as a Mild Oxidant and a Reagent. *J. Org. Chem.* **2019**, *84*, 12992–13002.
- (40) Li, D.; Shen, X.; Lei, J. Metal-Free Iodine/TEMPO-Mediated Aerobic Oxidative Ugi-Type Multicomponent Reactions with Tertiary Amines. *J. Org. Chem.* **2020**, *85*, 2466–2475.
- (41) Pan, N.; Ling, J.; Zapata, R.; Pulicani, J.-P.; Grimaud, L.; Vitale, M. R. Electrochemical TEMPO-Catalyzed Multicomponent C(Sp^3)-H α -Carbamoylation of Free Cyclic Secondary Amines. *Green Chem.* **2019**, *21*, 6194–6199.
- (42) Drouet, F.; Masson, G.; Zhu, J. Ugi Four-Component Reaction of Alcohols: Stoichiometric and Catalytic Oxidation/MCR Sequences. *Org. Lett.* **2013**, *15*, 2854–2857.
- (43) Khan, I. A.; Saxena, A. K. Parikh-Doering Oxidation-Dehydration-Ugi Cyclization Cascade in the Development of Lactams from Formidoalkanols (3 > Chain Length > 7). *Tetrahedron* **2012**, *68*, 294–299.
- (44) Moni, L.; Banfi, L.; Basso, A.; Carcone, L.; Rasparini, M.; Riva, R. Ugi and Passerini Reactions of Biocatalytically Derived Chiral Aldehydes: Application to the Synthesis of Bicyclic Pyrrolidines and of Antiviral Agent Telaprevir. *J. Org. Chem.* **2015**, *80*, 3411–3428.
- (45) Moni, L.; Banfi, L.; Basso, A.; Martino, E.; Riva, R. Diastereoselective Passerini Reaction of Biobased Chiral Aldehydes: Divergent Synthesis of Various Polyfunctionalized Heterocycles. *Org. Lett.* **2016**, *18*, 1638–1641.
- (46) Dagar, N.; Sen, P. P.; Roy, S. R. Electrifying Sustainability on Transition Metal-Free Modes: An Eco-Friendly Approach for the Formation of C-N Bonds. *ChemSusChem* **2021**, *14*, 1229–1257.
- (47) Yan, M.; Kawamata, Y.; Baran, P. S. Synthetic Organic Electrochemical Methods Since 2000: On the Verge of a Renaissance. *Chem. Rev.* **2017**, *117*, 13230–13319.
- (48) Kärkäs, M. D. Electrochemical Strategies for C-H Functionalization and C-N Bond Formation. *Chem. Soc. Rev.* **2018**, *47*, 5786–5865.
- (49) Wiebe, A.; Gieshoff, T.; Möhle, S.; Rodrigo, E.; Zirbes, M.; Waldvogel, S. R. Electrifying Organic Synthesis. *Angew. Chem., Int. Ed.* **2018**, *57*, 5594–5619.
- (50) Echeverría, P.-G.; Delbrayelle, D.; Letort, A.; Nomertin, F.; Perez, M.; Petit, L. The Spectacular Resurgence of Electrochemical Redox Reactions in Organic Synthesis. *Aldrichimica Acta* **2018**, *51*, 3–19.
- (51) Hammerich, O.; Speiser, B. *Organic Electrochemistry*, 5th ed.; CRC Press: New York, 2016.
- (52) Nutting, J. E.; Rafiee, M.; Stahl, S. S. Tetramethylpiperidine *N*-Oxyl (TEMPO), Phthalimide *N*-Oxyl (PINO), and Related *N*-Oxyl Species: Electrochemical Properties and Their Use in Electrocatalytic Reactions. *Chem. Rev.* **2018**, *118*, 4834–4885.
- (53) Gerken, J. B.; Pang, Y. Q.; Lauber, M. B.; Stahl, S. S. Structural Effects on the PH-Dependent Redox Properties of Organic Nitroxyls: Pourbaix Diagrams for TEMPO, ABNO, and Three TEMPO Analogs. *J. Org. Chem.* **2018**, *83*, 7323–7330.
- (54) Taitt, B. J.; Bender, M. T.; Choi, K. S. Impacts of the Regeneration Pathways of the Oxoammonium Cation on Electrochemical Nitroxyl Radical-Mediated Alcohol Oxidation. *ACS Catal.* **2020**, *10* (1), 265–275.