

Pd-Catalyzed Regioselective Cyclopropanation of 2-Substituted 1,3-Dienes

Agonist Kastrati, Vincent Jaquier, Michele Garbo, Céline Besnard, and Clément Mazet*

Cite This: *ACS Org. Inorg. Au* 2023, 3, 291–298

Read Online

ACCESS |




Metrics & More



Article Recommendations



Supporting Information



ABSTRACT: A Pd-catalyzed 3,4-regioselective cyclopropanation of 2-substituted 1,3-dienes by decomposition of diazo esters is reported. The vinylcyclopropanes generated are isolated in practical chemical yields with high levels of regioselectivity but low diastereoselectivity. The system operates under mild reaction conditions, is scalable, and tolerates various sensitive functional groups. A series of original postcatalytic derivatizations is presented to highlight the synthetic potential of the catalytic method.

KEYWORDS: palladium catalysis, conjugated dienes, selective catalysis, cyclopropanation, vinylcyclopropanes

Vinylcyclopropanes (VCPs) are highly prevalent structural motifs in natural and synthetic bioactive molecules.^{1–5} Owing to their propensity to ring-open in the presence of transition-metal catalysts, their reactivity has been widely studied and they are now frequently used as platforms for further transformations.^{6–12} Retrosynthetically, the transition-metal-catalyzed cyclopropanation of dienes based on diazo-alkane decomposition ranks among the most direct routes for their preparation.^{13,14} In practice, while this is certainly true for symmetrical substrates where both alkenes are equivalent, the situation is more contrasted for unsymmetrical 1,3-dienes. Indeed, the highly enantio- and diastereoselective Cu-catalyzed cyclopropanation of 2,5-dimethyl-2,4-hexadiene (DMHD) for the production of pyrethroids was brought to industrial scale by Aratani and his group at Sumimoto Co., in the 1980s (Figure 1A).² Today, it still constitutes one of the most significant achievements of selective homogeneous catalysis. In comparison, until recently, the cyclopropanation of 1,3-dienes featuring two distinct alkenes was considered of poor synthetic utility because of the low levels of regio- and diastereoselectivity obtained with the transition-metal catalysts typically used for simple olefins.^{15–17} Two recent studies have begun to address these shortcomings. Our group has shown that, using readily available diazo esters, the chiral Cu-bisoxazoline system catalyzes the cyclopropanation of 2-substituted 1,3-dienes with excellent levels of regio- and enantioselectivity but modest *trans/cis* selectivity (from 1:1 to 2:1) (Figure 1B). Because the ester stereocenter (C5) is controlled by the chiral ligand, the lack of diastereocontrol at C2 was circumvented by engaging the VCP mixtures in a subsequent Rh-catalyzed stereoconvergent intermolecular (5 + 2) cycloaddition with a variety of alkynes. Overall, this sequential approach yielded 7-membered rings with

high levels of enantiopurity.¹⁸ Concurrently to this study, Uyeda and co-workers reported a unique dinuclear Ni catalyst for the cyclopropanation of branched dienes using silylated diazo-alkanes. Cyclopropanation occurred exclusively at the most substituted double bond ($rr_{(1,2/3,4)} > 20:1$), and the corresponding racemic VCPs were isolated with moderate to excellent levels of diastereoselectivity (Figure 1C). Quite notably, an unusual diradical mechanism distinct from the classical (2 + 1) cycloaddition involving $M = CR_2$ intermediates was established.¹⁹ Following these advances, we sought to identify a complementary catalytic system for the regioselective cyclopropanation of the terminal alkene in 2-substituted 1,3-dienes (Figure 1D). We report herein the results of our investigations in this direction.

The renewed interest in Ni and Pd catalysis for the development of cyclopropanation reactions prompted us to initiate our investigations with the evaluation of group X transition-metal precatalysts using branched diene **1a** as a model substrate and ethyl diazo acetate **2a** (Table 1).^{20–26} Whereas no reaction was observed with several nickel sources, most of the Pd(II) precatalysts yielded diethyl fumarate and diethyl maleate (entries 1–3 and 5–7). Cyclopropanation occurred to a marginal extent using $Pd(OAc)_2$ but with excellent 3,4-regioselectivity (12% conv., $rr_{3/4} > 20:1$; entry 4). The use of

Received: June 7, 2023

Revised: July 6, 2023

Accepted: July 6, 2023

Published: July 25, 2023







Figure 1. (A) Cu-catalyzed enantioselective cyclopropanation of DMHD. (B) Cu-catalyzed 1,2-regioselective and enantioselective cyclopropanation of branched dienes. (C) Ni-catalyzed 1,2-regioselective, and diastereoselective cyclopropanation of 2-substituted 1,3-dienes. (D) Pd-catalyzed 3,4-regioselective cyclopropanation of 2-substituted 1,3-dienes.

$\text{Pd}_2(\text{dba})_3$ (>98% purity based on Ananikov's method)²⁷ led to a noticeable increase in catalytic activity, a similarly high level of regiocontrol, but essentially no diastereocontrol (entry 8). While catalytic inhibition was observed when P- or N-based ligands were employed, 64% conv. in **3a** (*rr*_{3/4} > 20:1) was achieved in tetrahydrofuran (THF) after a brief solvent survey (entries 9–12). The commercially available [(NHC)Pd(0)] complexes **C**₂ and **C**₃ displayed significant reactivity and selectivity but did not outcompete $\text{Pd}_2(\text{dba})_3$ (entries 13 and 14). In the absence of Lewis acid, no polymerization of ethyl diazo acetate was observed with **C**₂ and **C**₃.^{28,29} Finally, we found that the optimal results were obtained with **C**₄, an underused though readily available Pd(0) precursor (69% conv., *rr*_{3/4} > 20:1, *trans/cis* 1:1; entry 14).³⁰ Under these conditions, the less reactive diazomalonate **2b** enabled the regioselective cyclopropanation of **1a** into VCP **3b** in 47% conversion (*rr*_{3/4} > 20:1), a result that could not be improved at higher temperature (entries 19 and 20).


The generality of the optimized protocol was subsequently evaluated with a representative selection of 2-substituted 1,3-dienes **1a–n** using ethyl diazo acetate **2a** (Figure 2). Substrates containing an electron-poor, an electron-neutral, an electron-rich aromatic ring, or a heteroaromatic substituent delivered the vinylcyclopropanes in usually high yield, low *cis/trans* ratio but with consistently excellent 3,4-regioselectivity. Only the 3-thiophene derivative **3ia** was isolated in low yield (38% yield). *Ortho*-substitution was well-tolerated (**3da–3fa**). Primary, secondary, and even sterically demanding tertiary aliphatic derivatives were cyclopropanated with similar catalytic efficiency (**3ja–3na**). Among the diverse functional groups that were accommodated, the perfect chemo- and regioselectivity observed for substrates featuring a 1,2-(*Z*)-disubstituted alkenes (**3na**), as well as trisubstituted alkenes (**3fa**, **3ka**), is particularly noticeable.

The optimized reaction conditions using **C**₄ were next applied to other classes of 1,3-dienes (Figure 3). We found that terminal diene **1o** underwent cyclopropanation with a very high level of

Table 1. Reaction Optimization^a

entry	catalyst	2	solvent	conv. (%) ^b	rr (3,4/1,2) ^b	trans/cis ^b
1	Ni(cod) ₂	2a	toluene	nr ^c		
2	Ni(cod)(DQ)	2a	toluene	nr		
3	NiCl ₂ (PPh ₃) ₂	2a	toluene	nr		
4	Pd(OAc) ₂	2a	toluene	12	>20:1	1.4:1.0
5	PdCl ₂ (cod)	2a	toluene	nr		
6	(CPhos)Pd(G3)	2a	toluene	nr		
7	C ₁	2a	toluene	nr		
8	Pd ₂ (dba) ₃	2a	toluene	37	>20:1	1.3:1.0
9	Pd ₂ (dba) ₃ /L ₁	2a	toluene	9	>20:1	2.0:1.0
10	Pd ₂ (dba) ₃ /PCy ₃ ^d	2a	toluene	<5	nd ^e	nd
11	Pd ₂ (dba) ₃	2a	CH ₂ Cl ₂	31	>20:1	1.5:1.0
12	Pd ₂ (dba) ₃	2a	THF	64	>20:1	1.2:1.0
13	C ₂	2a	THF	38	>20:1	2.0:1.0
14	C ₃	2a	THF	41	>20:1	1.2:1.0
15	C ₄	2a	THF	69	>20:1	1.0:1.0
16	C ₄ /L ₁	2a	THF	nr		
17	C ₄ /PCy ₃ ^d	2a	THF	18	>20:1	1.0:1.0
18	C ₃	2b	THF	nr		
19	C ₄	2b	THF	47	>20:1	na ^g
20 ^f	C ₄	2b	THF	23	>20:1	na

^aReaction conditions: **1a** (0.1 mmol), **2a–b** (0.12–0.15 mmol).



^bDetermined by ¹H NMR using an internal standard. ^cNo reaction. ^d10 mol % of PCy₃. ^eNot determined. ^fAt 60 °C. ^gNot applicable.

1,2-regioselectivity, affording **30a** in 86% yield (*trans/cis* 1:1). In contrast, symmetrical dienes such as **1p** and **1q** were less reactive, delivering **3pa** and **3qa/3qa'** in 17% and 39% yields, respectively. The robustness of the cyclopropanation protocol was confirmed by successfully conducting the model reaction between **1a** and **2a** on a gram scale (Figure 4). Gratifyingly, the combined yield for this experiment was slightly improved, and, more importantly, we showed that both diastereoisomers could be separated by standard chromatographic purification affording 530 mg of *trans*-**3aa** and 560 mg of *cis*-**3aa**.

The synthetic utility of the VCP was demonstrated through a series of comparative postcatalytic derivatizations

using *trans*-**3aa** and *cis*-**3aa** or the corresponding primary alcohols *trans*-**5aa** and *cis*-**5aa** prepared following a standard reduction procedure (see the Supporting Information for details). We initiated our investigations by evaluating two protocols recently reported by the Marek group for the cyclopropanation and epoxidation of densely substituted alkenyl cyclopropyl carbynols.³¹ Our interest stemmed from the fact that—to the best of our knowledge—the substitution pattern of the VCP generated by the Pd-catalyzed cyclopropanation of 2-substituted 1,3-dienes had not been explored in previous studies.^{32–34} While we did not observe product formation using diiodomethane for the Zn-mediated Simmons–Smith–




Figure 2. Scope of the Pd-catalyzed 3,4-regioselective cyclopropanation of 2-substituted 1,3-dienes (0.5 mmol scale). Regio- and diastereoselectivity determined by ^1H NMR. Yield after purification.

Furukawa cyclopropanation of either *trans*-**5aa** or *cis*-**5aa**, reactivity was restored with the use of chloriodomethane (Figure 5A).³⁵ Biscyclopropyl carbinol *trans*-**6aa** was obtained in a 53% yield after purification by column chromatography starting from *trans*-**5aa**. In contrast, when *cis*-**5aa** was subjected to similar reaction conditions, cyclopropanation of the C=C bond was accompanied by competitive O–H insertion, a feature that is commonly observed with transition metals.^{36,37} The biscyclopropyl carbinol *cis*-**6aa** and the biscyclopropyl methyl ether *cis*-**7aa** thus generated could be separated by column chromatography and isolated in 28% and 48% yields, respectively. While no reaction occurred when attempting to perform a V-catalyzed cyclopropanation of *trans*-**5aa** using

*t*BuOOH as an oxidant, alkenyl cyclopropyl carbinol *cis*-**5aa** afforded **8aa** as a single diastereoisomer (Figure 5B).^{38–40} Unexpectedly, benzoylation of **8aa** using 3,5-dinitrobenzoyl chloride led to the diastereoselective formation of the 3-oxabicyclo[3.1.0]hexane derivative **9aa**, generated through intramolecular $\text{S}_{\text{N}}2$ ring-opening of the epoxide by the pendant alcohol functionality and subsequent benzoylation. The relative stereochemistry of the three contiguous stereocenters in **8aa** and **9aa** was assigned by growing crystals of suitable quality for X-ray analysis of the latter. Finally, we found that Cu-catalyzed protoboration of both *trans*-**3aa** and *cis*-**3aa** led to the formation of the same ring-opened polyfunctional allyl boronate ester (*E*)-**10aa** in good yields and with excellent level of stereocontrol (*E*/




Figure 3. Pd-catalyzed cyclopropanation of differently substituted 1,3-dienes (0.5–1.0 mmol scale). Regio- and diastereoselectivity determined by ¹H NMR. ^aNot determined.




Figure 4. Large-scale experiment. Regio- and diastereoselectivity determined by ¹H NMR.

Z > 20:1) (Figure 5C).^{41–44} Quite notably, the use of morpholine trifluoroacetic acid salt appeared crucial to avoid deborylation, a phenomenon we observed using various alcohols as proton sources. It is worth noting that the stereoconvergent nature of this unprecedented Cu-catalyzed protoboration alleviates the lack of stereocontrol of the Pd-catalyzed 3,4-regioselective cyclopropanation of branched dienes and obviates the need to separate the *cis* and *trans* VCPs.

In conclusion, we have developed a 3,4-regioselective Pd-catalyzed cyclopropanation of 2-substituted 1,3-dienes using readily available diazo esters. The vinylcyclopropanes generated are isolated in practical chemical yields with exquisite regioselectivity and low diastereoselectivity. The method operates under mild reaction conditions, is compatible with a wide number of potentially sensitive functional groups, and can be performed on a gram scale, thus allowing separation of the *cis* and *trans* isomers. A series of postcatalytic derivatizations served


to highlight some of the intrinsic reactivity differences between these diastereoisomeric structures. Among these, an original stereoconvergent Cu-catalyzed ring-opening protoboration mitigated the modest level of stereocontrol of the catalytic cyclopropanation. Current efforts in our laboratory are directed toward understanding the origin of the high levels of regioselectivity obtained in the Pd-catalyzed 3,4-cyclopropanation of branched 1,3-dienes.

EXPERIMENTAL SECTION


General Cyclopropanation Procedure

In a N₂-filled glovebox, a 15% solution of ethyl diazo acetate in toluene (**2a** or **2b**, 0.75 mmol, 1.50 equiv) was added at once to a Schlenk flask containing the appropriate diene **1a–n** (0.5 mmol, 1.00 equiv) and precatalyst **C**₄ (5 mol %) in THF (5 mL, 0.1 M) at room temperature. The reaction mixture was stirred at 25 °C for 30 min. The Schlenk was taken out of the glovebox, and the reaction was quenched by dilution

(A) Simmons-Smith-Furukawa cyclopropanation



(B) V-catalyzed epoxidation



(C) Cu-catalyzed borylative ring-opening




Figure 5. (A) Simmons–Smith–Furukawa cyclopropanations, (B) V-catalyzed epoxidations, and (C) Cu-catalyzed borylative ring-opening reactions.

using THF (5 mL). The crude mixture was concentrated under reduced pressure and adsorbed in silica using CH_2Cl_2 . Purification by flash chromatography using the appropriate eluent yielded the desired vinylcyclopropanes (VCPs).

■ ASSOCIATED CONTENT

Data Availability Statement

The data underlying this study are available in the published article and its Supporting Information.

Supporting Information

The Supporting Information is available free of charge at <https://pubs.acs.org/doi/10.1021/acsorginorgau.3c00024>.

Experimental procedures and characterization of all new compounds and X-ray data for compound **9aa** (PDF)

Accession Codes

CCDC 2266910 contains the supplementary crystallographic data for this paper. These data can be obtained free of charge via www.ccdc.cam.ac.uk/data_request/cif, or by emailing data_request@ccdc.cam.ac.uk, or by contacting The Cambridge Crystallographic Data Centre, 12 Union Road, Cambridge CB2 1EZ, UK; fax: +44 1223 336033.

AUTHOR INFORMATION

Corresponding Author

Clément Mazet – Department of Organic Chemistry, University of Geneva, 1211 Geneva, Switzerland; orcid.org/0000-0002-2385-280X; Email: clement.mazet@unige.ch

Authors

Agonist Kastrati – Department of Organic Chemistry, University of Geneva, 1211 Geneva, Switzerland

Vincent Jaquier – Department of Organic Chemistry, University of Geneva, 1211 Geneva, Switzerland

Michele Garbo – Department of Organic Chemistry, University of Geneva, 1211 Geneva, Switzerland

Céline Besnard – Laboratory of Crystallography, University of Geneva, 1211 Geneva, Switzerland; orcid.org/0000-0001-5699-9675

Complete contact information is available at:

<https://pubs.acs.org/doi/10.1021/acsorginorgau.3c00024>

Author Contributions

CRedit: **Agonist Kastrati** formal analysis (lead), methodology (lead), writing-review & editing (supporting); **Vincent Jaquier** formal analysis (lead), methodology (lead), writing-review & editing (supporting); **Michele Garbo** data curation (lead), methodology (lead), writing-review & editing (supporting); **Céline Besnard** data curation (lead), formal analysis (lead); **Clément Mazet** conceptualization (lead), formal analysis (supporting), funding acquisition (lead), investigation (supporting), methodology (supporting), project administration (lead), resources (lead), supervision (lead), writing-original draft (lead), writing-review & editing (lead).

Notes

The authors declare no competing financial interest.

ACKNOWLEDGMENTS

Financial support was provided by the University of Geneva and the Swiss National Foundation (grants 200020_175489 and 200021_188490). Stéphane Rosset (University of Geneva) was acknowledged for measuring HRMS. Marion Pupier (University of Geneva) was acknowledged for assistance with NMR analyses. Dr. Sylvain Taillemaud (University of Geneva) was acknowledged for fruitful scientific discussions.

REFERENCES

- (1) Aratani, T.; Yoneyoshi, Y.; Nagase, T. Asymmetric synthesis of chrysanthemic acid. An application of copper carbenoid reaction. *Tetrahedron Lett.* **1975**, *16*, 1707–1710.
- (2) Aratani, T. Catalytic asymmetric synthesis of cyclopropanecarboxylic acids: an application of chiral copper carbenoid reaction. *Pure Appl. Chem.* **1985**, *57*, 1839–1844.
- (3) Doyle, M. P.; Protopopova, M. N. New Aspects of Catalytic Asymmetric Cyclopropanation. *Tetrahedron* **1998**, *54*, 7919–7946.
- (4) Suenobu, K.; Itagaki, M.; Nakamura, E. Reaction Pathway and Stereoselectivity of Asymmetric Synthesis of Chrysanthemate with the Aratani C₁-Symmetric Salicylaldimine–Copper Catalyst. *J. Am. Chem. Soc.* **2004**, *126*, 7271–7280.
- (5) Talele, T. T. The “cyclopropyl fragment” is a versatile player that frequently appears in preclinical/clinical drug molecules. *J. Med. Chem.* **2016**, *59*, 8712–8756.
- (6) Rubin, M.; Rubina, M.; Gevorgyan, V. Transition Metal Chemistry of Cyclopropenes and Cyclopropanes. *Chem. Rev.* **2007**, *107*, 3117–3179.
- (7) Jiao, L.; Yu, Z.-X. Vinylcyclopropane Derivatives in Transition-Metal-Catalyzed Cycloadditions for the Synthesis of Carbocyclic Compounds. *J. Org. Chem.* **2013**, *78*, 6842–6848.
- (8) Souillart, L.; Cramer, N. Catalytic C–C Bond Activations via Oxidative Addition to Transition Metals. *Chem. Rev.* **2015**, *115*, 9410–9464.
- (9) Fumagalli, G.; Stanton, S.; Bower, J. F. Recent Methodologies That Exploit C–C Single-Bond Cleavage of Strained Ring Systems by Transition Metal Complexes. *Chem. Rev.* **2017**, *117*, 9404–9432.
- (10) Meazza, M.; Guo, H.; Rios, R. Synthetic applications of vinyl cyclopropane opening. *Org. Biomol. Chem.* **2017**, *15*, 2479–2490.
- (11) Pirenne, V.; Muriel, B.; Waser, J. Catalytic Enantioselective Ring-Opening Reactions of Cyclopropanes. *Chem. Rev.* **2021**, *121*, 227–263.
- (12) Wang, J.; Blaszczyk, S. A.; Li, X.; Tang, W. Transition Metal-Catalyzed Selective Carbon–Carbon Bond Cleavage of Vinylcyclopropanes in Cycloaddition Reactions. *Chem. Rev.* **2021**, *121*, 110–139.
- (13) Lebel, H.; Marcoux, J.-F.; Molinaro, C.; Charette, A. B. Stereoselective Cyclopropanation Reactions. *Chem. Rev.* **2003**, *103*, 977–1050.
- (14) Davies, H. M. L.; Manning, J. R. Catalytic C–H Functionalization by Metal Carbenoid and Nitrenoid Insertion. *Nature* **2008**, *451*, 417–424.
- (15) Doyle, M. P.; Dorow, R. L.; Tamblyn, W. H.; Buhro, W. E. Regioselectivities of Cyclopropanation Reactions with Ethyl Diazoacetate. *Tetrahedron Lett.* **1982**, *23*, 2161–2264.
- (16) Doyle, M. P.; Dorow, R. L.; Buhro, W. E.; Griffin, J. H.; Tamblyn, W. H.; Trudell, M. L. Stereoselectivity of Catalytic Cyclopropanation Reactions. Catalyst Dependence in Reactions of Ethyl Diazoacetate with Alkenes. *Organometallics* **1984**, *3*, 44–52.
- (17) Murelli, R. P.; Catalán, S.; Gannon, M. P.; Snapper, M. L. Ruthenium-Catalyzed Tandem Enyne-Cross Metathesis-Cyclopropanation: Three-Component Access to Vinyl Cyclopropanes. *Tetrahedron Lett.* **2008**, *49*, 5714–5717.
- (18) Garbo, M.; Besnard, C.; Guenée, L.; Mazet, C. Access to Optically Active 7-Membered Rings by a 2-Step Synthetic Sequence: Cu Catalyzed Stereoselective Cyclopropanation of Branched 1,3-Dienes/Rh-Catalyzed Stereoconvergent [5 + 2] Cycloaddition. *ACS Catal.* **2020**, *10*, 9604–9611.
- (19) Maity, A. K.; Kalb, A. E.; Zeller, M.; Uyeda, C. A Dinickel Catalyzed Cyclopropanation without the Formation of a Metal Carbene Intermediate. *Angew. Chem., Int. Ed.* **2021**, *60*, 1897–1902.
- (20) Tomilov, Y. V.; Bordakov, V. G.; Dolgii, I. E.; Nefedov, O. M. Reaction of diazoalkanes with unsaturated compounds. Cyclopropanation of olefins by diazomethane in presence of palladium compounds. *Bull. Acad. Sci. USSR, Div. Chem. Sci.* **1984**, *3*, 533–538.
- (21) Denmark, S. E.; Stavenger, R. A.; Faucher, A.-M.; Edwards, J. P. Cyclopropanation with Diazomethane and Bis(oxazoline)palladium(II) Complexes. *J. Org. Chem.* **1997**, *62*, 3375–3389.
- (22) Chen, S.-F.; Ma, J.; Wang, J.-B. Palladium-catalyzed cyclopropanation of electron-deficient olefins with aryl diazocarbonyl compounds. *Tetrahedron Lett.* **2008**, *49*, 6781–6783.
- (23) Martín, C.; Molina, F.; Alvarez, E.; Belderrain, T. R. Stable N-Heterocyclic Carbene (NHC)–Palladium(0) Complexes as Active

Catalysts for Olefin Cyclopropanation Reactions with Ethyl Diazoacetate. *Chem. - Eur. J.* **2011**, *17*, 14885–14895.

(24) Zhou, Y.-Y.; Uyeda, C. Reductive Cyclopropanations Catalyzed by Dinuclear Nickel Complexes. *Angew. Chem., Int. Ed.* **2016**, *55*, 3171–3175.

(25) Sakurai, S.; Inagaki, T.; Kodama, T.; Yamanaka, M.; Tobisu, M. Palladium-Catalyzed Siloxycyclopropanation of Alkenes Using Acylsilanes. *J. Am. Chem. Soc.* **2022**, *144*, 1099–1105.

(26) Inagaki, T.; Sakurai, S.; Yamanaka, M.; Tobisu, M. Palladium-Catalyzed Silylacylation of Allenes Using Acylsilanes. *Angew. Chem., Int. Ed.* **2022**, *61*, No. e202202387.

(27) Zaleskiy, S. S.; Ananikov, V. P. $\text{Pd}_2(\text{dba})_3$ as a Precursor of Soluble Metal Complexes and Nanoparticles: Determination of Palladium Active Species for Catalysis and Synthesis. *Organometallics* **2012**, *31*, 2302–2309.

(28) Ihara, E.; Ishiguro, Y.; Yoshida, N.; Hiraren, T.; Itoh, T.; Inoue, K. (N-Heterocyclic Carbene) Pd/Borate Initiating Systems for Polymerization of Ethyl Diazoacetate. *Macromolecules* **2009**, *42*, 8608–8610.

(29) Zhukhovitskiy, A. V.; Kobylanskiy, I. J.; Thomas, A. A.; Evans, A. M.; Delaney, C. P.; Flanders, N. C.; Denmark, S. E.; Dichtel, W. R.; Toste, F. D. A Dinuclear Mechanism Implicated in Controlled Carbene Polymerization. *J. Am. Chem. Soc.* **2019**, *141*, 6473–6478.

(30) Itoh, K.; Ueda, F.; Hirai, K.; Ishii, Y. New mixed olefin complexes of palladium (0). A stabilization of the $\text{Pd}^{(0)}$ state with a combination of electron-donating and electron-withdrawing olefins. *Chem. Lett.* **1977**, *6*, 877–880.

(31) Cohen, A.; Siddaraju, Y.; Marek, I. Directed Diastereoselective Cyclopropanation and Epoxidation of Alkenyl Cyclopropyl Carbinol Derivatives. *Org. Lett.* **2022**, *24*, 8322–8325.

(32) Simmons, H. E.; Smith, R. D. A New Synthesis of Cyclopropanes from Olefins. *J. Am. Chem. Soc.* **1958**, *80*, 5323–5324.

(33) Charette, A. B.; Beauchemin, A. Simmons-Smith Cyclopropanation Reaction. *Org. React.* **2001**, *58*, 1–415.

(34) Nakamura, M.; Hirai, A.; Nakamura, E. Reaction Pathways of the Simmons–Smith Reaction. *J. Am. Chem. Soc.* **2003**, *125*, 2341–2350.

(35) Denmark, S. E.; Edwards, J. P. A Comparison of (Chloromethyl)- and (Iodomethyl)zinc Cyclopropanation Reagents. *J. Org. Chem.* **1991**, *56*, 6974–6981.

(36) *Reactive Intermediate Chemistry*; Moss, R. A.; Platz, M. S.; Jones, M., Jr., Eds.; John Wiley & sons, Inc.: Hoboken, NJ, 2004.

(37) Zhu, S.-F.; Zhou, Q.-L. Transition-Metal-Catalyzed Enantioselective Heteroatom–Hydrogen Bond Insertion Reactions. *Acc. Chem. Res.* **2012**, *45*, 1365–1377.

(38) Ligtenbarg, A. G. J.; Hage, R.; Feringa, B. L. Catalytic Oxidations by Vanadium Complexes. *Coord. Chem. Rev.* **2003**, *237*, 89–101.

(39) Mihelich, E. D.; Daniels, K.; Eickhoff, D. J. Vanadium-Catalyzed Epoxidations. 2. Highly Stereoselective Epoxidations of Acyclic Homoallylic Alcohols Predicted by a Detailed Transition-State Model. *J. Am. Chem. Soc.* **1981**, *103*, 7690–7692.

(40) When the epoxidation was conducted with *m*CPBA, we observed a *dr* > 10:1 based on ^1H NMR analysis of the crude reaction mixture.

(41) For related (seminal) work on 1,3-dienes, see: Sasaki, Y.; Zhong, C.; Sawamura, M.; Ito, H. Copper(I)-Catalyzed Asymmetric Monoborylation of 1,3-Dienes: Synthesis of Enantioenriched Cyclic Homoallyl- and Allylboronates. *J. Am. Chem. Soc.* **2010**, *132*, 1226–1227.

(42) Sebelius, S.; Olsson, V. J.; Szabó, K. J. Palladium Pincer Complex Catalyzed Substitution of Vinyl Cyclopropanes, Vinyl Aziridines, and Allyl Acetates with Tetrahydroxydiboron. An Efficient Route to Functionalized Allylboronic Acids and Potassium Trifluoro(allyl)borates. *J. Am. Chem. Soc.* **2005**, *127*, 10478–10479.

(43) Chen, C.; Wang, H.; Li, T.; Lu, D.; Li, J.; Zhang, X.; Hong, X.; Lu, Z. Cobalt-Catalyzed Asymmetric Sequential Hydroboration/Isomerization/Hydroboration of 2-Aryl Vinylcyclopropanes. *Angew. Chem., Int. Ed.* **2022**, *61*, No. e202205619.

(44) Lu, D.; Chen, C.; Zheng, L.; Ying, J.; Lu, Z. Regio- and Stereoselective Cobalt-Catalyzed Hydroboration of Vinylcyclopro-

panes to Access Homoallylic Boronates *Organometallics* **2023**, DOI: 10.1021/acs.organomet.2c00592.