LEMMA

CLASSE SUPERIOR EM ATIVOS

BIFIDOBACTERIUM INFANTIS

Bifidobacterium é uma bactéria em forma de bastonete anaeróbias, o que significa que não pode crescer na presença de oxigênio. É normalmente encontrado nos intestinos e é capaz de quebrar açúcares em ácido láctico. Esta bactéria é também capaz de usar alguns compostos "não digeríveis" plantas, incluindo algumas proteínas e celulose, para o combustível. Sua capacidade de usar muitos compostos diferentes como o alimento lhe permite prosperar no trato digestivo e competir com bactérias potencialmente prejudiciais para o espaço e os recursos no intestino.

Bifidobacterium Infantis é um membro da família da bifidobactérias, uma estirpe de bactérias que é normalmente encontrada no intestino humano. Uma vez que estas bactérias normalmente não causam infecções, elas ajudam a manter a flora intestinal saudável.

Propriedades e Benefícios

- Síndrome do Intestino Irritável;
- Prevenção da Enterocolite Necrosante;
- Tratamento da Diarréia do Viajante e Colite Ulcerativa;
- Redução dos sintomas de Infecções Pulmonares em crianças;
- Ajuda a prevenir a diarreia causada pelo uso de antibióticos.

Rua Dr. Arthur Costacurta, 550 – Área Industrial.
Jardinópolis / SP – CEP 14.680-000
Escritório: 11 5044 7234 – São Paulo / SP
www.lemma.com.br - lemma@lemma.com.br

LEMMA

CLASSE SUPERIOR EM ATIVOS

Posologia Recomendada

As doses típicas usadas em estudos **variaram de 1 a 2 bilhões de UFC/dia.**

A dosagem de Probioticos varia dependendo do produto e indicação específica. Não existe consenso quanto ao número mínimo de microrganismos que devem ser ingeridos para obter um efeito benéfico.

Tipicamente, um probiótico deve conter várias bactérias e milhões de microrganismos para aumentar a probabilidade de colonização intestinal adequada.

Referências:

1. EZAKI, T. HASHIMOTO, Y.& YABUUCHI, E(1989). Fluorometric deoxyribonucleic acid-deoxyribonucleic acid hybridization in microdilution wells as an alternative to membrane ®lter hybridization in which radioisotopes are used to determine genetic relatedness among bacterial strains. *Int J Syst Bacteriol* 39, 224±229.

2. HIRAISHI, A. IWASAKI, M.& SHINJO, H.(2000). Terminal restriction pattern analysis of 16S rRNA genes for the characterization of bacterial communities of activated sludge. *J Biosci Bioeng* 90, 148±156.

3. JIAN, W.ZHU, L.J.DONG, X.(2001) New approach to phylogenetic analysis of the genus *Bi®dobacterium* based on partial HSP60 gene sequences. *Int J Syst Evol Microbiol* 51, 1633±1638.

4. LAUER, E.&KANDLER, O.(1983). DNA-DNA homology, murein types, and enzyme patterns in the type strains of the genus *Bi®dobacterium*. *Syst Appl Microbiol* 4, 42±64.

5. LIN, AW.USERA, M.A.,BARRETT, T.J.&GOLSBY, R.A.(1996). Application of random ampli®ed polymorphic DNA analysis to differentiate strains of *salmonella*. *J Clin Microbiol* 34, 870±876.

6. MANGIN, I.BOURGET, N., BOUHNIK, Y.BISETTI, N.,SIMONET, J.M&DECARIS, .(1994) Identi®cation of *Bi®dobacterium* strains by rRNA gene restriction patterns. *Appl Environ Microbiol* 60, 1451±1458.

7. MARTIN, P. Identi®cation and typing of *Streptomyces* strains : evaluation of interspeci®c, intraspeci ®c and intraclonal differences by RAPD ®ngerprinting. *Res Microbiol* 151

Rua Dr. Arthur Costacurta, 550 – Área Industrial.
Jardinópolis / SP – CEP 14.680-000
Escritório: 11 5044 7234 – São Paulo / SP
www.lemma.com.br - lemma@lemma.com.br