

Review of PDF and R

Emanuele Coradin

March 2024

Contents

1	Pairing and ordering objects	1
2	Probability distributions in R	1
2.1	Binomial distribution	1
2.2	Geometric Distribution	4

1 Pairing and ordering objects

We can identify different useful tools to compute the numbers of different pairings as orderings that we can apply to a sequence of length r taken from a dictionary of n objects.

2 Probability distributions in R

R provides almost all the standard PDFs that we could wish. The name convention prescribes: `*d` for the pdf `*p` for the cumulative density function (cdf) `*q` for the quantile function `*r` to sample a random number from the distribution.

Now we will experiment with them with same simple exercise.

2.1 Binomial distribution

The Binomial probability function is described by: $P(X = k) = \binom{n}{k} \cdot p^k \cdot (1 - p)^{n-k}$

First of all let's define a palette.

```
color_vector <- c("#CC0000", # Wine-like color
                  "#0072B2", # Strong blue
                  "#CC79A7", # Muted purple
                  "#D55E00", # Vermilion
                  "#009E73", # Bluish green
                  "#56B4E9", # Sky blue
                  "#E69F00") # Yellow-orange
```

```
# Parameters for the binomial distribution
```

```
n <- c(5, 10, 15, 20) # number of trials
linetype <- 2-(1:length(n))%2
p <- 1./2. # probability of success
```

```
# Generate values for x (number of successes)
x <- 0:20
```

```

#define a function to prepare the plots
plot_pdf <- function(N) {dbinom(x, N, p)}
binomial_pdf <- sapply(n, plot_pdf)

# create the first straight line plot
plot(x,binomial_pdf[,1], type = "s", lwd = 2, col = color_vector[1],
     xlab = "Number of Successes", ylab = "Probability", lty = linetype[1])


# create all the others
for (i in 2:length(n)) {
  lines(x, binomial_pdf[, i], col=color_vector[i], type='s', lwd=2, lty = linetype[i])
}

grid() # Add grid

# Create legend labels with strings "n = "
legend_labels <- paste("n =", n)
# Add a legend
legend("topright", legend = legend_labels, col = color_vector[1:length(n)], lty = linetype, lwd = 2)
# Add a title
title("Theoretical Binomial Distribution P = 0.5")

```

Theoretical Binomial Distribution P = 0.5



```

# Parameters for the binomial distribution

n <- c(5, 10, 15, 20) # number of trials
linetype <- 2-(1:length(n))%2
p <- 1./5. # probability of success

```

```

# Generate values for x (number of successes)
x <- 0:20

#define a function to prepare the plots
plot_pdf <- function(N) {dbinom(x, N, p)}
binomial_pdf <- sapply(n, plot_pdf)

# create the first straight line plot
plot(x,binomial_pdf[,1], type = "s", lwd = 2, col = color_vector[1],
      xlab = "Number of Successes", ylab = "Probability", lty = linetype[1])


# create all the others
for (i in 2:length(n)) {
  lines(x, binomial_pdf[, i], col=color_vector[i], type='s', lwd=2, lty = linetype[i])
}

grid() # Add grid

# Create legend labels with strings "n = "
legend_labels <- paste("n =", n)
# Add a legend
legend("topright", legend = legend_labels, col = color_vector[1:length(n)], lty = linetype, lwd = 2)
# Add a title
title("Theoretical Binomial Distribution P = 0.2")

```

Theoretical Binomial Distribution P = 0.2



2.2 Geometric Distribution

The Geometric distribution can be described by: $P(X = k) = (1 - p)^{k-1} \cdot p$

Now let's visualize it.

```
# Parameters for the binomial distribution

p <- c(1./2., 1/4, 1/10, 10**(-3)) # probability of success
linetype <- 2-(1:length(p))%2

# Generate values for x (number of successes)
x <- 0:20

#define a function to prepare the plots
plot_pdf <- function(P) {dgeom(x, P)}
binomial_pdf <- sapply(p, plot_pdf)

# create the first straight line plot
plot(x,binomial_pdf[,1], type = "s", lwd = 2, col = color_vector[1],
      xlab = "Number of Successes", ylab = "Probability", lty = linetype[1], log="y")

# create all the others
for (i in 2:length(p)) {
  lines(x, binomial_pdf[, i], col=color_vector[i], type='s', lwd=2, lty = linetype[i])
}

grid() # Add grid

# Create legend labels with strings "p = "
legend_labels <- paste("p =", p)
# Add a legend
legend("topright", legend = legend_labels, col = color_vector[1:length(p)], lty = linetype, lwd = 2)
# Add a title
title("Theoretical Geometric distribution")
```

Theoretical Geometric distribution

