



XGBoost



Prediction of L -th tree


each check is
an inequality
for one feature



$g(x)$: function mapping
sample x to
leaf j

Prediction of t -th tree $\Rightarrow g_t(x) = w_t(q(x))$

each check is
an inequality
for one feature




$q(x)$: function mapping
sample x to
leaf j

Questions

- Q_1 • how to find best split ($x^i < v$) for every node?
- Q_2 • when to stop splitting and set a leaf?
- Q_3 • why a linear combination of tree's predictions? ($G = \sum_{t=1}^T g_t$)

Loss function

$$C(\{x_m\}, G) = \sum_{m=1}^N \underbrace{\ell(y_m, \hat{y}_m)}_{\text{convex loss function}} + \sum_{t=1}^T \Omega(\theta_t)$$



convex
loss function

$N \geq m$ samples

e.g. square deviation $\frac{1}{2}(y - \hat{y})^2$

• cross entropy for classification $y = (0, 1)$

Loss function

$$C(\{x_m\}, G) = \sum_{m=1}^N \ell(y_m, \hat{y}_m) + \sum_{t=1}^T \Omega(g_t)$$

regularization
for every tree $t \leq T$

$$\Omega(g_t) = \gamma \cdot J_t + \underbrace{\frac{\lambda}{2} \|w_t\|^2}_{\text{Ridge on } w\text{'s}}$$

"cost"
of one
leaf

leaves

($J_t = 4$ in the example)

Loss function

$$C(\{x_m\}, G) = \sum_{m=1}^N \ell(y_m, \hat{y}_m) + \sum_{t=1}^T \Omega(\theta_t)$$

build C iteratively at every "time" t

$t=1$

$C_1 \rightarrow C_2 \rightarrow \dots \rightarrow C_{t-1} \rightarrow C_t \rightarrow \dots \rightarrow C_T \equiv C$

$t \gg 1$: perturbation expansion, Taylor

Loss function, up to time τ

$$C_{\tau}(\{x_m\}, G) = \sum_{m=1}^N \ell(y_m, \hat{y}_m^{(\tau)}) + \sum_{t=1}^{\tau} \Omega(g_t)$$

$$\hat{y}_m^{(\tau)} = \hat{y}_m^{(\tau-1)} + g_t(x_m)$$

prediction
up to tree
 $\tau-1$

addition to
prediction
from tree
 τ

Taylor
(answers)
 R_3



$$= C_{\tau-1} + \Delta C_{\tau}$$

Contribution to loss function from tree τ

Taylor expansion

$$\Delta C_\tau = \sum_m \left[\left. \partial_z \ell(y_m, z) \right|_{z = \hat{y}_m^{(\tau-1)}} g_\tau(x_m) + \frac{1}{2} \left. \partial_z^2 \ell(y_m, z) \right|_{z = \hat{y}_m^{(\tau-1)}} g_\tau(x_m)^2 \right] + \mathcal{L}(g_\tau)$$

def. a_m, b_m


$$\equiv \sum_m \left[a_m g_\tau(x_m) + \frac{1}{2} b_m g_\tau(x_m)^2 \right] + \mathcal{L}(g_\tau)$$

⚠ neglecting terms $O(g_t^3)$

Contribution to loss function from tree τ

Taylor expansion

$$\Delta C_{\tau} = \sum_m \left[a_m g_{\tau}(x_m) + \frac{1}{2} b_m g_{\tau}(x_m)^2 \right] + \mathcal{O}(g_{\tau})$$



all samples sent by tree τ (via $q_{\tau}(x_m)$)
to the same leaf j
will have the same $g_{\tau}(x_m) = W_{\tau}(j)$

Contribution to loss function from tree τ

Taylor expansion

$$\Delta C_\tau = \sum_m \left[a_m g_\tau(x_m) + \frac{1}{2} b_m g_\tau(x_m)^2 \right] + \Omega(g_\tau)$$

all samples sent by tree τ (via $g_\tau(x_m)$)
to the same leaf j

will have the same $g_\tau(x_m) = W_\tau(j)$

rewrite

$$\sum_{\substack{m \\ \text{samples}}} \dots = \sum_{\substack{j=1 \\ \text{leaves}}}^{J_\tau} \sum_{m \in \mathcal{I}_\tau(j)} \dots \quad \parallel \quad \begin{array}{l} \text{Indicator} \\ \mathcal{I}_\tau(j) = \{m \mid g_\tau(x_m) = j\} \end{array}$$

Contribution to loss function from tree τ

Taylor expansion

$$\Delta C_{\tau} = \sum_m \left[a_m g_{\tau}(x_m) + \frac{1}{2} b_m g_{\tau}(x_m)^2 \right] + \underbrace{\Omega(g_{\tau})}_{\text{regularization}}$$

$$= \sum_{j=1}^{J_{\tau}} \left[\underbrace{\left(\sum_{m \in I_{\tau}(j)} a_m \right)}_{A_{\tau}(j)} w_{\tau}(j) + \frac{1}{2} \underbrace{\left(\sum_{m \in I_{\tau}(j)} b_m \right)}_{B_{\tau}(j)} w_{\tau}(j)^2 + \underbrace{\lambda}_{\text{regularization}} w_{\tau}(j)^2 \right] + \underbrace{\gamma J_{\tau}}_{\text{regularization}}$$


Contribution to loss function from tree τ

$$\Delta C_{\tau} = \sum_{j=1}^{J_{\tau}} \left[A_{\tau}(j) w_{\tau}(j) + \frac{1}{2} (B_{\tau}(j) + \lambda) w_{\tau}(j)^2 \right] + \gamma J_{\tau}$$

\forall leaf j , parabola

$$\frac{B+\lambda}{2} w^2 + Aw + \gamma J$$

$$B+\lambda > 0$$




$$\text{best } w_{\tau}^*(j) = - \frac{A_{\tau}(j)}{B(j) + \lambda}$$

Contribution to loss function from tree τ

$$\Delta C_{\tau} = -\frac{1}{2} \sum_{j=1}^{J_{\tau}} \frac{A_{\tau}(j)^2}{B_{\tau}(j) + \lambda} + \gamma J_{\tau}$$

optimized
with Newton's
method (w^*)

⚠ very quick!
no gradient descent



$$\text{best } w_{\tau}^*(j) = -\frac{A_{\tau}(j)}{B_{\tau}(j) + \lambda}$$

Questions

Q₁

- how to find best split ($x^i < v$) for every node?

Q₂

- when to stop splitting and set a leaf?




there is a huge amount of possible combinations (9, 9)

weights tree
 structure

Practical implementation of XGBoost


for many samples ($N \gg 1$) with many features ($L \gg 1$)

- partition each feature in percentiles, v^i




Split


v^i



VS



L



R

Is the original leaf on the left better or worse than the two leaves on the right?

Practical implementation of XGBoost

Greedy algorithm
in every current leaf \Rightarrow


A, B
?

try many splits
(V percentile)

Practical implementation of XGBoost

Greedy algorithm

in every current leaf \Rightarrow



try many splits
(V percentile)

choose the one providing smallest ΔC_{SPLIT}

$$\Delta C_{\text{SPLIT}} = -\frac{1}{2} \left(\frac{A_L^2}{B_L + \lambda} + \frac{A_R^2}{B_R + \lambda} - \frac{A^2}{B + \lambda} \right) + \gamma$$

L/R split


all together \Rightarrow


$Q_1 \checkmark$

- start from a root node
 - iterate split for every node until
 - a) it is not convenient (all $\Delta C_{\text{split}} > 0$)
 - b) conditions max depth, etc., are met
- $\Rightarrow Q_2 \checkmark$



root





this was too easy

normally the 1st tree does not solve it all, and the $\hat{y}_m - y_m$ difference is the starting point for the next tree

Example of A, B , for square deviation loss f.

(\hat{y}_m up to previous
free)

$$\ell(y_m, \hat{y}_m) = \frac{1}{2} (\hat{y}_m - y_m)^2$$

$$a_m = \left. \partial_z \ell(y_m, z) \right|_{z=\hat{y}_m} = \hat{y}_m - y_m \quad \Rightarrow \quad A(j) = \sum_{m \in \mathcal{I}(j)} (\hat{y}_m - y_m)$$

$$b_m = \left. \partial_z^2 \ell(y_m, z) \right|_{z=\hat{y}_m} = 1 \quad \Rightarrow \quad B(j) = \sum_{m \in \mathcal{I}(j)} 1 =$$

= # samples sent to leaf j by $g(x)$