RANDOMIZED CONTROLLED TRIAL · DIGITAL MENTAL HEALTH

# Delivering Cognitive Behavior Therapy to Young Adults With Symptoms of Depression and Anxiety Using a Fully Automated Conversational Agent (Woebot): A Randomized Controlled Trial

**Kathleen Kara Fitzpatrick, PhD[1]*, Alison Darcy, PhD[2]*, Molly Vierhile, BA[1]**

[1] Stanford School of Medicine, Department of Psychiatry and Behavioral Sciences, Stanford, CA, United States
[2] Woebot Labs Inc., San Francisco, CA, United States
* Contributed equally. Corresponding author: Alison Darcy, PhD

Received 29 March 2017 · Revised 5 May 2017 · Accepted 22 May 2017 · Published 6 June 2017
**Abridged from JMIR Mental Health 2017;4(2):e19, doi:10.2196/mental.7785 · © the authors, CC BY 4.0**

## ABSTRACT

**Background** Web-based cognitive-behavioral therapeutic (CBT) apps have demonstrated efficacy but are characterized by poor adherence. Conversational agents may offer a convenient, engaging way of getting support at any time.

**Objective** The objective of the study was to determine the feasibility, acceptability, and preliminary efficacy of a fully automated conversational agent to deliver a self-help program for college students who self-identify as having symptoms of anxiety and depression.

**Methods** In an unblinded trial, 70 individuals age 18-28 years were recruited online from a university community social media site and were randomized to receive either 2 weeks (up to 20 sessions) of self-help content derived from CBT principles in a conversational format with a text-based conversational agent (Woebot) (n=34) or were directed to the National Institute of Mental Health ebook, "Depression in College Students," as an information-only control group (n=36). All participants completed Web-based versions of the 9-item Patient Health Questionnaire (PHQ-9), the 7-item Generalized Anxiety Disorder scale (GAD-7), and the Positive and Negative Affect Scale at baseline and 2-3 weeks later (T2).

**Results** Participants were on average 22.2 years old (SD 2.33), 67% female (47/70), mostly non-Hispanic (93%, 54/58), and Caucasian (79%, 46/58). Participants in the Woebot group engaged with the conversational agent an average of 12.14 (SD 2.23) times over the study period. No significant differences existed between the groups at baseline, and 83% (58/70) of participants provided data at T2 (17% attrition). Intent-to-treat univariate analysis of covariance revealed a significant group difference on depression such that those in the Woebot group significantly reduced their symptoms of depression over the study period as measured by the PHQ-9 ($F$=6.47; $P$=.01) while those in the information control group did not. In an analysis of completers, participants in both groups significantly reduced anxiety as measured by the GAD-7 ($F_{1,54}$= 9.24; $P$=.004). Participants' comments suggest that process factors were more influential on their acceptability of the program than content factors mirroring traditional therapy.

**Conclusions** Conversational agents appear to be a feasible, engaging, and effective way to deliver CBT.

**Keywords** conversational agents; mobile mental health; mental health; chatbots; depression; anxiety; college students; digital health

**Trial registration** None. The trial involved a nonclinical population of college students and was considered exempt from registration in a public trials registry; it is reported with the CONSORT-EHEALTH checklist.

## Introduction

Up to 74% of mental health diagnoses have their first onset before the age of 24 (Kessler et al., 2007). Depression and anxiety symptoms are particularly common among college students, with more than half reporting symptoms of anxiety and depression in the previous year that were so severe they had difficulty functioning (Zivin et al., 2009). In addition, epidemiological data

suggest that mental health problems are both increasing in prevalence and severity (Hunt & Eisenberg, 2010). However, up to 75% of the college students that need them do not access clinical services (Hunt & Eisenberg, 2010). While the reasons for this are varied, the ubiquity of free or inexpensive mental health services on campuses suggests that service availability and cost are not primary barriers to care (Hunt & Eisenberg, 2010). Like non-college populations, stigma is considered the primary barrier to accessing psychological health services.

Overcoming problems of stigma has been traditionally considered a major benefit of Internet-delivered and more recently mobile mental health interventions. In recent years, there has been an explosion of interest and development of such services to either supplement existing mental health treatments or expand limited access to quality mental health services (Bakker et al., 2016). This development is matched by great patient demand with about 70% showing interest in using mobile apps to self-monitor and self-manage their mental health (Torous et al., 2014). Internet interventions for anxiety and depression have empirical support (Spek et al., 2007) with outcomes comparable to therapist-delivered cognitive behavioral therapy (CBT) (Andersson & Cuijpers, 2009; Barak et al., 2008). Yet, despite demonstrated efficacy, they are characterized by relatively poor adoption and adherence. One review found a median minimal completion rate of 56% (Donkin et al., 2013). A hypothesized reason for this lack of adherence is the loss of the human interactional quality that in-person CBT retains.

With recent advancements in voice recognition, conversational interfaces (ie, those that use natural language as inputs and outputs) have begun to emerge. Conversational agents (such as Apple's Siri or Amazon's Alexa) may be a more natural medium through which individuals engage with technology. Humans respond and converse with nonhuman agents in ways that mirror emotional and social discourse dynamics when discussing behavioral health (Bickmore et al., 2005) and their capacity to act as first responders has already been evaluated (Miner et al., 2016). Theoretically, conversational interfaces may be better positioned than visually oriented mobile apps to deliver structured, manualized therapies because in addition to delivering therapeutic content, they can mirror therapeutic process. Indeed, Bickmore et al. (2005) demonstrated that a carefully designed health-related conversational agent could establish a therapeutic relationship with adults attempting to increase exercise. The intervention was an embodied conversational agent, that is, it was designed with a graphical face to mirror human interactions that are typically face-to-face.

Thus, the objective of this study was to assess the feasibility of delivering CBT in a conversational interface via an automated bot in a way that facilitates engagement and reduction in symptoms. The current study compared outcomes from 2 weeks of a CBT-oriented conversational agent (Woebot), or an information control group (National Institute of Mental Health's [NIMH] ebook) in a nonclinical college population. We hypothesized that conversation with a therapeutic process-oriented conversational agent would lead to greater improvement in symptoms relative to the information control group. We also hypothesized that receiving psychoeducational material in a conversational manner would be more acceptable to those who received it.

## Methods

### Recruitment and Procedure

Potential participants were recruited using a flyer posted on social media websites targeting a US university community for students who self-identified as experiencing symptoms of depression and anxiety. Inclusion criteria included age 18 and over (screened at the first level via checkbox confirmation) and able to read English (implied). To guard against compromise, for example from malicious bots, all potential participants were sent an email requesting that they respond denoting their confirmation. Confirmed participants were randomized via computer algorithm that automatically generated a number between 0 and 1. Participants with numbers <0.5 were allocated to receive a direct link to begin chatting with Woebot in an instant messenger app, and participants with numbers >0.5 were sent a link to NIMH's ebook on depression among college students (National Institute of Mental Health, 2017), after completion of online baseline questionnaires. Because the randomization allocation occurred algorithmically, allocation concealment was in place. However, the condition to which each participant was allocated was not masked for the service providers (Woebot Labs). After approximately 2 weeks (T2), participants were contacted again to complete a second set of questionnaires online. Participants were offered a prorated incentive of US $10 per completed assessment (US $20 for completion of both assessments).

Since this trial involved a nonclinical population of college students, it was considered exempt from registration in a public trials registry. See Multimedia Ap-

pendix 1 for the study's CONSORT-EHEALTH checklist (Eysenbach & CONSORT-EHEALTH Group, 2011).

### Woebot

Woebot is an automated conversational agent designed to deliver CBT in the format of brief, daily conversations and mood tracking. Woebot is used within an instant messenger app that is platform agnostic and can be used either on a desktop or mobile device. Each interaction begins with a general inquiry about context (eg, "What's going on in your world right now?"), and mood (eg, "How are you feeling?") with responses provided as word or emoji images to represent affect in that moment. After gathering mood data, participants are presented with core concepts related to CBT by link to short video, or by way of short "word games" designed to facilitate teaching participants about cognitive distortions. The first day included an "onboarding" process that introduced the bot, adding that while the bot may seem like a person, it is closer to a "choose your own adventure self-help book" and therefore not fully capable of understanding what the needs of the user may be. The bot also briefly explained CBT and notified the user that while a psychologist was "keeping an eye on things" (ie, monitoring), this was not happening in real time and thus the service should not be used as a replacement for therapy. In addition, participants were encouraged to call 911 for emergencies.

The bot's conversational style was modeled on human clinical decision making and the dynamics of social discourse. Psychoeducational content was adapted from self-help for CBT (Burns, 1980, 2006; Towery, 2016). Aside from CBT content, the bot was created to include the following therapeutic process-oriented features:

- **Empathic responses:** The bot replied in an empathic way appropriate to the participants' inputted mood.
- **Tailoring:** Specific content is sent to individuals depending on mood state. For example, a participant indicating that they feel anxious is offered in-vivo assistance with the anxious event.
- **Goal setting:** The conversational agent asked participants if they had a personal goal that they hoped to achieve over the 2-week period.
- **Accountability:** To facilitate a sense of accountability, the bot set expectations of regular check-ins and followed up on earlier activities, for example, on the status of the stated goal.
- **Motivation and engagement:** To engage the individual in daily monitoring, the bot sent one personalized message every day or every other day to initiate a conversation (ie, prompting).
- **Reflection:** The bot also provided weekly charts depicting each participant's mood over time. Each graph was sent with a brief description of the data to facilitate reflection.

**THE BOT'S MESSAGES QUOTED IN THE METHODS**

CHECK-IN · CONTEXT
What's going on in your world right now?

CHECK-IN · MOOD
How are you feeling?

EMPATHIC RESPONSE · ENDORSED LONELINESS
I'm so sorry you're feeling lonely. I guess we all feel a little lonely sometimes

EMPATHIC RESPONSE · EXCITEMENT
Yay, always good to hear that!

REFLECTION · WEEKLY MOOD CHART
Overall, your mood has been fairly steady, though you tend to become tired after periods of anxiety. It looks like Tuesday was your best day.

### Information Control Condition

In the information control condition, participants were directed to the NIMH resources section and specifically, a free publication entitled "Depression in College Students" (National Institute of Mental Health, 2017).

### Measures

#### *The Patient Health Questionnaire-9*

The Patient Health Questionnaire (PHQ-9) (Kroenke et al., 2001) is a 9-item, self-report questionnaire that assesses the frequency and severity of depressive symptomatology within the previous 2 weeks. Each of the 9 items is based on the Diagnostic and Statistical Manual of Mental Disorders, 4th edition (DSM-IV) criteria for major depressive disorder and can be scored on a 0 (not at all) to 3 (nearly every day) scale.

#### *Generalized Anxiety Disorder-7*

The Generalized Anxiety Disorder 7-item scale (GAD-7) (Spitzer et al., 2006) is a valid, brief self-report tool to assess the frequency and severity of anxious thoughts and behaviors over the past 2 weeks. Based on the DSM-IV diagnostic criteria for GAD, the scores of all 7 items range from 0 (not at all) to 3 (nearly every day). Therefore, the total score ranges from 0-21.

#### *Positive and Negative Affect Schedule*

The Positive and Negative Affect Schedule (PANAS) (Watson et al., 1988) is a 20-item self-report measure of current positive and negative affect. Items are scored on a 1 (very slightly or not at all) to 5 (extremely) scale, with higher scores representing higher affect. Positive and negative affect are summed independent of each other with possible scores from 10-50.

### Statistical Analysis

Statistical power calculations using analysis of covariance (ANCOVA) revealed that a sample size of 70 would have sufficient (80%) power to detect a moderate-large effect size (Cohen *d*=0.4) for depression, reported by a meta-analysis of Internet-delivered treatments for adult depression and anxiety (Andersson & Cuijpers, 2009), with alpha at 5%.

To determine whether any significant differences between groups existed at baseline, independent *t* tests were conducted on continuous baseline variables (eg, age, PHQ-9, GAD-7, and PANAS), and chi-square analyses were conducted on categorical or nominal variables (gender, race, ethnicity). Univariate effects of group membership on T2 outcomes were examined using between-subjects ANCOVA adjusting for baseline measures. Cohen *d* effect sizes were calculated to examine the magnitude of between-group differences. All subjects were included in intention-to-treat (ITT) analyses. Prior to conducting these analyses, the multiple imputation procedure in SPSS v. 23 was used to handle missing data assumed to be missing at random. As secondary subgroup analyses, we conducted completer analyses using 2x2 repeated measures analysis of variance (ANOVA) to explore main and interaction effects.

### Ethics and Informed Consent

The study was reviewed and approved by Stanford School of Medicine's Institutional Review Board. Participants indicated their consent to the terms of the study via checkbox on an information sheet. As additional safety measures, participants in the Woebot group who denoted long-standing depression, suicidality, or self-harm were automatically provided with helpline numbers and a crisis text line number, and were encouraged to call 911 in emergencies.

## Results

**Figure 1** shows the participant flow throughout the study. A total of 204 registrations were received between January 31 and February 20, 2017, and all registrants were asked to confirm their interest by return email. A total of 115 responded to this email, though 45 of these were deemed bot-generated (eg, email addresses with unusual almost identical formats and identical responses) and were deemed ineligible. The resultant sample of N=70 were randomized via computer algorithm to receive either a direct link to begin chatting with Woebot (n=34) in an instant messenger app, or NIMH's ebook on depression among college students (National Institute of Mental Health, 2017) (n=36), after completion of online questionnaires at baseline.

### Attrition

Of the randomized participants, 83% (58/70) went on to provide partial or complete data at T2 representing an overall attrition rate of 17%. Attrition was not equal between the arms and was greater among the information control group (31% vs 9%; $\chi^2_1$=5.16; *P*=.023). However, independent *t* tests and chi-square analyses failed to detect evidence of significant differences at baseline between those who dropped out of the study versus those who did not on age ($t_{68}$=1.18; *P*=.24); GAD-7 ($t_{68}$=1.28; *P*=.89); PHQ-9 ($t_{68}$=.63; *P*=.59); PANAS positive ($t_{68}$=.79; *P*=.43) and negative ($t_{68}$=.02; *P*=.98) affect scores; or on gender ($\chi^2_1$=1.75; *P*=.18) or ethnicity ($\chi^2_1$=.066; *P*=.79).

### Participant Demographics

**Table 1** shows the demographic information and baseline scores on clinical variables for those with data from the entire sample (N=58). Participants were an average of 22.2 years old (SD 2.33) and over two-thirds female. Participants were mostly non-Hispanic (93%, 54/58), 79% Caucasian (46/58), with 7% (4/58) Asian, 9% (5/58) more than one race, 2% (2/58) African American, and 2% (2/58) Native American/Alaskan Native.

In terms of baseline characteristics, nearly half (46%, 32/69) of the sample was in the moderately-severe or severe range of depression at baseline as measured by the PHQ-9, while three-quarters (74%, 52/70) were in the severe range for anxiety as measured by the GAD-7.

### Preliminary Efficacy

**Table 2** shows the results of the primary ITT analyses conducted on the entire sample. Univariate ANCOVA revealed a significant treatment effect on depression revealing that those in the Woebot group significantly reduced PHQ-9 score while those in the information control group did not ($F_{1,48}$=6.03; *P*=.017). This represented a moderate between-groups effect size (*d*=0.44). This effect is robust after Bonferroni correction for multiple comparisons (*P*=.04). No other significant between-group differences were observed on anxiety or affect.





**Figure 1.** Participant flow through the trial (CONSORT diagram).

Redrawn from Figure 1 of the original; the Analysis row follows the Statistical Analysis section. The text reports 58 participants with T2 data.

**Table 1.** Demographic and clinical variables of participants at baseline.

| | Information control | Woebot |
|---|---|---|
| **Scale, mean (SD)** | | |
| Depression (PHQ-9) | 13.25 (5.17) | 14.30 (6.65) |
| Anxiety (GAD-7) | 19.02 (4.27) | 18.05 (5.89) |
| Positive affect | 26.19 (8.37) | 25.54 (9.58) |
| Negative affect | 28.74 (8.92) | 24.87 (8.13) |
| Age, mean (SD) | 21.83 (2.24) | 22.58 (2.38) |
| **Gender, n (%)** | | |
| Male | 4 (7) | 7 (21) |
| Female | 20 (55) | 27 (79) |
| **Ethnicity, n (%)** | | |
| Latino/Hispanic | 2 (8) | 2 (6) |
| Non-Latino/Hispanic | 22 (92) | 32 (94) |
| Caucasian | 18 (75) | 28 (82) |
| Non-Caucasian | 6 (25) | 6 (18) |

Participants with data at T2 (N=58), as printed in the original Table 1.

### Completer Analysis

As a secondary analysis, to explore whether any main effects existed, 2x2 repeated measures ANOVAs were conducted on the primary outcome variables (with the exception of PHQ-9) among completers only. A significant main effect was observed on GAD-7 ($F_{1,54}$=9.24; $P$=.004) suggesting that completers experienced a significant reduction in symptoms of anxiety between baseline and T2, regardless of the group to which they were assigned with a within-subjects effect size of $d$=0.37. No main effects were observed for positive ($F_{1,50}$=.001; $P$=.951; $d$=0.21) or negative affect ($F_{1,50}$=.06; $P$=.80; $d$=0.003) as measured by the PANAS.

### Use and Acceptability

Participants in the Woebot condition checked in with the bot (defined as at least providing context and mood information) an average of 12.14 times (SD 2.23; median 12; range 8-18) over the 2-week period, with almost all check-ins occurring on unique days.

While ratings indicated that both conditions were acceptable (above 3/5), participants in the Woebot condition reported significantly higher levels of satisfaction both overall (4.3 versus 3.4; $t_{48}$=3.99; $P$<.001) and with content (4.0 versus 3.4; $t_{48}$=2.30; $P$=.02), and they reported a significantly greater amount of emotional awareness as a result of using the bot (3.3 versus 2.7; $t_{47.06}$=2.38; $P$=.021) than the information control group.

**Table 2.** Results of ITT analysis of entire sample on primary outcomes in the study at T2.

| | Information-only control | | Woebot | | *F* | *P* | *d*[c] |
|---|---|---|---|---|---|---|---|
| | T2[a] | 95% CI[b] | T2[a] | 95% CI[b] | | | |
| PHQ-9 | 13.67 (.81) | 12.07-15.27 | 11.14 (0.71) | 9.74-12.32 | 6.03 | .017 | 0.44 |
| GAD-7 | 16.84 (.67) | 15.52-18.56 | 17.35 (0.60) | 16.16-18.13 | 0.38 | .581 | 0.14 |
| PANAS positive affect | 26.02 (1.45) | 23.17-28.86 | 26.88 (1.29) | 24.35-29.41 | 0.17 | .707 | 0.02 |
| PANAS negative affect | 27.53 (1.42) | 24.73-30.32 | 25.98 (1.24) | 23.54-28.42 | 0.91 | .912 | 0.344 |

[a]Baseline=pooled mean (standard error). [b]95% confidence interval. [c]Cohen *d* shown for between-subjects effects using means and standard errors at Time 2.

### Qualitative Results

**Figure 2** shows a thematic map of participants' responses to the question "What was the best thing about your experience using Woebot?" Two major themes emerged in respect to this question: process and content. In the process theme, the subthemes that emerged were accountability from daily check-ins (noted by 9 participants); the empathy that the bot showed, or other factors relating to his "personality" (n=7); and the learning that the bot facilitated (n=12), which in turn was divided into further subthemes of emotional insight (n=5), general insight (n=5), and insights about cognitions (n=2).

**Figure 3** illustrates a thematic map of participants' responses to the question: "What was the worst thing about your experience with Woebot?" Three themes emerged: process violations (n=15), technical problems (n=8), and problems with content (n=8). By far the most common subtheme to emerge among the process violations related to the limitations in natural conversation such as the bot not being able to understand some responses or getting confused when unexpected answers were provided by participants (n=10), and 2 individuals noted that the conversations could get repetitive.

## Discussion

### Principal Results

To our knowledge this is the first randomized trial of a nonembodied text-based conversational agent designed for therapeutic use. The objective of the study was to explore whether a fully automated conversational agent based on CBT principals could deliver a therapeutic experience to college students over a 2-week period.

The study confirmed that after 2 weeks, those in the Woebot group experienced a significant reduction in depression, thus our hypothesis was partially supported. Woebot was associated with a high level of engagement with most individuals using the bot nearly every day and was generally viewed more favorably than the information-only comparison.

### Limitations

There are several methodological weaknesses that limit the generalizability of the findings. As a feasibility study,





**Figure 2.** Best features of the Woebot experience: themes of the answers to "What was the best thing about your experience using Woebot?"

Number of participants per theme. Redrawn as bars from the thematic map in Figure 3 of the original; numbers as printed there.





**Figure 3.** Least favored experiences: themes of the answers to "What was the worst thing about your experience of using Woebot?"

Number of participants per theme. Redrawn as bars from the thematic map in Figure 4 of the original; numbers as printed there.

we recruited a limited number of participants to receive a relatively short intervention, and no follow-up data were available to assess whether gains were sustained. The small number of participants meant that a formal mediator analysis was not possible, thus we cannot formally test a theorized relationship between engagement and outcome in this context of conversational agents. The study should be replicated with more participants, a longer dose, and a follow-up period to investigate if findings persist. Nonetheless, the relatively strong comparison group can be viewed as a strength of the study. Indeed, the relative strength of the control group was illustrated by the fact that individuals providing data in that group saw a similar reduction in anxiety as those who received Woebot, which supports the literature that suggests minimal passive psychoeducation alone can reduce symptoms of psychological distress (Donker et al., 2009). Nonetheless, the choice of control group was somewhat limiting for two main reasons. First, it may have contributed to the high attrition rate since an ebook is not designed for multiple or recurring sessions. It also did not introduce any CBT-specific material, thus it was not possible to evaluate whether the conversational delivery mediated symptom reduction, rather than the CBT content that the bot delivered.

Finally, the study was conducted in a New York area university community population and since we did not formally assess digital divide factors such as socioeconomic status, findings may be limited in their generalizability.

### Conclusions

While results should be viewed with some caution and the findings need to be replicated, this study nonetheless demonstrates that a text-based conversational agent designed to mirror therapeutic process has the potential to offer an alternative and engaging method of delivering CBT for some 10 million college students in the United States who experience debilitating anxiety and depression.

## CONFLICTS OF INTEREST

The second author (AMD) is the founder of a commercial entity Woebot Labs Inc. (formerly, the Life Ninja Project) that created the intervention (Woebot) that is the subject of this trial and therefore has financial interest in that company. Woebot Labs Inc. covered the cost of participant incentives, though Standford [sic] made the payments.

## ABBREVIATIONS

**ANCOVA** analysis of covariance
**ANOVA** analysis of variance
**DSM-IV** Diagnostic and Statistical Manual of Mental Disorders, 4th edition
**GAD-7** Generalized Anxiety Disorders scale
**ITT** intention to treat
**NIMH** National Institute of Mental Health
**PANAS** Positive and Negative Affect Scale
**PHQ-9** Patient Health Questionnaire scale
**T2** time 2

## REFERENCES

Andersson, G., & Cuijpers, P. (2009). Internet-based and other computerized psychological treatments for adult depression: a meta-analysis. *Cognitive Behaviour Therapy*, 38(4), 196–205. https://doi.org/10.1080/16506070903318960

Bakker, D., Kazantzis, N., Rickwood, D., & Rickard, N. (2016). Mental Health Smartphone Apps: Review and Evidence-Based Recommendations for Future Developments. *JMIR Mental Health*, 3(1), e7. https://doi.org/10.2196/mental.4984

Barak, A., Hen, L., Boniel-Nissim, M., & Shapira, N. (2008). A Comprehensive Review and a Meta-Analysis of the Effectiveness of Internet-Based Psychotherapeutic Interventions. *Journal of Technology in Human Services*, 26(2–4), 109–160. https://doi.org/10.1080/15228830802094429

Bickmore, T., Gruber, A., & Picard, R. (2005). Establishing the computer-patient working alliance in automated health behavior change interventions. *Patient Education and Counseling*, 59(1), 21–30. https://doi.org/10.1016/j.pec.2004.09.008

Burns, D. D. (1980). *Feeling good: The new mood therapy*. Harper Collins.

Burns, D. D. (2006). *When panic attacks*. Harmony.

Donker, T., Griffiths, K. M., Cuijpers, P., & Christensen, H. (2009). Psychoeducation for depression, anxiety and psychological distress: a meta-analysis. *BMC Medicine*, 7, 79. https://doi.org/10.1186/1741-7015-7-79

Donkin, L., Hickie, I. B., Christensen, H., Naismith, S. L., Neal, B., Cockayne, N. L., & Glozier, N. (2013). Rethinking the dose-response relationship between usage and outcome in an online intervention for depression: randomized controlled trial. *Journal of Medical Internet Research*, 15(10), e231. https://doi.org/10.2196/jmir.2771

Eysenbach, G. & CONSORT-EHEALTH Group. (2011). CONSORT-EHEALTH: Improving and Standardizing Evaluation Reports of Web-based and Mobile Health Interventions. *Journal of Medical Internet Research*, 13(4), e126. https://doi.org/10.2196/jmir.1923

Hunt, J., & Eisenberg, D. (2010). Mental health problems and help-seeking behavior among college students. *Journal of Adolescent Health*, 46(1), 3–10. https://doi.org/10.1016/j.jadohealth.2009.08.008

Kessler, R. C., Amminger, G. P., Aguilar-Gaxiola, S., Alonso, J., Lee, S., & Üstün, T. B. (2007). Age of onset of mental disorders: a review of recent literature. *Current Opinion*

*in Psychiatry*, 20(4), 359–364. https://doi.org/10.1097/YCO.0b013e32816ebc8c

Kroenke, K., Spitzer, R. L., & Williams, J. B. W. (2001). The PHQ-9: validity of a brief depression severity measure. *Journal of General Internal Medicine*, 16(9), 606–613. https://doi.org/10.1046/j.1525-1497.2001.016009606.x

Miner, A. S., Milstein, A., Schueller, S., Hegde, R., Mangurian, C., & Linos, E. (2016). Smartphone-Based Conversational Agents and Responses to Questions About Mental Health, Interpersonal Violence, and Physical Health. *JAMA Internal Medicine*, 176(5), 619–625. https://doi.org/10.1001/jamainternmed.2016.0400

National Institute of Mental Health. (2017). *Depression in college students*. https://www.nimh.nih.gov/health/publications/depression-and-college-students/index.shtml

Spek, V., Cuijpers, P., Nyklícek, I., Riper, H., Keyzer, J., & Pop, V. (2007). Internet-based cognitive behaviour therapy for symptoms of depression and anxiety: a meta-analysis. *Psychological Medicine*, 37(3), 319–328. https://doi.org/10.1017/S0033291706008944

Spitzer, R. L., Kroenke, K., Williams, J. B. W., & Löwe, B. (2006). A brief measure for assessing generalized anxiety disorder: the GAD-7. *Archives of Internal Medicine*, 166(10), 1092–1097. https://doi.org/10.1001/archinte.166.10.1092

Torous, J., Chan, S. R., Yee-Marie Tan, S., Behrens, J., Mathew, I., Conrad, E. J., Hinton, L., Yellowlees, P., & Keshavan, M. (2014). Patient Smartphone Ownership and Interest in Mobile Apps to Monitor Symptoms of Mental Health Conditions: A Survey in Four Geographically Distinct Psychiatric Clinics. *JMIR Mental Health*, 1(1), e5. https://doi.org/10.2196/mental.4004

Towery, J. (2016). *The anti-depressant book: A practical guide for teens and young adults to overcome depression and stay healthy*. Jacob Towery.

Watson, D., Clark, L. A., & Tellegen, A. (1988). Development and validation of brief measures of positive and negative affect: the PANAS scales. *Journal of Personality and Social Psychology*, 54(6), 1063–1070. https://doi.org/10.1037//0022-3514.54.6.1063

Zivin, K., Eisenberg, D., Gollust, S. E., & Golberstein, E. (2009). Persistence of mental health problems and needs in a college student population. *Journal of Affective Disorders*, 117(3), 180–185. https://doi.org/10.1016/j.jad.2009.01.001

## ABOUT THIS EDITION

Abridged from Fitzpatrick KK, Darcy A, Vierhile M. Delivering Cognitive Behavior Therapy to Young Adults With Symptoms of Depression and Anxiety Using a Fully Automated Conversational Agent (Woebot): A Randomized Controlled Trial. JMIR Mental Health 2017;4(2):e19. doi:10.2196/mental.7785. © Kathleen Kara Fitzpatrick, Alison Darcy, Molly Vierhile. Originally published in JMIR Mental Health (http://mental.jmir.org), 06.06.2017. This is an open-access article distributed under the terms of the Creative Commons Attribution License (https://creativecommons.org/licenses/by/4.0/), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work, first published in JMIR Mental Health, is properly cited.

Changes: sections, paragraphs and sentences cut; the original Figure 2 and Multimedia Appendix 1 left out; citations converted from numbered to APA author–date; Figure 1 redrawn as a CONSORT diagram and Figures 3 and 4 redrawn as bar charts (Figures 2 and 3 here); the bot's quoted messages gathered in a box. Set in Lora and Nunito Sans (SIL OFL) for the Postext Cookbook.