


Geometric combinatorics of paths and deformations of convex polytopes

Germain Poullot




- Mon site : “Germain Poullot” dans Google
- Onglet “Petit jeu”
- Suivez les indications, mot de passe : 0000
- ⇒ Amusez-vous !

Only in French, sorry...

Merci Guillaume !!!

Geometric combinatorics of paths and deformations of convex polytopes

Germain Poullot





Institut de Mathématiques
de
Jussieu-Paris Rive Gauche

Directeurs / Advisors

Arnaud PADROL

Vincent PILAUD

Jury

Fu LIU

Jesús DE LOERA

Martina JUHNKE-KUBITZKE

Lionel POURNIN

Frédéric MEUNIER

Vic REINER

1 What is “Combinatorics of Polytopes”?

2 Generalized permutohedra

- Deformations
- Submodular Cone
- Ongoing work


3 Max-slope Pivot Polytopes

- Max-slope pivot rule
- Poset of slopes
- Pivot rule polytope of products of simplices

What is “Combinatorics of Polytopes”?



Definition

Polytope: convex hull of finitely many points in \mathbb{R}^n





Definition

Polytope: convex hull of finitely many points in \mathbb{R}^n
bounded intersection of finitely many half-spaces in \mathbb{R}^n





Definition

Polytope: convex hull of finitely many points in \mathbb{R}^n
bounded intersection of finitely many half-spaces in \mathbb{R}^n



Definition



Polytope: convex hull of finitely many points in \mathbb{R}^n
bounded intersection of finitely many half-spaces in \mathbb{R}^n



Polytopes

Definition



Polytope: convex hull of finitely many points in \mathbb{R}^n
bounded intersection of finitely many half-spaces in \mathbb{R}^n



Polytopes

Definition

Polytope: convex hull of finitely many points in \mathbb{R}^n
bounded intersection of finitely many half-spaces in \mathbb{R}^n



Representing polytopes




Tetrahedron
Fire




Hexahedron
Earth



Octahedron
Air



Dodecahedron
the Universe



Icosahedron
Water

Representing polytopes




Tetrahedron
Fire



Hexahedron
Earth



Octahedron
Air



Dodecahedron
the Universe



Icosahedron
Water



Representing polytopes




Tetrahedron
Fire



Hexahedron
Earth




Octahedron
Air



Dodecahedron
the Universe




Icosahedron
Water



Definition

Face: $P^c := \{x \in \mathbb{R}^n ; \langle x, c \rangle = \max_{y \in P} \langle y, c \rangle\}$



P

Definition


Face: $P^c := \{x \in \mathbb{R}^n ; \langle x, c \rangle = \max_{y \in P} \langle y, c \rangle\}$



P

Definition




Face: $P^c := \{x \in \mathbb{R}^n ; \langle x, c \rangle = \max_{y \in P} \langle y, c \rangle\}$



P




Face lattice

Face lattice: poset of inclusions of faces




Face lattice

Face lattice: poset of inclusions of faces




Face lattice

Face lattice: poset of inclusions of faces




Face lattice

Face lattice: poset of inclusions of faces



Face lattice


Face lattice: poset of inclusions of faces



Definition

Normal cone of a face F : $\mathcal{N}_P(F) := \{\mathbf{c} ; P^c = F\}$

Normal fan \mathcal{N}_P : collection of $\mathcal{N}_P(F)$ for F face of P




P

\mathcal{N}_P

Definition

Normal cone of a face F : $\mathcal{N}_P(F) := \{\mathbf{c} ; P^c = F\}$

Normal fan \mathcal{N}_P : collection of $\mathcal{N}_P(F)$ for F face of P



P




\mathcal{N}_P


Definition

Normal cone of a face F : $\mathcal{N}_P(F) := \{\mathbf{c} ; P^c = F\}$

Normal fan \mathcal{N}_P : collection of $\mathcal{N}_P(F)$ for F face of P



P




\mathcal{N}_P


Definition

Normal cone of a face F : $\mathcal{N}_P(F) := \{\mathbf{c} ; P^c = F\}$

Normal fan \mathcal{N}_P : collection of $\mathcal{N}_P(F)$ for F face of P



P



\mathcal{N}_P

Combinatorics of Polytopes

One way: Take a polytope → combinatorial info (e.g. face lattice)

Combinatorics of Polytopes

One way: Take a polytope → combinatorial info (e.g. face lattice)

Other way: Take a combinatorial family → embed it as a polytope

Combinatorics of Polytopes

One way: Take a polytope → combinatorial info (e.g. face lattice)

Other way: Take a combinatorial family → embed it as a polytope

Example (Permutohedron)

$$\Pi_n = \text{conv} \left\{ \begin{pmatrix} \sigma(1) \\ \vdots \\ \sigma(n) \end{pmatrix} ; \sigma \text{ permutation of } \{1, \dots, n\} \right\}$$


Combinatorics of Polytopes

One way: Take a polytope → combinatorial info (e.g. face lattice)

Other way: Take a combinatorial family → embed it as a polytope

Example (Permutahedron)

$$\Pi_n = \text{conv} \left\{ \begin{pmatrix} \sigma(1) \\ \vdots \\ \sigma(n) \end{pmatrix} ; \sigma \text{ permutation of } \{1, \dots, n\} \right\}$$




Combinatorics of Polytopes

One way: Take a polytope → combinatorial info (e.g. face lattice)

Other way: Take a combinatorial family → embed it as a polytope

Example (Permutahedron)

$$\Pi_n = \text{conv} \left\{ \begin{pmatrix} \sigma(1) \\ \vdots \\ \sigma(n) \end{pmatrix} ; \sigma \text{ permutation of } \{1, \dots, n\} \right\}$$




Combinatorics of Polytopes

One way: Take a polytope → combinatorial info (e.g. face lattice)

Other way: Take a combinatorial family → embed it as a polytope

Example (Permutahedron)

$$\Pi_n = \text{conv} \left\{ \begin{pmatrix} \sigma(1) \\ \vdots \\ \sigma(n) \end{pmatrix} ; \sigma \text{ permutation of } \{1, \dots, n\} \right\}$$




Combinatorics of Polytopes

One way: Take a polytope → combinatorial info (e.g. face lattice)

Other way: Take a combinatorial family → embed it as a polytope

Example (Permutahedron)

$$\Pi_n = \text{conv} \left\{ \begin{pmatrix} \sigma(1) \\ \vdots \\ \sigma(n) \end{pmatrix} ; \sigma \text{ permutation of } \{1, \dots, n\} \right\}$$





Combinatorics of Polytopes

One way: Take a polytope → combinatorial info (e.g. face lattice)

Other way: Take a combinatorial family → embed it as a polytope

Example (Permutahedron)

$$\Pi_n = \text{conv} \left\{ \begin{pmatrix} \sigma(1) \\ \vdots \\ \sigma(n) \end{pmatrix} ; \sigma \text{ permutation of } \{1, \dots, n\} \right\}$$




Generalized permutohedra

Coarsening: Choose maximal cones and merge them

Definition




Q is a *deformation* of P iff \mathcal{N}_Q coarsens \mathcal{N}_P .



Coarsening: Choose maximal cones and merge them

Definition




Q is a *deformation* of P iff \mathcal{N}_Q coarsens \mathcal{N}_P .




Coarsening: Choose maximal cones and merge them

Definition

Q is a *deformation* of P iff \mathcal{N}_Q coarsens \mathcal{N}_P .



coarsens



Definition

Braid fan: arrangement of hyperplanes $H_{i,j} := \{\mathbf{x} ; x_i = x_j\}$

Braid fan



Definition

Braid fan: arrangement of hyperplanes $H_{i,j} := \{\mathbf{x} ; x_i = x_j\}$

Definition

Generalized permutohedron: deformation of Π_n

i.e. P generalized permutohedron iff \mathcal{N}_P coarsens braid fan



Braid fan



Definition

Braid fan: arrangement of hyperplanes $H_{i,j} := \{\mathbf{x} ; x_i = x_j\}$

Definition


Generalized permutohedron: deformation of Π_n

i.e. P generalized permutohedron iff \mathcal{N}_P coarsens braid fan



$\mathcal{P}(P)$: all the posets associated to faces of P

Deformations of Π_4



Permutahedron Π_4

Sequence of deformations of Π_4

Cone of deformations

Minkowski sum: $P + Q = \{p + q ; p \in P, q \in Q\}$

Theorem

If Q, R deformations of P , then: λQ deform. of P
 $Q + R$ deform. of P

Definition

Deformation cone: $\mathbb{DC}(P) := \{Q ; Q \text{ deformation of } P\}$ is a cone.

Cone of deformations


Minkowski sum: $P + Q = \{p + q ; p \in P, q \in Q\}$

Theorem

If Q, R deformations of P , then: λQ deform. of P
 $Q + R$ deform. of P

Definition

Deformation cone: $\mathbb{DC}(P) := \{Q ; Q \text{ deformation of } P\}$ is a cone.



Cone of deformations


Minkowski sum: $P + Q = \{p + q ; p \in P, q \in Q\}$

Theorem

If Q, R deformations of P , then: λQ deform. of P
 $Q + R$ deform. of P

Definition

Deformation cone: $\mathbb{DC}(P) := \{Q ; Q \text{ deformation of } P\}$ is a cone.



Cone of deformations


Minkowski sum: $P + Q = \{p + q ; p \in P, q \in Q\}$

Theorem

If Q, R deformations of P , then: $\begin{aligned} &\text{for all } \lambda > 0, \lambda Q \text{ deform. of } P \\ &Q + R \text{ deform. of } P \end{aligned}$

Definition

Deformation cone: $\mathbb{DC}(P) := \{Q ; Q \text{ deformation of } P\}$ is a cone.



Cone of deformations


Minkowski sum: $P + Q = \{p + q ; p \in P, q \in Q\}$

Theorem

If Q, R deformations of P , then: $\begin{aligned} &\text{for all } \lambda > 0, \lambda Q \text{ deform. of } P \\ &Q + R \text{ deform. of } P \end{aligned}$

Definition

Deformation cone: $\mathbb{DC}(P) := \{Q ; Q \text{ deformation of } P\}$ is a cone.



Cone of deformations


Minkowski sum: $P + Q = \{p + q ; p \in P, q \in Q\}$

Theorem

If Q, R deformations of P , then: $\begin{aligned} &\text{for all } \lambda > 0, \lambda Q \text{ deform. of } P \\ &Q + R \text{ deform. of } P \end{aligned}$

Definition

Deformation cone: $\mathbb{DC}(P) := \{Q ; Q \text{ deformation of } P\}$ is a cone.



Cone of deformations


Minkowski sum: $P + Q = \{p + q ; p \in P, q \in Q\}$

Theorem

If Q, R deformations of P , then: $\begin{aligned} &\text{for all } \lambda > 0, \lambda Q \text{ deform. of } P \\ &Q + R \text{ deform. of } P \end{aligned}$

Definition

Deformation cone: $\mathbb{DC}(P) := \{Q ; Q \text{ deformation of } P\}$ is a cone.



Cone of deformations


Minkowski sum: $P + Q = \{p + q ; p \in P, q \in Q\}$

Theorem

If Q, R deformations of P , then: $\begin{aligned} &\text{for all } \lambda > 0, \lambda Q \text{ deform. of } P \\ &Q + R \text{ deform. of } P \end{aligned}$

Definition

Deformation cone: $\mathbb{DC}(P) := \{Q ; Q \text{ deformation of } P\}$ is a cone.



Parametrization:

height vector:

$$\mathbf{h} = (h_r)_{r \text{ rays}}$$

Cone of deformations


Minkowski sum: $P + Q = \{p + q ; p \in P, q \in Q\}$

Theorem

If Q, R deformations of P , then: $\text{for all } \lambda > 0, \lambda Q \text{ deform. of } P$
 $Q + R \text{ deform. of } P$

Definition

Deformation cone: $\mathbb{DC}(P) := \{Q ; Q \text{ deformation of } P\}$ is a cone.



Parametrization:

height vector:

$$\mathbf{h} = (h_r)_{r \text{ rays}}$$

Wall-crossing inequalities:

linear ineq on \mathbf{h}

Cone of deformations


Minkowski sum: $P + Q = \{p + q ; p \in P, q \in Q\}$

Theorem

If Q, R deformations of P , then: $\text{for all } \lambda > 0, \lambda Q \text{ deform. of } P$
 $Q + R \text{ deform. of } P$

Definition

Deformation cone: $\mathbb{DC}(P) := \{Q ; Q \text{ deformation of } P\}$ is a cone.



Parametrization:

height vector:

$$\mathbf{h} = (h_r)_{r \text{ rays}}$$

Wall-crossing inequalities:

linear ineq on \mathbf{h}

Cone of deformations


Minkowski sum: $P + Q = \{p + q ; p \in P, q \in Q\}$

Theorem

If Q, R deformations of P , then: $\text{for all } \lambda > 0, \lambda Q \text{ deform. of } P$
 $Q + R \text{ deform. of } P$

Definition

Deformation cone: $\mathbb{DC}(P) := \{Q ; Q \text{ deformation of } P\}$ is a cone.



Parametrization:

height vector:

$$\mathbf{h} = (h_r)_{r \text{ rays}}$$

Wall-crossing inequalities:

linear ineq on \mathbf{h}

Cone of deformations


Minkowski sum: $P + Q = \{p + q ; p \in P, q \in Q\}$

Theorem

If Q, R deformations of P , then: $for\ all\ \lambda > 0,\ \lambda Q\ deform.\ of\ P$
 $Q + R\ deform.\ of\ P$

Definition

Deformation cone: $\mathbb{DC}(P) := \{Q ; Q\ deformation\ of\ P\}$ is a cone.



Parametrization:

height vector:

$$\mathbf{h} = (h_r)_{r \text{ rays}}$$

Wall-crossing inequalities:


linear ineq on \mathbf{h}

Definition

Submodular cone: deformation cone of the permutohedron Π_n

	$\mathbb{DC}(\Pi_n)$
Dim (no lineal)	$2^n - n - 1$
# facets	$\binom{n}{2} 2^{n-2}$
# rays	unknown!

Submodular Cone for Π_3



Definition

Submodular cone: deformation cone of the permutohedron Π_n

	$\mathbb{DC}(\Pi_n)$
Dim (no lineal)	$2^n - n - 1$
# facets	$\binom{n}{2} 2^{n-2}$
# rays	unknown!

Submodular Cone's faces

Definition

Submodular cone: deformation cone of the permutohedron Π_n

Theorem (Faces of $\mathbb{DC}(\mathbf{P})$)

If \mathbf{Q} deformation of \mathbf{P} , then $\mathbb{DC}(\mathbf{Q})$ is a face of $\mathbb{DC}(\mathbf{P})$.

	$\mathbb{DC}(\Pi_n)$
Dim (no lineal)	$2^n - n - 1$
# facets	$\binom{n}{2} 2^{n-2}$
# rays	unknown!

Submodular Cone's faces

Definition

Submodular cone: deformation cone of the permutohedron Π_n

Theorem (Faces of $\mathbb{DC}(P)$)

If Q deformation of P , then $\mathbb{DC}(Q)$ is a face of $\mathbb{DC}(P)$.

	$\mathbb{DC}(\Pi_n)$	$\mathbb{DC}(\text{Asso}_n)$
Dim (no lineal)	$2^n - n - 1$	$\binom{n}{2}$
# facets	$\binom{n}{2} 2^{n-2}$	$\binom{n}{2}$
# rays	unknown!	$\binom{n}{2}$ is simplicial!

Submodular Cone's faces

Definition

Submodular cone: deformation cone of the permutohedron Π_n

Theorem (Faces of $\mathbb{DC}(P)$)

If Q deformation of P , then $\mathbb{DC}(Q)$ is a face of $\mathbb{DC}(P)$.

	$\mathbb{DC}(\Pi_n)$	$\mathbb{DC}(\text{Asso}_n)$	$\mathbb{DC}(Z_G)$	$\mathbb{DC}(N_B)$
Dim (no lineal)	$2^n - n - 1$	$\binom{n}{2}$	N	N
# facets	$\binom{n}{2} 2^{n-2}$	$\binom{n}{2}$	E	E
# rays	unknown!	$\binom{n}{2}$ is simplicial!	X	X


My contribution - Graphical Zonotopes

$G = (V, E)$ a graph, $n = |V|$

Definition

Graphical zonotope $Z_G := \sum_{(i,j) \in E} [\mathbf{e}_i, \mathbf{e}_j]$

Z_G deformation of $\Pi_n \implies \mathbb{DC}(Z_G)$ is a face of $\mathbb{DC}(\Pi_n)$



The diagram illustrates the construction of a graphical zonotope Z_G from four vectors Δ_{12} , Δ_{23} , Δ_{34} , and Δ_{14} . The vectors are represented by arrows originating from vertex 1. They are combined using vector addition (+) to form the zonotope Z_G , which is shown as a hexagon.

$$\Delta_{12} + \Delta_{23} + \Delta_{34} + \Delta_{14} = Z_G$$

Theorem (Padrol, Pilaud, P., '23)

Explicit facet-description of $\mathbb{DC}(Z_G)$

Theorem (Padrol, Pilaud, P., '23)

Explicit facet-description of $\mathbb{DC}(Z_G)$

Corollary

$\dim \mathbb{DC}(Z_G) = \# \text{ cliques of } G$

$\# \text{ facets of } \mathbb{DC}(Z_G) = \sum_{(i,j) \in E} 2^{|\{k ; (i,k), (j,k) \in E\}|}$

Theorem (Padrol, Pilaud, P., '23)

Explicit facet-description of $\mathbb{DC}(Z_G)$

Corollary

$\dim \mathbb{DC}(Z_G) = \# \text{ cliques of } G$


$\# \text{ facets of } \mathbb{DC}(Z_G) = \sum_{(i,j) \in E} 2^{|\{k ; (i,k), (j,k) \in E\}|}$

Corollary

$\mathbb{DC}(Z_G)$ simplicial iff G without triangle

NB: Recover facet-description of $\mathbb{DC}(\Pi_n)$

My contribution - Graphical Zonotopes



My contribution - Nestohedra


Definition

Building set $B \subseteq 2^{[n]}$ with: $X_{1,2} \in B, X_1 \cap X_2 \neq \emptyset \Rightarrow X_1 \cup X_2 \in B$

Definition

Nestohedron $N_B := \sum_{X \in B} \Delta_X$ where $\Delta_X = \text{conv}\{\mathbf{e}_i ; i \in X\}$

N_B deformation of $\Pi_n \implies \mathbb{DC}(N_B)$ is a face of $\mathbb{DC}(\Pi_n)$



My contribution - Nestohedra

Elementary blocks $X \in \varepsilon(B)$ iff X is not a union

Maximal block $\mu(X) := \max\{Y \in B ; Y \subsetneq X\}$

Theorem (Padrol, Pilaud, P., '23)

Explicit facet description of $\mathbb{DC}(N_B)$

My contribution - Nestohedra

Elementary blocks $X \in \varepsilon(B)$ iff X is not a union

Maximal block $\mu(X) := \max\{Y \in B ; Y \subsetneq X\}$

Theorem (Padrol, Pilaud, P., '23)

Explicit facet description of $\mathbb{DC}(N_B)$

Corollary

$\dim \mathbb{DC}(N_B) = |B| - \# \text{ singletons}$

$\# \text{ facets of } \mathbb{DC}(N_B) = |\varepsilon(B)| + \sum_{X \in B \setminus \varepsilon(B)} \binom{|\mu(X)|}{2}$

My contribution - Nestohedra

Elementary blocks $X \in \varepsilon(B)$ iff X is not a union

Maximal block $\mu(X) := \max\{Y \in B ; Y \subsetneq X\}$

Theorem (Padrol, Pilaud, P., '23)

Explicit facet description of $\mathbb{DC}(N_B)$

Corollary

$$\dim \mathbb{DC}(N_B) = |B| - \# \text{ singletons}$$


$$\# \text{ facets of } \mathbb{DC}(N_B) = |\varepsilon(B)| + \sum_{X \in B \setminus \varepsilon(B)} \binom{|\mu(X)|}{2}$$

Corollary

$\mathbb{DC}(N_B)$ simplicial iff B has no non-elementary block with 3 maximal subblocks


NB: Recover facet-description of $\mathbb{DC}(\Pi_n)$

My contribution - Nestohedra




Definition

Hypergraphic pol $P_H := \sum_{x \in H} \Delta_x$ with $H \subseteq 2^{[n]}$




Definition

Quotientopes: Minkowski sum of shard polytopes




Max-slope Pivot Polytopes

Linear optimization




Simplex method




Max-slope pivot rule

Linear optimization in dimension 2 (simplex method):




Max-slope pivot rule

Linear optimization in dimension 2 (simplex method):




Max-slope pivot rule

Linear optimization in dimension 2 (simplex method):




Max-slope pivot rule

Linear optimization in dimension 2 (simplex method):



Max-slope pivot rule


Linear optimization in dimension 2 (simplex method): **EASY !**



Convention: choose upper


Max-slope pivot rule

Optimization in higher dimension: make it 2-dimensional !




Max-slope pivot rule

Optimization in higher dimension: make it 2-dimensional !




Max-slope pivot rule

Optimization in higher dimension: make it 2-dimensional !




Max-slope pivot rule

Optimization in higher dimension: make it 2-dimensional !




Max-slope pivot rule

Optimization in higher dimension: make it 2-dimensional !



Max-slope pivot rule


Optimization in higher dimension: make it 2-dimensional !



Max-slope pivot rule: take (improving) neighbor with best slope


Max-slope pivot rule

Optimization in higher dimension: make it 2-dimensional !




Max-slope pivot rule: take (improving) neighbor with best slope



Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex




Coherent paths of the d -simplex




Coherent paths of the d -simplex



Coherent paths of the d -simplex





ω




ω
 c


Coherent paths of the d -simplex



Coherent paths of the d -simplex




ω




$\omega \uparrow$
 c


Coherent paths of the d -simplex




ω





Coherent paths of the d -simplex





ω





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex





Coherent paths of the d -simplex




Coherent paths of the d -simplex



Coherent paths of the d -simplex




Monotone path polytope



Coherent monotone path: path obtained via max-slope pivot rule

Monotone path fan: Fan with $\omega \sim \omega'$ iff same path

Monotone path polytope



Coherent monotone path: path obtained via max-slope pivot rule


Monotone path fan: Fan with $\omega \sim \omega'$ iff same path

Theorem (Billera, Sturmfels, '92)

The monotone path fan is polytopal.

Monotone path polytope $\Sigma_c(P)$: dual to monotone path fan

Monotone path polytope



Coherent monotone path: path obtained via max-slope pivot rule

Monotone path fan: Fan with $\omega \sim \omega'$ iff same path

Theorem (Billera, Sturmfels, '92)


The monotone path fan is polytopal.

Monotone path polytope $\Sigma_c(P)$: dual to monotone path fan


$$\Sigma_c(\Delta_d) = \text{Cube}_{d-1}$$

$$\Sigma_c(\text{Cube}_d) = \Pi_d$$



Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex




Coherent arborescences of the d -simplex




Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





ω





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex





Coherent arborescences of the d -simplex




Coherent arborescences of the d -simplex



Coherent arborescences of the d -simplex




(Max-slope) pivot polytope



Coherent arborescence:
arborescence obtained via
max-slope pivot rule

Pivot rule fan:
 $\omega \sim \omega'$ iff same arborescence.

(Max-slope) pivot polytope




Coherent arborescence:
arborescence obtained via
max-slope pivot rule

Pivot rule fan:
 $\omega \sim \omega'$ iff same arborescence.

Pivot rule fan refines monotone
path fan

(Max-slope) pivot polytope



Coherent arborescence:
arborescence obtained via
max-slope pivot rule

Pivot rule fan:
 $\omega \sim \omega'$ iff same arborescence.


Pivot rule fan refines monotone
path fan

Theorem (Black, De Loera, Lütjeharms, Sanyal '22)

The pivot rule fan is polytopal.

(Max-slope) pivot polytope Π_c : dual to the pivot rule fan

(Max-slope) pivot polytope



Coherent arborescence:
arborescence obtained via
max-slope pivot rule

Pivot rule fan:
 $\omega \sim \omega'$ iff same arborescence.

Pivot rule fan refines monotone
path fan

Theorem (Black, De Loera, Lütjeharms, Sanyal '22)

The pivot rule fan is polytopal.

(Max-slope) pivot polytope Π_c : dual to the pivot rule fan

$$\Sigma_c(\Delta_d) = \text{Cube}_{d-1}$$

$$\Pi_c(\Delta_d) = \text{Asso}_d$$

Link with generalized permutohedra?

Theorem (Black, De Loera, Lütjeharms, Sanyal '23+)

$$\Pi_c(\Delta_d) \simeq \text{Asso}_d \quad \text{for all } c$$

Link with generalized permutohedra?

Theorem (Black, De Loera, Lütjeharms, Sanyal '23+)

$$\Pi_c(\Delta_d) \simeq \text{Asso}_d \quad \text{for all } c$$

Theorem (Black, De Loera, Lütjeharms, Sanyal '22)

$$\Pi_c(\text{Cube}_d) \simeq \Pi_d \quad \text{for all } c$$

Link with generalized permutohedra?

Theorem (Black, De Loera, Lütjeharms, Sanyal '23+)

$$\Pi_c(\Delta_d) \simeq \text{Asso}_d \quad \text{for all } c$$

Theorem (Black, De Loera, Lütjeharms, Sanyal '22)

$$\Pi_c(\text{Cube}_d) \simeq \Pi_d \quad \text{for all } c$$

Conjecture (Pilaud, Sanyal)

$$\Pi_c(\Delta_{d_1} \times \Delta_{d_2}) \simeq \text{Asso}_{d_1} \star \text{Asso}_{d_2} \quad \star \text{shuffle product}$$

Link with generalized permutohedra?

Theorem (Black, De Loera, Lütjeharms, Sanyal '23+)

$$\Pi_c(\Delta_d) \simeq \text{Asso}_d \quad \text{for all } c$$

Theorem (Black, De Loera, Lütjeharms, Sanyal '22)

$$\Pi_c(\text{Cube}_d) \simeq \Pi_d \quad \text{for all } c$$

Conjecture (Pilaud, Sanyal)

$$\Pi_c(\Delta_{d_1} \times \Delta_{d_2}) \simeq \text{Asso}_{d_1} \star \text{Asso}_{d_2} \quad \star \text{shuffle product}$$

Challenge 1: Prove the conjecture!

Link with generalized permutohedra?

Theorem (Black, De Loera, Lütjeharms, Sanyal '23+)

$$\Pi_c(\Delta_d) \simeq \text{Asso}_d \quad \text{for all } c$$

Theorem (Black, De Loera, Lütjeharms, Sanyal '22)

$$\Pi_c(\text{Cube}_d) \simeq \Pi_d \quad \text{for all } c$$


Conjecture (Pilaud, Sanyal)

$$\Pi_c(\Delta_{d_1} \times \Delta_{d_2}) \simeq \text{Asso}_{d_1} \star \text{Asso}_{d_2} \quad \star \text{shuffle product}$$


Challenge 1: Prove the conjecture!

Challenge 2: Give **geometric** proofs!

Case of the d -simplex




Case of the d -cube



Slope comparisons

Max-slope pivot rule: take (improving) neighbor with best slope

For ω , what is important?




Slope comparisons

Max-slope pivot rule: take (improving) neighbor with best slope

For ω , what is important?

$$\text{Slopes: } \tau_\omega(u, v) = \frac{\langle \omega, u - v \rangle}{\langle c, u - v \rangle}$$



Slope comparisons


Max-slope pivot rule: take (improving) neighbor with best slope

For ω , what is important?

$$\text{Slopes: } \tau_\omega(u, v) = \frac{\langle \omega, u - v \rangle}{\langle c, u - v \rangle}$$

Slope vector:

$$\theta(\omega) = (\tau_\omega(u, v))_{uv \in E(P)}$$



Slope comparisons

Max-slope pivot rule: take (improving) neighbor with best slope


For ω , what is important?

$$\text{Slopes: } \tau_\omega(u, v) = \frac{\langle \omega, u - v \rangle}{\langle c, u - v \rangle}$$

Slope vector:

$$\theta(\omega) = (\tau_\omega(u, v))_{uv \in E(P)}$$

$\theta : \mathbb{R}^d \rightarrow \mathbb{R}^m$, injective linear map



Slope comparisons

Max-slope pivot rule: take (improving) neighbor with best slope

For ω , what is important?


$$\text{Slopes: } \tau_\omega(u, v) = \frac{\langle \omega, u - v \rangle}{\langle c, u - v \rangle}$$

Slope vector:

$$\theta(\omega) = (\tau_\omega(u, v))_{uv \in E(P)}$$

$\theta : \mathbb{R}^d \rightarrow \mathbb{R}^m$, injective linear map

What is **really** important??



Slope comparisons

Max-slope pivot rule: take (improving) neighbor with best slope

For ω , what is important?


$$\text{Slopes: } \tau_\omega(u, v) = \frac{\langle \omega, u - v \rangle}{\langle c, u - v \rangle}$$

Slope vector:

$$\theta(\omega) = (\tau_\omega(u, v))_{uv \in E(P)}$$

$\theta : \mathbb{R}^d \rightarrow \mathbb{R}^m$, injective linear map

What is **really** important?? The comparisons of slopes!



Slope comparisons

Max-slope pivot rule: take (improving) neighbor with best slope

For ω , what is important?

$$\text{Slopes: } \tau_{\omega}(u, v) = \frac{\langle \omega, u - v \rangle}{\langle c, u - v \rangle}$$


Slope vector:

$$\theta(\omega) = (\tau_{\omega}(u, v))_{uv \in E(P)}$$

$\theta : \mathbb{R}^d \rightarrow \mathbb{R}^m$, injective linear map

What is **really** important?? The comparisons of slopes!

Compare coordinates of $\theta(\omega)$



Slope comparisons

Max-slope pivot rule: take (improving) neighbor with best slope

For ω , what is important?

Slopes: $\tau_\omega(u, v) = \frac{\langle \omega, u - v \rangle}{\langle c, u - v \rangle}$

Slope vector:


$$\theta(\omega) = (\tau_\omega(u, v))_{uv \in E(P)}$$

$\theta : \mathbb{R}^d \rightarrow \mathbb{R}^m$, injective linear map




What is **really** important?? The comparisons of slopes!

Compare coordinates of $\theta(\omega)$

Where is $\theta(\omega)$ in the braid fan \mathcal{B}_m ?






Case of the d -cube






Too many edges

Case of the d -cube




Too many edges, **but**
parallelism saves us!

Case of the d -cube




Too many edges, **but**
parallelism saves us!

Geometric proof of
 $\Pi_c(\text{Cube}_d) = \Pi_d$




Case of the d -simplex




Too many edges

Case of the d -simplex




Too many edges, **but** affine independence saves us!

Case of the d -simplex



Too many edges, **but** affine independence saves us!

Geometric proof of $\Pi_c(\Delta_d) = \text{Asso}_d$



Shuffle: (E, \leq) and (F, \preceq) posets, then \trianglelefteq is a shuffle when:

group set : $E \sqcup F$

relations : all relations of \leq ; all relations of \preceq ;

for each $e \in E, f \in F$, choose if $e \trianglelefteq f$ or $e \trianglerighteq f$
(+ transitive closure)

Shuffle: (E, \leq) and (F, \preceq) posets, then \trianglelefteq is a shuffle when:

groud set : $E \sqcup F$

relations : all relations of \leq ; all relations of \preceq ;

for each $e \in E, f \in F$, choose if $e \trianglelefteq f$ or $e \trianglerighteq f$
(+ transitive closure)

Theorem (Chapoton, Pilaud '22)

P, Q : *generalized permutohedra*.

Exists polytope $P \star Q$ s.t.

$$\mathcal{P}(P \star Q) = \{ \text{all shuffles between } \leq \in \mathcal{P}(P) \text{ and } \preceq \in \mathcal{P}(Q) \}$$

Combine parallelism & affine independence:

Theorem

For $\Delta_{d_1} \times \cdots \times \Delta_{d_r}$, all (generic) direction:

$$\Pi_c(\Delta_{d_1} \times \cdots \times \Delta_{d_r}) \simeq \text{Asso}_{d_1} \star \cdots \star \text{Asso}_{d_r}$$

Product of simplices

Combine parallelism & affine independence:

Theorem

For $\Delta_{d_1} \times \cdots \times \Delta_{d_r}$, all (generic) direction:

$$\Pi_c(\Delta_{d_1} \times \cdots \times \Delta_{d_r}) \simeq \text{Asso}_{d_1} \star \cdots \star \text{Asso}_{d_r}$$

Example

- (a) $\Pi_c(\square_d) \simeq \text{Perm}_d$
- (b) $\Pi_c(\square_m \times \Delta_n) \simeq (m, n)\text{-multiplihedron}$
- (c) $\Pi_c(\Delta_m \times \Delta_n) \simeq (m, n)\text{-constrainahedron}$

Ongoing work


- 1) For which P , $\Pi_c(P)$ is a generalized permutohedron?
→ a priori, only products of simplices, but no proof
- 2) Is $\Pi_c(P)$ projection of a generalized permutohedron?
→ pivot fan sent inside $\text{Im}(\theta) \cap \mathcal{B}_m$
- 3) When $\Pi_c(P)$ and $\Pi_c(Q)$ **not** generalized permutohedra, what happen to $\Pi_c(P \times Q)$?
→ not equivalent to $\Pi_c(P) \star \Pi_c(Q)$, but "embeds" in it

What I have presented

Contents

Introduction	5
1 Preliminaries	
1.1 Partially ordered sets	11
1.2 Polytopes	12
1.2.1 Simplex	15
1.2.2 Cube	15
1.2.3 Permutohedron	15
1.2.4 Associahedron	18
1.3 Linear programming	20
2 Deformations of polytopes and generalized permutohedra	24
2.1 Deformations of polytopes	24
2.2 Deformation cones of graphical zonotopes	28
2.2.1 Graphical zonotopes	28
2.2.2 Graphical deformation cones	29
2.2.3 The facets of graphical deformation cones	33
2.2.4 Simplicial graphical deformation cones	36
2.2.5 Perspectives and open questions	37
2.3 Deformation cones of nestohedra	38
2.3.1 Deformation cones of graphical nested fans	39
2.3.2 Deformation cones of arbitrary nested fans	47
2.3.3 Simplicial deformation cones and interval building sets	61
2.3.4 Perspectives and open questions	63
3 Max-slope pivot rule polytopes	65
3.1 Max-slope pivot rule and max-slope pivot polytope	65
3.2 Max-slope pivot polytope of cyclic polytopes	70
3.2.1 Cyclic associahedra and the intrinsic degree	71
3.2.2 Realization sets and universal arborescences	74
3.2.3 Pivot polytopes of cyclic polytopes of dimension 2 and 3	86
3.2.4 Perspectives and open questions	92
3.3 Max-slope pivot polytopes of products of polytopes	95
3.3.1 Max-slope pivot polytopes of the cube and the simplex	99
3.3.2 Max-slope pivot polytope of a product of simplices	102
3.3.3 Perspectives and open questions	106
4 Fiber polytopes	108
4.1 Preliminaries on fiber polytopes	108
4.2 Monotone path polytopes of the hypersimplices	111
4.2.1 Monotone paths polytopes in general	111
4.2.2 A necessary criterion for coherent paths on $\Delta(n, k)$	119
4.2.3 Sufficiency of this criterion in the case $\Delta(n, 2)$	119
4.2.4 Counting the number of coherent monotone paths on $\Delta(n, 2)$	127
4.2.5 Perspectives and open questions	130
4.3 Fiber polytopes for the projection from $\text{Cyc}_d(t)$ to $\text{Cyc}_2(t)$	133
4.3.1 Bijection between triangulations and non-crossing arborescences	133
4.3.2 Fiber polytopes for the projection $\text{Cyc}_d(t) \xrightarrow{\sim} \text{Cyc}_2(t)$	135
4.3.3 Realization sets and universal triangulations for $\text{Cyc}_d(t) \xrightarrow{\sim} \text{Cyc}_2(t)$	138
4.3.4 Perspectives and open questions	143
A A Vandermonde-like determinant	146

Thank you!





The tool: submodular dependancies

Notations: $Sx = S \cup \{x\}$, $(f_x)_{x \subseteq [n]}$ canonical basis of $\mathbb{R}^{2^{[n]}}$

Definition

Submodular vector $n(S, u, v) = f_{Suv} - f_{Su} - f_{Sv} + f_S$
for $u, v \in S \subseteq [n]$





The tool: submodular dependancies

Notations: $Sx = S \cup \{x\}$, $(f_x)_{x \subseteq [n]}$ canonical basis of $\mathbb{R}^{2^{[n]}}$

Definition

Submodular vector $n(S, u, v) = f_{Suv} - f_{Su} - f_{Sv} + f_S$
for $u, v \in S \subseteq [n]$



The tool: submodular dependancies


Lemma (Submodular normal)

$\mathbf{n}(S, u, v)$ are the facet's normals of $\mathbb{DC}(\Pi_n)$

Lemma (Cubic relation)

$u, v, x \notin S \subseteq [n]$

$$\mathbf{n}(S_{uvx}, u, v) + \mathbf{n}(S_{ux}, u, x) = \mathbf{n}(S_{uv}, u, v) + \mathbf{n}(S_{uvx}, u, x)$$



NB: Cubic relations generates all relations of submodular vectors

The tool: submodular dependancies


Lemma (Submodular normal)

$\mathbf{n}(S, u, v)$ are the facet's normals of $\mathbb{DC}(\Pi_n)$

Lemma (Cubic relation)

$u, v, x \notin S \subseteq [n]$

$$\mathbf{n}(S_{uvx}, u, v) + \mathbf{n}(S_{ux}, u, x) = \mathbf{n}(S_{uv}, u, v) + \mathbf{n}(S_{uvx}, u, x)$$



NB: Cubic relations generates all relations of submodular vectors

The tool: submodular dependancies


Lemma (Submodular normal)

$\mathbf{n}(S, u, v)$ are the facet's normals of $\mathbb{DC}(\Pi_n)$

Lemma (Cubic relation)

$u, v, x \notin S \subseteq [n]$

$$\mathbf{n}(S_{uvx}, u, v) + \mathbf{n}(S_{ux}, u, x) = \mathbf{n}(S_{uv}, u, v) + \mathbf{n}(S_{uvx}, u, x)$$



NB: Cubic relations generates all relations of submodular vectors

The tool: submodular dependancies



Lemma (Submodular normal)

$\mathbf{n}(S, u, v)$ are the facet's normals of $\mathbb{DC}(\Pi_n)$

Lemma (Cubic relation)

$u, v, x \notin S \subseteq [n]$

$$\mathbf{n}(S_{uvx}, u, v) + \mathbf{n}(S_{ux}, u, x) = \mathbf{n}(S_{uv}, u, v) + \mathbf{n}(S_{uvx}, u, x)$$



NB: Cubic relations generates all relations of submodular vectors

Monotone path polytope and pivot rule polytope

Let $P \subset \mathbb{R}^d$ be a polytope.

Max-slope pivot rule: $A^\omega(v) = \operatorname{argmax} \left\{ \frac{\langle \omega, u - v \rangle}{\langle c, u - v \rangle}; u \text{ impr. neig. of } v \right\}$.

Coherent monotone path: A monotone path that can be obtained via the max-slope pivot rule.

Monotone path polytope $\Sigma_c(P)$ [?]: Fiber polytope of $P \xrightarrow{\pi} Q$ with Q a segment. (Can be seen as a Minkowski sum of sections of P .)
The vertices of $\Sigma_c(P)$ are all coherent monotone paths.

Coherent arborescence: An arborescence that can be obtained via the max-slope pivot rule.

Pivot rule polytope $\Pi_c(P)$: Polytope which vertices are all coherent arborescences.


$$\Pi_c(P) = \operatorname{conv} \left\{ \sum_{v \neq v_{opt}} \frac{1}{\langle c, A(v) - v \rangle} (A(v) - v); A \text{ coherent arbo. of } P \right\}$$

Monotone path polytope and pivot rule polytope

Coherent arborescence: An arborescence that can be obtained via the max-slope pivot rule.

Pivot rule polytope $\Pi_c(P)$: Polytope which vertices are all coherent arborescences. Can also be seen as a Minkowski sum of sections:

$$\sum_{v \in V(P)} (\text{section between } v \text{ and its improving neighbors})$$



Mimicking the case of the d -cube

Idea 1:

Fix a polytope P , and direction c , n vertices, m edges.

$\theta : \mathbb{R}^d \rightarrow \mathbb{R}^m$ sends the pivot fan inside $\text{Im}(\theta) \cap \mathcal{B}_m$

Problem: This is not a braid fan as $d \ll m \dots$

If m' classes of parallelism:

$\bar{\theta} : \mathbb{R}^d \rightarrow \mathbb{R}^{m'}$ sends the pivot fan inside $\text{Im}(\theta) \cap \mathcal{B}_{m'}$

Problem: This is not a braid fan as $d \ll m' < m \dots$

We need to go lower dimensional!


Adapted slope map

Idea 2:

Fix a polytope P , direction c ,
 n vertices, m edges.

Fix A arborescence:

$$\vartheta_A(\omega) = (\tau_\omega(u, A(u)) ; u \text{ vertex})$$



ϑ_A : linear, injective, $\mathbb{R}^d \rightarrow \mathbb{R}^{n-1}$

but if ω does not capture A , then $\vartheta_A(\omega)$ have no meaning...

Adapted slope map: $\vartheta(\omega) = \vartheta_{A^\omega}(\omega)$

i.e. take ω and look at the slope of the edges it selects.

Case of the d -simplex

$$d = n - 1 \iff P \text{ is a simplex}$$

For Δ_d : $\vartheta : \mathbb{R}^d \rightarrow \mathbb{R}^d$ piece-wise linear, $\ker \vartheta = \{\mathbf{0}\} \Rightarrow$ bijection
 ϑ sends the pivot fan of Δ_d inside \mathcal{B}_d .

