


How many generalized permutahedra are there?

Germain Poullot with Georg Loho & Arnau Padrol



24 March 2026, Combinatorial CoWorkspace
arXiv:2510.03177

How many generalized permutahedra are there?

Infinitely many!

How many generalized permutahedra are there?

Infinitely many!

Well, we can still gather them in finitely many classes.

Infinitely many!

Well, we can still gather them in finitely many classes.

Too many?

How many generalized permutahedra are there?

Infinitely many!

Well, we can still gather them in finitely many classes.

Too many?

Yes, but let's try to get a slightly more precise answer, shall we?

1 Generalized permutohedra, submodular cone

- 4 definitions
- Deformations

2 GP-sums and equality sets

- $\text{SC}_{n+1} \simeq \text{SC}_n^2$ more or less
- Drawing $\mathcal{L}(\text{SC}_n)^2$ inside $\mathcal{L}(\text{SC}_{n+1})$

3 Applications (now, we can count stuffs)

- Rays
- f -vector, total number of faces, non-simplicial faces

Generalized permutohedra, submodular cone

Generalized permutohedra

Generalized n -permutohedra are equivalently:

- Polytopes $\subseteq \mathbb{R}^n$ with edge directions $\mathbf{e}_i - \mathbf{e}_j$

Generalized permutohedra


Generalized n -permutohedra are equivalently:

- Polytopes $\subseteq \mathbb{R}^n$ with edge directions $\mathbf{e}_i - \mathbf{e}_j$
- Minkowski summand of the *permutohedron*

$$\Pi_n := \text{conv}\left((\sigma(1), \dots, \sigma(n)) ; \sigma \in \mathcal{S}_n\right)$$

Example (Permutahedron)



$$\Pi_n = \text{conv} \left\{ \begin{pmatrix} \sigma(1) \\ \vdots \\ \sigma(n) \end{pmatrix} ; \sigma \text{ permutation of } \{1, \dots, n\} \right\}$$



Permutahedron

Example (Permutahedron)


$$\Pi_n = \text{conv} \left\{ \begin{pmatrix} \sigma(1) \\ \vdots \\ \sigma(n) \end{pmatrix} ; \sigma \text{ permutation of } \{1, \dots, n\} \right\}$$



Minkowski summands

Definition (Minkowski sum)


P, Q polytopes, $P + Q := \{\mathbf{p} + \mathbf{q} \ ; \ \mathbf{p} \in P, \ \mathbf{q} \in Q\}$



Minkowski summands

Definition (Minkowski sum)


P, Q polytopes, $P + Q := \{\mathbf{p} + \mathbf{q} \ ; \ \mathbf{p} \in P, \ \mathbf{q} \in Q\}$



Minkowski summands

Definition (Minkowski sum)


P, Q polytopes, $P + Q := \{\mathbf{p} + \mathbf{q} \ ; \ \mathbf{p} \in P, \ \mathbf{q} \in Q\}$



Minkowski summands

Definition (Minkowski sum)

P, Q polytopes, $P + Q := \{\mathbf{p} + \mathbf{q} \ ; \ \mathbf{p} \in P, \ \mathbf{q} \in Q\}$



Generalized n -permutohedra are equivalently:

- Polytopes $\subseteq \mathbb{R}^n$ with edge directions $\mathbf{e}_i - \mathbf{e}_j$
- Minkowski summand of the *permutohedron*

$$\Pi_n := \text{conv}\left((\sigma(1), \dots, \sigma(n)) ; \sigma \in \mathcal{S}_n\right)$$

Generalized n -permutohedra are equivalently:

- Polytopes $\subseteq \mathbb{R}^n$ with edge directions $\mathbf{e}_i - \mathbf{e}_j$
- Minkowski summand of the *permutohedron*

$$\Pi_n := \text{conv}\left((\sigma(1), \dots, \sigma(n)) ; \sigma \in \mathcal{S}_n\right)$$

- Polytopes whose normal fan coarsens the *braid fan*

$$\text{Braid}_n := (\{\mathbf{x} ; x_i = x_j\} ; 1 \leq i < j \leq n)$$


Fan coarsening

$\mathcal{N}_P$: Normal fan of polytope P


Coarsening: Choose maximal cones and merge them

Theorem

Q is a Minkowski summand of P iff $\mathcal{N}_Q$ coarsens $\mathcal{N}_P$.



summand of






Fan coarsening

$\mathcal{N}_P$: Normal fan of polytope P

Coarsening: Choose maximal cones and merge them

Theorem

Q is a Minkowski summand of P iff $\mathcal{N}_Q$ coarsens $\mathcal{N}_P$.





Fan coarsening

$\mathcal{N}_P$: Normal fan of polytope P


Coarsening: Choose maximal cones and merge them

Theorem

Q is a Minkowski summand of P iff $\mathcal{N}_Q$ coarsens $\mathcal{N}_P$.




coarsens



Braid fan



Definition

Braid fan: arrangement of hyperplanes $H_{i,j} := \{\mathbf{x} ; x_i = x_j\}$



Definition

Braid fan: arrangement of hyperplanes $H_{i,j} := \{\mathbf{x} ; x_i = x_j\}$



Generalized permutohedra

Generalized n -permutohedra are equivalently:

- Polytopes $\subseteq \mathbb{R}^n$ with edge directions $\mathbf{e}_i - \mathbf{e}_j$
- Minkowski summand of the *permutohedron*

$$\Pi_n := \text{conv}\left((\sigma(1), \dots, \sigma(n)) ; \sigma \in \mathcal{S}_n\right)$$

- Polytopes whose normal fan coarsens the *braid fan*

$$\text{Braid}_n := (\{\mathbf{x} ; x_i = x_j\} ; 1 \leq i < j \leq n)$$

Generalized permutohedra

Generalized n -permutohedra are equivalently:

- Polytopes $\subseteq \mathbb{R}^n$ with edge directions $\mathbf{e}_i - \mathbf{e}_j$
- Minkowski summand of the *permutohedron*

$$\Pi_n := \text{conv}\left((\sigma(1), \dots, \sigma(n)) ; \sigma \in \mathcal{S}_n\right)$$

- Polytopes whose normal fan coarsens the *braid fan*

$$\text{Braid}_n := (\{\mathbf{x} ; x_i = x_j\} ; 1 \leq i < j \leq n)$$

- Set functions $f : 2^{[n]} \setminus \{\emptyset, [n]\} \rightarrow \mathbb{R}$ which are *submodular*, i.e.

$$f(A) + f(B) \geq f(A \cap B) + f(A \cup B)$$

Generalized permutohedra

Generalized n -permutohedra are equivalently:

- Polytopes $\subseteq \mathbb{R}^n$ with edge directions $\mathbf{e}_i - \mathbf{e}_j$
- Minkowski summand of the *permutahedron*

$$\Pi_n := \text{conv}\left((\sigma(1), \dots, \sigma(n)) ; \sigma \in \mathcal{S}_n\right)$$


- Polytopes whose normal fan coarsens the *braid fan*

$$\text{Braid}_n := (\{\mathbf{x} ; x_i = x_j\} ; 1 \leq i < j \leq n)$$

- Set functions $f : 2^{[n]} \setminus \{\emptyset, [n]\} \rightarrow \mathbb{R}$ which are *submodular*, i.e.

$$f(A) + f(B) \geq f(A \cap B) + f(A \cup B)$$

Submodular cone $\text{SC}_n := \{\text{all generalized } n\text{-permutohedra}\}$



$$P, P' \in \mathbb{SC}_n$$

P' is a *deformation* of P , denoted $P' \trianglelefteq P$, iff:

- P' is a Minkowski summand of P , i.e. $\exists R, \lambda > 0, \lambda P = P' + R$

$$P, P' \in \mathbb{SC}_n$$

P' is a *deformation* of P , denoted $P' \trianglelefteq P$, iff:

- P' is a Minkowski summand of P , i.e. $\exists R, \lambda > 0, \lambda P = P' + R$
- $\mathcal{N}_{P'}$ coarsens $\mathcal{N}_P$

$$P, P' \in \mathbb{SC}_n$$

P' is a *deformation* of P , denoted $P' \trianglelefteq P$, iff:

- P' is a Minkowski summand of P , i.e. $\exists R, \lambda > 0, \lambda P = P' + R$
- $\mathcal{N}_{P'}$ coarsens $\mathcal{N}_P$
- $\text{modular}(f') \subseteq \text{modular}(f)$ where
 $\text{modular}(f) := \{(A, B) ; f(A) + f(B) = f(A \cap B) + f(A \cup B)\}$


$$P, P' \in \mathbb{SC}_n$$

P' is a *deformation* of P , denoted $P' \trianglelefteq P$, iff:

- P' is a Minkowski summand of P , i.e. $\exists R, \lambda > 0, \lambda P = P' + R$
- $\mathcal{N}_{P'}$ coarsens $\mathcal{N}_P$
- $\text{modular}(f') \subseteq \text{modular}(f)$ where
 $\text{modular}(f) := \{(A, B) ; f(A) + f(B) = f(A \cap B) + f(A \cup B)\}$

P and P' are *equivalent*, denoted $P \sim P'$, iff $P' \trianglelefteq P$ and $P \trianglelefteq P'$

Deformations of Π_4



Permutahedron Π_4

Sequence of deformations of Π_4

What's the goal?

How many generalized permutahedra are there?

$\mathbb{SC}_n$ is a polyhedral cone, $\mathcal{L}(\mathbb{SC}_n)$ *its face lattice*.

What's the goal?

How many generalized permutahedra are there?

$\mathbb{SC}_n$ is a polyhedral cone, $\mathcal{L}(\mathbb{SC}_n)$ *its face lattice*.

faces of $\mathbb{SC}_n$ are gen. permutahedra.

What's the goal?

How many generalized permutahedra are there?

$\mathbb{SC}_n$ is a polyhedral cone, $\mathcal{L}(\mathbb{SC}_n)$ *its face lattice*.

Interior faces of $\mathbb{SC}_n$ are classes of equivalent gen. permutahedra.

What's the goal?

How many generalized permutahedra are there?

$\mathbb{SC}_n$ is a polyhedral cone, $\mathcal{L}(\mathbb{SC}_n)$ *its face lattice*.

Interior faces of $\mathbb{SC}_n$ are classes of equivalent gen. permutahedra.

$\mathcal{L}(\mathbb{SC}_n)$ is the set of classes of gen. permutahedra, ordered by $\trianglelefteq$.

What's the goal?

How many generalized permutahedra are there?

$\mathbb{SC}_n$ is a polyhedral cone, $\mathcal{L}(\mathbb{SC}_n)$ *its face lattice*.

Interior faces of $\mathbb{SC}_n$ are classes of equivalent gen. permutahedra.

$\mathcal{L}(\mathbb{SC}_n)$ is the set of classes of gen. permutahedra, ordered by $\trianglelefteq$.

I want to enumerate $\mathcal{L}(\mathbb{SC}_n)$.

What's the goal?

How many generalized permutahedra are there?

$\mathbb{SC}_n$ is a polyhedral cone, $\mathcal{L}(\mathbb{SC}_n)$ *its face lattice*.

Interior faces of $\mathbb{SC}_n$ are classes of equivalent gen. permutahedra.

$\mathcal{L}(\mathbb{SC}_n)$ is the set of classes of gen. permutahedra, ordered by $\trianglelefteq$.

I want to enumerate $\mathcal{L}(\mathbb{SC}_n)$.

I want to parameterise $\mathcal{L}(\mathbb{SC}_{n+1})$ via $\mathcal{L}(\mathbb{SC}_n)$

What's the goal?

How many generalized permutahedra are there?

$\mathbb{SC}_n$ is a polyhedral cone, $\mathcal{L}(\mathbb{SC}_n)$ *its face lattice*.

Interior faces of $\mathbb{SC}_n$ are classes of equivalent gen. permutahedra.

$\mathcal{L}(\mathbb{SC}_n)$ is the set of classes of gen. permutahedra, ordered by $\trianglelefteq$.

I want to enumerate $\mathcal{L}(\mathbb{SC}_n)$.

I want to parameterise $\mathcal{L}(\mathbb{SC}_{n+1})$ via $\mathcal{L}(\mathbb{SC}_n)^2$.

What's the goal?

How many generalized permutahedra are there?

$\mathbb{SC}_n$ is a polyhedral cone, $\mathcal{L}(\mathbb{SC}_n)$ its face lattice.

Interior faces of $\mathbb{SC}_n$ are classes of equivalent gen. permutahedra.

$\mathcal{L}(\mathbb{SC}_n)$ is the set of classes of gen. permutahedra, ordered by $\trianglelefteq$.

I want to enumerate $\mathcal{L}(\mathbb{SC}_n)$.


I want to parameterise $\mathcal{L}(\mathbb{SC}_{n+1})$ via $\mathcal{L}(\mathbb{SC}_n)^2$.

$\dim \mathbb{SC}_n = 2^n - 1$, $\#_{\text{facets}} \mathbb{SC}_n = \binom{n}{2} 2^{n-2}$, $\#_{\text{rays}} \mathbb{SC}_n = \text{unknown}$


Some faces are “known”: associahedron, graphical zonotopes (arXiv:2111.12422), nestohedra (arXiv:2109.09200), etc.

GP-sums and equality sets

First, embed $P \in \mathbb{SC}_{n+1}$ inside $\mathbb{R}^n$: forget last coordinate.
 Now, edges directions are $\mathbf{e}_i$ and $\mathbf{e}_i - \mathbf{e}_j$ for $1 \leq i < j \leq n$




First, embed $P \in \mathbb{SC}_{n+1}$ inside $\mathbb{R}^n$: forget last coordinate.
 Now, edges directions are $\mathbf{e}_i$ and $\mathbf{e}_i - \mathbf{e}_j$ for $1 \leq i < j \leq n$



P^+ , P^- : **opposite** faces, min/max $\mathbf{1}$ on P are $\in \mathbb{SC}_n$.

First, embed $P \in \mathbb{SC}_{n+1}$ inside $\mathbb{R}^n$: forget last coordinate.
 Now, edges directions are $\mathbf{e}_i$ and $\mathbf{e}_i - \mathbf{e}_j$ for $1 \leq i < j \leq n$




P^+ , P^- : **opposite** faces, min/max $\mathbf{1}$ on P are $\in \mathbb{SC}_n$.

Theorem (Frank '11)

*Deformed permutahedra are **uniquely** determined by their pair of top and bottom faces.*

Thus, $P \mapsto (P^+, P^-)$ is a bijection:

First, embed $P \in \mathbb{SC}_{n+1}$ inside $\mathbb{R}^n$: forget last coordinate.
 Now, edges directions are $\mathbf{e}_i$ and $\mathbf{e}_i - \mathbf{e}_j$ for $1 \leq i < j \leq n$




P^+ , P^- : **opposite** faces, min/max $\mathbf{1}$ on P are $\in \mathbb{SC}_n$.

Theorem (Frank '11)

Deformed permutahedra are **uniquely** determined by their pair of top and bottom faces.

Thus, $P \mapsto (P^+, P^-)$ is a bijection: $\mathbb{SC}_{n+1} \simeq (\text{roughly } \mathbb{SC}_n^2)$

First, embed $P \in \mathbb{SC}_{n+1}$ inside $\mathbb{R}^n$: forget last coordinate.
 Now, edges directions are $\mathbf{e}_i$ and $\mathbf{e}_i - \mathbf{e}_j$ for $1 \leq i < j \leq n$



P^+ , P^- : **opposite** faces, min/max $\mathbf{1}$ on P are $\in \mathbb{SC}_n$.

Theorem (Frank '11)

*Deformed permutahedra are **uniquely** determined by their pair of top and bottom faces.*


Thus, $P \mapsto (P^+, P^-)$ is a bijection: $\mathbb{SC}_{n+1} \simeq (\text{roughly } \mathbb{SC}_n^2)$

What's the reciprocal bijection?

Theorem (Frank '11)

Deformed permutahedra are **uniquely** determined by their pair of top and bottom faces: $P \mapsto (P^+, P^-)$ is a bijection.


GP-sum: $Q \boxdot P := (Q + \mathbb{R}^+) \cap (P + \mathbb{R}^-)$



Theorem (Frank '11)


Deformed permutahedra are **uniquely** determined by their pair of top and bottom faces: $P \mapsto (P^+, P^-)$ is a bijection.

GP-sum: $Q \boxdot P := (Q + \mathbb{R}^+) \cap (P + \mathbb{R}^-)$







Reciprocal of $P \mapsto (P^+, P^-)$ is the map $(P, Q) \mapsto Q \boxdot P$

SC_3 drawn with GP-sums




GP-sums

GP-sum exists iff $P \subseteq (Q + \mathbb{R}^+)$ and $Q \subseteq (P + \mathbb{R}^-)$




GP-sums

GP-sum exists iff $P \subseteq (Q + \mathbb{R}^+)$ and $Q \subseteq (P + \mathbb{R}^-)$
 iff for all $i \in [n]$, $\sigma \in S_n$, we have $P_i^\sigma \leq Q_i^\sigma$



GP-sums


GP-sum exists iff $P \subseteq (Q + \mathbb{R}^+)$ and $Q \subseteq (P + \mathbb{R}^-)$
 iff for all $i \in [n], \sigma \in S_n$, we have $P_i^\sigma \leq Q_i^\sigma$



But: I have " $\text{SC}_{n+1} \simeq \text{SC}_n^2$ ". I'd like " $\mathcal{L}(\text{SC}_{n+1}) \simeq \mathcal{L}(\text{SC}_n)^2$ "...

GP-sums



GP-sum exists iff $P \subseteq (Q + \mathbb{R}^+)$ and $Q \subseteq (P + \mathbb{R}^-)$
 iff for all $i \in [n], \sigma \in S_n$, we have $P_i^\sigma \leq Q_i^\sigma$



But: I have " $SC_{n+1} \simeq SC_n^2$ ". I'd like " $\mathcal{L}(SC_{n+1}) \simeq \mathcal{L}(SC_n)^2$ "...

GP-sums


GP-sum exists iff $P \subseteq (Q + \mathbb{R}^+)$ and $Q \subseteq (P + \mathbb{R}^-)$
 iff for all $i \in [n], \sigma \in S_n$, we have $P_i^\sigma \leq Q_i^\sigma$



But: I have " $SC_{n+1} \simeq SC_n^2$ ". I'd like " $\mathcal{L}(SC_{n+1}) \simeq \mathcal{L}(SC_n)^2$ "...

GP-sums

GP-sum exists iff $P \subseteq (Q + \mathbb{R}^+)$ and $Q \subseteq (P + \mathbb{R}^-)$
 iff for all $i \in [n], \sigma \in S_n$, we have $P_i^\sigma \leq Q_i^\sigma$



But: I have " $SC_{n+1} \simeq SC_n^2$ ". I'd like " $\mathcal{L}(SC_{n+1}) \simeq \mathcal{L}(SC_n)^2$ "...

Poset of valid equality sets

$$P, Q \in \mathbb{SC}_n$$

Definition

Equality set: $\mathbf{E}(P, Q) := \{(i, \sigma) \in [n] \times \mathcal{S}_n ; Q_i^\sigma = P_i^\sigma\}$

Poset of valid equality sets:

$$\mathcal{E}(P, Q) := \{\mathbf{E}(P', Q') ; P', Q' \in \mathbb{SC}_n, P' \sim P \text{ and } Q' \sim Q\}$$

Poset of valid equality sets

$$P, Q \in \mathbb{SC}_n$$

Definition


Equality set: $\mathbf{E}(P, Q) := \{(i, \sigma) \in [n] \times \mathcal{S}_n ; Q_i^\sigma = P_i^\sigma\}$

Poset of valid equality sets:

$$\mathcal{E}(P, Q) := \{\mathbf{E}(P', Q') ; P', Q' \in \mathbb{SC}_n, P' \sim P \text{ and } Q' \sim Q\}$$

A GP-sum is *generic*, denoted $Q \square P$, if $\mathbf{E}(P, Q) = \emptyset$.

SC_3 with equality sets



Poset of valid equality sets

$$P, Q \in \mathbb{SC}_n$$

Definition

Equality set: $\mathbf{E}(P, Q) := \{(i, \sigma) \in [n] \times \mathcal{S}_n ; Q_i^\sigma = P_i^\sigma\}$

Poset of valid equality sets:

$$\mathcal{E}(P, Q) := \{\mathbf{E}(P', Q') ; P', Q' \in \mathbb{SC}_n, P' \sim P \text{ and } Q' \sim Q\}$$

A GP-sum is *generic*, denoted $Q \square P$, if $\mathbf{E}(P, Q) = \emptyset$.

Poset of valid equality sets

$$P, Q \in \mathbb{SC}_n$$

Definition

Equality set: $\mathbf{E}(P, Q) := \{(i, \sigma) \in [n] \times \mathcal{S}_n ; Q_i^\sigma = P_i^\sigma\}$

Poset of valid equality sets:

$$\mathcal{E}(P, Q) := \{\mathbf{E}(P', Q') ; P', Q' \in \mathbb{SC}_n, P' \sim P \text{ and } Q' \sim Q\}$$

A GP-sum is *generic*, denoted $Q \square P$, if $\mathbf{E}(P, Q) = \emptyset$.

Theorem (Loho–Padrol–P., 25+)

$Q' \square P' \sim Q \square P$ iff $P' \sim P$, $Q' \sim Q$ and $\mathbf{E}(P', Q') = \mathbf{E}(P, Q)$.

Poset of valid equality sets

$$P, Q \in \mathbb{SC}_n$$

Definition

Equality set: $\mathbf{E}(P, Q) := \{(i, \sigma) \in [n] \times \mathcal{S}_n ; Q_i^\sigma = P_i^\sigma\}$

Poset of valid equality sets:

$$\mathcal{E}(P, Q) := \{\mathbf{E}(P', Q') ; P', Q' \in \mathbb{SC}_n, P' \sim P \text{ and } Q' \sim Q\}$$

A GP-sum is *generic*, denoted $Q \square P$, if $\mathbf{E}(P, Q) = \emptyset$.

Theorem (Loho–Padrol–P., 25+)


$Q' \square P' \sim Q \square P$ iff $P' \sim P$, $Q' \sim Q$ and $\mathbf{E}(P', Q') = \mathbf{E}(P, Q)$.



Corollary (Loho–Padrol–P., 25+)

The faces of $\mathbb{SC}_{n+1}$ are in bijection with triples $(P, Q, \mathbf{E})$ where P and Q are taken up to equivalence and $\mathbf{E} \in \mathcal{E}(P, Q)$.

$\mathbb{DC}(P)$: class of $P' \sim P$, i.e. faces of $\mathbb{SC}_n$

$$\mathcal{L}(\mathbb{SC}_n)^2 \simeq \{\mathbb{DC}(Q \square P) ; P, Q \in \mathbb{SC}_n^2\}$$






Applications (now, we can count stuffs)

Rays


Rays of $\mathbb{SC}_n$ = indecomposable generalized permutohedra
= bottom-most elements in:



How to get rays of $\mathbb{SC}_{n+1}$?

Rays

Rays of $\mathbb{SC}_n$ = indecomposable generalized permutohedra
= bottom-most elements in:




How to get rays of $\mathbb{SC}_{n+1}$?

Start with $Q \square P$,

Rays

Rays of $\mathbb{SC}_n$ = indecomposable generalized permutohedra
= bottom-most elements in:




How to get rays of $\mathbb{SC}_{n+1}$?

Start with $Q \square P$, then move P and Q around to increase $\mathbf{E}(P, Q)$,

Rays

Rays of $\mathbb{SC}_n$ = indecomposable generalized permutohedra
= bottom-most elements in:




How to get rays of $\mathbb{SC}_{n+1}$?

Start with $Q \square P$, then move P and Q around to increase $\mathbf{E}(P, Q)$,
then **hope** for touching the bottom...

Rays

Rays of $\mathbb{SC}_n$ = indecomposable generalized permutohedra
= bottom-most elements in:




How to get rays of $\mathbb{SC}_{n+1}$?

Start with $Q \square P$, then move P and Q around to increase $\mathbf{E}(P, Q)$,
then **hope** for touching the bottom...


To increase hopes, start with $P, Q \in \text{rays of } \mathbb{SC}_n$!

Rays


$P, Q \in \text{rays of } \mathbb{SC}_n$, i.e. deformations are translations & dilations.



$E = \emptyset$




bigger E




biggest E

Rays


$P, Q \in \text{rays of } \mathbb{SC}_n$, i.e. deformations are translations & dilations.



$E = \emptyset$



bigger E




biggest E

Problem: Sometimes, you'll need to dilate by 0 to get a ray.

Rays

$P, Q \in \text{rays of } \mathbb{SC}_n$, i.e. deformations are translations & dilations.



Problem: Sometimes, you'll need to dilate by 0 to get a ray.

Lemma (Loho–Padrol–P., 25+)

Technical lemma about non-0 dilations that you wouldn't read.

Theorem (Loho–Padrol–P., 25+)

If $P_1, \dots, P_s$ are pair-wise fertile rays of $\mathbb{SC}_n$, then you can construct explicitly $R_1, \dots, R_{s^2}$ pair-wise fertile rays of $\mathbb{SC}_{n+1}$.

Number of rays of $\mathbb{SC}_n$

t_n := number of rays of $\mathbb{SC}_n$.

Theorem ((approximately) Nguyen, '86)

The number t_n of rays of $\mathbb{SC}_n$ satisfies

$$2^{2^n/n^{3/2}} \leq t_n$$

Number of rays of $\mathbb{SC}_n$

t_n := number of rays of $\mathbb{SC}_n$.

Theorem ((approximately) Nguyen, '86)

The number t_n of rays of $\mathbb{SC}_n$ satisfies

$$2^{2^n/n^{3/2}} \leq t_n$$

Theorem (Loho–Padrol–P., '25+)

The number t_n of rays of $\mathbb{SC}_n$ satisfies

$$2^{2^{n-2}} \leq t_n$$

N.B.: By Upper Bound Theorem: $t_n \leq n^{2^n}$.

Precisely: $n - 2 \leq \log_2 \log_2 t_n \leq n + \log_2 \log_2 n + 1$

Lemma (Loho–Padrol–P., 25+)

$$\dim \mathbb{DC}(Q \square P) = \dim \mathbb{DC}(P) + \dim \mathbb{DC}(Q) + 1$$

Lemma (Loho–Padrol–P., 25+)

$$\dim \mathbb{DC}(Q \square P) = \dim \mathbb{DC}(P) + \dim \mathbb{DC}(Q) + 1$$

$F_n(X) := \sum_k f(n, k) X^k$ where $f(n, k) :=$ number k -faces of $\mathbb{SC}_n$

Corollary (Loho–Padrol–P., 25+)

$$F_{n+1}(X) \succeq_{\text{coeff.-wise}} X \cdot F_n(X)^2$$

Lemma (Loho–Padrol–P., 25+)

$$\dim \mathbb{DC}(Q \square P) = \dim \mathbb{DC}(P) + \dim \mathbb{DC}(Q) + 1$$

$F_n(X) := \sum_k f(n, k) X^k$ where $f(n, k) :=$ number k -faces of $\mathbb{SC}_n$

Corollary (Loho–Padrol–P., 25+)

$$F_{n+1}(X) \succeq_{\text{coeff.-wise}} X \cdot F_n(X)^2$$

$$\text{Hence, } F_{n+1}(X) \succeq_{\text{coeff.-wise}} F_4^{2^{n-4}}$$

Lemma (Loho–Padrol–P., 25+)

$$\dim \mathbb{DC}(Q \square P) = \dim \mathbb{DC}(P) + \dim \mathbb{DC}(Q) + 1$$

$F_n(X) := \sum_k f(n, k) X^k$ where $f(n, k) :=$ number k -faces of $\mathbb{SC}_n$

Corollary (Loho–Padrol–P., 25+)

$$F_{n+1}(X) \succeq_{\text{coeff.-wise}} X \cdot F_n(X)^2$$

$$\text{Hence, } F_{n+1}(X) \succeq_{\text{coeff.-wise}} F_4^{2^{n-4}}$$

$$\text{And, } \#_{\text{faces}} \mathbb{SC}_n \geq 22\,108^{2^{n-4}}$$

Lemma (Loho–Padrol–P., 25+)

$\mathbb{DC}(P)$ *not simplicial* $\Rightarrow \mathbb{DC}(Q \square P)$ *not simplicial* for any $Q \in \mathbb{SC}_n$.

Lemma (Loho–Padrol–P., 25+)

$\mathbb{DC}(P)$ not simplicial $\Rightarrow \mathbb{DC}(Q \square P)$ not simplicial for any $Q \in \mathbb{SC}_n$.

Corollary (Loho–Padrol–P., 25+)


Number of non-simplicial faces of $\mathbb{SC}_n \geq \prod_{p < n} \#_{\text{faces}} \mathbb{SC}_p$

In particular, $\mathbb{SC}_n$ is “very far” from being simplicial, so we should be “very far” for the Upper Bound Theorem equality case.

Furthermore:

- $\mathcal{L}(\mathbb{SC}_n)^2 \simeq \{\mathbb{DC}(Q \square P); P, Q \in \mathbb{SC}_n^2\}$ is the face figure of $[0, 1]^n$.
- $\mathcal{E}(P, P)$ is iso. to the interval of $\mathcal{L}(\mathbb{SC}_{n+1})$ between P and $P \square P$.
In particular, $\mathcal{E}(\Pi_n, \Pi_n)$ is a face figure.
- In general, $\mathcal{E}(P, Q)$ is a co-graded, co-atomistic, join-semi-lattice with a geometric interpretation.
- Notion of independence polytope Ind_P for all generalized permutohedra, similar to matroid polytopes, with:
 Ind_P indecomposable iff P indecomposable, and
 $\text{Ind}_{P+Q} = \text{Ind}_P + \text{Ind}_Q$

Thank you!





Tools: submodular dependencies

Notations: $Sx = S \cup \{x\}$, $(\mathbf{f}_x)_{x \subseteq [n]}$ canonical basis of $\mathbb{R}^{2^{[n]}}$

Definition

Submodular vector $\mathbf{n}(S, u, v) = \mathbf{f}_{Suv} - \mathbf{f}_{Su} - \mathbf{f}_{Sv} + \mathbf{f}_S$
for $u, v \in S \subseteq [n]$





Tools: submodular dependencies

Notations: $Sx = S \cup \{x\}$, $(\mathbf{f}_x)_{x \subseteq [n]}$ canonical basis of $\mathbb{R}^{2^{[n]}}$

Definition

Submodular vector $\mathbf{n}(S, u, v) = \mathbf{f}_{Suv} - \mathbf{f}_{Su} - \mathbf{f}_{Sv} + \mathbf{f}_S$
for $u, v \in S \subseteq [n]$




Lemma (Submodular normal)

$\mathbf{n}(S, u, v)$ are the facet's normals of $\mathbb{DC}(\Pi_n)$

Lemma (Cubic relation)

$u, v, x \notin S \subseteq [n]$

$$\mathbf{n}(Suvx, u, v) + \mathbf{n}(Sux, u, x) = \mathbf{n}(Suv, u, v) + \mathbf{n}(Suvx, u, x)$$



NB: Cubic relations generates all relations of submodular vectors


Lemma (Submodular normal)

$\mathbf{n}(S, u, v)$ are the facet's normals of $\mathbb{DC}(\Pi_n)$

Lemma (Cubic relation)

$u, v, x \notin S \subseteq [n]$

$$\mathbf{n}(Suvx, u, v) + \mathbf{n}(Sux, u, x) = \mathbf{n}(Suv, u, v) + \mathbf{n}(Suvx, u, x)$$



NB: Cubic relations generates all relations of submodular vectors


Lemma (Submodular normal)

$\mathbf{n}(S, u, v)$ are the facet's normals of $\mathbb{DC}(\Pi_n)$

Lemma (Cubic relation)

$u, v, x \notin S \subseteq [n]$

$$\mathbf{n}(Suvx, u, v) + \mathbf{n}(Sux, u, x) = \mathbf{n}(Suv, u, v) + \mathbf{n}(Suvx, u, x)$$



NB: Cubic relations generates all relations of submodular vectors



Lemma (Submodular normal)

$\mathbf{n}(S, u, v)$ are the facet's normals of $\mathbb{DC}(\Pi_n)$

Lemma (Cubic relation)

$u, v, x \notin S \subseteq [n]$


$$\mathbf{n}(Suvx, u, v) + \mathbf{n}(Sux, u, x) = \mathbf{n}(Suv, u, v) + \mathbf{n}(Suvx, u, x)$$




NB: Cubic relations generates all relations of submodular vectors

Fertility


(P, Q) is *fertile* iff there exists $i \in [n]$ such that for all $\tau \in S_n$, either $P_i^\tau \neq \min_{\sigma \in S_n} P_i^\sigma$ or $Q_i^\tau \neq \max_{\sigma \in S_n} Q_i^\sigma$




$i = 1$



$i = 2$



$i = 3$



Multiple choice questions (sometimes several answer possible)

A sum of 2 co-planar triangles can have	3 edges	4 edges	5 edges	6 edges
$\mathcal{N}_{P+Q} = \text{--- of } \mathcal{N}_P, \mathcal{N}_Q$	union	intersection	common refinement	product
Edge directions of $\text{GP} \in$	$\{\mathbf{e}_i\}_i$	$\{\mathbf{e}_i + \mathbf{e}_j\}_{i,j}$	$\{\mathbf{e}_i - \mathbf{e}_j\}_{i,j}$	$\{\mathbf{e}_i \pm \mathbf{e}_j\}_{i,j}$
Z_G has no hexagonal face iff G is	a tree	K_3 -free	K_4 -free	complete
--- form a basis of $\text{Vect}(\mathbb{SC}_n)$	$\{\Delta_X\}_{X \subseteq [n]}$	nestohedra	shard polytopes	matroid polytopes
$\dim \mathbb{SC}_3 =$	3	4	5	11
Group of symmetries of $\mathbb{SC}_n$:	Free_n	$\mathcal{S}_n$	$\mathcal{S}_n \times \mathbb{Z}_2$	$\mathcal{S}_n \times \mathbb{Z}_n$
# 3-dim indecomposable GP:	3	5	7	37

1. Show that $\text{Vert}(P + Q) \subseteq \{\mathbf{u} + \mathbf{v} ; \mathbf{u} \in \text{Vert}(P), \mathbf{v} \in \text{Vert}(Q)\}$. Find an example with strict inclusion. Prove that $\mathcal{N}_{P+Q}$ is the common refinement of $\mathcal{N}_P$ and $\mathcal{N}_Q$.
2. Find all the ways to write the 2-dimensional permutahedron as a Minkowski sum of indecomposable polytopes.
3. Prove that $\mathbf{h}_{P+Q} = \mathbf{h}_P + \mathbf{h}_Q$ and $\ell_{P+Q} = \ell_P + \ell_Q$. What are $\mathbf{h}_{P+t}$ and ℓ_{P+t} ?
4. Prove that a triangle is Minkowski indecomposable. Deduce that all simplicial polytopes are Minkowski indecomposable.

5. Take vectors $\mathbf{s}, \mathbf{s}', \mathbf{r}_1, \dots, \mathbf{r}_{d-1}$ such that $C = \text{cone}(\mathbf{s}, \mathbf{r}_1, \dots, \mathbf{r}_{d-1})$ and $C' = \text{cone}(\mathbf{s}', \mathbf{r}_1, \dots, \mathbf{r}_{d-1})$ are simplicial cones which intersect on their proper face $C \cap C' = \text{cone}(\mathbf{r}_1, \dots, \mathbf{r}_{d-1})$. Show that there exist $\alpha_{\mathbf{s}}, \alpha_{\mathbf{s}'} > 0$ and $\alpha_{\mathbf{r}_i} \in \mathbb{R}$ such that:

$$\alpha_{\mathbf{s}}\mathbf{s} + \alpha_{\mathbf{s}'}\mathbf{s}' + \sum_i \alpha_{\mathbf{r}_i}\mathbf{r}_i = \mathbf{0}$$

6. Compute (and draw) the deformation cone of a parallelogram (use the edge-lengths point of view). What about a parallelepiped? Show that there is a **unique** way to write a parallelepiped P as a sum of Minkowski indecomposable polytopes.

7. Show that $\mathbb{SC}_n$ has $\binom{n}{2}2^{n-2}$ facets. Give bounds on its number of rays.

8. Show that graphical zonotopes are generalized permutahedra. Show that nestohedra are generalized permutahedra. Show that matroid polytopes are generalized permutahedra. Which are indecomposable?

9. If G is triangle-free, then $\mathbb{DC}(Z_G)$ is simplicial (see lecture). Each face of $\mathbb{DC}(Z_G)$ is associated to a polytope: which polytope?

10. The *weighted graphical zonotope* of a graph $G = (V, E)$ and weight (on edges) $\omega : E \rightarrow \mathbb{R}_+^*$ is $Z_{G,\omega} := \sum_{(i,j) \in E} \omega(i,j)[\mathbf{e}_i, \mathbf{e}_j]$. Show that: $\{\text{zonotopes}\} \cap \{\text{generalized permutahedra}\} = \{\text{weighted graphical zonotopes}\}$. Show that the cone of weighted graphical zonotopes is a section (i.e. intersection with a linear sub-space) of the submodular cone, such that the rays of this section are rays of $\mathbb{SC}_n$. What is the dimension of this section?

11. Prove the cubic relations hold. Show that cubic relations generates all linear relations between submodular vectors.