


Minkowski indecomposability of frameworks

Germain Poullot with Arnau Padrol
ArXiv: 2512.05307




31 March 2026

- 1 What are polytopes and frameworks?
 - Two definitions of polytopes
 - Faces and graphs
- 2 Deformations (weak Minkowski summands)
 - Minkowski decomposition
 - Cone of deformations
- 3 Graph of edge dependencies
 - Triangle, parallelograms
 - Graph of edge dependencies
 - Pinning closure and our strongest theorem
- 4 Applications
 - First applications
 - New indecomposable generalized permutahedra
 - 0/1-polytopes
 - Platonic, Archimedean, and Jhonson solids
 - Products, parallelogramic Minkowski sums
 - More...

What are polytopes and frameworks?



Definition

Polytope: convex hull of finitely many points in $\mathbb{R}^n$





Definition

Polytope: convex hull of finitely many points in $\mathbb{R}^n$
bounded intersection of finitely many half-spaces in $\mathbb{R}^n$





Definition

Polytope: convex hull of finitely many points in $\mathbb{R}^n$
bounded intersection of finitely many half-spaces in $\mathbb{R}^n$





Definition

Polytope: convex hull of finitely many points in $\mathbb{R}^n$
bounded intersection of finitely many half-spaces in $\mathbb{R}^n$





Definition

Polytope: convex hull of finitely many points in $\mathbb{R}^n$
bounded intersection of finitely many half-spaces in $\mathbb{R}^n$




Definition

Polytope: convex hull of finitely many points in $\mathbb{R}^n$
bounded intersection of finitely many half-spaces in $\mathbb{R}^n$



Definition

Face: $P^c := \{ \mathbf{x} \in \mathbb{R}^n ; \langle \mathbf{x}, \mathbf{c} \rangle = \max_{\mathbf{y} \in P} \langle \mathbf{y}, \mathbf{c} \rangle \}$




P



Definition

Face: $P^c := \{ \mathbf{x} \in \mathbb{R}^n ; \langle \mathbf{x}, \mathbf{c} \rangle = \max_{\mathbf{y} \in P} \langle \mathbf{y}, \mathbf{c} \rangle \}$




P



Definition

Face: $P^c := \{ \mathbf{x} \in \mathbb{R}^n ; \langle \mathbf{x}, \mathbf{c} \rangle = \max_{\mathbf{y} \in P} \langle \mathbf{y}, \mathbf{c} \rangle \}$




P




Face lattice

Face lattice: poset of inclusions of faces




Face lattice

Face lattice: poset of inclusions of faces




Face lattice

Face lattice: poset of inclusions of faces




Face lattice

Face lattice: poset of inclusions of faces



Face lattice

Face lattice: poset of inclusions of faces



Graph of polytopes *versus* framework

Definition

Framework $\mathcal{F} = (V, E, \phi)$ where (V, E) is a graph, and $\phi : V \rightarrow \mathbb{R}^d$ its *realization*.

Graph of polytopes *versus* framework

Definition

Framework $\mathcal{F} = (V, E, \phi)$ where (V, E) is a graph, and $\phi : V \rightarrow \mathbb{R}^d$ its *realization*.

Examples:

- $\mathcal{F}(P)$: vertex-edge graph of P with ϕ its realization

Graph of polytopes *versus* framework

Definition

Framework $\mathcal{F} = (V, E, \phi)$ where (V, E) is a graph, and $\phi : V \rightarrow \mathbb{R}^d$ its *realization*.

Examples:

- $\mathcal{F}(P)$: vertex-edge graph of P with ϕ its realization
- Plane embedding ϕ of a planar graph (V, E)

Definition

Framework $\mathcal{F} = (V, E, \phi)$ where (V, E) is a graph, and $\phi : V \rightarrow \mathbb{R}^d$ its *realization*.

Examples:

- $\mathcal{F}(P)$: vertex-edge graph of P with ϕ its realization
- Plane embedding ϕ of a planar graph (V, E)
- $V =$ affinely indep. points $\phi_1, \dots, \phi_n \in \mathbb{R}^d$; and $E =$ cycle on V

Graph of polytopes *versus* framework

Definition

Framework $\mathcal{F} = (V, E, \phi)$ where (V, E) is a graph, and $\phi : V \rightarrow \mathbb{R}^d$ its *realization*.

Examples:

- $\mathcal{F}(P)$: vertex-edge graph of P with ϕ its realization
- Plane embedding ϕ of a planar graph (V, E)
- $V =$ affinely indep. points $\phi_1, \dots, \phi_n \in \mathbb{R}^d$; and $E =$ cycle on V

Lemma

The cycle space of a polytopal framing is generated by 2-dimensional cycles.


Deformations (weak Minkowski summands)

Definition

P, Q polytopes. *Minkowski sum*:

$$P + Q = \{ \mathbf{p} + \mathbf{q} \ ; \ \mathbf{p} \in P, \ \mathbf{q} \in Q \}$$

N.B. $\text{Vert}(P + Q) \subseteq \text{Vert}(P) + \text{Vert}(Q)$




Minkowski sum

Definition

P, Q polytopes. *Minkowski sum*:


$$P + Q = \{ \mathbf{p} + \mathbf{q} \ ; \ \mathbf{p} \in P, \ \mathbf{q} \in Q \}$$

N.B. $\text{Vert}(P + Q) \subseteq \text{Vert}(P) + \text{Vert}(Q)$




Sum of frameworks $\mathcal{F} + \mathcal{F}'$ with same graph has realization $\phi + \phi'$.


Minkowski sum




Minkowski sum



Minkowski sum



Minkowski sum



Definition

Q is a *Minkowski summand*, a.k.a. *deformation*, of P when there exists R and $\lambda > 0$ with:

$$\lambda P = Q + R$$

Definition

Q is a *Minkowski summand*, a.k.a. *deformation*, of P when there exists R and $\lambda > 0$ with:

$$\lambda P = Q + R$$

Deformation cone: $\mathbb{DC}(P) = \{Q ; Q \text{ is a deformation of } P\}$

Indecomposable: deformations of P are translated-dilations of P

Definition

Q is a *Minkowski summand*, a.k.a. *deformation*, of P when there exists R and $\lambda > 0$ with:

$$\lambda P = Q + R$$

Deformation cone: $\mathbb{DC}(P) = \{Q ; Q \text{ is a deformation of } P\}$

Indecomposable: deformations of P are translated-dilations of P

What is the best way to write P as a Minkowski sum ?

- With the fewest number of (indecomposable) summands ?
- With the (indecomposable) summands of smallest dimension ?
- Respecting some symmetries ?
- ...

Definition

Q is a *Minkowski summand*, a.k.a. *deformation*, of P when there exists R and $\lambda > 0$ with:

$$\lambda P = Q + R$$

Deformation cone: $\mathbb{DC}(P) = \{Q ; Q \text{ is a deformation of } P\}$


Indecomposable: deformations of P are translated-dilations of P

What is the best way to write P as a Minkowski sum ?

- With the fewest number of (indecomposable) summands ?
- With the (indecomposable) summands of smallest dimension ?
- Respecting some symmetries ?
- ...



$\implies$ What is the structure of $\mathbb{DC}(P)$?

Minkowski summands





Observation

If $P = Q + R$, then the edges of P “are” edges of Q or of R .
 $\Rightarrow$ I can write deformations of P using edge-length vectors.




Observation

If $P = Q + R$, then the edges of P “are” edges of Q or of R .
 $\Rightarrow$ I can write deformations of P using edge-length vectors.




Observation

If $P = Q + R$, then the edges of P “are” edges of Q or of R .
 $\Rightarrow$ I can write deformations of P using edge-length vectors.




Observation

If $P = Q + R$, then the edges of P “are” edges of Q or of R .
 $\Rightarrow$ I can write deformations of P using edge-length vectors.



Observation


If $P = Q + R$, then the edges of P “are” edges of Q or of R .
 $\Rightarrow$ I can write deformations of P using edge-length vectors.



Remark:

As P, Q, R have “same edges”, we also have $\mathcal{F}(P) = \mathcal{F}(Q) + \mathcal{F}(R)$.

Deformations of 3-dim permutahedron



Permutahedron Π_4

Sequence of deformations of Π_4

Edge-length deformation cone

Theorem

Q deformation of $P \Leftrightarrow \mathcal{F}(Q)$ deformation of $\mathcal{F}(P)$
 $\Leftrightarrow$ same edge-directions, but different lengths

Definition

Deformation cone: $\mathbb{DC}(\mathcal{F}) = \{\mathcal{G} ; \mathcal{G} \text{ same graph and edge-dir } \mathcal{F}\}$


Edge-length deformation cone

Theorem

Q deformation of $P \Leftrightarrow \mathcal{F}(Q)$ deformation of $\mathcal{F}(P)$
 $\Leftrightarrow$ same edge-directions, but different lengths

Definition

Deformation cone: $\mathbb{DC}(\mathcal{F}) = \{\mathcal{G} ; \mathcal{G} \text{ same graph and edge-dir } \mathcal{F}\}$



Parametrization:

edge-length vector:

$$\ell = (\ell_e)_{e \text{ edge}}$$


Edge-length deformation cone

Theorem

Q deformation of $P \Leftrightarrow \mathcal{F}(Q)$ deformation of $\mathcal{F}(P)$
 $\Leftrightarrow$ same edge-directions, but different lengths

Definition

Deformation cone: $\mathbb{DC}(\mathcal{F}) = \{\mathcal{G} ; \mathcal{G} \text{ same graph and edge-dir } \mathcal{F}\}$



Parametrization:

edge-length vector:

$$\ell = (\ell_e)_{e \text{ edge}}$$


Edge-length deformation cone

Theorem

Q deformation of $P \Leftrightarrow \mathcal{F}(Q)$ deformation of $\mathcal{F}(P)$
 $\Leftrightarrow$ same edge-directions, but different lengths

Definition

Deformation cone: $\mathbb{DC}(\mathcal{F}) = \{\mathcal{G} ; \mathcal{G} \text{ same graph and edge-dir } \mathcal{F}\}$



Parametrization:

edge-length vector:

$$\ell = (\ell_e)_{e \text{ edge}}$$


Edge-length deformation cone

Theorem

Q deformation of $P \Leftrightarrow \mathcal{F}(Q)$ deformation of $\mathcal{F}(P)$
 $\Leftrightarrow$ same edge-directions, but different lengths

Definition

Deformation cone: $\mathbb{DC}(\mathcal{F}) = \{\mathcal{G} ; \mathcal{G} \text{ same graph and edge-dir } \mathcal{F}\}$



Parametrization:

edge-length vector:

$$\ell = (\ell_e)_{e \text{ edge}}$$


Edge-length deformation cone

Theorem

Q deformation of $P \Leftrightarrow \mathcal{F}(Q)$ deformation of $\mathcal{F}(P)$
 $\Leftrightarrow$ same edge-directions, but different lengths

Definition

Deformation cone: $\mathbb{DC}(\mathcal{F}) = \{\mathcal{G} ; \mathcal{G} \text{ same graph and edge-dir } \mathcal{F}\}$



Parametrization:

edge-length vector:

$$\ell = (\ell_e)_{e \text{ edge}}$$

Cycle equations:

linear equations on ℓ

$$\ell_e \geq 0 \text{ for all } e \text{ edge}$$


Edge-length deformation cone

Theorem

Q deformation of $P \Leftrightarrow \mathcal{F}(Q)$ deformation of $\mathcal{F}(P)$
 $\Leftrightarrow$ same edge-directions, but different lengths

Definition

Deformation cone: $\mathbb{DC}(\mathcal{F}) = \{\mathcal{G} ; \mathcal{G} \text{ same graph and edge-dir } \mathcal{F}\}$



Parametrization:

edge-length vector:


$$\ell = (\ell_e)_{e \text{ edge}}$$

Cycle equations:

linear equations on ℓ

$$\ell_e \geq 0 \text{ for all } e \text{ edge}$$

Cycle equations



For a cycle in $\mathcal{F}$:

$$\sum_{e \in \text{cycle}} \mathbf{u}_e = \mathbf{0} \quad , \quad \mathbf{u}_e \text{ edge vector}$$

hence

$$\sum_e \ell_e \mathbf{u}_e = \mathbf{0}$$

Summary on $\mathbb{DC}(P)$

$\mathbb{DC}(P)$		
Q	ℓ	h
Minkowski summands	edge-lengths	heights on rays
$Q_1 + Q_2$	$\ell_1 + \ell_2$	$h_1 + h_2$
Dilation λQ	$\lambda \ell$	λh
Translations	Has been fixed	Lineal
<i>complicated</i>	edge directions Cycle equations V -description	normal fan $\mathcal{N}_P$ Wall-crossing ineq. H -description
Polytope algebra	Weight algebra	Polynomial algebra

$\mathbb{DC}(P)$ is a ray = P Minkowski indecomposable

$\mathbb{DC}(P)$ is simplicial cone = P has **unique** Minkowski decomposition

Graph of edge dependencies

Cycle equations of a triangle, quadrilateral


Goal: Solve quickly $\{\text{cycle equations}\} \cap \{\text{positive variables}\}$.

Cycle equations of a triangle, quadrilateral

Goal: Solve quickly $\{\text{cycle equations}\} \cap \{\text{positive variables}\}$.

Cycle equations of a triangle, quadrilateral

Goal: Solve quickly $\{\text{cycle equations}\} \cap \{\text{positive variables}\}$.



Triangle:


3 variables (= lengths of the edges)

2 equations (= 1 equation in dimension 2)

$\Rightarrow \dim(\text{space of solutions}) = 1$

Cycle equations of a triangle, quadrilateral

Goal: Solve quickly $\{\text{cycle equations}\} \cap \{\text{positive variables}\}$.



Triangle:

3 variables (= lengths of the edges)


2 equations (= 1 equation in dimension 2)

$\Rightarrow \dim(\text{space of solutions}) = 1$

$\Rightarrow$ Indecomposable

Cycle equations of a triangle, quadrilateral

Goal: Solve quickly $\{\text{cycle equations}\} \cap \{\text{positive variables}\}$.



Triangle:

3 variables (= lengths of the edges)

2 equations (= 1 equation in dimension 2)


$\Rightarrow \dim(\text{space of solutions}) = 1$

$\Rightarrow$ Indecomposable

$\Rightarrow$ If I know the length of 1 edge, I know the length of the others.

Cycle equations of a triangle, quadrilateral

Goal: Solve quickly $\{\text{cycle equations}\} \cap \{\text{positive variables}\}$.



Triangle:

3 variables (= lengths of the edges)

2 equations (= 1 equation in dimension 2)


$\Rightarrow \dim(\text{space of solutions}) = 1$

$\Rightarrow$ Indecomposable

$\Rightarrow$ If I know the length of 1 edge, I know the length of the others.

Cycle equations of a triangle, quadrilateral

Goal: Solve quickly $\{\text{cycle equations}\} \cap \{\text{positive variables}\}$.



Triangle:

3 variables (= lengths of the edges)

2 equations (= 1 equation in dimension 2)


$\Rightarrow \dim(\text{space of solutions}) = 1$

$\Rightarrow$ Indecomposable

$\Rightarrow$ If I know the length of 1 edge, I know the length of the others.

Cycle equations of a triangle, quadrilateral

Goal: Solve quickly $\{\text{cycle equations}\} \cap \{\text{positive variables}\}$.



Triangle:

3 variables (= lengths of the edges)

2 equations (= 1 equation in dimension 2)

$\Rightarrow \dim(\text{space of solutions}) = 1$

$\Rightarrow$ Indecomposable


$\Rightarrow$ If I know the length of 1 edge, I know the length of the others.

Graph of edge dependencies


Graph of edge dependencies $ED(\mathcal{F})$:

nodes: edges of framework $\mathcal{F}$


arcs: link two *dependent* edges, i.e. I can deduce the length of one from the length of the other, using the cycle equations



(a) Triangle




(b) Quadrilateral




(c) Trapezoid




(d) Parallelogram




(e) Cube



(f) Prism



(g) Hemicube



(h) Pyramid

Problem 1:

How to find arcs of $ED(\mathcal{F})$.

The two problems

Problem 1:

How to find arcs of $ED(\mathcal{F})$.

Problem 2:

If I have enough arcs in $ED(\mathcal{F})$, what can I conclude on $\mathbb{DC}(\mathcal{F})$ or on the indecomposability of $\mathcal{F}$?

The two problems

Problem 1:

How to find arcs of $ED(\mathcal{F})$.

Problem 2:

If I have enough arcs in $ED(\mathcal{F})$, what can I conclude on $\mathbb{DC}(\mathcal{F})$ or on the indecomposability of $\mathcal{F}$?

A Gale's criterion, '54

All 2-faces of P are triangles $\Rightarrow P$ indecomposable.

The two problems

Problem 1:

How to find arcs of $ED(\mathcal{F})$.

Problem 2:

If I have enough arcs in $ED(\mathcal{F})$, what can I conclude on $\mathbb{DC}(\mathcal{F})$ or on the indecomposability of $\mathcal{F}$?

A Gale's criterion, '54

All 2-faces of P are triangles $\Rightarrow P$ indecomposable.



"triangles" $\rightarrow$ arcs in $ED(\mathcal{F}(P))$

"all 2-faces" $\rightarrow$ enough arcs in $ED(\mathcal{F}(P))$

Problem 1: creating edges

In $ED(\mathcal{F})$, two nodes e, f ($\in$ edges of $\mathcal{F}$) are linked if, e.g.:



- in a common triangle (not necessarily a 2-face),



Problem 1: creating edges

In $ED(\mathcal{F})$, two nodes e, f ($\in$ edges of $\mathcal{F}$) are linked if, e.g.:



- in a common triangle (not necessarily a 2-face),
- are opposite non-parallel in a trapezoid,



Problem 1: creating edges

In $ED(\mathcal{F})$, two nodes e, f ($\in$ edges of $\mathcal{F}$) are linked if, e.g.:



- in a common triangle (not necessarily a 2-face),
- belong to a cycle of edges which spans a full-dim space,
- are opposite non-parallel in a trapezoid,



Problem 1: creating edges

In $ED(\mathcal{F})$, two nodes e, f ($\in$ edges of $\mathcal{F}$) are linked if, e.g.:



- in a common triangle (not necessarily a 2-face),
- belong to a cycle of edges which spans a full-dim space,
- belong to an indecomposable sub-graph of $\mathcal{F}$,
- are opposite non-parallel in a trapezoid,



Problem 1: creating edges

In $ED(\mathcal{F})$, two nodes e, f ($\in$ edges of $\mathcal{F}$) are linked if, e.g.:

- in a common triangle (not necessarily a 2-face),
- belong to a cycle of edges which spans a full-dim space,
- belong to an indecomposable sub-graph of $\mathcal{F}$,
- are opposite non-parallel in a trapezoid,
- are in a sub-graph of $\mathcal{F}$ which projects down to indecomposable,





Problem 1: creating edges

In $ED(\mathcal{F})$, two nodes e, f ($\in$ edges of $\mathcal{F}$) are linked if, e.g.:

- in a common triangle (not necessarily a 2-face),
- belong to a cycle of edges which spans a full-dim space,
- belong to an indecomposable sub-graph of $\mathcal{F}$,
- are opposite non-parallel in a trapezoid,
- are in a sub-graph of $\mathcal{F}$ which projects down to indecomposable,
- ... (there is a nicer way to write these properties)

+ implicit edges



Problem 2: consequences of $ED(\mathcal{F})$

Theorem (Padrol–P. '25+)

If $ED(\mathcal{F})$ is connected, then $\mathcal{F}$ is indecomposable

Problem 2: consequences of $ED(\mathcal{F})$

Theorem (Padrol–P. '25+)

If $ED(\mathcal{F})$ is connected, then $\mathcal{F}$ is indecomposable

Dependent $S \subseteq V$: for all $u, v \in S$, there is a path (in graph of $\mathcal{F}$) of dependent edges between u and v .

Theorem (Padrol–P. '25+)

If there exists a set S of dependent vertices such that every facet contains a vertex of S , then $\mathcal{F}$ is indecomposable.

Problem 2: consequences of $ED(\mathcal{F})$

Theorem (Padrol–P. '25+)

If $ED(\mathcal{F})$ is connected, then $\mathcal{F}$ is indecomposable

Dependent $S \subseteq V$: for all $u, v \in S$, there is a path (in graph of $\mathcal{F}$) of dependent edges between u and v .

Theorem (Padrol–P. '25+)

If there exists a set S of dependent vertices such that every facet contains a vertex of S , then $\mathcal{F}$ is indecomposable.

Corollaries: Gale '54, Shepard '63, Kallay '82, McMullen '87, Yost–Przesławski '08 & '16 criteria...

flat: set of vertices inducing a *connected* sub-framework of $\mathcal{F}$

$$\text{dir}_{\mathcal{F}}(F) = \text{lin}\{\phi_u - \phi_v ; u, v \in F\}$$

flat: set of vertices inducing a *connected* sub-framework of $\mathcal{F}$

$$\text{dir}_{\mathcal{F}}(F) = \text{lin}\{\phi_u - \phi_v ; u, v \in F\}$$

A collection of flats $\mathcal{C}$ *pins* a vertex v iff:

$$v \in \bigcap_{F \in \mathcal{C}} F \quad \text{and} \quad \bigcap_{F \in \mathcal{C}} \text{dir}_{\mathcal{F}}(F) = \{\mathbf{0}\}$$

flat: set of vertices inducing a *connected* sub-framework of $\mathcal{F}$

$$\text{dir}_{\mathcal{F}}(F) = \text{lin}\{\phi_u - \phi_v ; u, v \in F\}$$

A collection of flats $\mathcal{C}$ *pins* a vertex v iff:

$$v \in \bigcap_{F \in \mathcal{C}} F \quad \text{and} \quad \bigcap_{F \in \mathcal{C}} \text{dir}_{\mathcal{F}}(F) = \{\mathbf{0}\}$$

Definition

$$S \subseteq V$$

flat: set of vertices inducing a *connected* sub-framework of $\mathcal{F}$

$$\text{dir}_{\mathcal{F}}(F) = \text{lin}\{\phi_u - \phi_v ; u, v \in F\}$$

A collection of flats $\mathcal{C}$ *pins* a vertex v iff:

$$v \in \bigcap_{F \in \mathcal{C}} F \quad \text{and} \quad \bigcap_{F \in \mathcal{C}} \text{dir}_{\mathcal{F}}(F) = \{\mathbf{0}\}$$

Definition

$$S \subseteq V$$

$$S_0 = S$$

$$S_{i+1} = S_i \cup \{v \in V ; v \text{ pinned by collection of flats} \\ \text{having each } \geq 1 \text{ vertex in } S_{i-1}\}$$

flat: set of vertices inducing a *connected* sub-framework of $\mathcal{F}$

$$\text{dir}_{\mathcal{F}}(F) = \text{lin}\{\phi_u - \phi_v ; u, v \in F\}$$

A collection of flats $\mathcal{C}$ *pins* a vertex v iff:

$$v \in \bigcap_{F \in \mathcal{C}} F \quad \text{and} \quad \bigcap_{F \in \mathcal{C}} \text{dir}_{\mathcal{F}}(F) = \{\mathbf{0}\}$$

Definition


$$S \subseteq V$$

$$S_0 = S$$

$$S_{i+1} = S_i \cup \{v \in V ; v \text{ pinned by collection of flats} \\ \text{having each } \geq 1 \text{ vertex in } S_{i-1}\}$$

$$\text{pinning closure: } \text{cl}(S) = \max_i S_i$$

Pinning closure



This is indecomposable

Our strongest theorem

Theorem (Padrol–P. '25+)

If there is $S \subseteq V$, and $X_1, \dots, X_r$ dependent sets of edges with

- *any two vertices of S are connected via $\bigcup_i X_i$, and*
- *$\text{cl}(S) = V$,*

then: $\dim \mathbb{DC}(\mathcal{F}) \leq r$.

Corollary (Padrol–P. '25+)

If there is S such that:

- *S is a dependent set of vertices*
- *$\text{cl}(S) = V$,*

then: $\mathcal{F}$ is indecomposable.

Applications

Theorem (Padrol–P. '25+)

If there is $S \subseteq V$, and $X_1, \dots, X_r$ dependent sets of edges with

- any two vertices of S are connected via $\bigcup_i X_i$, and*
- $\text{cl}(S) = V$,*


then: $\dim \mathbb{DC}(\mathcal{F}) \leq r$.

Theorem (Padrol-P. '25+)


If there is $S \subseteq V$, and $X_1, \dots, X_r$ dependent sets of edges with

- any two vertices of S are connected via $\bigcup_i X_i$, and*
- $\text{cl}(S) = V$,*


then: $\dim \mathbb{DC}(\mathcal{F}) \leq r$.



(a) Triang. cupola




(b) Sum of 2 triangles



(c) Chiseled cube

$\dim \mathbb{DC}(\mathcal{P}) = 2$, only one Minkowski decomposition (in 2 terms)

First applications



$\dim \mathbb{DC}(P) = 2$, this is the only Minkowski decomposition

New indecomposable generalized permutahedra

Generalized permutahedra: edge directions are $\mathbf{e}_i - \mathbf{e}_j$ for $i \neq j$

Edmonds problem '70: find all indecomposable gene. permut.

New indecomposable generalized permutahedra


Generalized permutahedra: edge directions are $\mathbf{e}_i - \mathbf{e}_j$ for $i \neq j$

Edmonds problem '70: find all indecomposable gene. permut.

The *graphical zonotope* of $G = (V, E)$ is: $Z_G = \sum_{i,j \in E} [\mathbf{e}_i, \mathbf{e}_j]$

Theorem (Padrol-P. '25+)

For all complete bipartite graphs $K_{n,m}$ (except $n = m = 2$), it is possible to truncate 1 or 2 vertices of $Z_{K_{n,m}}$ to obtain an indecomposable generalized permutahedron.



New indecomposable generalized permutahedra


Generalized permutahedra: edge directions are $\mathbf{e}_i - \mathbf{e}_j$ for $i \neq j$

Edmonds problem '70: find all indecomposable gene. permut.

The *graphical zonotope* of $G = (V, E)$ is: $Z_G = \sum_{i,j \in E} [\mathbf{e}_i, \mathbf{e}_j]$

Theorem (Padrol-P. '25+)

For all complete bipartite graphs $K_{n,m}$ (except $n = m = 2$), it is possible to truncate 1 or 2 vertices of $Z_{K_{n,m}}$ to obtain an indecomposable generalized permutahedron.



This gives $2 \left\lfloor \frac{n-1}{2} \right\rfloor$ new indecomposable gene. permutahedra

New indecomposable generalized permutahedra


Generalized permutahedra: edge directions are $\mathbf{e}_i - \mathbf{e}_j$ for $i \neq j$

Edmonds problem '70: find all indecomposable gene. permut.

The *graphical zonotope* of $G = (V, E)$ is: $Z_G = \sum_{i,j \in E} [\mathbf{e}_i, \mathbf{e}_j]$

Theorem (Padrol-P. '25+)

For all complete bipartite graphs $K_{n,m}$ (except $n = m = 2$), it is possible to truncate 1 or 2 vertices of $Z_{K_{n,m}}$ to obtain an indecomposable generalized permutahedron.



This gives $2 \left\lfloor \frac{n-1}{2} \right\rfloor$ new indecomposable gene. permutahedra

Loho-Padrol-P.'25: we create $2^{2^{n-2}}$ indecomposable gene. permut.

Corollary (Padrol–P. '25+)

If all 2-faces of P are triangles and parallelograms, then there is a unique way to write P as a sum of indecomposable polytopes (and we know the which ones).

Corollary (Padrol-P. '25+)

If all 2-faces of P are triangles and parallelograms, then there is a unique way to write P as a sum of indecomposable polytopes (and we know the which ones).

0/1-polytope: polytopes whose vertices have coordinates 0s and 1s.

Corollary

Let P be a 0/1-polytope.

There is a unique way to write P as a Minkowski sum of indecomposable polytopes: it is a product of 0/1-polytopes

Corollary (Padrol–P. '25+)

If all 2-faces of P are triangles and parallelograms, then there is a unique way to write P as a sum of indecomposable polytopes (and we know the which ones).

0/1-polytope: polytopes whose vertices have coordinates 0s and 1s.

Corollary

Let P be a 0/1-polytope.

There is a unique way to write P as a Minkowski sum of indecomposable polytopes: it is a product of 0/1-polytopes

Easy proof: matroid pol. indecomposable $\Leftrightarrow$ matroid connected


See Higashitani–Padrol–Sanyal for more (soon to come).

Platonic, Archimedean, and Jhonson solids


Platonic solids: 3-dim., 1 type of regular polygons, vertex transitive

Archimedean solids: 3-dim., regular polygons, vertex transitive


Jhonson solids: 3-dim., regular polygons




(a) J_6




(b) J_{32}



(c) J_{33}



(d) J_{34}




(e) A_8

Platonic, Archimedean, and Jhonson solids


Platonic solids: 3-dim., 1 type of regular polygons, vertex transitive

Archimedean solids: 3-dim., regular polygons, vertex transitive


Jhonson solids: 3-dim., regular polygons




(a) J_6




(b) J_{32}



(c) J_{33}



(d) J_{34}



(e) A_8

Theorem (P. '26+)

$\dim \mathbb{DC}(P)$ for all Platonic, Archimedean, and Jhonson solids:
2 infinite families + 110 sporadic solids

Products, parallelogramic Minkowski sums

Theorem (Padrol–P. '25+)

$$\mathbb{DC}(P \times Q) \simeq \mathbb{DC}(P) \times \mathbb{DC}(Q)$$

Products, parallelogramic Minkowski sums


Theorem (Padrol–P. '25+)

$$\mathbb{DC}(P \times Q) \simeq \mathbb{DC}(P) \times \mathbb{DC}(Q)$$


P, Q are in *parallelogramic position* if no edge of P is in a plane spanned by the 2-faces of Q

Theorem (Padrol–P. '25+)

P, Q in *parallelogramic position*: $\mathbb{DC}(P + Q) \simeq \mathbb{DC}(P) \times \mathbb{DC}(Q)$.



(a) Diminished trapezohedron




(b) Gyrobifastigium

Other applications:

- “autonomous & dependent” parts of $ED(\mathcal{F})$ give rays of $\mathbb{DC}(\mathcal{F})$
- edge decomposability implies unique decomposability property
- stack/delete vertices on polytopes, then study indecomposability
- you can deal with zonotopes whose 2-faces are parallelograms
- accelerate the computation $\mathbb{DC}(\mathcal{F})$ using combinatorial data
- re-write $\mathbb{DC}(P)$ as a 2-dimensional problem
- ...

Remark:

Be careful, $ED(\mathcal{F})$ is not always sufficient for recovering $\mathbb{DC}(\mathcal{F})$.



Merci !

Thank you!

Bonus slides...

What else do I do?

Submodular cone: “Many rays of the submodular cone”

Linear programming: “Vertices of the monotone path polytopes of hypersimplices” & “Pivot polytopes of products of simplices and shuffles of associahedra”

Random polytopes: “Unimodality of the number of paths per length on polytopes: Examples, counter-examples, and central limit theorem”

Ehrhart theory, triangulations: “Ehrhart non-positivity and unimodular triangulations for classes of s-lecture hall simplices”

3D polytopes, framing lattices, polytope/weight algebra, hypergraphs, Gray codes, hypergraphs...

Anything that might contain a polytope somewhere is interesting !!