



Theoretical Physics

---

# Diagrammatic Monte Carlo

---

Blomquist, Emil  
emilbl@kth.se

Master of Science Thesis  
Supervisor: Egor Babaev

July 18, 2017

## Abstract

...

## Sammanfattning

...

Throughout this text  $\hbar = 1$  has been used.

# Contents

<b>1</b>	<b>Introduction</b>	<b>5</b>
1.1	Many-particle physics . . . . .	5
<b>2</b>	<b>Background Material</b>	<b>6</b>
2.1	A few words about finite temperature formalism . . . . .	6
2.1.1	Green's function . . . . .	6
2.2	Derivation of the Fröhlich Hamiltonian . . . . .	6
2.2.1	Classical picture . . . . .	6
2.2.2	Quantisation . . . . .	11
2.3	Feynman diagrams . . . . .	13
2.4	Dyson equation . . . . .	13
2.5	Monte Carlo Simulation . . . . .	13
2.5.1	Detailed balance and ergodicity . . . . .	13
2.5.2	Metropolis-Hastings algorithm . . . . .	13
<b>3</b>	<b>Diagrammatic Monte Carlo</b>	<b>14</b>
3.1	General idea . . . . .	14
3.2	Generalisation of stochastic integration (better name needed) . . . . .	14
3.2.1	Example 1 . . . . .	14
3.2.2	Example 2 . . . . .	14
3.2.3	Example 3 . . . . .	14
3.3	Polaron implementation . . . . .	14
3.3.1	Update function 1 . . . . .	15
3.3.2	Update function 2 . . . . .	15
3.3.3	Update function 3 . . . . .	15
3.4	Sample self energy . . . . .	15
3.4.1	Divergent diagram . . . . .	15
3.5	Boldification . . . . .	15
<b>4</b>	<b>Results</b>	<b>16</b>
4.1	Analytical predictions . . . . .	16
4.2	Numerics . . . . .	16
4.3	Discussion . . . . .	16
<b>5</b>	<b>Summary and Conclusions</b>	<b>17</b>

<b>Appendix A (make this go away, also Appendix A -&gt; Appendix)</b>	<b>18</b>
A.1 Heavy theory from the finite temperature formalism . . . . .	18
A.2 Derivation of the finite temperature electron Green's function . . . . .	18
A.3 Derivation of the finite temperature phonon Green's function . . . . .	18

# Chapter 1

## Introduction

Here we shall summarise what this chapter is about

### 1.1 Many-particle physics

# Chapter 2

## Background Material

Here we shall summarise what this chapter is about

### 2.1 A few words about finite temperature formalism

#### 2.1.1 Green's function

fit to exponential for large  $\tau$ 's etc.

G and D derived in the appendix

### 2.2 Derivation of the Fröhlich Hamiltonian

What will happen to a dielectric medium if one introduces a charged particle, and how will this particle react to changes in the dielectric medium? These are questions which Herbert Fröhlich answered in his 1954 paper *Electrons in lattice fields*[1]. In this section we will outline the original derivation but also look into some details relevant to us.

#### 2.2.1 Classical picture

The idea is that we introduce a free electron into a dielectric crystal, e.g. rock salt. This electron will deform the lattice and hence polarise the crystal. This deformation may be described by a displacement vector for each lattice point. However, to simplify matters, we treat this displacement vector as a continuous vector field.

If we first consider a steady state situation, the polarisation is determined solely by the dielectric permittivity  $\epsilon_r$ . Introducing the electric displacement field  $\mathbf{D} = \mathbf{E} + 4\pi\mathbf{P}$  (CGS units) for which the source is the free electron, this may be thought of as the external

electrical field. If our electron is at position  $\mathbf{r}_e$ , the  $\mathbf{D}$ -field at a position  $r$  is given by the usual

$$\mathbf{D}(\mathbf{r}, \mathbf{r}_e) = -\nabla \frac{Q}{|\mathbf{r} - \mathbf{r}_e|} \quad (2.1)$$

so that

$$\nabla \cdot \mathbf{D}(\mathbf{r}, \mathbf{r}_e) = 4\pi q \delta(\mathbf{r} - \mathbf{r}_e). \quad (2.2)$$

Here  $Q$  is the charge of the electron and the second equality is to be thought of as valid inside of an integral. In agreement with the definition of the  $\mathbf{D}$ -field we may also define the electric field due to the bound charges as  $\mathbf{E}_{\text{bound}} = -4\pi\mathbf{P}$  since  $\mathbf{D} = \mathbf{E} + 4\pi\mathbf{P} = \mathbf{E} - \mathbf{E}_{\text{bound}} \equiv \mathbf{E}_{\text{free}}$ .

The interaction energy, which is minimised when the  $\mathbf{D}$ -field is parallel to the polarisation, is given by (why!? Dipole something?)

$$E_{\text{int}} = - \int_V \mathbf{D}(\mathbf{r}, \mathbf{r}_e) \cdot \mathbf{P}(\mathbf{r}) d^3r. \quad (2.3)$$

By utilising the scalar potential of the bound electric field  $\mathbf{E}_{\text{bound}}(r) = -\nabla\Phi(\mathbf{r})$  along with the quantities defined above, this interaction energy may be rewritten as

$$\begin{aligned} E_{\text{int}} &= -\frac{1}{4\pi} \int_{\Omega} \mathbf{D}(\mathbf{r}, \mathbf{r}_e) \cdot \nabla\Phi(\mathbf{r}) d^3r \\ &= -\frac{1}{4\pi} \int_{\Omega \setminus V_\varepsilon} \mathbf{D}(\mathbf{r}, \mathbf{r}_e) \cdot \nabla\Phi(\mathbf{r}) d^3r - \frac{1}{4\pi} \int_{V_\varepsilon} \mathbf{D}(\mathbf{r}, \mathbf{r}_e) \cdot \nabla\Phi(\mathbf{r}) d^3r \\ &= -\frac{1}{4\pi} \int_{\Omega \setminus V_\varepsilon} \nabla \cdot [\Phi(\mathbf{r})\mathbf{D}(\mathbf{r}, \mathbf{r}_e)] d^3r + \underbrace{\frac{1}{4\pi} \int_{\Omega \setminus V_\varepsilon} \Phi(\mathbf{r}) \nabla \cdot \mathbf{D}(\mathbf{r}, \mathbf{r}_e) d^3r}_0 + \underbrace{\frac{q}{4\pi} \int_{V_\varepsilon} \frac{1}{r^2} \frac{\partial\Phi(\mathbf{r} + \mathbf{r}_e)}{\partial r} d^3r}_{\rightarrow 0 \text{ as } \varepsilon \rightarrow 0} \\ &= \frac{Q}{4\pi} \int_{-\partial V'_\varepsilon} \Phi(\mathbf{r} + \mathbf{r}_e) \nabla \frac{1}{r} \cdot d^2\mathbf{r} \\ &= Q \Phi(\mathbf{r}_e) \end{aligned} \quad (2.4)$$

Here we have both assumed that  $\Phi(\mathbf{r})$  has a continuous first derivative as well as  $\Phi(\mathbf{r})\mathbf{D}(\mathbf{r}, \mathbf{r}_e) \simeq 0$  on the boundary  $\mathbf{r} \in \partial\Omega$ . In order to use the divergence theorem the singularity at  $\mathbf{r}_e$  has been isolated in a spherical volume  $V_\varepsilon$  centered at  $\mathbf{r}_e$  with radius  $\varepsilon \rightarrow 0$ .

Next we bring back the time dependence and no longer consider a situation which is in steady state. Since each lattice point in our crystal is occupied by a ion, the dynamic in our system is characterised by two time scales. That is, the time it takes to displace the bound electrons relative their nuclei (deformation of ion) and the time it takes to displace



the nucleus relative the lattice (deformation of lattice structure). Since the electrons are significantly lighter than their nucleus, the ion deformation time should thus be much smaller than the time it takes to deformation the lattice. Denoting these times as  $t_{\text{uv}}$  and  $t_{\text{ir}}$  respectively we thus have that  $t_{\text{uv}} \ll t_{\text{ir}}$  which corresponds to  $\omega_{\text{uv}} \gg \omega_{\text{ir}}$ . The subscripts of course indicate that frequencies lie in the ultraviolet and infrared region respectively. The total deformation is due to these two deformation so that  $\mathbf{P} = \mathbf{P}_{\text{ir}} + \mathbf{P}_{\text{uv}}$ .




Figure 2.1: Illustration of the lattice and the two types of deformation. In the two images with polarisation, depict the cumulative polarisation as well. Also there should be subcaptions!

It is further reasonable to assume that each of these contributions behave as a driven harmonic oscillator, that is

$$\ddot{\mathbf{P}}_{\text{ir}}(\mathbf{r}) + \omega_{\text{ir}}^2 \mathbf{P}_{\text{ir}}(\mathbf{r}) = \frac{\mathbf{D}(\mathbf{r}, \mathbf{r}_e)}{\gamma}, \quad \ddot{\mathbf{P}}_{\text{uv}}(\mathbf{r}) + \omega_{\text{uv}}^2 \mathbf{P}_{\text{uv}}(\mathbf{r}) = \frac{\mathbf{D}(\mathbf{r}, \mathbf{r}_e)}{\delta}. \quad (2.5)$$

Here  $\gamma$  and  $\delta$  are constants to be determined in what follows. First we must once again quickly consider a static situation. Here the static dielectric constant  $\epsilon$  would give us  $\mathbf{D} = \epsilon \mathbf{E}$  which we could use together with the previously stated definition of the  $\mathbf{D}$ -field to relate  $\mathbf{P}$  and  $\mathbf{D}$  as

$$4\pi \mathbf{P}(\mathbf{r}) = (1 - 1/\epsilon) \mathbf{D}(\mathbf{r}, \mathbf{r}_e). \quad (2.6)$$

Next we once again turn back time and utilise the high frequency dielectric constant  $\epsilon_\infty$ , defined by  $\mathbf{D} = \epsilon_\infty \mathbf{E}$ , under the assumption that the frequency  $\omega_\infty$  of the external field satisfies  $\omega_{uv} \gg \omega_\infty \gg \omega_{ir}$ . The latter is simply implying that the free electron in our crystal is moving slowly (Fast enough for the IR polarisation not to follow, right!?). This tells us that the UV-part of the polarisation will follow the changes in the  $\mathbf{D}$ -field nearly adiabatically whilst the IR-part wont have time adapt to the changes and may thus be thought of as a negligible constant field in comparison, that is,  $\mathbf{P} \simeq \mathbf{P}_{uv}$ . Using this together with the relation of  $\mathbf{D}$  and  $\mathbf{E}$  through  $\epsilon_\infty$  we find, in similarity to (2.6),

$$4\pi \mathbf{P}_{uv}(\mathbf{r}) = (1 - 1/\epsilon_\infty) \mathbf{D}(\mathbf{r}, \mathbf{r}_e). \quad (2.7)$$

Taking the difference between (2.6) and (2.7) we find a similar equation for  $\mathbf{P}_{ir}$

$$4\pi \mathbf{P}_{ir}(\mathbf{r}) = (1/\epsilon_\infty - 1/\epsilon) \mathbf{D}(\mathbf{r}, \mathbf{r}_e). \quad (2.8)$$

Noting that  $\ddot{\mathbf{P}}_{uv} \approx \omega_\infty^2 \mathbf{P}_{uv} \ll \omega_{uv}^2 \mathbf{P}_{uv}$  and assuming that  $\mathbf{P}_{ir}$  is seemingly constant at timescales of length  $1/\omega_{ir}$ , equation (2.9) simplifies into (in the original derivation they simply compared using steady state where the time derivatives obviously are zero, but this must be equivalent due to  $\omega_{uv} \gg \omega_\infty \gg \omega_{ir}$ ?)

$$\omega_{ir}^2 \mathbf{P}_{ir}(\mathbf{r}) = \frac{\mathbf{D}(\mathbf{r}, \mathbf{r}_e)}{\gamma}, \quad \omega_{uv}^2 \mathbf{P}_{uv}(\mathbf{r}) = \frac{\mathbf{D}(\mathbf{r}, \mathbf{r}_e)}{\delta}. \quad (2.9)$$

Comparing this equation to (2.7) and (2.8) we find the value of  $\gamma$  and  $\delta$  in terms of know material properties

$$\frac{1}{\gamma} = \frac{\omega_{ir}^2}{4\pi} \left( \frac{1}{\epsilon_\infty} - \frac{1}{\epsilon} \right), \quad \frac{1}{\delta} = \frac{\omega_{uv}^2}{4\pi} \left( 1 - \frac{1}{\epsilon_\infty} \right). \quad (2.10)$$

Having obtained the equation of motion for the two types of polarisation, our next task is to find the one for the free electron. Since the electron had low in kinetic energy, we may without hesitation treat it in the non relativistic limit. The only force acting on the electron is due to the bound electric field, that is  $\mathbf{F}(\mathbf{r}_e) = q \mathbf{E}_{\text{bound}}(\mathbf{r}_e) = -q \nabla \Phi(\mathbf{r}_e)$ . Thus  $m \ddot{\mathbf{r}}_e = \dot{\mathbf{p}}_e = -q \nabla \Phi(\mathbf{r}_e)$  where  $m$  is an effective mass which in general is different from the electronic mass  $m_e$ . By treating  $\mathbf{p}_e$ ,  $\mathbf{P}_{uv}(\mathbf{r})$  and  $\mathbf{P}_{ir}(\mathbf{r})$  as variables, we may

by combining all the equation of motions derived above together with expression for the interaction energy (2.4) form the Lagrangian

$$L = \frac{\gamma}{2} \int \left[ \dot{\mathbf{P}}_{\text{ir}}^2(\mathbf{r}) - \omega_{\text{ir}}^2 \mathbf{P}_{\text{ir}}^2(\mathbf{r}) \right] d^3r + \frac{\delta}{2} \int \left[ \dot{\mathbf{P}}_{\text{uv}}^2(\mathbf{r}) - \omega_{\text{uv}}^2 \mathbf{P}_{\text{uv}}^2(\mathbf{r}) \right] d^3r + \int \mathbf{D}(\mathbf{r}, \mathbf{r}_e) \cdot [\mathbf{P}_{\text{ir}}(\mathbf{r}) + \mathbf{P}_{\text{uv}}(\mathbf{r})] d^3r + \frac{\mathbf{P}_e^2}{2m} \quad (2.11)$$

and by a Legendre transformation, the Hamiltonian

$$H = \frac{\gamma}{2} \int \left[ \dot{\mathbf{P}}_{\text{ir}}^2(\mathbf{r}) + \omega_{\text{ir}}^2 \mathbf{P}_{\text{ir}}^2(\mathbf{r}) \right] d^3r + \frac{\delta}{2} \int \left[ \dot{\mathbf{P}}_{\text{uv}}^2(\mathbf{r}) + \omega_{\text{uv}}^2 \mathbf{P}_{\text{uv}}^2(\mathbf{r}) \right] d^3r - \int \mathbf{D}(\mathbf{r}, \mathbf{r}_e) \cdot [\mathbf{P}_{\text{ir}}(\mathbf{r}) + \mathbf{P}_{\text{uv}}(\mathbf{r})] d^3r + \frac{\mathbf{P}_e^2}{2m}. \quad (2.12)$$

Since the UV-part of the polarisation follows the external field nearly adiabatically, the dynamic solution  $\mathbf{P}_{\text{uv}}$  is approximatively the statical one with respect to current  $\mathbf{D}$ -field. Because of this  $\mathbf{P}_{\text{uv}}$  will only contribute to the Hamiltonian with a constant and may thus be removed from the theory giving rise only to an energy shift.

Next we observe that the integrals in the expressions above are divergent since  $\mathbf{P}_{\text{ir}}$  according to (2.7) and (2.8) are proportional to the  $\mathbf{D}$ -field which is singular at  $\mathbf{r} = \mathbf{r}_e$ . However this divergence is a resulted from us disregarding the atomic structure when interpreting the displacements as a continuous vector field. To resolve this one should express the polarisation in a Fourier series and omit terms with a wave length smaller than that of the lattice constant. Hence we need not to worry about this divergence.

Further more it is useful to introduce here a complex vector field  $\mathbf{B}$ ,

$$\mathbf{B}(\mathbf{r}) = \sqrt{\frac{\gamma\omega_{\text{ir}}}{2}} \left[ \mathbf{P}_{\text{ir}}(\mathbf{r}) + \frac{i}{\omega_{\text{ir}}} \dot{\mathbf{P}}_{\text{ir}}(\mathbf{r}) \right], \quad \mathbf{B}^\dagger(\mathbf{r}) = \sqrt{\frac{\gamma\omega_{\text{ir}}}{2}} \left[ \mathbf{P}_{\text{ir}}(\mathbf{r}) - \frac{i}{\omega_{\text{ir}}} \dot{\mathbf{P}}_{\text{ir}}(\mathbf{r}) \right]. \quad (2.13)$$

Here we have also dropped the subscript of  $\omega_{\text{ir}}$  since  $\omega_{\text{uv}}$  no longer appears in our theory. The reasons to why one prefer to work with  $\mathbf{B}$  rather than  $\mathbf{P}_{\text{ir}}$  will be evident later. With this our now non-divergent Hamiltonian becomes,

$$H = \frac{1}{2} m \dot{\mathbf{r}}_e^2 + \omega \int \mathbf{B}^\dagger(\mathbf{r}) \cdot \mathbf{B}(\mathbf{r}) d^3r - \sqrt{\frac{1}{2\gamma\omega}} \int \mathbf{D}(\mathbf{r}, \mathbf{r}_e) \cdot [\mathbf{B}(\mathbf{r}) + \mathbf{B}^\dagger(\mathbf{r})] d^3r. \quad (2.14)$$

Using a periodic boundary condition over a cube of volume  $V = l^3$ ,  $\mathbf{B}$  might be expressed as

$$\mathbf{B}(\mathbf{r}) = \sum_{\mathbf{q}} \frac{\mathbf{q}}{q} b_{\mathbf{q}} \frac{e^{i\mathbf{q}\cdot\mathbf{r}}}{\sqrt{V}}, \quad \mathbf{B}^\dagger(\mathbf{r}) = \sum_{\mathbf{q}} \frac{\mathbf{q}}{q} b_{\mathbf{q}}^\dagger \frac{e^{-i\mathbf{q}\cdot\mathbf{r}}}{\sqrt{V}} \quad (2.15)$$

with

$$\mathbf{q}_i = \frac{2\pi}{l} n_i; \quad n_i = 0, \pm 1, \pm 2, \dots, \pm N_{\text{cut}} \quad (2.16)$$

Here  $i = x, y, z$ ,  $|\mathbf{q}| = q$  and the cutoff magnitude  $q_{\text{cut}} = 2\pi N_{\text{cut}}/l$  is inversely proportional to the lattice constant in accordance to what was discussed earlier ( $a \lesssim \lambda = 2\pi/k \rightarrow k \lesssim a$  right!?) (mention  $q_{\text{cut}}$  since we are only looking at the IR-part). Also defining a scalar potential for the IR-part of the polarisation as  $4\pi\mathbf{P}_{\text{ir}} = \nabla\Phi_{\text{ir}}$  we may, by using equation (2.4), express the interaction energy between  $\mathbf{D}$  and  $\mathbf{P}_{\text{ir}}$  as  $Q\Phi_{\text{ir}}(\mathbf{r}_e)$ . Using (2.13, 2.15) together with  $\Phi_{\text{ir}}$  the expression for the Hamiltonian becomes

$$H = \frac{\mathbf{p}_e^2}{2m} + \omega \sum_{\mathbf{q}} b_{\mathbf{q}}^\dagger b_{\mathbf{q}} + i 4\pi Q \sqrt{\frac{1}{2\gamma\omega V}} \sum_{\mathbf{q}} \frac{1}{q} (b_{\mathbf{q}}^\dagger e^{-i\mathbf{q}\cdot\mathbf{r}_e} - b_{\mathbf{q}} e^{i\mathbf{q}\cdot\mathbf{r}_e}). \quad (2.17)$$

Introduce  $V(q)$  as well as  $\alpha$ .

Here it seems as if the interaction part of the Hamiltonian is ill behaved due to the  $\mathbf{q} = \mathbf{0}$  term of the sum. This is a result from the periodic boundary condition imposed on our system. Originally the system contained only one free electron. With a periodic boundary condition however, the system now consists of an infinite number of copies of the cube with volume  $V$ , each containing a free electron. These electrons interact with one another due to the long range Coulomb repulsion. To avoid this behaviour when periodically continuing the system, one should introduce in each cube a charge density of  $-Q/V$ . Doing this would give rise to an attractive Coulomb force which for each electron would cancel the Coulomb repulsion. Such an interaction would exactly cancel the ill behaved  $\mathbf{q} = \mathbf{0}$  term in the interaction part of the Hamiltonian since the only difference would be the sign of the charge, i.e  $Q \rightarrow -Q$ .

## 2.2.2 Quantisation

Having sorted out these details we are ready to quantise the Hamiltonian. To do so the first step is to symmetrise  $b_{\mathbf{q}}^\dagger b_{\mathbf{q}} = (b_{\mathbf{q}}^\dagger b_{\mathbf{q}} + b_{\mathbf{q}} b_{\mathbf{q}}^\dagger)/2$  (why? source needed.). Next up, using[2]

$$\dot{q} = \frac{\delta H}{\delta p}, \quad \dot{p} = -\frac{\delta H}{\delta q} \quad (2.18)$$

together with (2.9, 2.12) it is easy to verify that  $\mathbf{P}_{\text{ir}}$  and  $\gamma\dot{\mathbf{P}}_{\text{ir}}$  or equivalent,  $(b_{\mathbf{q}}^\dagger + b_{\mathbf{q}})/\sqrt{2\gamma\omega}$  and  $i(b_{\mathbf{q}}^\dagger - b_{\mathbf{q}})\sqrt{\gamma\omega/2}$  for each  $\mathbf{q}$ , are conjugate variables (how do we arrive at the final

set of conjugate variables? Using some sort of transformation perhaps? Try find in the book by Egor.). By promoting to operators  $b_{\mathbf{q}} \rightarrow \hat{b}_{\mathbf{q}}$ ,  $\mathbf{r}_e \rightarrow \hat{\mathbf{r}}_e$ ,  $\mathbf{p}_e \rightarrow \hat{\mathbf{p}}_e$  and imposing Bose statistics (How do we know this? spinless?) onto the conjugate variables,

$$i\delta_{\mathbf{q},\mathbf{k}} = \frac{i}{2}[\hat{b}_{\mathbf{q}}^\dagger + \hat{b}_{\mathbf{q}}, \hat{b}_{\mathbf{k}}^\dagger - \hat{b}_{\mathbf{k}}]/2 \quad (2.19)$$

we obtain  $[b_{\mathbf{q}}, b_{\mathbf{k}}^\dagger] = \delta_{\mathbf{q},\mathbf{k}}$  and  $[b_{\mathbf{q}}, b_{\mathbf{k}}] = 0$  (really, they follow?). Using the first of these to rewrite  $(\hat{b}_{\mathbf{q}}^\dagger \hat{b}_{\mathbf{q}} + \hat{b}_{\mathbf{q}} \hat{b}_{\mathbf{q}}^\dagger)/2 = \hat{b}_{\mathbf{q}}^\dagger \hat{b}_{\mathbf{q}} + \frac{1}{2}$  the Hamiltonian, in quantised form, becomes

$$\hat{H} = \frac{\hat{\mathbf{p}}_e^2}{2m} + \omega \sum_{\mathbf{q}} \hat{b}_{\mathbf{q}}^\dagger \hat{b}_{\mathbf{q}} + i4\pi Q \sqrt{\frac{1}{2\gamma\omega V}} \sum_{\mathbf{q}} \frac{1}{q} \left( \hat{b}_{\mathbf{q}}^\dagger e^{-i\mathbf{q}\cdot\hat{\mathbf{r}}_e} - \hat{b}_{\mathbf{q}} e^{i\mathbf{q}\cdot\hat{\mathbf{r}}_e} \right). \quad (2.20)$$

Here we have omitted the zero point energy  $\omega \sum_{\mathbf{q}} \frac{1}{2}$  of the oscillating ions. At this stage it has become evident that  $\hat{b}_{\mathbf{q}}^\dagger$  and  $\hat{b}_{\mathbf{q}}$  may be recognised[1] as the creation and annihilation operator of a harmonic oscillator of mass  $M$ , with position and momentum coordinates

$$\hat{q}_{\mathbf{k}} = \sqrt{\frac{1}{2M\omega}} (\hat{b}_{\mathbf{k}}^\dagger + \hat{b}_{\mathbf{k}}), \quad \hat{p}_{\mathbf{k}} = i\sqrt{\frac{M\omega}{2}} (\hat{b}_{\mathbf{k}}^\dagger - \hat{b}_{\mathbf{k}}). \quad (2.21)$$

The polarisation field  $\mathbf{P}_{\text{ir}}$  is in this manner being represented by a set of harmonic oscillators, all of them having the same frequency  $\omega$ .

Having sorted out the quantisation of the  $\mathbf{B}$ -field we turn our attention towards the electron. In order to later expand the interacting theory using Feynman diagrams, we wish to represent the free electron by a many-body Fock state instead of having it represented by a single-body Hilbert state. In order to do this the second quantisation formalism need to be utilised. Using the momentum basis as the electronic many-body Fock occupation state with corresponding creation and annihilation operators  $\hat{a}_{\mathbf{k}}^\dagger$  and  $\hat{a}_{\mathbf{k}}$  respectively, the translation of a one-body operator from first-quantisation to second-quantisation is[3]

$$\hat{O}_2 = \sum_{\mathbf{k},\mathbf{p}} \langle \mathbf{k} | O_1 | \mathbf{p} \rangle \hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{p}}. \quad (2.22)$$

Here  $\hat{O}_2$  is the second-quantised operator and  $O_1$  is the corresponding first-quantised operator sandwiched between the single-particle Hilbert states  $|\mathbf{k}\rangle$ ,  $|\mathbf{p}\rangle$ . Mention that these momenta once again are discretised according to . Thus

$$\frac{\hat{\mathbf{p}}_e^2}{2m} \rightarrow \sum_{\mathbf{p}} \frac{p^2}{2m} \hat{a}_{\mathbf{p}}^\dagger \hat{a}_{\mathbf{p}}, \quad e^{-i\mathbf{q}\cdot\hat{\mathbf{r}}_e} \rightarrow \sum_{\mathbf{p}} \hat{a}_{\mathbf{p}-\mathbf{q}}^\dagger \hat{a}_{\mathbf{p}}, \quad e^{i\mathbf{q}\cdot\hat{\mathbf{r}}_e} \rightarrow \sum_{\mathbf{p}} \hat{a}_{\mathbf{p}+\mathbf{q}}^\dagger \hat{a}_{\mathbf{p}} \quad (2.23)$$

$$\begin{aligned}
e^{-i\mathbf{q}\cdot\hat{\mathbf{r}}_e} &\rightarrow \sum_{\mathbf{k},\mathbf{p}} \langle \mathbf{k} | e^{-i\mathbf{q}\cdot\hat{\mathbf{r}}_e} | \mathbf{p} \rangle \hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{p}} \\
&= \sum_{\mathbf{k},\mathbf{p}} \langle \mathbf{k} | \left[ \int |\mathbf{x}\rangle \langle \mathbf{x}| d^3x \right] e^{-i\mathbf{q}\cdot\hat{\mathbf{r}}_e} \left[ \int |\mathbf{y}\rangle \langle \mathbf{y}| d^3y \right] | \mathbf{p} \rangle \hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{p}} \\
&= \sum_{\mathbf{k},\mathbf{p}} \hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{p}} \int d^3x \frac{e^{-i\mathbf{k}\cdot\mathbf{x}}}{\sqrt{V}} e^{-i\mathbf{q}\cdot\mathbf{x}} \frac{e^{i\mathbf{p}\cdot\mathbf{x}}}{\sqrt{V}} \\
&= \sum_{\mathbf{k},\mathbf{p}} \hat{a}_{\mathbf{k}}^\dagger \hat{a}_{\mathbf{p}} \delta(\mathbf{k} + \mathbf{q} - \mathbf{p}) \\
&= \sum_{\mathbf{p}} \hat{a}_{\mathbf{p}-\mathbf{q}}^\dagger \hat{a}_{\mathbf{p}}
\end{aligned} \tag{2.24}$$

We substitute this into the Hamiltonian, and make in the interaction term use of the fact that the summation over  $\mathbf{q}$  is symmetric, to finally obtain

$$\hat{H} = \sum_{\mathbf{p}} \frac{p^2}{2m} \hat{a}_{\mathbf{p}}^\dagger \hat{a}_{\mathbf{p}} + \omega \sum_{\mathbf{q}} \hat{b}_{\mathbf{q}}^\dagger \hat{b}_{\mathbf{q}} + \sum_{\mathbf{q},\mathbf{p}} V(q) \left( \hat{b}_{\mathbf{q}}^\dagger - \hat{b}_{-\mathbf{q}} \right) \hat{a}_{\mathbf{p}-\mathbf{q}}^\dagger \hat{a}_{\mathbf{p}}. \tag{2.25}$$

Take the limit  $l \rightarrow \infty$ . Perhaps do this where we apply the algorithm? So at the Feynman diagrams Section?

what is the interpretation of the  $\mathbf{B}$ -field. Oscillating displacements in the lattice  $\rightarrow$  phonons.

make the reader pay attention to the phonons being dispersionless. Discuss why this model is still a pretty accurate (optical phonons)

## 2.3 Feynman diagrams

## 2.4 Dyson equation

## 2.5 Monte Carlo Simulation

### 2.5.1 Detailed balance and ergodicity

### 2.5.2 Metropolis-Hastings algorithm

# Chapter 3

## Diagrammatic Monte Carlo

Here we shall summarise what this chapter is about

### 3.1 General idea

### 3.2 Generalisation of stochastic integration (better name needed)

#### 3.2.1 Example 1

#### 3.2.2 Example 2

#### 3.2.3 Example 3

### 3.3 Polaron implementation

Here we will demonstrate what happens when we are using  $T = 0$  and not having any free electrons in the system at the beginning.

### 3.3.1 Update function 1

### 3.3.2 Update function 2

### 3.3.3 Update function 3

## 3.4 Sample self energy

### 3.4.1 Divergent diagram

The first order proper self energy diagram is proportional to  $1/\sqrt{\tau}$  for small  $\tau$  and thus diverges when  $\tau \rightarrow 0$ . This combined with a discretised histogram becomes a problem for the bins in the neighbourhood of  $\tau = 0$  since they acquire more mass than they are supposed to. In order to fix this we could use a much finer discretisation or calculate the value for these bins somehow else.

The value of the diagram for  $\mathbf{p} = \mathbf{0}$  and  $\Delta G = 0$  is

$$S^{(1)}(\tau) = \frac{\alpha}{\sqrt{\pi\tau}} e^{(\mu-\omega)\tau} \quad (3.1)$$

The value of the bins in our histogram is

$$\tilde{V}_i \propto \frac{1}{\Delta\tau} \int_{\tau_i}^{\tau_{i+1}} S^{(1)}(t) dt = \frac{\alpha}{\Delta\tau \sqrt{\omega - \mu}} [\text{erf}(\sqrt{\omega - \mu} \sqrt{\tau_{i+1}}) - \text{erf}(\sqrt{\omega - \mu} \sqrt{\tau_i})] \quad (3.2)$$

but the true value should be

$$V_i \propto S^{(1)}\left(\frac{\tau_{i+1} + \tau_i}{2}\right) \quad (3.3)$$

For all bins which fulfil  $\tilde{V}_i - V_i > \epsilon$  we instead of using DMC utilise ordinary stochastic integration using MC for the fixed times  $(\tau_{i+1} + \tau_i)/2$ .

## 3.5 Boldification



# Chapter 4

## Results

Here we shall summarise what this chapter is about

### 4.1 Analytical predictions

### 4.2 Numerics

### 4.3 Discussion

# Chapter 5

## Summary and Conclusions

Here we shall summarise what this chapter is about

# Appendix A

(make this go away, also Appendix A  
-> Appendix)

- A.1 Heavy theory from the finite temperature formalism
- A.2 Derivation of the finite temperature electron Green's function
- A.3 Derivation of the finite temperature phonon Green's function

# References

- [1] Fröhlich H. Electrons in lattice fields. *Advances in Physics*. 1954;3(11):325–361.  
Available from: <http://dx.doi.org/10.1080/00018735400101213>.
- [2] Boris V Svistunov NVP Egor S Babaev. *Superfluid States of Matter*. CRC Press; 2015.
- [3] Fetter AL, Walecka JD. *Quantum Theory of Many-Particle Systems*. Dover Publications; 2003.