

Poisson distribution


From Wikipedia, the free encyclopedia

In probability theory and statistics, the **Poisson distribution** (French pronunciation [pwaʒɔ̃]; in English usually /ˈpɔːzɒn/), named after French mathematician Siméon Denis Poisson, is a discrete probability distribution that expresses the probability of a given number of events occurring in a fixed interval of time and/or space if these events occur with a known average rate and independently of the time since the last event.^[1] The Poisson distribution can also be used for the number of events in other specified intervals such as distance, area or volume.


For instance, an individual keeping track of the amount of mail they receive each day may notice that they receive an average number of 4 letters per day. If receiving any particular piece of mail doesn't affect the arrival times of future pieces of mail, i.e., if pieces of mail from a wide range of sources arrive independently of one another, then a reasonable assumption is that the number of pieces of mail received per day obeys a Poisson distribution.^[2] Other examples that may follow a Poisson: the number of phone calls received by a call center per hour or the number of decay events per second from a radioactive source.

Contents

- 1 Introduction to the Poisson distribution
 - 1.1 Examples
 - 1.2 Assumptions: When is the Poisson distribution an appropriate model?
 - 1.3 Probability of events for a Poisson distribution
 - 1.3.1 Examples of probability for Poisson distributions
 - 1.3.2 Once in an interval events: The special case of $\lambda = 1$ and $k = 0$
 - 1.4 Examples that violate the Poisson assumptions
 - 1.5 Poisson regression and negative binomial regression
- 2 History



The horizontal axis is the index k , the number of occurrences. λ is the expected number of occurrences. The vertical axis is the probability of k occurrences given λ . The function is defined only at integer values of k . The connecting lines are only guides for the eye.



The horizontal axis is the index k , the number of occurrences. The CDF is discontinuous at the integers of k and flat everywhere else because a variable that is Poisson distributed takes on only integer values.

Parameters	$\lambda > 0$ (real)
-------------------	----------------------

- 3 Definition
- 4 Properties
 - 4.1 Mean
 - 4.2 Median
 - 4.3 Higher moments
 - 4.4 Sums of Poisson-distributed random variables
 - 4.5 Other properties
 - 4.6 Poisson races
- 5 Related distributions
- 6 Occurrence
 - 6.1 Law of rare events
 - 6.2 Poisson point process
 - 6.3 Other applications in science
- 7 Generating Poisson-distributed random variables
- 8 Parameter estimation
 - 8.1 Maximum likelihood
 - 8.2 Confidence interval
 - 8.3 Bayesian inference
 - 8.4 Simultaneous estimation of multiple Poisson means
- 9 Bivariate Poisson distribution
- 10 Computer software for the Poisson distribution
 - 10.1 Poisson distribution using R
 - 10.2 Poisson distribution using Excel
 - 10.3 Poisson distribution using Python (SciPy)
 - 10.4 Poisson distribution using Mathematica
- 11 See also
- 12 Notes
- 13 Further reading
- 14 References

Introduction to the Poisson distribution

The Poisson distribution is popular for modelling the number of times an event occurs in an interval of time or space.

Examples

The Poisson distribution may be useful to model events such as

- The number of meteors greater than 1 meter diameter that strike earth in a year
- The number of patients arriving in an emergency room between 10 and 11 pm

Support	$k \in \mathbb{N} \cup 0$;
pmf	$\frac{\lambda^k e^{-\lambda}}{k!}$
CDF	$\frac{\Gamma(\lfloor k+1 \rfloor, \lambda)}{\lfloor k \rfloor!}$, or $e^{-\lambda} \sum_{i=0}^{\lfloor k \rfloor} \frac{\lambda^i}{i!}$, or $Q(\lfloor k+1 \rfloor, \lambda)$ (for $k \geq 0$, where $\Gamma(x, y)$ is the incomplete gamma function, $\lfloor k \rfloor$ is the floor function, and Q is the regularized gamma function)
Mean	λ
Median	$\approx \lfloor \lambda + 1/3 - 0.02/\lambda \rfloor$
Mode	$\lfloor \lambda \rfloor - 1, \lfloor \lambda \rfloor$
Variance	λ
Skewness	$\lambda^{-1/2}$
Ex. kurtosis	λ^{-1}
Entropy	$\lambda[1 - \log(\lambda)] + e^{-\lambda} \sum_{k=0}^{\infty} \frac{\lambda^k \log(k!)}{k!}$ (for large λ) $\frac{1}{2} \log(2\pi e \lambda) - \frac{1}{12\lambda} - \frac{1}{24\lambda^2} - \frac{19}{360\lambda^3} + O\left(\frac{1}{\lambda^4}\right)$
MGF	$\exp(\lambda(e^t - 1))$
CF	$\exp(\lambda(e^{it} - 1))$
PGF	$\exp(\lambda(z - 1))$
Fisher information	$\frac{1}{\lambda}$