

Ancestral Metabolic Architecture Lost in European Neolithic Lineages Explains Global Type 2 Diabetes Disparities

Emilio García-Morán

17 May 2026

Target journal: Genome Medicine

emilio.garcia.moran@uva.es · ORCID: 0000-0002-2487-6686

Data & code: <https://github.com/EmilioGarciaMoran/EastWestDM>

Zenodo DOI: 10.5281/zenodo.20260486

Competing interests: None declared · Funding: None declared

Abstract

Type 2 diabetes (T2D) disproportionately affects East Asian (EAS), admixed American (AMR), and African-ancestry populations compared to non-Finnish Europeans (NFE). We propose that this disparity reflects not differential pathology but the application of a clinical reference framework calibrated on one post-Neolithic European metabolic architecture to populations whose equally successful evolutionary trajectories diverged prior to — and were never subject to — the selective pressures that drove European metabolic remodelling.

We analysed 168 Eastern and 67 Western variants from four archaic and early modern human genomes — Denisovan D17, Ust'-Ishim (~45 ka), Vindija Neandertal (~44 ka), and Ötzi (~5.3 ka) — against gnomAD v4. The ancestral NCDN variant (chr1:35,565,741) is present at ~70% in sub-Saharan African populations (San 92%, Biaka 91%), confirming African ancestral origin predating the Out-of-Africa dispersal. Vindija Neandertal already carries Western regulatory

variants prior to agriculture (~44 ka), indicating that the European divergence began in pre-Neolithic temperate environments. Unadmixed Native American populations (Surui, Karitiana: 100%; Pima: 77%; Maya: 67%) and EAS (78%) carry the AMA at or above the African baseline, reflecting enrichment consistent with cold-environment selection during the Beringian migration (~15,000 ya). NFE Europeans carry the variant at 3% — a 23-fold reduction from the African baseline consistent with Neolithic counter-selection. A single upstream variant at chr11:18,411,103 (C>T) acts as a coordinated cis-eQTL suppressing LDHC expression across seven tissues in GTEx v8 (skeletal muscle $p = 3.71 \times 10^{-64}$, NES = -0.935; independent replication in eQTLGen $n = 31,684$, $Z = 75.96$, $p = 3.27 \times 10^{-310}$). SLC2A3 (GLUT3) remains invariant across 430,000 years of hominin evolution, establishing brain glucose transport as a non-negotiable evolutionary constraint. African Americans carry intermediate frequencies (ASW 49%, ACB 73%) consistent with post-colonial admixture superimposed on the African ancestral baseline. These findings reframe global T2D disparities as a consequence of clinical reference frameworks derived predominantly from European-ancestry cohorts, with immediate implications for ancestry-informed diagnostic thresholds and pharmacogenomic stratification.

Keywords: type 2 diabetes · ancestral metabolic architecture · LDHA · LDHC · eQTL · paleogenomics · Beringia · Neolithic transition · precision medicine · admixture

Introduction

The global burden of type 2 diabetes is distributed with striking population specificity. East Asian and admixed American populations show higher prevalence, earlier onset, greater severity at equivalent body mass index, and differential response to dietary and pharmacological interventions compared to Europeans. African-ancestry populations share this elevated burden and are additionally underrepresented in genome-wide association studies and pharmacogenomic trials. The three population groups most poorly served by current clinical frameworks — EAS, AMR, and African-ancestry — thus share a common feature: systematic underperformance of risk models calibrated on European reference cohorts. This constitutes a systematic clinical bias: diagnostic thresholds, BMI cutoffs, and therapeutic guidelines calibrated on European-ancestry populations consistently underperform in EAS, AMR, and African-ancestry groups not because of differential disease biology, but because they measure the wrong metabolic baseline.

The thrifty genotype hypothesis posited a single ancestral adaptation to feast-famine cycles. The thrifty phenotype hypothesis shifted causality to developmental programming. Neither framework explains the shared elevated burden across three geographically and historically distinct population groups, nor does either account for why non-Finnish Europeans — who also experienced prehistoric food scarcity — specifically do not share it.

We propose a framework centred on a universal ancestral metabolic architecture — present in Africa at ~70% frequency prior to the Out-of-Africa dispersal — that subsequently diverged along two distinct post-Out-of-Africa trajectories under radically different selective pressures.

The clinical consequences are a product of the encounter between these trajectories and a medical reference standard calibrated, by historical circumstance, on only one of them.

The metabolic architectures documented here represent independent adaptive solutions of equivalent fitness. The European lineage, beginning with pre-agricultural Neandertals in temperate environments and consolidated by the Neolithic transition, optimised for carbohydrate processing from plant-based substrates. The Siberian and Beringian lineage optimised for maximal glycolytic efficiency under cold stress and caloric scarcity, with brain glucose supply as the inviolable constraint in both cases. The clinical problem arises not from the architectures themselves but from their encounter with a globally uniform food environment and clinical reference frameworks that, by historical circumstance, were calibrated on only one of them.

Results

Eastern and Western variant discovery

Comparison of D17 Denisovan and Ust'-Ishim (~45 ka, Siberia) genomes against gnomAD v4 identified 168 high-confidence Eastern variants after C→T/G→A damage filtering (mapDamage 2.2.1; PMDtools threshold 3). Comparison of Vindija Neandertal and Ötzi genomes identified 67 Western variants. Pathway analysis revealed Eastern enrichment in glycolysis regulators (PFKM, LDHA, LDHC), adipose biology (ADIPOQ, FASN), and cold thermosensation (TRPA1); Western enrichment in lipid metabolism (ACACA), thermogenesis (PPARGC1A), and plant-based nutrient processing. Eighty-six percent of Eastern variants showed symmetric polarisation toward AMR/EAS populations (Fisher $p = 3.2 \times 10^{-18}$), with LDHA/LDHC regulatory regions showing 21.4-fold enrichment.

SLC2A3 invariance across 430,000 years

Pileup analysis of five Sima de los Huesos specimens (PRJEB10597; remapped to hg38; mean depth 1.18×) at SLC2A3 exonic and promoter positions reveals no fixed differences relative to modern humans at 11 canonical positions examined. Combined with conservation data from Vindija, D17, and Ust'-Ishim, SLC2A3 has remained functionally invariant for at least 430,000 years across hominin lineages, establishing brain glucose transport as a non-negotiable evolutionary constraint around which both metabolic architectures were independently organised.

LDHA/LDHC cis-eQTL across seven tissues

GTEX v8 analysis identified chr11:18,411,103 (C>T; rs7951953) — an Eastern-specific upstream variant 1,228 bp from the LDHC transcription start site — as a coordinated cis-eQTL suppressing LDHC expression in seven tissues simultaneously (Table 1). Effect sizes ranged from NES = -0.935 (skeletal muscle) to NES = -0.620 (whole blood), all in the same negative

direction. The Eastern-specific allele (C>T) is associated with lower LDHC expression in all seven tissues, consistent with a regulatory variant that suppresses LDHC transcription, shifting metabolic flux toward oxidative phosphorylation — thermogenically advantageous under cold stress, potentially impairing metabolic flexibility under caloric excess. Independent replication in eQTLGen ($n = 31,684$; $Z = 75.96$; $p = 3.27 \times 10^{-310}$) confirms this as among the most statistically robust regulatory signals in the human genome.

Table 1: LDHA/LDHC cis-eQTL across GTEx v8 tissues (chr11:18,411,103 C>T). NES: normalised effect size (negative = lower LDHC expression in Eastern allele carriers). eQTLGen: independent whole-blood replication cohort.

Tissue	p -value	NES	n
Skeletal muscle	3.71×10^{-64}	−0.935	706
Subcutaneous adipose	1.46×10^{-59}	−0.881	581
Visceral adipose	1.11×10^{-41}	−0.812	469
Whole blood	7.39×10^{-30}	−0.620	670
Pancreas	7.33×10^{-26}	−0.751	305
Liver	9.12×10^{-24}	−0.734	208
Brain cortex	2.70×10^{-22}	−0.698	205
eQTLGen (blood)	3.27×10^{-310}	$Z = 75.96$	31,684

Independent fine-mapping in Open Targets Genetics confirmed 11_18411103_C_T as a cis-eQTL credible set variant for LDHC and LDHA across 13 independent molQTL studies (SuSiE fine-mapping), including amygdala ($p = 1.95 \times 10^{-18}$), skin, induced pluripotent stem cells, ascending aorta, frontal cortex, sigmoid colon, cerebellum, and omental fat pad, with consistent fine-mapping confidence across all tissues. This constitutes independent replication of the LDHC/LDHA cis-regulatory signal across GTEx v8 (7 tissues), eQTLGen ($n = 31,684$), and Open Targets molQTL (13 studies).

The brain cortex eQTL ($p = 2.70 \times 10^{-22}$) alongside SLC2A3 invariance establishes a critical dissociation: brain glucose *transport* was conserved for 430,000 years, while brain glucose *metabolism* retains population-specific regulatory variation.

Global frequency distribution: an ancestral variant with asymmetric selective history

Query of the gnomAD v3.1.2 HGDP+1000G joint callset at the NCDN locus (chr1:35,565,741) revealed a global frequency distribution consistent with an ancestral allele present prior to the Out-of-Africa dispersal, subsequently subject to asymmetric selection across lineages (Table 2, Figures 5–6).

Sub-Saharan African populations carry the variant at a mean frequency of 70.4% (range 49–92%), establishing the ancestral baseline. San and Biaka Pygmy populations show the

highest frequencies (91.7% and 90.9% respectively), consistent with long-term maintenance of the ancestral state. EAS (78%) and unadmixed Native American populations (Surui/Karitiana: 100%; Pima: 77%; Maya: 67%) show frequencies at or above the African baseline. NFE Europeans show near-complete loss of the variant (3%). African Americans carry intermediate frequencies (ASW: 49%; ACB: 73%) consistent with variable European admixture superimposed on the African ancestral baseline. The independent replication SNP at chr1:35,542,512 shows consistent directional effect across all populations.

Table 2: Global frequency of ancestral NCDN variant (chr1:35,565,741) by population. Sources: HGDP, 1000 Genomes Project, gnomAD v3.1.2.

Population	<i>n</i>	Frequency	Selective context
San (Kalahari)	12	92%	Ancestral African baseline
Biaka Pygmy (Congo)	44	91%	Ancestral African baseline
Mbuti Pygmy (Congo)	24	83%	Ancestral African baseline
Surui (Amazonia)	14	100%	Consistent with cold-environment enrichment
Karitiana (Amazonia)	20	100%	Consistent with cold-environment enrichment
EAS (gnomAD)	5,174	78%	Consistent with cold-environment enrichment
Pima (N. Mexico)	22	77%	Consistent with cold-environment enrichment
Yoruba (Nigeria)	40	70%	Ancestral African baseline
ACB (African Caribbean)	188	73%	Admixture dilution
Maya (Mesoamerica)	36	67%	Consistent with cold-environment enrichment
AFR gnomAD (global)	41,322	56%	Pan-African mean
Peruvian (Andes)	170	53%	Low European admixture

Population	n	Frequency	Selective context
ASW (African American)	104	49%	Admixture dilution
Mexican (MXL, 1000G)	126	39%	Post-1521 European admixture
AMR gnomAD	15,244	31%	Post-1521 European admixture
Colombian (CLM, 1000G)	188	29%	Post-1521 European admixture
NFE (gnomAD)	67,976	3%	Neolithic counter-selection

Discussion

Two adaptive solutions of equivalent fitness

We propose a terminological framework that clarifies the evolutionary asymmetry documented here. The **Ancestral Metabolic Architecture (AMA)** — characterised by high NCDN variant frequency, LDHC suppression across peripheral tissues, enhanced anaerobic glycolytic flux, and preserved brain glucose priority — represents the metabolic baseline present in Africa prior to the Out-of-Africa dispersal and maintained in EAS, unadmixed Native American, and sub-Saharan African populations. The **European Post-Neolithic Metabolic Architecture (EPMA)** — characterised by near-complete loss of AMA variants (NCDN 3%), ACACA lipogenesis, and PPARGC1A thermogenesis — emerged in European lineages beginning with pre-agricultural Neandertals in temperate environments and was consolidated by the Neolithic agrarian transition. The clinical disparity arises not from AMA pathology but from the universal application of EPMA-calibrated diagnostic and therapeutic standards to populations in whom AMA predominates.

The AMA and EPMA represent independent evolutionary solutions of equivalent adaptive success — not differential pathology. The EPMA lineage optimised for carbohydrate processing from plant-based substrates: ACACA lipogenesis, PPARGC1A thermogenesis, PPAR adipogenesis calibrated for cereal-derived glucose. The AMA lineage optimised for maximal glycolytic efficiency under cold stress and caloric scarcity, with brain glucose supply as the inviolable constraint in both cases. The clinical problem arises not from the architectures themselves but from their encounter with a globally uniform food environment and clinical reference frameworks that, by historical circumstance, were calibrated predominantly on EPMA-carrying populations.

The European divergence began before agriculture

Vindija Neandertal (~44 ka) already carries Western regulatory variants in the absence of cereals, domesticated animals, or settled lifestyles — predating the Neolithic transition by approximately 35,000 years. The temperate European environment, characterised by moderate cold stress and reliance on megafaunal fat as caloric substrate, appears to have initiated counter-selection against the AMA prior to the Neolithic. The Neolithic transition accelerated a process already underway. Ötzi (5,300 ya) represents not the origin of the EPMA but its completion.

Whether the ancestral NCDN architecture was present in Sima de los Huesos hominin lineages (430 ka) and subsequently lost during the European divergence remains undetermined. Coverage at chr1:35,565,741 in SH specimens was insufficient for allelic inference (0 reads; mean genome-wide depth 1.18×). This question represents a compelling target for higher-coverage ancient European genome sequencing.

From regulatory variant to glycolytic mechanism

The identification of chr11:18,411,103 (C>T) as a coordinated cis-eQTL suppressing LDHC expression across seven metabolically relevant tissues converts this population genomic observation into a tractable molecular mechanism. LDHC encodes the muscle/testis isoform of lactate dehydrogenase, catalysing the final step of anaerobic glycolysis. Although classically described as testis-enriched, LDHC shows documented ectopic expression in skeletal muscle, liver, and brain under metabolic stress conditions, and its co-regulation with LDHA in glycolytically active tissues has been reported in multiple transcriptomic datasets. The coordinated suppression of LDHC alongside LDHA regulatory variants is therefore consistent with a systemic glycolytic programme rather than a tissue-restricted phenomenon. Its coordinated downregulation across muscle, adipose, pancreas, liver, and brain cortex — all at negative NES — is consistent with a pleiotropic regulatory programme rather than a tissue-specific effect. In calorically abundant environments, LDHC suppression shifts the pyruvate/lactate equilibrium toward oxidative phosphorylation — thermogenically advantageous under Arctic conditions, potentially impairing metabolic flexibility under caloric excess.

Regulatory ghost adaptation

LDHA missense variants shared between D17 and Ust'-Ishim are absent from gnomAD v4, indicating active purifying selection on coding changes. Yet the surrounding regulatory landscape persists at high frequency across most human populations outside Europe. We term this pattern *regulatory ghost adaptation*: coding changes eliminated by purifying selection, regulatory architecture co-selected with them persisting in non-coding space, visible only through eQTL analysis. The European loss of this architecture — rather than its presence elsewhere — is the evolutionary anomaly requiring explanation.

The Beringia bottleneck: cold selection as amplifier, not origin

The peopling of the Americas did not introduce a novel metabolic architecture — it exported and amplified an ancestral African one. Unadmixed Native American populations carry the Eastern NCDN variant at 67–100%, at or above the African baseline, consistent with enrichment during the Arctic cold-environment bottleneck (~15,000 ya). The stepwise frequency reduction through populations of increasing European admixture — Peruvians (53%), Mexicans (39%), AMR gnomAD (31%), Colombians (29%) — constitutes a quantitative genomic record of post-1521 demographic change. The same dilution mechanism operates in African Americans (ASW: 49%; ACB: 73%), where post-colonial European admixture has superimposed the EPMA low-frequency architecture on the African ancestral baseline. The ~24% frequency difference between ASW and ACB reflects differential admixture proportions, not differential selection.

Three clinically undercharacterised populations, one shared mechanism

The populations most poorly served by European-calibrated clinical thresholds share a single genomic feature: they carry the AMA that Europeans specifically lost through EPMA selection. SGLT2 inhibitors, GLP-1 receptor agonists, and thiazolidinediones were validated predominantly in European-ancestry cohorts, and direct pharmacogenomic evidence in AMA-carrying populations is currently limited. The LDHC cis-eQTL signal in pancreas ($p = 7.33 \times 10^{-26}$), skeletal muscle ($p = 3.71 \times 10^{-64}$), and liver ($p = 9.12 \times 10^{-24}$) identifies specific tissues where differential glycolytic flux may alter pharmacodynamic response. Stratification by LDHA/LDHC regulatory haplotype — rather than continental ancestry label — would recover this signal in admixed populations currently misclassified by European-reference panels. The BMI thresholds documented by Caleyachetty et al. are not arbitrary measurement failures: they are the epidemiological shadow of the evolutionary divergence documented here.

The human singularity and its metabolic price

SLC2A3 invariance across 430,000 years establishes brain glucose priority as the non-negotiable constraint around which all hominin metabolic architectures were organised. The periphery adapted — differently in different lineages, under different selective pressures, across different timescales. The brain never compromised. The clinical consequence is that the organ most dependent on stable glucose supply is protected by the most conserved gene in our dataset, while the organs that regulate glucose availability — muscle, liver, pancreas, adipose — carry the full burden of evolutionary divergence.

Limitations

This analysis is limited by available high-coverage archaic genomes. The conservative C→T/G→A damage filter may remove genuine variants; PMDtools-based filtering gave

comparable directional results. Coverage at the NCDN locus in Sima de los Huesos specimens was insufficient for allelic inference (0 reads; mean depth 1.18 $\times$). Native American and African HGDP sample sizes are small ($n = 12\text{--}44$); replication in the PAGE study, Arctic Biobank, and Mexico City Prospective Study is required before clinical translation. The eQTLGen replication cohort consists of whole blood; however, consistency of negative NES across all seven GTEx tissues supports a pleiotropic regulatory programme. The Neolithic counter-selection hypothesis requires formal testing with time-series ancient DNA spanning the European Mesolithic-Neolithic transition. CRISPR functional validation of the LDHA/LDHC regulatory variant is needed to establish causality. Finally, these findings do not imply deterministic ancestry-specific pathology. They reflect population-level differences in regulatory metabolic architecture shaped by historical selection pressures and demographic processes, and should be interpreted in that evolutionary and epidemiological context rather than as fixed biological categories.

Methods

Archaic genome processing and variant calling

Five Sima de los Huesos specimens (PRJEB10597; Meyer et al. 2016) were remapped to hg38 using AdapterRemoval ($-\text{qualitymax } 93$) $\rightarrow$ BWA-MEM ($-\text{k16}$) $\rightarrow$ Picard MarkDuplicates $\rightarrow$ mapDamage 2.2.1 ($-\text{rescale, MCMC}$). Variants were called with GATK HaplotypeCaller. D17 Denisovan, Ust'-Ishim, Vindija Neandertal, and Ötzi genomes were processed with identical parameters. C $\rightarrow$ T and G $\rightarrow$ A variants were excluded at positions with PMDtools score < 3 .

Population frequency analysis

gnomAD v4 allele frequencies were extracted via the gnomAD REST API. Population-specific frequencies for HGDP and 1000 Genomes populations were extracted via remote bcftools query from the gnomAD v3.1.2 HGDP+TGP joint callset. Symmetric polarisation was assessed using Fisher's exact test with Bonferroni correction. African population frequencies were extracted for 15 sub-Saharan populations including deep-branching lineages (San, Biaka Pygmy, Mbuti Pygmy).

eQTL analysis

GTEx v8 single-tissue eQTL data were queried for all Eastern variants within 1 Mb of LDHC (ENSG00000166796). The eQTLGen cis-eQTL dataset (2019-12-11 release; $n = 31,684$; FDR < 0.05) was queried for independent replication. Effect size direction (NES) was compared across all tissues to assess pleiotropic consistency.

SLC2A3 conservation analysis

SLC2A3 exonic positions were queried in all archaic genomes using samtools mpileup (minimum base quality 20, minimum mapping quality 25). Positions covered at $1\times$ depth were considered for conservation analysis.

Statistical analysis and figure generation

All analyses were performed in R v4.5.2 (ggplot2, dplyr) and Python 3.11 (matplotlib, geopandas, pandas). All code is available at <https://github.com/EmilioGarciaMoran/EastWestDM>.

Data Availability

All analysis code, variant lists, population frequency tables, and figure generation scripts are available at: <https://github.com/EmilioGarciaMoran/EastWestDM> (DOI: 10.5281/zenodo.20260486). Raw archaic genome data: PRJEB10597 (Sima de los Huesos). gnomAD: <https://gnomad.broadinstitute.org>. GTEx: <https://gtexportal.org>. eQTLGen: <https://eqtlgen.org>.

Figure Legends

Figure 1. East-West variant frequency polarisation. 86% of Eastern variants enriched in AMR/EAS vs NFE (Fisher $p = 3.2 \times 10^{-18}$). Bars: proportion of Eastern (blue) and Western (red) variants in each population group.

Figure 2. Pathway enrichment. Eastern: glycolysis (PFKM, LDHA, LDHC), adipose (ADIPOQ, FASN), cold thermosensation (TRPA1). Western: lipogenesis (ACACA), thermogenesis (PPARGC1A).

Figure 3. SLC2A3 invariance across 430,000 years. Pileup at 11 canonical positions across five Sima de los Huesos specimens, D17, Ust'-Ishim, Vindija, Ötzi. No fixed differences observed.

Figure 4. LDHA/LDHC cis-eQTL. Panel A: NES by tissue (GTEx v8). Panel B: eQTLGen replication ($n = 31,684$; $Z = 75.96$; $p = 3.27 \times 10^{-310}$). Panel C: regional association plot chr11:18.35–18.55 Mb.

Figure 5. Global frequency gradient of ancestral NCDN variant (chr1:35,565,741). Bars ordered by frequency. Colours: deep-branching African (dark red), unadmixed Native American/EAS (red), admixed African American/AMR (amber), European (blue). Open circles: replication at chr1:35,542,512.

Figure 6. World map of ancestral NCDN variant frequency. Circle size proportional to allele frequency; colour scale: blue (NFE 3%) to red (Surui/Karitiana 100%). Arrows: Out-of-Africa

(~70 ka, black); Beringia cold-environment enrichment (~15 ka, dark red); Neolithic counter-selection in European lineages (~8–5 ka, blue); post-colonial admixture events (dashed grey). Source: HGDP + 1000 Genomes + gnomAD v3.1.2.

“The periphery adapted; the brain never compromised.”

\begin{center} EastWestDM v7.1 · García-Morán 2026 · Universidad de Valladolid · DOI:
10.5281/zenodo.20260486 \end{center}