

A74 EXERCISES: Stokes parameters and moving charges (2 & 3)

1. Consider the Stoke's parameters for monochromatic plane waves (Eq 2.40; note that 2.47 is the more general forms for quasi-monochromatic plane waves). Note that the ωt part doesn't appear in these equations by definition (and I think also by virtue of reality: except in interferometry, we're always measuring a time average of a light wave, not recording a specific phase).
 - (a) Just by considering the geometry and the verbal definitions given for the Stoke's parameters at the beginning of the lecture notes, what are the Stokes parameters for horizontally (along 0 degrees) polarized light and vertically (along 90 degrees) polarized light?
 - (b) Consider the formulation of the electric field that uses β and χ . What must β and χ be for the E fields in the two scenarios above? (There are actually a combination of options). Then, use Eq 2.40 to determine the Stokes parameters, confirming your answer to part a).
 - (c) Consider the formulation of the electric field that uses complex numbers (Eq. 2.35 + 2.36). For each of the two scenarios, what is the complex vector E field? (The answer is simple).
2. Consider the Lienard-Wiechart potentials, the potential fields for a moving charge (R&L Eq 3.7). Show that the scaling of the potentials with κ results in a beaming effect. Do this by considering **qualitatively** how κ changes when you consider the directions parallel, perpendicular, and anti-parallel to the velocity.
3. Show that the electric field (R&L Eq 3.9a) becomes Coulomb's law in the limit that $v \ll c$ and zero acceleration.