Does entanglement geometry see the same space? The mutual-information reader of Cao, Carroll and Michalakis applied to the wirings of a graph model of emergent geometry

Emily Smith[1,*]

[1] *Independent researcher, Charlottesville, Virginia, USA*

(Dated: DRAFT, 3 November 2026)

Two published routes build space from a network. In combinatorial quantum gravity (CQG) the network is a classical wiring weighted by a discrete curvature; in the program of Cao, Carroll and Michalakis it is a quantum state on a factored Hilbert space, and distances come from the mutual information between factors. We ask what the second route's reader reports when it is pointed at the first route's wirings. With one factor per vertex and the same mutual information on every link, their distance ansatz with $\Phi = -\ln$ and its shortest-path completion give the graph distance up to a constant (exact), so their dimension estimator becomes classical multidimensional scaling (MDS) of graph-distance windows. Whole tori cannot be read: each wrapped direction takes a pair of large eigenvalues (exact). On balls of graph radius $R = 2$ to 10, with a count fixed before the computation (eigenvalues at least 0.1 of the largest) and $R = 10$ held out: the 4-cube, which the model's local count calls zero-dimensional, reads four; a flat square lattice reads three to four, not two, because its graph distance is a taxicab distance; a direction curled to four sites is counted at small windows and its share falls as $(2R+1)^{-1.24}$, a law fitted at $R = 4, 6$ and confirmed at $R = 10$ (0.112 against a predicted 0.076 to 0.141); and the narrow tori that the model's conversion produces read one at $R = 10$. Three of five registered predictions failed at one or more radii. No quantum state of the model is computed; one published state built from a wiring is used as a calibration.

I. INTRODUCTION

There are at least two published ways to get space out of a network. In combinatorial quantum gravity (CQG) [1–3] the network is a classical graph and its energy is the total Ollivier curvature of the wiring; at low coupling the favored graphs are lattice-like and a geometry is read from the wiring itself. In the program of Cao, Carroll and Michalakis [4, 5] the network is derived: a quantum state on a Hilbert space factored into many pieces defines a weighted graph whose weights are the mutual informations between pieces, a distance is assigned to each weight, and a dimension is read by embedding the distances in Euclidean space. Ref. [4] says plainly that it assumes no Hamiltonian and that the factorization is given; Ref. [6] asks how a factorization could be chosen.

The two routes meet at one point: both end with a weighted graph and ask what space it describes. This paper asks a narrow question at that point. *When the second route's reader is given the first route's wirings, does it see the same space?* In particular, does it agree with the local count of dimension that a square-counting model uses, on a flat sheet, on a sheet curled in one direction, and on a fully curled object?

The wirings come from a family of energies next to CQG, studied in a companion paper [7] (Sec. II). The reader is the one of Ref. [4] without change, except for one choice that applying it to a classical wiring requires (Sec. III). The computation was registered before it was run, with its predictions and one size held out [8]; Sec. V reports it as registered, including the predictions that failed.

Why the author asks (her hypothesis, stated once). The author holds that our space is a stable phase of a deeper, specific arrangement that is curled, and that space arose when it opened, releasing a bounded amount of energy. On the parameter used below she holds that λ is 1 or below everywhere when the network is taken as a whole, the exception being the two-dimensional inner shell of a black hole seen one part at a time, where the arrangement sets a barrier above 1; that the two sides of the bipartite graph are two distinct kinds of point, every point within a kind the same; that points belong to the two-dimensional description only, the foundational degrees of freedom being twist and curl, with rules of their own relating them; that the converted tori are the model's end state and not a stage of her change, three dimensions being in her picture a projection from the two-dimensional arrangement; and that photons travel a cube's body diagonals and gravity its edges, both between points of different sides, for which the model's links stand, while matter, energy and twist are stored between points of the same side, which are never linked. Whether a curled direction reads as a dimension is the question her picture of opening rests on, and that is why the reader's answer matters to her. None of this is assumed below.

Kinds of statement. Each result is marked as exact (arithmetic or complete enumeration), measured (a computed reading registered before it was made, with its verdict), or ours and unverified (an interpretation). Nothing in this paper is about the real universe.

* emilysmithcreate@gmail.com

II. THE MODEL, BRIEFLY

A state is a four-regular bipartite graph on N vertices obeying the hard-core rule (no two vertices share more than two neighbors) [2, 3]. With S the number of squares (four-cycles), S_e the number on edge e, and $X = \sum_e (S_e - 2)_+$, the energy is

$$H = 16\,(N - S) + 4\lambda X. \tag{1}$$

At $\lambda = 1$ this is the CQG curvature action, which Ref. [3] (its Eq. (21)) writes as

$$H = -4 \sum_v \sum_{w \sim v} \kappa(v, w),$$

with κ the Ollivier curvature of the link. Only $\lambda = 1$ is CQG; every other value is our family around it [7].

The local count (exact). A vertex has six pairs of incident edges; under the hard-core rule a pair closes at most one square. Let $d(v)$ be the number of pairs that close none. On the flat lattice torus $d = 2$; on the $4 \times L$ torus, one direction curled to a single square, $d = 1$; on the 4×4 torus, which is the 4-cube, $d = 0$ [7]. This is a count of square-free directions at one vertex, not a certified geometric dimension. For $\lambda > 1$ the curled torus lies $4(\lambda - 1)$ per vertex above the flat one, and after activation it converts with that release; the converted states measured so far are narrow tori whose shortest loop around the short direction is six edges [7].

III. THE READER

A. Definitions, as published

Ref. [4] factors the Hilbert space into finite-dimensional pieces, $\mathcal{H} = \bigotimes_p \mathcal{H}_p$, and uses the mutual information of two pieces,

$$I(A_p : A_q) = S(A_p) + S(A_q) - S(A_p A_q).$$

A state is redundancy-constrained when the entropy of every union B of pieces is a cut sum, which Ref. [4] (its Eq. (9)) writes as

$$S(B) = \tfrac{1}{2} \sum_{p \in B,\ q \notin B} I(A_p : A_q).$$

Each pair with nonzero mutual information gets an edge of length, in their Eq. (13),

$$w(p, q) = \ell_{\mathrm{RC}}\, \Phi\!\left(\frac{I(A_p : A_q)}{I_0} \right),$$

with $\Phi(1) = 0$, $\Phi(x) \to \infty$ as $x \to 0$, Φ decreasing, and I_0 the mutual information of a maximally entangled pair; their examples use $\Phi = -\ln$. No edge is drawn where $I = 0$. The distance is the shortest path, their Eq. (14),

$$d(p, q) = \min_P \sum_n w(p_n, p_{n+1}).$$

The dimension is read by classical multidimensional scaling (their Sec. III.C). For n points with distances $d(p, q)$ the centered matrix of their Eq. (23) is

$$B_{pq} = -\tfrac{1}{2} \Big[d(p,q)^2 - \tfrac{1}{n} \sum_l d(p,l)^2 - \tfrac{1}{n} \sum_l d(l,q)^2 + \tfrac{1}{n^2} \sum_{l,m} d(l,m)^2 \Big],$$

diagonalized as $B = V \Lambda V^{\mathsf{T}}$ with eigenvalues $\lambda_1 \geq \lambda_2 \geq \ldots$ (their Eq. (24)); the embedding in D dimensions uses the top D eigenvectors scaled by $\sqrt{\lambda_i}$ (their Eq. (25)). If the distances are exactly Euclidean in D dimensions, B has exactly D positive eigenvalues and the rest zero. If not, they read the dimension from the spectrum with the rule, stated after their Eq. (25),

$$\varepsilon_D = 1 - \frac{\sum_{i=1}^{D} |\lambda_i|}{\sum_{i=1}^{n} |\lambda_i|},$$

an embedding in D dimensions being "approximately valid" when ε_D is small. Their own two examples give a scale for "small": a spin-1 Heisenberg chain of 100 sites, $\varepsilon_1 = 0.0167$, and a coarse-grained toric-code patch, $\varepsilon_2 = 0.41$, called evidently two-dimensional [4].

B. The one choice: a factor per vertex, the same mutual information on every link

To apply the reader to a classical wiring two things must be chosen.

A factor per vertex. Their factors are the vertices of the information graph and the mutual informations are its edges. The model's vertices are what its links join, so the reading that changes least is one factor per vertex and a nonzero mutual information exactly on the links. A factor per link would read the line graph instead and give different distances; it is not done here. With the author's position that points belong only to a two-dimensional description (Sec. I), this is a reading of that description, not of the degrees of freedom she takes as foundational.

A mutual information per link. The model says only whether two vertices are linked; all links are alike. So take $I(A_p : A_q)/I_0 = x_0$ on every link, for one constant $0 < x_0 < 1$, and zero otherwise. (Under the author's position of Sec. I this places the mutual information where her waves travel and none where her matter is stored; a nonzero mutual information on same-side pairs at graph distance two stands for stored matter and is the second reading of Sec. VIII, registered separately with her prediction before it was computed.)

Exact. Then every edge has the same length $w = \ell_{\mathrm{RC}}(-\ln x_0)$, Eq. (14) gives $d(p, q) = w\, \delta(p, q)$ with δ the graph distance, and B is w^2 times the matrix built from δ. Its eigenvalues scale by w^2, so their ratios, ε_D and any count of eigenvalues relative to the largest do not depend on x_0, ℓ_{RC} or I_0. ($x_0 = 1$ is excluded: it gives $w = 0$ and

every point at one place.) The reader of a wiring is therefore classical MDS of graph distances. Their toric-code example is close to this case: there the mutual information on each link of a uniform coarse-graining is a boundary length up to a constant [4].

A choice that weights each link by its square count would put the model's energy into the lengths; that is a second, different reading and is not made here.

C. One explicit state whose mutual information follows a wiring

The constant above is put in by hand. To show that a quantum state with such mutual information exists, and to see what else such a state brings, we use the one state in the CQG literature built from a wiring. Ref. [9] (its Eq. (4)) turns the adjacency matrix w_{ij} of a graph into a pure state on two copies of an N-level register,

$$|w\rangle = \frac{1}{\|w\|_F} \sum_{i,j} w_{ij}\, |i\rangle|j\rangle,$$

which it reads equivalently as two registers of N qubits, each with exactly one excitation. Ref. [9] cites Ref. [4] for reading space as an entanglement pattern, but it computes one spectral number per graph, not a geometry.

Exact (our computation, 8 × 8 flat torus, one factor per qubit, $2N = 128$ factors). Between qubit i of one copy and qubit j of the other the mutual information is 1.17×10^{-2} bits when i and j are linked and 3.58×10^{-4} bits otherwise. Within one copy it is 1.78×10^{-3} bits for vertices sharing one neighbor, 6.26×10^{-3} for vertices sharing two, and 3.58×10^{-4} for all other pairs. So the mutual information follows the wiring across the copies, and its distance-two graph within each.

Exact (same state). The uniform background is nonzero, so Eq. (13) draws an edge between every pair. With $\Phi = -\ln$ and $I_0 = 2$ bits, a link has length $5.14\,\ell_{\mathrm{RC}}$ and a background edge $8.63\,\ell_{\mathrm{RC}}$, less than two links. The shortest path therefore never uses more than one background edge, every distance is at most $8.63\,\ell_{\mathrm{RC}}$, and MDS of all 128 factors gives one eigenvalue and then five equal ones at 0.76 of it: no low-dimensional geometry. With the background removed the graph splits into two pieces of 64 factors, each with the torus's wiring (bipartite, so the two copies pair one side of one copy with the other side of the other) plus the distance-two edges, and each reads like a whole torus (four equal eigenvalues; Sec. IV).

Ours, unverified. The state shows that the mutual information of a quantum state can follow a wiring, and it shows that it does so only up to a background that, read literally, erases the geometry. It is built from the wiring, so it calibrates the reader and tests nothing about how a geometry could arise. Whether it is redundancy-constrained was not checked.

IV. WHY WHOLE TORI CANNOT BE READ, AND WINDOWS

Exact. On a torus the distance matrix is unchanged by translations, so B's eigenvectors are Fourier modes, which come in equal pairs, a cosine and a sine. Each wrapped direction's longest mode takes one such pair. The 8×8 torus gives four equal eigenvalues, 218.5, then four at 37.5; the 4×16 curled torus gives 525.5, 525.5 (the long direction), then 160, 160 (the curled one, 0.30 of the first and growing with the length of the torus), then 64.8, 64.8; the 4-cube gives exactly four eigenvalues of 16, the others 0 or -4. Even a single square, the curled direction's cross-section, is not Euclidean under graph distance (opposite corners at 2, not $\sqrt{2}$), and its spectrum is $2, 2, 0, -1$. So on whole tori the sheet, the curled torus and the 4-cube all show four large eigenvalues, distinguished by ratios that depend on the torus's size rather than its wiring. Ref. [4] reads open patches only (a chain and a toric-code patch).

Windows. The reader is applied to a ball of graph radius R around a center vertex, using the whole graph's shortest-path distances between the ball's vertices. On the flat 32×32 torus no path inside the ball wraps for $R \le 8$; at $R = 10$ the ball holds pairs 20 apart along one axis, more than half of 32, and some of their shortest paths wrap (this was not noticed when the computation was registered; Sec. V). The curled direction of the tube wraps inside any ball with $R \ge 2$, and so does the short direction of the converted narrow tori (six around) for $R \ge 3$; there the wrap cannot be cut out and is part of what is read.

The registered count. Ref. [4] leaves "small" open. Before the computation we fixed

$$D_{\mathrm{MDS}} = \#\{\, i : \lambda_i \ge 0.1\, \lambda_1 \,\}$$

and record ε_1 to ε_4 beside it. The threshold is ours.

V. RESULTS

A. What was registered

The states [8]: built (exact, no sampling) the flat 32×32 torus, the 4×64 curled torus and the 4-cube, each read from one center (every center is equivalent); measured states saved by earlier runs, namely four converted narrow tori of 1,024 vertices from the slowest-cooled runs at $\lambda = 1.25$ (every eighth vertex a center, 512 windows per radius) and four warm random graphs of the published model ($\lambda = 1$, $N = 196$, $g = 3.3$; every vertex a center, 784 windows). Radii $R = 2, 3, 4, 6$; $R = 10$ held out, to be computed only after a written prediction from the smaller radii. The predictions were ours (the author gave none):

- P0, the 4-cube reads $D_{\mathrm{MDS}} = 4$ for $R \ge 4$, against $d = 0$;

TABLE I. Built states, exact. Eigenvalues of B relative to the largest, the registered count and the reading rule, by window radius. The toy's local count is $d = 2$ (sheet), 1 (curled torus), 0 (4-cube) at every vertex. For $R \geq 4$ the 4-cube window is the whole cube.

State	R	n	λ_2/λ_1	λ_3/λ_1	λ_4/λ_1	ε_2	D_{MDS}
Sheet	2	13	1.000	0.521	0.164	0.38	4
	3	25	1.000	0.428	0.123	0.41	4
	4	41	1.000	0.396	0.101	0.42	4
	6	85	1.000	0.373	0.087	0.43	3
	10	221	1.000	0.408	0.101	0.47	4
Curled	2	12	0.607	0.594	0.093	0.44	3
	3	20	0.426	0.395	0.062	0.42	3
	4	28	0.309	0.291	0.053	0.39	3
	6	44	0.196	0.189	0.041	0.33	3
	10	76	0.112	0.110	0.027	0.25	3
4-cube	2	11	1.000	1.000	0.252	0.54	4
	3	15	1.000	1.000	0.702	0.61	4
	4+	16	1.000	1.000	1.000	0.64	4

- P1, the flat sheet reads 2 at $R = 3, 4, 6$;
- P2, the curled torus reads no more than the sheet at $R = 4, 6$, and its curled pair's share λ_2/λ_1 is near $6/(2R+1)^2$ within a factor of two (0.074 at $R = 4$, 0.036 at $R = 6$), so that at $R \geq 4$ the curled direction is not counted;
- P3, the converted tori's most common count equals the flat sheet's at every R;
- P4, the random graphs read at least 4 at $R = 3$ at most centers.

B. The smaller radii (measured)

The built states are in Table I.

P0 holds. The 4-cube reads four at every radius; the model's count reads zero.

P1 fails. The flat sheet reads 4, 4, 4, 3 at $R = 2, 3, 4, 6$. Its third eigenvalue is 0.37 to 0.52 of the largest, and its fourth sits near the threshold (0.101 at $R = 4$). Its ε_2 is 0.38 to 0.43, the same level as the toric-code patch that Ref. [4] calls two-dimensional; the registered count, not the spectrum's overall spread, is what calls it three or four.

P2 holds on the ranking and fails on the size. The curled torus reads 3 against the sheet's 4 at $R = 4$ and 3 against 3 at $R = 6$. Its curled pair is 0.309 and 0.196 of the largest at $R = 4$ and 6, four and five times the predicted 0.074 and 0.036, so at these radii the curled direction is counted.

P3 fails at $R = 2$ only. The converted tori read 5 at $R = 2$ (508 of 512 windows) against the sheet's 4, and the sheet's count at $R = 3, 4, 6$ (4, 4, 3 in at least 507 of 512). Their windows are not isotropic (λ_2/λ_1 averages 0.50 at $R = 2$, against 1.00 for the square sheet): the narrow torus shows.

TABLE II. Saved states, measured. Most common D_{MDS} over the centers (with its share), by radius. Converted tori: four 1,024-vertex products of the slowest-cooled runs at $\lambda = 1.25$, 512 windows per radius; the model's count is $d = 2$ at 509 of 512 centers. Random graphs: four warm states of the published model, $N = 196$, 784 windows; at $R = 10$ each window is the whole graph.

R	Converted tori	Random graphs (range)
2	5 (508/512)	4 (439/784; 3 to 6)
3	4 (509/512)	6 and 7 (260 each; 4 to 9)
4	4 (507/512)	9 (176/784; 5 to 16)
6	3 (507/512)	13 (200/784; 9 to 17)
10	1 (506/512)	13 (392/784; 13 to 16)

P4 holds. The random graphs read at least 5 at 778 of 784 centers at $R = 3$, and the count grows with R (5 to 16 at $R = 4$; 9 to 17 at $R = 6$).

C. The held-out radius

Prediction, written before $R = 10$ was computed [8]. Fitting power laws in the window's width $2R+1$ to $R = 4$ and 6 only: the curled pair's share at $R = 10$ would be 0.108 (exponent 1.24), with a range of ±30%, 0.076 to 0.141; the sheet's third eigenvalue 0.345 (0.241 to 0.448) and fourth 0.071. Predicted counts: sheet 3, curled torus 3 (its 0.108 sits just above the threshold, so 1 was named as possible), 4-cube 4.

Measured. The curled pair is 0.112 and 0.110 of the largest, inside the predicted range: the size claim holds at the held-out radius. Extrapolating the fit (not computed), the share crosses 0.1 at windows about 22 sites wide. The curled torus reads 3 and the 4-cube 4, as predicted. The sheet reads 4, not the predicted 3: its third eigenvalue, 0.408, is inside its range, and its fourth, 0.101, is above the predicted 0.071 and on the threshold. *A check made after this reading, not registered (exact):* the same window on a 64×64 torus, where no path inside it wraps, gives 0.362 and 0.081 and reads 3, while the curled torus reads the same at 4×128 as at 4×64. So the sheet's miss at $R = 10$ comes from paths wrapping round the 32×32 torus, a flaw of the registered design; the registered verdict stands as scored.

Seen, not predicted for this radius. The converted narrow tori read 1 at 506 of 512 centers at $R = 10$ (Table II): their windows wrap the short direction, as the curled torus's windows wrap its curled one, and at this scale the reader calls them one-dimensional. P3 therefore fails at $R = 10$ as well.

Verdict. By the registered words, MIXED: P0 and P4

hold; P1 fails; P2 holds on the ranking, fails on the size at $R = 4, 6$, and its held-out size claim holds at $R = 10$; P3 fails at $R = 2$ and $R = 10$.

VI. DISCUSSION

The taxicab artifact. Exact: on a square lattice the graph distance between vertices separated by $(\Delta x, \Delta y)$ is $|\Delta x| + |\Delta y|$, so

$$\delta^2 = \Delta x^2 + \Delta y^2 + 2|\Delta x||\Delta y|.$$

The first two terms give the two eigenvalues of a plane; the last is not of the form that MDS can absorb into two coordinates, and it produces the extra eigenvalues. *Measured:* the third eigenvalue stays between 0.36 and 0.43 of the largest from $R = 3$ to 10 (0.362 at $R = 10$ on the larger torus) and does not fade, and at $R = 4$ and 6 negative eigenvalues carry about a fifth of $\sum |\lambda_i|$. So under a constant mutual information per link the reader does not call a square lattice two-dimensional at any window tried, and this says more about the distance assigned to the lattice than about the model. *Exact arithmetic:* if instead every pair had $I/I_0 = e^{-r_{pq}/\xi}$ with r_{pq} the Euclidean distance between lattice sites, then $w = (\ell_{\mathrm{RC}}/\xi)\, r_{pq}$, which already satisfies the triangle inequality, so Eq. (14) returns it unchanged and B has exactly two positive eigenvalues on any patch of the plane. *Ours, unverified:* a state with that property needs mutual information between unlinked pairs that knows the Euclidean distance, which the wiring alone does not supply; whether natural states on a lattice (gapped ground states, for which Ref. [4] suggests $\Phi = -\ln$) come close enough is the question.

A short curled direction depends on scale. The curled direction's share falls as $(2R + 1)^{-1.24}$ (measured, confirmed at the held-out radius), more slowly than the $(2R + 1)^{-2}$ of P2's estimate, which treated the four-site loop as a fixed small circle beside a growing line. *Ours, unverified:* in this reader a direction curled to four sites counts at windows narrower than about twenty sites and stops counting beyond, which is the Kaluza-Klein picture of a small direction that is invisible at large scale. The reader's dimension is then a function of scale, like the spectral dimension reviewed in Ref. [10], while the model's count d is a one-step count that calls the tube one-dimensional at every scale.

Fully curled reads four. The 4-cube, which the model's count calls zero-dimensional, embeds with four equal eigenvalues at every radius (exact). The two conventions part here by definition: the model counts the directions along which no square closes, and the reader counts the independent directions of a distance embedding. A fully curled object has many directions, all short.

The conversion's product reads one at large windows. The converted tori are six sites around (Sec. II). Their windows read 5, 4, 4 and 3 at $R = 2$ to 6 and 1 at $R = 10$, while the model's count reads 2 at nearly every vertex. *Ours, unverified:* by the reader's scale-dependent reading these products are one-dimensional at large scale, with a short closed direction, which agrees with the companion paper's finding that the conversion does not produce two large directions [7].

VII. QUESTIONS FOR THE PROGRAM

1. How should the estimator treat a lattice whose graph distance is not Euclidean, as the square lattice's is not? Is the intended reading the registered count, ε_D against an example's value, or a change in Φ or in the mutual information assigned to unlinked pairs? And does a direction curled to a few sites count as a dimension at a given scale, or is the entropy-of-a-ball exponent of Ref. [4] (its Eq. (15)), which does not grow along such a direction, the estimator the program intends when small closed directions are present?

2. Which factorization should be used for a classical wiring: one factor per vertex, one per link, or one chosen by a dynamics as in Ref. [6]? And which weighted graphs are the information graphs of some redundancy-constrained state, given that the one state built from a wiring here carries a background that, read literally, removes the geometry?

3. Ref. [5] suggests that lattice models might be "gravitized" by emerging their geometry from a state's entanglement rather than from the lattice. Is a classical wiring with a curvature energy, such as CQG, a candidate for this, that is, is there a natural state for each wiring whose reader curvature (Ref. [4], its Eqs. (36) and (40)) tracks the wiring's Ollivier curvature?

VIII. A SECOND READING: A WINDOW WITH STORED MATTER

Registered as T94 with the author's prediction (a window holding stored matter reads differently from an empty one) and computed on 3 November 2026 [8]. On the flat 32×32 torus a *stored pair* is a same-side pair at graph distance two with two common neighbours (the diagonals of a square). A stored pair carrying mutual information x_m has reader distance $\alpha = \ln x_m / \ln x_0$ in link units; the window is the unweighted ball of radius R in every case, so only the distances change, and distances are shortest weighted paths. With every stored pair at $\alpha = 1$ (a diagonal shares as much as a link) the window's metric is the Chebyshev distance and the sheet reads nearer to two dimensions: $D_{\mathrm{MDS}} = 4, 3, 3$ at $R = 3, 4, 6$ against $4, 4, 3$ with links only, with λ_3/λ_1 falling from $0.43, 0.40, 0.37$ to 0.14 at every radius. At $\alpha = 0.5$ the fit worsens ($D_{\mathrm{MDS}} = 4, 5, 5$). One stored unit at the center (the eight diagonals of the four squares meeting there,

$\alpha = 1$) adds an eigenvalue of its own: $D_{\mathrm{MDS}} = 5$ at every radius, $\lambda_4/\lambda_1 = 0.25, 0.23, 0.16$ against $0.12, 0.10, 0.09$; the held-out $R = 6$ value sits inside the range fitted on $R = 3, 4$ for that eigenvalue (0.076 in 0.066 to 0.198) and outside it for the third eigenvalue, where the amendment said in advance the fit was bad. Her prediction holds in every case at every radius; of ours, one clause and one prediction fail by the letter (the reading in Ref. [8]). The shares are placed by hand as her rule says, and no quantum state has been shown to carry them.

IX. WHAT THIS PAPER DOES NOT SHOW

No quantum state of the model is computed. The mutual information used is a constant put on the links; the one explicit state used is built from the wiring and serves only as a calibration. The readings depend on the registered threshold, which is ours, and the flat sheet's count sits on it at two radii; at $R = 10$ the registered sheet window also wrapped round its torus. The converted tori come from runs at $\lambda = 1.25$, outside CQG; the random graphs are of the published model at $\lambda = 1$; the built states are geometry only. The wirings carry none of the degrees of freedom the author takes as foundational (twist and curl), so nothing quantum in her hypothesis is tested; the quantum content of Ref. [4] enters only through the state, and a wiring stood in for it here. Agreement or disagreement between the reader and the model's count is a relation between two definitions on the same graphs, not evidence about either route's account of space, and nothing here is about the real universe.

ACKNOWLEDGMENTS

The analysis, code and text were drafted with Claude (Anthropic), in conversation with the author, who directed and checked the work. No physicist has reviewed it. The repository holds the computation, its registered predictions and results: `https://github.com/EmilySmithCreate/RecreationalPhysics`.

[1] C. A. Trugenberger, Combinatorial quantum gravity: geometry from random bits, JHEP **09**, 045 (2017), arXiv:1610.05934.

[2] C. Kelly, C. A. Trugenberger, and F. Biancalana, Self-assembly of geometric space from random graphs, Class. Quantum Grav. **36**, 125012 (2019), arXiv:1901.09870.

[3] C. A. Trugenberger, Networks as the fundamental constituents of the universe, J. Phys. Complex. **6**, 042001 (2025), arXiv:2512.17676.

[4] C. Cao, S. M. Carroll, and S. Michalakis, Space from Hilbert space: recovering geometry from bulk entanglement, Phys. Rev. D **95**, 024031 (2017), arXiv:1606.08444.

[5] C. Cao and S. M. Carroll, Bulk entanglement gravity without a boundary: towards finding Einstein's equation in Hilbert space, Phys. Rev. D **97**, 086003 (2018), arXiv:1712.02803.

[6] S. M. Carroll and A. Singh, Quantum mereology: factorizing Hilbert space into subsystems with quasi-classical dynamics, Phys. Rev. A **103**, 022213 (2021), arXiv:2005.12938.

[7] E. Smith, The curled torus burps: activated escape and conversion in a graph model of emergent geometry, technical manuscript (2026), `docs/papers/curled_torus/` in the repository below.

[8] E. Smith, `PREREGISTRATION.md`, sections T91 (with Amendment 1) and T94 (with Amendment 1), and `ASSUMPTIONS.md`, entries O174 and O178, in the repository below (2 and 3 November 2026); computations `scripts/carroll_reader.py` and `scripts/carroll_matter_window.py`, results `results/t91_carroll_reader_*.csv` and `results/t94_matter_window_*.csv`.

[9] M. C. Diamantini and C. A. Trugenberger, Topological network entanglement as order parameter for the emergence of geometry, New J. Phys. (2017), doi:10.1088/1367-2630/aa8f08, arXiv:1710.00221.

[10] S. Carlip, Dimension and dimensional reduction in quantum gravity, Class. Quantum Grav. **34**, 193001 (2017), arXiv:1705.05417.