

Metastable curled tori and their front-driven decay in combinatorial quantum gravity

Emily Smith^{1,*}

¹*Independent researcher, Charlottesville, Virginia, USA*

(Dated: DRAFT, 23 September 2026)

Combinatorial quantum gravity assigns to a four-regular bipartite graph an energy built from its total Ollivier curvature, which splits into a global term that rewards squares and a local term that removes the reward for a third square on an edge. We treat the coefficient λ of the local term as a control parameter; $\lambda = 1$ is the published model. For $\lambda \geq 1$ the flat lattice torus is the ground state exactly, and a $4 \times L$ torus, one direction curled to a single square, lies $4(\lambda - 1)$ per vertex above it. At $\lambda = 1.25$ the curled torus is metastable and decays at fixed coupling by a memoryless nucleation followed by a single front: at half conversion 98.8 to 99.8% of vertices belong to one ordered state or the other, and the released energy is exactly $4(\lambda - 1)$ per vertex. The barrier is a single local move. In a sealed system a seed of 12 energy units starts the change at every size from 48 to 192 vertices and nothing smaller does, while the release grows with the system. Sealed, the change completes only if its surroundings can absorb the release, and it leaves one defect of fixed size however large the torus. [λ scan: one sentence, pending.] The release vanishes at $\lambda = 1$. Every verdict quoted was pre-registered.

I. INTRODUCTION

In combinatorial quantum gravity (CQG) [1–3] geometry emerges from random graphs driven by a discrete Ricci curvature. The question studied so far is whether geometry forms out of the random phase sharply or smoothly: a first-order transition with hysteresis when only the global term is kept [4], a continuous or hybrid one with the full action [2, 3]. A different kind of change has not, as far as we have found, been studied in this model: one *ordered* arrangement turning into another. Such changes matter for the proposal that dark matter consists of long-lived regions of the graph stuck in a different discrete arrangement [5]. We study the simplest case the model offers, a torus with one direction curled, opening into the flat torus. It exists as a metastable state only for $\lambda > 1$; we map it along λ and follow its decay.

II. MODEL AND EXACT RESULTS

States are four-regular bipartite graphs obeying the hard-core rule (no two vertices share more than two neighbours) [2, 3]. With S the number of squares, S_e the squares on edge e and $X = \sum_e (S_e - 2)_+$,

$$H = 16(N - S) + 4\lambda X = 4 \sum_e [(2 - S_e) + \lambda(S_e - 2)_+]. \quad (1)$$

At $\lambda = 1$, H is the total combinatorial curvature of Refs. [2, 3]; λ multiplies the local (saturation) term. The hard-core rule limits $S_e \leq 3$. Three exact consequences follow.

Ground state. For $\lambda \geq 1$ every term in the sum is non-negative, so $H \geq 0$, with $H = 0$ for any graph with two

squares on every edge: the flat lattice torus. For $\lambda > 1$ nothing else reaches zero.

Ladder. The $4 \times L$ torus has $S = 5N/4$ and $X = N$ (the edges of each wrap-around 4-cycle carry three squares), so $H/N = 4(\lambda - 1)$; the 4-cube has $H/N = 8(\lambda - 1)$. Each curled direction costs $4(\lambda - 1)$ per vertex. We call $d(v)$, the number of pairs of edges at v that close no square, the local dimension: 2 on the flat torus, 1 on the curled torus, 0 on the 4-cube.

Metastability window. Enumerating every valid graph at $N = 18$ (26 classes up to relabelling) [8], states above the ground state from which every single move costs energy exist for $1 < \lambda < 1.6$, and at neither $\lambda = 1$ nor any $\lambda \geq 1.6$ (steps of 0.05 up to 1.7, coarser to 30).

The way out. The Markov chain swaps one partner each between two vertices of the same side (an edge switch preserving the bipartition; Metropolis; the hard-core rule is enforced) [8]. From the perfect $4 \times L$ torus the cheapest moves are of two kinds, counted exactly: move A ($\Delta S = -2$, $\Delta X = -4$; cost $32 - 16\lambda$; $3N$ moves) and move B ($\Delta S = -4$, $\Delta X = -10$; cost $64 - 40\lambda$; $2N$ moves). A sweep is $2N$ attempts drawn from $4N^2$ equally likely proposals, so A is offered 3 times and B twice per sweep, independent of N . The mean time to the first exit is therefore predicted without free parameters,

$$\tau(\lambda, g) = [3e^{-(32-16\lambda)/g} + 2e^{-(64-40\lambda)/g}]^{-1}, \quad (2)$$

with A the cheaper for $\lambda < 4/3$.

III. DECAY AT FIXED COUPLING

At $\lambda = 1.25$ the barrier is 12. Over twelve conditions ($g = 1.4$ to 2.5 , $N = 64$ and 144 , 16 runs each), the measured mean waiting time against the move-A term of Eq. (2) has a plain mean ratio of 0.98 and a weighted mean of 0.88 ± 0.06 , every ratio between 0.71 and 1.62, over a ninety-fold range of times; a fit of the

* emilysmithcreate@gmail.com

barrier at $N = 144$ gives 12.1 ± 0.9 [Fig. 1(a)]. Including move B lowers the predicted waiting time by 15% at $g = 1.5$; which of the two describes the data better is open (Sec. V).

The decay itself was pre-registered [8]: at $\lambda = 1.25$, $g = 1.5$, thirty decays at each of $N = 64, 96, 144, 192$, and the adjacency read when the conversion fraction first crossed 25, 50 and 75%. The predictions were a memoryless waiting time (coefficient of variation between 0.7 and 1.3), at least 80% of vertices at $d \in \{1, 2\}$ at half conversion, and the converted vertices in one piece holding at least 70% of them. All three hold at every size and on two independent sets of seeds (Table I): the change starts at one place and runs as a single front, with the two orders on either side of it and almost nothing in between [Fig. 1(b)]. The mean waiting time does not fall with N , as a local barrier offered at an N -independent rate per sweep requires.

The released energy is exact. After the change, 24 to 28 of 30 decays per size rest on the flat torus with release $4(\lambda - 1)$ per vertex to within 1%. The others settle on metastable defected sheets, read from their wiring: most often one column of the torus left curled, four vertices at $d = 1$ with $S = N + 1$, $X = 6$, costing $24\lambda - 16$ (14 at $\lambda = 1.25$); more rarely combinations of small defects of 8 to 45 units. The pre-registered verdict, a two-state change, holds at $N = 64, 96$ and 192 ; at $N = 144$ one decay changed state inside its final measuring window and the size was set aside by the energy gate as written, with all three predictions holding there too.

For contrast: at the same $\lambda = 1.25$ the change out of the random phase shows one hump in the energy distribution at every coupling, size and replica we ran, with any latent heat below 1.28 per vertex at $N = 100$ and falling with N [8]. In this model the sharp change is between orders.

IV. SEALED SYSTEM

To follow the released energy we replace the heat bath by Creutz demons [6]: C stores, each never negative; each attempted move picks one store to pay for it or be paid, and is refused if that store cannot pay. H plus the stores' contents is conserved exactly, checked to the last unit in every run, and the stores' mean reads the temperature.

A local, size-independent threshold. With one store

TABLE I. The decay of the $4 \times L$ torus at $\lambda = 1.25$, $g = 1.5$ (first run / independent rerun, thirty decays each).

N	mean wait	CV	$d \in \{1, 2\}$ at 50%	largest piece
64	871 / 660	0.91 / 0.84	0.990 / 0.998	0.98 / 0.99
96	913 / 928	1.10 / 1.02	0.990 / 0.988	0.95 / 0.96
144	856 / 743	1.01 / 0.76	0.997 / 0.997	0.92 / 0.96
192	1039 / -	0.87 / -	0.995 / -	0.85 / -

holding a seed s and nothing else, the torus converted in 8 of 8 runs for $s \geq 12$ and in 0 of 8 for $s \leq 11.5$ (seeds 4 to 16), at each of $N = 48, 64, 96, 144, 192$ (320 runs). Twelve is the cost of move A. With nothing to spare the torus never converts, although the flat torus is lower. The cost of starting is fixed while the release, $4(\lambda - 1)N$, grows with the system.

The surroundings must absorb the release. With the seed in one of C stores ($N = 64, 96, 192$; twenty runs each; 420 runs): the converted region melts into the random phase in 17 to 20 of 20 runs for $C \leq N/16$ and in 9 to 16 of 20 at $C = N/8$; at $C = N/4$ most runs end partly converted or partly melted; for $C \geq N/2$ the flat torus results in 18 to 20 of 20. The crossover scales with N , as a release of $4(\lambda - 1)N$ into C stores against a melting coupling near 3.5 predicts ($C^* \approx N/3.5$). No run stalled with both orders present and the bath below the melting coupling: the curled torus has no coexistence temperature with the flat one.

One leftover, however large. In the coldest box ($C = 2N$), the number of leftover defects per converted torus is 0.90 ± 0.31 , 0.85 ± 0.37 , 1.05 ± 0.22 and 0.95 ± 0.22 at $N = 64, 96, 192, 288$ (twenty runs each; slope 0.0003 ± 0.0003 per vertex); the defect is the curled column above (14 units), occasionally a two-vertex twist (20 units). Its position relative to where the change started is indistinguishable from uniform.

V. DEPENDENCE ON λ

[PENDING: the pre-registered map at $\lambda = 1.05$ to 1.45, all four sizes, thirty decays each. Exact inputs already fixed: the barrier $\min(32 - 16\lambda, 64 - 40\lambda)$, the waiting time of Eq. (2) (8330 sweeps at $\lambda = 1.05$ down to 22 at 1.45 for $g = 1.5$), the release $4(\lambda - 1)$ per vertex, and the curled column's share of the release, $(24\lambda - 16)/[4(\lambda - 1)N]$, which is 22% at $\lambda = 1.25$ and 72% at 1.05 for $N = 64$. Table and one figure.]

VI. DISCUSSION

At $\lambda = 1$, the published model, the flat, curled and 4-cube tori are degenerate and the change releases nothing; the $N = 4p^2$ sizes recommended for the random-to-lattice problem admit neither of the latter two. The sharp change between two orders described here therefore belongs to the family on the $\lambda > 1$ side, [and the map of Sec. V shows how it approaches $\lambda = 1$]. The same questions can be asked at $\lambda = 1$ of regions stuck in a different discrete arrangement inside a curved background [5]: whether such a region is metastable, and whether it gives way by a front with a definite release. The tools used here (exact barrier counting, front tracking, sealed baths) apply unchanged.

Not claimed: a first-order transition in the thermodynamic sense (this is the decay of a metastable state at




FIG. 1. (a) Mean waiting time of the perfect $4 \times L$ torus at $\lambda = 1.25$ against inverse coupling, sixteen runs per point, with Eq. (2) using move A only (solid) and moves A and B (dashed); nothing is fitted. (b) Share of vertices at each local dimension d when half the torus has converted, averaged over thirty decays per size.

fixed coupling, and “sharp” is used in the pre-registered sense above); anything at sizes beyond 288; why the cold box leaves exactly one defect. All runs label vertices; treating them as interchangeable [7] weights each state by its number of symmetries, which is $2N$ for the curled torus and 2 one move away, so the escape would be slower by a factor of about N , with the release unchanged.

METHODS AND DATA

Simulations use single-edge switches that preserve the bipartition, with Metropolis acceptance; the chain reproduces exact averages computed by enumerating every valid graph at $N = 16$ and 18, and the published square

density of Ref. [2] (Fig. 8a) at $N = 160$ to rms 0.005. Seeds are fixed in configuration files; every result file carries its configuration and software versions. Predictions and analysis rules were committed before the runs they concern, with every amendment recorded. Code, data, configurations and pre-registrations: [8].

ACKNOWLEDGMENTS

Code, analysis and text were drafted with Claude (Anthropic), directed and checked by the author; every kernel is tested against an independent computation before use. We thank C. A. Trugenberger for correspondence.

-
- [1] C. A. Trugenberger, J. High Energy Phys. **09** (2017) 045, arXiv:1610.05934.
 - [2] C. Kelly, C. A. Trugenberger, and F. Biancalana, Class. Quantum Grav. **36**, 125012 (2019), arXiv:1901.09870.
 - [3] C. A. Trugenberger, *Networks as the fundamental constituents of the universe*, arXiv:2512.17676.
 - [4] A. Gorsky and O. Valba, Phys. Rev. D **103**, 106013 (2021), arXiv:2101.04072.

- [5] C. A. Trugenberger, *Dark matter and dark energy in combinatorial quantum gravity*, arXiv:2409.09385.
- [6] M. Creutz, Phys. Rev. Lett. **50**, 1411 (1983).
- [7] K. Betre and N. Lewis, *Dynamical quantum multigraphs*, arXiv:2509.08296 (2025).
- [8] E. Smith, RecreationalPhysics, <https://github.com/EmilySmithCreate/RecreationalPhysics>.