THE STORY BEHIND THE CALCULATION

Two ways to read a space

A map drawn from shared quantum information,
pointed at a small model world

Emily Smith Independent researcher

3 November 2026 • Plain-language edition of the technical draft of the same date [1]

There is more than one published way to get space out of something that is not space. This paper takes two of them and asks a plain question: when they look at the same shapes, do they see the same space?




Figure 1. The question in one picture. SKETCH, NOT DATA Reading 1 belongs to the small model world this project studies; Reading 2 is a published method from quantum physics. Chapters 2 and 3 explain each, and Chapter 6 compares them.

Who is saying what

The project keeps four kinds of statement apart, and this edition tags them where it matters, so that a sentence is never quietly promoted from one kind to another.

- EXACT Arithmetic, or a complete listing of every possibility: true whatever a run does.
- MEASURED A computation whose rules and predictions were written down before it ran, with its verdict. MEASURED, EXPLORATORY means a computation that was not written down first, or a closer look at results already read.
- OUR READING, UNVERIFIED A reading of a result, offered by the author and her AI assistants and checked by no physicist.
- THE AUTHOR'S CLAIM The author's hypothesis about reality, stated as hers.
- PUBLISHED Other researchers' published results.

Numbers in square brackets point to the sources listed at the end.

1 THE PLACE WE START

Where does space come from, and how would we know it when we saw it?

The author. Emily Smith is a software engineer, not a physicist, and also a startup founder and a foster mom. Her recreational time is limited and the computing bills are her own. Fittingly, the project repository is called RecreationalPhysics. She works, on the side, to map her intuition about how reality is organized onto what humanity knows about the universe, and to share what she learns with other curious people.

Her hypothesis. THE AUTHOR'S CLAIM Space is one settled arrangement of something deeper, which she calls X because nobody knows what it is. X is a specific arrangement, curled up tight, and the space we live in arose when it opened, releasing a definite amount of energy. So for her it matters whether a curled direction counts as a direction at all, and to whom.

The approach. Nothing in a computer can prove a hypothesis about reality. What a computer can do is take a small world whose rules are written out in full, in which space is an arrangement and not a given, and ask precise questions of it. The project's first paper [7] did that with one such world. This one borrows a tool from a different corner of physics, a way of reading space out of quantum information, and points it at that world's shapes. The calculation, and what it was expected to show, were written down before it ran, with one size held back to test the predictions.

The parts of this paper. Chapter 2 introduces the two published ideas that meet here. Chapter 3 explains the borrowed tool. Chapter 4 states the question and the one choice needed to ask it. Chapter 5 says how the shapes were read, Chapter 6 what was found, Chapter 7 what it means, and Chapter 8 what is next, including the questions this raises for the tool's authors.

A toy. Physicists call a small world of that kind a **toy model**: simple enough that every consequence of its rules can be counted or simulated, and not to be mistaken for the real thing. Ours is **the toy** from here on.

2 TWO PUBLISHED IDEAS

A network that curves, and space read from quantum information

The first idea: a network whose energy is curvature

PUBLISHED Carlo Trugenberger, Christy Kelly and Fabio Biancalana built a model of space with no space in it [5, 6]: only **dots** and **links**, with no positions. A dot has no place, only neighbors. The dots come in two kinds, black and white, and a link always joins a black dot to a white one. Every dot has exactly four links, and no two dots share more than two neighbors. Four links that close a loop make a **square**. The list of which dots are linked to which is the network's **wiring**.

PUBLISHED The energy is counted link by link from squares: a link that borders exactly two squares is free, a missing square costs 16 units, and a third square on a link costs 4. With those prices the energy is a network version of the curvature in Einstein's theory of gravity. This is **the published model**. The toy keeps all of it and turns one price into a knob: a third square costs 4λ, where λ (lambda) can

be turned up or down. Only $\lambda = 1$ is the published model.




Figure 2. The toy's two ingredients for this paper. EXACT Left: at a dot, four links make six pairs; a pair at a right angle closes a square, and a pair that runs straight through closes none. The **toy's count** d is the number of pairs that close no square. Right: three shapes, each a grid whose edges are joined both ways round, so none has an end. The sheet is a fat doughnut, long both ways. The tube is a thin ring: its thin way round is only four steps, so walking round it is itself a square, and that direction no longer counts. The knot is curled both ways. (The doughnut and ring are drawn bent; the network itself has no positions and is bent nowhere.)

Two words are kept apart from here on. **Flat** is about each dot's surroundings: every link on exactly two squares, the wiring of a plain grid, at zero energy. **Wide** is about the whole shape: how many steps it takes to walk round it and come back. The sheet is flat and wide; the tube is neither.

EXACT With the knob above 1, the tube sits $4(\lambda - 1)$ units per dot above the flat sheet, and gives that much off if it opens. The author calls that release **the burp**. MEASURED Paper 1 [7] watched tubes open at $\lambda = 1.25$. What they opened into was flat at every dot but still narrow: a narrower doughnut whose shortest way round is six steps instead of four. Those end states are called **the opened tubes** here. This paper reads some of them. THE AUTHOR'S CLAIM In the author's picture the opened tube is the toy's end state and not a stage of her change: when the deeper ingredients are not laid out as a plane, two dimensions need not pass through a narrow doughnut on the way to three. Her three dimensions are a projection from the two-dimensional arrangement, a result of the burp's energy wiring. The projection ends inside a black hole, where the arrangement can project again, and that closing is her strange loop.

Her positions on the toy. THE AUTHOR'S CLAIM The author holds that, taken as a whole, no network she uses has λ above 1: it is 1 or below everywhere. The exception is the two-dimensional inner shell of a black hole, looked at one part at a time, where the arrangement sets a barrier above 1. She holds that the black dots and the white dots are two distinct kinds, and that within a kind every dot is the same. And she holds that the flat sheet of dots is a simplified stand-in for the two-dimensional picture only. The deeper ingredients are two degrees of freedom, twist and curl, with their own rules for how they relate to each other and to themselves; they are not two directions to move along. None of this is assumed below, and none of it is tested here.

The second idea: space read from quantum information

PUBLISHED ChunJun Cao, Sean Carroll and Spyridon Michalakis start somewhere else entirely [2, 3]. In quantum physics, everything that can be known about a system is packed into one object, its **state**.

Split a system into parts, and the parts of a state can be tied together so that learning about one tells you something about another, with nothing in between: that tie is **entanglement**. Their idea is to begin with a quantum state and no space at all, measure how strongly each pair of parts is tied, and read a geometry off those ties. Their recipe for the reading is called **the reader** here. They say plainly that they assume no energy rule, and that the split into parts is taken as given [2]. Their second paper goes on toward Einstein's equation [3].

The two ideas meet at one point. Each ends with a list of which things are close to which, and each asks what space that list describes.

The point to keep: The toy is a network of black and white dots whose energy is a curvature; it counts the directions at a dot, d: two on the flat sheet, one on the tube, none on the knot. Carroll and his colleagues read space from how strongly the parts of a quantum state are tied. This paper hands the toy's shapes to their reader.

3 THE TOOL

How do you draw a map from a table of shared information?

The reader works in four steps (Figure 3).

Step 1: shared information. PUBLISHED The strength of the tie between two parts is measured as **shared information**: how much knowing one part tells you about the other, counted in bits. Two coins that always land the same way share one bit: see one and you know the other. Two unrelated coins share none. Physicists call it mutual information.

Step 2: sharing becomes distance. PUBLISHED The more two parts share, the closer they are put. Each pair that shares anything gets a direct line whose length is set by how much it shares, and a pair that shares nothing gets no direct line. In their examples the length is minus the natural logarithm of the share, measured against the most two parts can share:

Share, as a fraction of the most possible	all of it	a half	a tenth	a thousandth
Length of the direct line	0	0.69	2.3	6.9

The distance between any two parts is then the shortest chain of direct lines between them.

Step 3: a table of distances becomes a map. PUBLISHED The back of an old road atlas has a mileage chart: the driving distance between every pair of cities, and nothing else. From that chart alone you could redraw the map, putting each city where its distances to all the others fit best. The reader does exactly this with the parts. The standard recipe is called multidimensional scaling.

Step 4: count the directions. PUBLISHED A mileage chart for cities on flat ground fits a flat page perfectly: two directions, north and south, east and west, and nothing left over. The recipe reports how much of the map's **spread** lies along each of its directions, largest first; for cities on flat ground, two directions carry all of it. When the distances do not fit any page exactly, a third and fourth direction carry some spread too, and someone has to decide how much counts. Their papers leave that

open; their own examples call a chain of 100 quantum parts one-dimensional and a patch of a known two-dimensional quantum system two-dimensional [2]. For this paper the rule was fixed before any calculation: a direction counts if its spread is at least a tenth of the largest. The tenth is a choice, not a law. EXACT The spread is a squared quantity, so a tenth of the spread is about a third of the reach: a direction counts if the dots reach along it at least about a third as far as they reach along the longest direction. The number of directions that pass is **the reader's count**.




Figure 3. Carroll's reader in four steps. PUBLISHED Steps 1 to 3 are the published recipe [2]. SKETCH, NOT DATA The numbers are made up for the drawing. The cutoff in step 4, a tenth of the largest spread, was chosen for this paper before any calculation; the published papers leave it open.

The point to keep: Shared information becomes distance, a full table of distances becomes a map, and the number of directions the map needs is the reader's count. Where the map does not fit a page exactly, the count depends on a cutoff, which here is a tenth of the largest spread, fixed in advance.

4 THE QUESTION

What does the reader see when it is handed a wiring?

The toy has no quantum state; it has a wiring. To hand it to the reader, one choice is needed.

The one choice

Give each dot one quantum part, and give every linked pair the same share of information, and every unlinked pair none. The toy itself says nothing more: in it, two dots are linked or they are not, and every link is like every other.

THE AUTHOR'S CLAIM In the author's picture this is where photons and gravitational waves travel: between dots of different kinds. In a cube of her three-dimensional picture, gravity runs along the edges and light along the body diagonals, and both join a black corner to a white one; the toy's links are its only black-to-white connections, so they stand for both. Matter, energy and twist are stored between dots of the same kind, which are never linked. So the one choice puts the shared information where her waves travel, and nothing where her matter is stored.

EXACT Then every direct line has the same length, and the distance between two dots is simply the number of steps along links between them, times that length. The map's directions all scale together, so the reader's count does not depend on how large the share is. With this choice, the reader is just a map drawn from step counts.

Does a quantum state like that exist?

PUBLISHED One published state is built from a wiring: Marcelo Diamantini and Carlo Trugenberger turned the link list of a network into a quantum state on two copies of its dots [4]. EXACT Computed for this paper on an 8 by 8 flat sheet: a linked pair shares 0.0117 bits, about 33 times what every other pair shares, 0.00036 bits. So the sharing follows the wiring.

EXACT But that faint share between every pair matters. Carroll's rule gives every pair that shares anything a direct line, and a faint share still makes a line shorter than two links:

One link (a linked pair's share)	length 5.14
Two links in a row	length 10.28
A direct line between any other pair (the faint share)	**length 8.63**

So no pair is ever more than 8.63 apart, every pair looks close, and the map comes out as one direction followed by five at about three quarters of it: no shape at all. Subtract the faint background first, and the wiring's map returns. OUR READING, UNVERIFIED So a quantum state whose sharing follows a wiring exists, but only after a step, subtracting the background, that Carroll's papers do not take. The state is built from the wiring, so it checks the reader; it does not test how a geometry could arise.

Given the toy's shapes, does Carroll's reader count the same directions as the toy does:
two on the sheet, one on the tube, none on the knot?

Why we went here. Her picture of the birth of space is a curled direction opening. If an independent reader, built in quantum physics with no knowledge of the toy, did not see a curled direction as different from an open one, the toy's count would be a convention of the toy and nothing more.

5 WHAT WE TRIED

Why not read the whole doughnut?

Every way round counts twice

EXACT A loop cannot be laid along a line. Eight dots in a ring, each one step from the next, need a page: placed on a circle they fit, and a circle takes two directions. So every way round that a shape wraps takes a pair of directions on the map (Figure 4, left). The flat sheet wraps two ways, so whole it reads four; the tube wraps two ways, so it reads four; the knot reads four. Read whole, every doughnut looks much the same: four directions each.

Windows

So the reader is given a **window** instead: one dot, the center, and every dot within R steps of it, with their distances taken through the whole network. A window is what someone living at the center could reach on foot within R steps. Its width is $2R+1$ dots across. Windows were read at $R = 2$, 3, 4 and 6; the widest, $R = 10$, was held back, and computed only after a prediction for it had been written from the smaller ones.




Figure 4. Why whole doughnuts cannot be read, and what is read instead. EXACT Left: a ring cannot be laid on a line without stretching some step, so the map spends two directions on every way round. Right: a window is a diamond of dots around one center (Chapter 6 says why it is a diamond and not a circle). The reader maps only those dots.

The shapes, and what was expected

Three shapes were built exactly: a flat sheet 32 by 32 dots, a tube 4 around and 64 along, and the knot. Two kinds were taken from earlier runs: four opened tubes, each of 1,024 dots, read from 512 centers apiece; and four **random tangles**, warm networks of the published model in which the wiring is jumbled, 196 dots each, read from every dot. These sizes are conveniences, not special values.

Written down before any window was computed: the knot would read four; the flat sheet two; the tube no more than the sheet, with its curled direction too faint to count by $R = 4$; the opened tubes the

same as the sheet; the tangles four or more.

6 WHAT WE FOUND

Five shapes, read window by window

The knot reads four

EXACT At every window size the knot reads four directions, all with equal spread. The toy counts none. That was expected, and it is a difference of definitions, not a mistake on either side: the toy counts directions with room in them, and the reader counts how many directions a map needs. A knot has many directions, all short. The 16 dots of the knot are the corners of a four-dimensional cube, with that cube's step distances, and drawing 16 corners with those distances takes four axes. A knot of string points every way and goes nowhere.

The flat sheet reads three, sometimes four

EXACT The flat sheet reads four, four, four and three directions at $R = 2$, 3, 4 and 6. Its first two directions are equal, as a page's should be. But a third carries between about two fifths and a half of the largest spread at every size, and a fourth sits close to the cutoff: at $R = 4$ it is 0.101 of the largest, a hair above the tenth, which is why the count there is four and not three. MEASURED The prediction of two failed.

The reason is how distance is counted. In a city laid out in square blocks, a taxi cannot cut across: three blocks east and three north is six blocks of driving, while a crow flies about four and a quarter. Steps along a grid are taxi distances. The dots exactly three steps from the center do not form a circle but a diamond (Figure 5), and no flat page holds a diamond of equal distances, so the map reaches for extra directions.




Figure 5. Taxi distance is not straight-line distance. EXACT On a grid, steps along links are what a taxi drives in a city of square blocks. Equal step counts make a diamond, not a circle, and the map needs more than a page to hold it. The extra directions say more about the distance assigned to a grid than about the toy. EXACT If instead the sharing between every pair fell off with straight-line distance, the map of any patch of a flat sheet would need exactly two directions.

The tube's curled direction is seen up close and fades from afar

Think of a garden hose. An ant on it can walk round it as easily as along it; from across the yard, the hose is a line. The tube is a hose four dots round.

EXACT In the tube's windows the long direction carries the largest spread, and the curled direction shows up as a pair. Its share of the largest spread falls as the window widens: 0.61, 0.43, 0.31 and 0.20 at $R = 2$, 3, 4 and 6 (Figure 6). MEASURED So the ranking held: at $R = 4$ the tube read three directions against the sheet's four. But the prediction that the curled direction would be too faint to count by then failed: it was four times stronger than predicted.

MEASURED From $R = 4$ and 6 alone, a rule was fitted, the share falling as the window's width to the power -1.24, and written down before the widest window was computed: at $R = 10$, 0.108, with a range of 0.076 to 0.141. **The widest window measured 0.112.** The prediction held.




Figure 6. The hose seen from farther and farther away. Points at $R = 2$ to 6: EXACT. The fitted rule and its prediction: MEASURED, written down before the held-out window at $R = 10$ (width 21) was computed; the measured 0.112 lands inside the predicted range of 0.076 to 0.141. The dotted line is the first guess, which missed by a factor of four. Extended (not computed), the fitted rule crosses the cutoff at windows about 22 dots across.

The opened tubes read one direction from afar

MEASURED The opened tubes read five, four, four and three directions at $R = 2$, 3, 4 and 6, in nearly every one of their 512 windows. At $R = 10$ they read one, in 506 of 512. The prediction that they would read like the flat sheet failed at the smallest window and at the largest. OUR READING, UNVERIFIED The reason is the hose again: the opened tubes are narrow, their shortest way round only six steps, so a wide window wraps right round them, and from that far the reader calls them a line.

The random tangles read many

MEASURED The jumbled networks read four or more directions at $R = 3$ in 778 of 784 windows, and more as the window widens, up to 17, as expected: a tangle has no shape a page can hold.

One miss that was the window's fault

MEASURED At $R = 10$ the flat sheet was predicted to read three and read four, its fourth direction again right on the cutoff. EXACT A check made after the reading, not written down in advance, found

the cause: a window ten steps out holds pairs twenty steps apart, more than halfway round a sheet 32 dots wide, so some of their shortest paths go round the back of the doughnut. On a sheet 64 by 64 the same window reads three, as predicted. The verdict stays as scored; the fault was in the choice of sheet.

The point to keep: The knot reads four where the toy counts none. The flat sheet reads three or four, because grid steps are taxi distances. The tube's curled direction is counted up close and fades as the window widens, at the rate predicted in advance for the held-out window. The narrow opened tubes read as a line from afar. By the words fixed beforehand, the verdict is MIXED.

7 WHAT IT MEANS

Two readings that agree in part

Shape	The toy's count	The reader's count
Flat sheet	2	3 or 4, from taxi distance
Tube	1	3 up close; the curled pair fading as the window widens
Knot	0	4
The opened tubes	2 at nearly every dot	5 up close, falling to 1 in the widest window

OUR READING, UNVERIFIED Where they agree: up close, both rank the flat sheet above the tube. And both find that the burp left something narrow. The toy says so by the shape of the opened tube, flat at every dot but six steps round; the reader says so by calling it a line from afar. That matches paper 1's finding that the opening did not make two wide directions [7].

OUR READING, UNVERIFIED Where they differ: the knot, and, more deeply, what a direction is. The toy's count is taken at one dot and is the same at every scale: the tube is one-dimensional, up close or far away. The reader's count depends on how far it looks, like the hose. A direction curled to four dots counts in a window narrower than about twenty dots and stops counting beyond. That is an old idea in physics, a small curled direction invisible from far away, here showing up in a quantum reader without being put in.

OUR READING, UNVERIFIED For the author's picture, this is a modest point in her favor and a warning together. A curled direction is real to an outside reader: up close, it counts. But whether it counts depends on scale, so "how many directions" has no single answer until a scale is named.

Why we went here. Where two readings of one network agree, the agreement is a fact about the network; where they part, the parting shows what each definition is really measuring.

The point to keep: The two readings agree on the order of sheet and tube up close and on the narrowness of what the burp leaves; they part on the knot and on whether dimension depends on scale. This compares two definitions on the same shapes. It says nothing directly about the real universe.

8 WHAT COMES NEXT

Three questions for the reader's authors

Here is everything this paper asked, with the answer and the kind of answer it is.

The question	The answer	Kind
What does the one choice do?	With the same share on every link, distance becomes steps along links, and the share's size drops out.	EXACT
Can a quantum state carry the toy's wiring?	One published state does, once its faint background is subtracted.	EXACT
What does the reader see in the knot?	Four directions, where the toy counts none.	EXACT
In the flat sheet?	Three or four, where the toy counts two: grid steps are taxi distances.	EXACT
In the tube?	Its curled direction up close, fading as the window widens, at the predicted rate.	MEASURED
In the opened tubes?	One direction in the widest window.	MEASURED
Do the two readings agree?	In part.	OUR READING, UNVERIFIED

What this does not show. No quantum state of the toy was computed: the sharing was put on the links by hand, and the one real state used serves only as a check. The counts depend on the cutoff, a choice, and the flat sheet sits right on it at two window sizes. The opened tubes come from runs at $\lambda = 1.25$, outside the published model. The wirings handed to the reader carry none of the author's deeper ingredients, twist and curl, so nothing quantum in her picture is tested here: the quantum part of Carroll's method enters only through the state, and the toy supplied a wiring in its place. Nothing here is about the real universe.

A window with matter in it

In the author's picture matter, energy and twist are stored between dots of the same kind, so a share placed on those pairs stands for stored matter, and the question is whether a window holding some reads differently from a window of empty space. THE AUTHOR'S CLAIM Her prediction, recorded on 3 November before any design: it does.

MEASURED Registered and computed the same day, on the flat sheet, with the same windows as before so that only the distances change. A **stored pair** is two dots of the same kind two steps apart that share two neighbors: the two diagonals of a square. Give every stored pair of the sheet as much share as a link, and the sheet reads *nearer* to two dimensions, not further: three directions at $R = 4$ and 6 where empty space reads four and three, and the spurious third direction falls from about 0.4 of the largest to 0.14 at every window size. With every diagonal as short as a link, the distance in the window is the chess king's, which fits a flat map better than the taxi's does. Give the stored pairs *more* share

than a link and the map gets worse. Put one stored unit at the center only, the eight diagonals of the four squares that meet there, and the reader adds a direction of its own: five directions against empty space's four or three at every window size tried, up to 13 dots wide, with the held-out size predicted in advance from the two smaller ones. So the window with matter in it reads differently in every case, as she predicted.

OUR READING, UNVERIFIED Under her storage rule a sheet with matter stored everywhere is a better-behaved map for Carroll's reader than empty space is, and a single stored unit is a direction the reader counts. Whether shares between dots of the same kind are matter, the toy cannot say; the shares were put on by hand as her rule says.

What is next on our side

A state built from twist and curl. The reader was made for a quantum state, and the one state used here was built from a wiring. Carroll starts from a state of many parts with no geometry assumed, and reads the geometry out of how the parts share information. The author starts from an arrangement with rules. The two meet if the arrangement is read as a state: THE AUTHOR'S CLAIM the state she has in mind is the Klein bottle of one-sided loops, the inner shell of a black hole, whose parts would carry the allowed sets of curl, twist and spin in place of 0 and 1, with her rules for which sets may sit together tying the parts. Her rules as of 3 November: curl is continuous, from -1 to 1, with the two signs matter and antimatter at equal size; twist comes in thirds, which combine into a whole twist across several diagonals and meet light only at whole values, and in halves, which make one-sided matter, dark matter, the building blocks of the Klein bottle; only a full curl, a loop, stores energy stably; the dark-matter particle holds the most, can take no more curl or spin, and meets gravity but not light, since light travels the body diagonals and not the edges. No such state exists yet. OUR READING, UNVERIFIED Three published computers are its nearest relatives, and none gets the quantum part from counting: Gorard's Wolfram model follows every order of a rewriting rule at once and puts the phases in by hand; Lloyd's has quantum bits at its gates, so quantum mechanics is an input; 't Hooft's has one real state at a time and meets Bell's test by denying that the settings are freely chosen. Her "plays out all the options" sits between them, and which she means is hers to say. EXACT On the toy's complete class spaces at 16 and 18 dots, the count over paths of the toy's move and the count over renamings of an arrangement are not the same count and run opposite ways: the classes with the most renamings collect the fewest paths. So "all the options" and "versions" are two different counts in this toy.

The questions

The paper ends with three questions for Carroll and his colleagues, in plain words here.

1. **Taxi distance and short loops.** How should the reader treat a grid whose step distance is not straight-line distance: by a cutoff, by some measure of how badly the map fits, or by changing how much unlinked pairs share? And should a direction curled to a few dots count at a given scale, or is a different measure from their first paper [2], how the information inside a region grows with its size, the one they intend when small curled directions are present?
2. **Which parts.** Should a wiring be split into one quantum part per dot, as here, or one per link? And which tables of sharing can come from an allowed quantum state at all, given that the one state built from a wiring carries a background that, taken literally, erases the geometry?
3. **Gravity.** Their second paper suggests that a network model might gain gravity by growing its

geometry from entanglement instead of from the network [3]. Could a wiring with a curvature energy, like the published model, be such a case: is there a natural quantum state for each wiring whose curvature, as the reader sees it, follows the wiring's own?

The reader sees a curled direction up close.
What quantum state would make it see the same space?

Continuing into the technical paper

This edition follows the technical draft [1], which holds the definitions with their equation numbers, the full tables and the scoring of each prediction. Every number above is taken from it or from the project's record. Its sections run in the same order as the chapters here: the toy (Chapter 2), the reader and its one choice (3 and 4), windows (5), results (6) and discussion and questions (7 and 8).

A few translations for the next step

In the paper	What to keep in mind
Vertex, edge, graph	Dot, link, network.
Bipartite	The two kinds of dot, black and white; links join different kinds.
Mutual information	Shared information, in bits.
$\Phi = -\ln$	Carroll's rule turning a share into a length.
Multidimensional scaling (MDS)	Drawing the map from a table of distances.
Eigenvalues λ_i	The map's spread along each direction, largest first. (The paper's λ_i are not the toy's knob λ.)
D_{MDS}	The reader's count: directions whose spread is at least 0.1 of the largest. The 0.1 is a choice.
d	The toy's count: pairs of links at a dot that close no square.
Graph distance	Steps along links; on a grid, a taxi distance.
Ball of radius R	A window: every dot within R steps of the center.
Held out	Computed only after its prediction was written from the smaller windows.
MIXED	The verdict word fixed beforehand for an outcome in which some predictions held and some failed.

Where the evidence is recorded

The technical draft is in the folder `docs/papers/carroll_reader/` of the project repository, with the two checks made while writing it (the quantum state of Chapter 4 and the window check of Chapter 6). The predictions and their scoring are section T91 of `PREREGISTRATION.md`, with its amendment and addendum; the readings are entries O174 and O175 of `ASSUMPTIONS.md`; the computation is `scripts/carroll_reader.py`.

Disclosure

The author used Claude (Anthropic) to assist with code, analysis and writing, and is responsible for the result. This edition was drafted with Claude from the technical draft and the project's record. Its analogies and pictures introduce no new calculations or evidence. No physicist has reviewed any of it.

Sources

[1] E. Smith, *Does entanglement geometry see the same space? The mutual-information reader of Cao, Carroll and Michalakis applied to the wirings of a graph model of emergent geometry*, technical draft, 3 November 2026, `docs/papers/carroll_reader/` in the project repository.

[2] C. Cao, S. M. Carroll, and S. Michalakis, *Space from Hilbert space: recovering geometry from bulk entanglement*, Phys. Rev. D **95**, 024031 (2017), arXiv:1606.08444.

[3] C. Cao and S. M. Carroll, *Bulk entanglement gravity without a boundary: towards finding Einstein's equation in Hilbert space*, Phys. Rev. D **97**, 086003 (2018), arXiv:1712.02803.

[4] M. C. Diamantini and C. A. Trugenberger, *Topological network entanglement as order parameter for the emergence of geometry*, New J. Phys. (2017), doi:10.1088/1367-2630/aa8f08, arXiv:1710.00221.

[5] C. A. Trugenberger, *Networks as the fundamental constituents of the universe*, J. Phys. Complex. **6**, 042001 (2025), arXiv:2512.17676.

[6] C. Kelly, C. A. Trugenberger, and F. Biancalana, *Self-assembly of geometric space from random graphs*, Class. Quantum Grav. **36**, 125012 (2019), arXiv:1901.09870.

[7] E. Smith, *The curled torus burps: activated escape and conversion in a graph model of emergent geometry*, technical manuscript, 14 October 2026, with its plain-language edition, `docs/papers/curled_torus/` in the project repository.