

APPENDIX

Theorem 6 (Theorem 7 in [14]). *If x and y are confusable, then they have to be in one of the following forms.*

(A) $x = u\alpha\omega\beta v, y = u\bar{\alpha}\omega\bar{\beta}v$, where α and β are alternating sequences of length at least two.

(B) $x = u\alpha\bar{a}\gamma\bar{b}\beta v, y = u\alpha\bar{a}\gamma\bar{b}\bar{\beta}v$, where α, β and γ are alternating sequences. Here, α is of length at least two and ends with a , β is of length at least two and starts with b , and γ starts with \bar{a} and ends with \bar{b} .

Theorem 7 (Theorem 4.1 in [13]). *For two sequences $x = u\tilde{x}v$ and $y = u\tilde{y}v$ in Σ^n with $d_L(x, y) \geq 2$, if $|\mathcal{D}_2(x) \cap \mathcal{D}_2(y)| \geq 5$, then one of the following holds:*

(A) $\tilde{x} = a\omega b$ and $\tilde{y} = \bar{a}\omega\bar{b}$, where a and b are alternating sequences of length at least two and ω is a combination of periodic sequences. More precisely, suppose a ends with α and b starts with β' , then $\omega = u'\omega'v'$ and one of the following holds:

- u' is a run sequence starting with α , ω' is an alternating sequence of length at least one starting with $\bar{\alpha}$ and ending with β' , v' is a run sequence starting with $\bar{\beta}'$
- u' is a run sequence starting with $\bar{\alpha}$, ω' is an alternating sequence of length at least one starting with α and ending with β' , v' is a run sequence starting with β' .

(B) $\tilde{x} = a\bar{\alpha}\omega\bar{\beta}b, \tilde{y} = a\alpha\bar{\omega}\bar{\beta}b$, where a is an alternating sequence of length at least two ending with α , ω is an alternating sequence starting with $\bar{\alpha}$ and ending with $\bar{\beta}'$, and b is an alternating sequence of length at least two starting with β' .

Particularly, $|\mathcal{D}_2(x) \cap \mathcal{D}_2(y)| = 6$ if and only if $\tilde{x} = a\omega b$ and $\tilde{y} = \bar{a}\omega\bar{b}$, where a is an alternating sequence of length at least two ending with α , b is an alternating sequence of length at least two starting with β' , and one of the following holds:

- if ω starts with α' and ends with β' , ω is a alternating sequence of length at least two;
- if ω starts with α' and ends with $\bar{\beta}'$, ω is a alternating sequence of length at least three;
- if ω starts with $\bar{\alpha}'$ and ends with β' , ω is a alternating sequence of length at least three;
- if ω starts with $\bar{\alpha}'$ and ends with $\bar{\beta}'$, ω is a alternating sequence of length at least two;

Lemma 10 (Lemma 12(ii) in [14]). *Suppose that $x = \alpha r\beta$ and $y = \bar{\alpha} s\bar{\beta}$ and $d_L(x, y) \geq 2$. Set $\mathcal{S} = \mathcal{D}_2(x) \cap \mathcal{D}_2(y)$. If the first two bits of x are equal, i.e. $r = \alpha r'$, then $|\mathcal{S}| \leq 4$. Hence by symmetry, if $r = r'\beta$ or $s = \bar{\alpha} s'$ or $s = s'\bar{\beta}$, i.e., the first or last two bits of x or y are equal, then $|\mathcal{S}| \leq 4$.*

Lemma 7 Let x, y be two binary sequences of length n such that $d_L(x, y) \geq 3$, if $\mathcal{D}_3(x) \cap \mathcal{D}_3(y) \geq 18$, then $x = uas\omega^1 tbv, y = u\bar{a}\bar{s}\omega^2 \bar{t}\bar{b}v$, where

- u, v are the longest common prefix and suffix of x, y ,
- a, b, s, t are maximal alternating sequence of length at least 2.
- $\omega^1 \neq \omega^2$.

Proof: According Lemma 6, if $\mathcal{D}_3(x) \cap \mathcal{D}_3(y) \geq 18$, then they are of the form $x = uaw^1bv, y = u\bar{a}\bar{\omega}^2\bar{b}v$, such that

- u, v are the longest common prefix and suffix of x, y ;
- a, b are maximal alternating sequences with $|a|, |b| \geq 2$;
- ω^1 and ω^2 are two distinct sequences of length ℓ , where $\omega_1^1 = \omega_1^2, \omega_\ell^1 = \omega_\ell^2$.

Suppose a starts with i , ends with α , b starts with β ends with j . For the sequence x, y satisfying the above requirements, there are four possibilities as follows:

- 1) $x = uaa\omega_2^1 \dots \omega_{\ell-1}^1 \beta bv, y = u\bar{a}\bar{\alpha}\omega_2^2 \dots \omega_{\ell-1}^2 \bar{\beta} \bar{b}v$;
- 2) $x = uaa\omega_2^1 \dots \omega_{\ell-1}^1 \bar{\beta} bv, y = u\bar{a}\bar{\alpha}\omega_2^2 \dots \omega_{\ell-1}^2 \bar{\beta} \bar{b}v$;
- 3) $x = uaa\bar{\omega}_2^1 \dots \omega_{\ell-1}^1 \beta bv, y = u\bar{a}\bar{\alpha}\omega_2^2 \dots \omega_{\ell-1}^2 \bar{\beta} \bar{b}v$;
- 4) $x = uaa\bar{\omega}_2^1 \dots \omega_{\ell-1}^1 \bar{\beta} bv, y = u\bar{a}\bar{\alpha}\omega_2^2 \dots \omega_{\ell-1}^2 \bar{\beta} \bar{b}v$.

where (3)(4) are equivalent to (1)(2), so it is sufficient to discuss only (1)(2).

Denote $\omega_3^1 \dots \omega_\ell^1$ as c , $\omega_3^2 \dots \omega_\ell^2$ as d , $\mathcal{D}_3(a\omega^1 b) \cap \mathcal{D}_3(\bar{a}\omega^2 \bar{b})$ as \mathcal{S} . Without prejudice to generality, let $\omega_1^1 = \omega_1^2 = \alpha$, and We proceed with the following three cases.

Case (A): $\omega_2^1 = \alpha, \omega_2^2 = \alpha$

$$\begin{aligned} a\omega^1 b &= a \alpha \alpha c b = i \dots \alpha \alpha \alpha c \beta \dots j \\ \bar{a}\omega^2 \bar{b} &= \bar{a} \alpha \alpha d \bar{b} = \bar{i} \dots \bar{\alpha} \alpha \alpha d \bar{\beta} \dots \bar{j} \end{aligned}$$

Since a, b are alternating sequences, $\mathcal{S}_j^i = (a\alpha) \circ (\mathcal{D}_3(\alpha c \beta) \cap \mathcal{D}_1(d)) \circ (b_2 \dots b_{|b|})$. According to Lemma 4, $|\mathcal{S}_j^i| \leq 4$.

$\mathcal{S}_j^i = (a\alpha) \circ (\mathcal{D}_2(\alpha c) \cap \mathcal{D}_2(d\bar{\beta})) \circ (b_1 \dots b_{|b|-1})$. According to Lemma 1, $|\mathcal{S}_j^i| \leq 6$.

$\mathcal{S}_j^i = (a_2 \dots \alpha_{|a|}) \circ (\mathcal{D}_2(\alpha \alpha c \beta) \cap \mathcal{D}_2(\bar{\alpha} \alpha \alpha d)) \circ (b_2 \dots b_{|b|})$. We know that the first bit of the sequence of centers of the two error balls is different, then according to Lemma 10, $|\mathcal{S}_j^i| \leq 4$.

$\mathcal{S}_j^i = (a_2 \dots \alpha_{|a|}) \circ (\mathcal{D}_1(\alpha \alpha c) \cap \mathcal{D}_3(\bar{\alpha} \alpha \alpha d \bar{\beta})) \circ (b_1 \dots b_{|b|-1})$. The sequence in $\mathcal{D}_1(\alpha \alpha c) \cap \mathcal{D}_3(\bar{\alpha} \alpha \alpha d \bar{\beta})$ only can begin with α , $|\mathcal{D}_1(\alpha \alpha c) \cap \mathcal{D}_3(\bar{\alpha} \alpha \alpha d \bar{\beta})| = |\mathcal{D}_1(c) \cap \mathcal{D}_2(d\bar{\beta})|$, then according Lemma 5, $|\mathcal{S}_j^i| \leq 3$.

In this case, $\mathcal{D}_3(x) \cap \mathcal{D}_3(y) = |\mathcal{S}| \leq 17$.

Case (B): $\omega_2^1 = \alpha, \omega_2^2 = \bar{\alpha}$

$$\begin{aligned} a\omega^1 b &= a \alpha \alpha c b = i \dots \alpha \alpha \alpha c \beta \dots j \\ \bar{a}\omega^2 \bar{b} &= \bar{a} \alpha \bar{\alpha} d \bar{b} = \bar{i} \dots \bar{\alpha} \alpha \bar{\alpha} d \bar{\beta} \dots \bar{j} \end{aligned}$$

Since a, b are alternating sequences, $\mathcal{S}_j^i = a \circ (\mathcal{D}_3(\alpha \alpha c \beta) \cap \mathcal{D}_1(\bar{\alpha} d)) \circ (b_2 \dots b_{|b|})$. There is at most one sequence d that begins with α and two sequences $\mathcal{D}_1(c\beta) \cap \mathcal{D}_1(\bar{\alpha} d)$ that begins with $\bar{\alpha}$ in the set $\mathcal{D}_3(\alpha \alpha c \beta) \cap \mathcal{D}_1(\bar{\alpha} d)$ (It request $\mathcal{D}_3(\alpha \alpha c \beta) \cap \mathcal{D}_1(\bar{\alpha} d)$ begins with $\bar{\alpha}$ and Except for the common prefix and common suffix, the remainder of $c\beta$ and $\bar{\alpha} d$ are alternating sequences with different start symbols). Thus, $|\mathcal{S}_j^i| \leq 3$.

$\mathcal{S}_j^i = a \circ (\mathcal{D}_2(\alpha \alpha c) \cap \mathcal{D}_2(\bar{\alpha} d \bar{\beta})) \circ (b_1 \dots b_{|b|-1})$. According to Lemma 10, $|\mathcal{S}_j^i| \leq 4$.

$\mathcal{S}_j^i = (a_2 \dots \alpha_{|a|}) \circ (\mathcal{D}_2(\alpha \alpha c \beta) \cap \mathcal{D}_2(\bar{\alpha} \alpha \bar{\alpha} d)) \circ (b_2 \dots b_{|b|})$. According to Lemma 10, $|\mathcal{S}_j^i| \leq 4$.

$\mathcal{S}_j^i = (a_2 \dots \alpha_{|a|}) \circ (\mathcal{D}_1(\alpha \alpha c) \cap \mathcal{D}_3(\bar{\alpha} \alpha \alpha d \bar{\beta})) \circ (b_1 \dots b_{|b|-1})$. The sequence in $\mathcal{D}_1(\alpha \alpha c) \cap \mathcal{D}_3(\bar{\alpha} \alpha \alpha d \bar{\beta})$ only

can begin with α , $|\mathcal{D}_1(\alpha\alpha c) \cap \mathcal{D}_3(\bar{\alpha}\alpha\alpha d\bar{\beta})| = |\mathcal{D}_1(\alpha\alpha c) \cap \mathcal{D}_2(\alpha\alpha d\bar{\beta})|$, then according Lemma 5, $|\mathcal{S}_j^i| \leq 3$.

In sum, $\mathcal{D}_3(x) \cap \mathcal{D}_3(y) \leq 14$.

Case (C): $\omega_2^1 = \bar{\alpha}, \omega_2^2 = \alpha$

$$\begin{aligned} a\omega^1 b &= a\alpha\bar{\alpha}\omega_3^1 \cdots \omega_\ell^1 b = i \cdots \alpha\alpha\bar{\alpha}c\beta \cdots j \\ \bar{a}\omega^2 \bar{b} &= \bar{a}\alpha\alpha\omega_3^2 \cdots \omega_\ell^2 \bar{b} = \bar{i} \cdots \bar{\alpha}\alpha\alpha d\bar{\beta} \cdots \bar{j} \end{aligned}$$

Similarly to Case (A), $|\mathcal{S}_j^i| \leq 4, |\mathcal{S}_j^{\bar{i}}| \leq 4$. $\mathcal{S}_j^i = (a_1 \cdots \alpha_{|a|-1})\alpha \circ (\mathcal{D}_2(\bar{\alpha}c) \cap \mathcal{D}_2(d\bar{\beta})) \circ (b_1 \cdots b_{|b|-1})$. According to Lemma 1 and Theorem 6, $|\mathcal{S}_j^i| \leq 6$, and if $|\mathcal{S}_j^i| \geq 5$, then $\bar{\alpha}c, d\bar{\beta}$ satisfies one of the following two structures

- (i) $\bar{\alpha}c = s_1 t_1 t_2 t_3 s_2$, $d\bar{\beta} = s_1 \bar{t}_1 \bar{t}_2 \bar{t}_3 s_2$, or
- (ii) $\bar{\alpha}c = s_1 t_1 \bar{\gamma} t_2 \bar{\lambda} t_3 s_2$, $d\bar{\beta} = s_1 \bar{t}_1 \bar{\gamma} \bar{t}_2 \bar{\lambda} \bar{t}_3 s_2$,

where t_1, t_2, t_3 are alternating sequences of length at least two, and their lengths are denoted ℓ_1, ℓ_2, ℓ_3 respectively. In particular, in case (ii), t_1 ends with γ , t_2 starts with $\bar{\gamma}$ and ends with $\bar{\lambda}$, t_3 starts with λ .

In case (i), $\mathcal{S}_j^i = (a_2 \cdots \alpha_{|a|}) \circ (\mathcal{D}_1(\alpha\bar{\alpha}c) \cap \mathcal{D}_3(\bar{\alpha}\alpha\alpha d\bar{\beta})) \circ (b_1 \cdots b_{|b|-1})$, where $\mathcal{D}_1(\alpha\bar{\alpha}c) \cap \mathcal{D}_3(\bar{\alpha}\alpha\alpha d\bar{\beta}) = \mathcal{D}_1(\alpha s_1 t_1 t_2 t_3 s_2) \cap \mathcal{D}_3(\bar{\alpha}\alpha s_1 \bar{t}_1 \bar{t}_2 \bar{t}_3 s_2) = \mathcal{D}_1(\alpha s_1 t_1 t_2 t_3) \cap \mathcal{D}_3(\bar{\alpha}\alpha s_1 \bar{t}_1 \bar{t}_2 \bar{t}_3)$. For convenience, we suppose that t_3 ends with θ , and in this and subsequent proofs, we use $s_1[1], s_1[1, t]$ to denote the first bit of the sequence s_1 and subsequence formed by the first bit to the t -th bit respectively, then $\mathcal{D}_1(\alpha\alpha c) \cap \mathcal{D}_3(\bar{\alpha}\alpha\alpha d\bar{\beta})$ consists of the following parts:

- $\mathcal{D}_0(s_1 t_1 t_2 t_3) \cap (\bar{\alpha} \circ \mathcal{D}_2(\alpha\alpha s_1 \bar{t}_1 \bar{t}_2 \bar{t}_3[1, \ell_3 - 2]) \circ \theta)$, Note that we have known s begins with $\bar{\alpha}$, t_3 ends with θ .
- $\alpha \circ (\mathcal{D}_1(s_1 t_1 t_2 t_3[1, \ell_3 - 1]) \cap \mathcal{D}_1(\alpha s_1 \bar{t}_1 \bar{t}_2 \bar{t}_3[1, \ell_3 - 2])) \circ \theta$
- $\alpha \circ (\mathcal{D}_0(s_1 t_1 t_2 t_3[1, \ell_3 - 2]) \cap \mathcal{D}_2(\alpha s_1 \bar{t}_1 \bar{t}_2 \bar{t}_3[1, \ell_3 - 1])) \circ \bar{\theta}$

Firstly, denote $s_1 t_1 t_2 t_3[1, \ell_3 - 1]$ as \tilde{x} , $\alpha s_1 \bar{t}_1 \bar{t}_2 \bar{t}_3[1, \ell_3 - 2]$ as \tilde{y} and their length is ℓ . Since s_1 begins with $\bar{\alpha}$, t_3 is alternating sequence, then $\tilde{x}_1 \neq \tilde{y}_1$ and the last $\ell_3 - 2$ bits of x, y is equal. According to 2, $D_1(\tilde{x}) \cap D_1(\tilde{y}) = 2$ if and only if the first $\ell - \ell_3 + 2$ bits of \tilde{x}, \tilde{y} are alternating sequences beginning with different symbols. And since t_1 is alternating of length at least 2, then exists at least one bit $\tilde{x}_{|s_1|+2}$ equals to $\tilde{y}_{|s_1|+2}$, then $|\mathcal{D}_1(\tilde{x}) \cap \mathcal{D}_1(\tilde{y})| \leq 1$.

Next, we assume that there exists $|\mathcal{D}_0(s_1 t_1 t_2 t_3[1, \ell_3 - 2]) \cap \mathcal{D}_2(\alpha s_1 \bar{t}_1 \bar{t}_2 \bar{t}_3[1, \ell_3 - 1])| = 1$, then in order to get the same prefix $s_1 t_1[1, \ell_1 - 1]$, it is necessary to delete α in the beginning and $\bar{t}_1[1]$, since t_2 is alternating sequence, it is obviously that $t_2[1] = t_2[2]$ and then $t_1[\ell] t_2[1] \neq t_2[1] t_2[2]$, a contradiction. Thus, $(\alpha \circ (\mathcal{D}_0(s_1 t_1 t_2 t_3[1, \ell_3 - 2]) \cap \mathcal{D}_2(\alpha s_1 \bar{t}_1 \bar{t}_2 \bar{t}_3[1, \ell_3 - 1])) \circ \bar{\theta}) = 0$.

In sum, $|\mathcal{S}_j^i| \leq 2$. In case (ii), it can be derived similarly. So with all that, we have $|\mathcal{S}_j^i| + |\mathcal{S}_j^{\bar{i}}| \leq 8$.

$|\mathcal{S}_j^{\bar{i}}| = |\mathcal{D}_2(\alpha\bar{\alpha}c\beta) \cap \mathcal{D}_2(\bar{\alpha}\alpha\alpha d\bar{\beta})|$, according Theorem 3, if $|\mathcal{S}_j^{\bar{i}}| = 6$, then $\alpha\bar{\alpha}c\beta = t_1 t_2 t_3 s_2$, $\bar{\alpha}\alpha\alpha d\bar{\beta} = \bar{t}_1 \bar{t}_2 \bar{t}_3 s_2$, where $t_1 = \alpha\bar{\alpha}$, t_2, t_3 are alternating sequences and t_2 begins with α . We have:

$$\begin{aligned} x &= u i \cdots a_{|a|-1} \alpha \alpha \bar{\alpha} a t_2[2] \cdots t_2[\ell_2] t_3 s_2 b_2 \cdots j v \\ y &= u \bar{i} \cdots \bar{a}_{|a|-1} \bar{\alpha} \alpha \alpha t_2[2] \cdots t_2[\ell_2] \bar{t}_3 s_2 \bar{b}_2 \cdots \bar{j} v \end{aligned}$$

The subsequence obtained by deleting \bar{i}, \bar{j} from y is also obtained by deleting the one of the two last bits of t_2 and last bit of t_3 from x . It implies that $d_L(x, y) = 2$, a contradiction. Then $|\mathcal{S}_j^{\bar{i}}| \leq 5$.

Thus, if $\omega_2^1 = \bar{\alpha}, \omega_2^2 = \alpha$, then $\mathcal{D}_3(x) \cap \mathcal{D}_3(y) = |\mathcal{S}| \leq 4 + 8 + 5 = 17$.

Case (D): $\omega_2^1 = \bar{\alpha}, \omega_2^2 = \bar{\alpha}$

$$\begin{aligned} a\omega^1 b &= a\alpha\bar{\alpha}\omega_3^1 \cdots \omega_\ell^1 b = i \cdots \alpha\alpha\bar{\alpha}c\beta \cdots j \\ \bar{a}\omega^2 \bar{b} &= \bar{a}\alpha\bar{\alpha}\omega_3^2 \cdots \omega_\ell^2 \bar{b} = \bar{i} \cdots \bar{\alpha}\alpha\bar{\alpha}d\bar{\beta} \cdots \bar{j} \end{aligned}$$

In this case, if $\omega_{\ell-1}^1 \neq \omega_{\ell-1}^2$ is equivalent to Case (B) or Case (C); if $\omega_\ell^1 = \omega_\ell^2 = \omega_{\ell-1}^1 = \omega_{\ell-1}^2$ is equivalent to (B). Thus, when $\omega_1^1 = \omega_1^2 = \alpha$, $|\mathcal{S}| > 17$ holds only if $\omega_2^1 = \omega_2^2 = \bar{\alpha}$ and $\omega_\ell^1 = \omega_\ell^2, \omega_{\ell-1}^1 = \omega_{\ell-1}^2, \omega_{\ell-1}^1 = \omega_\ell^1$. There are two possible structures according as follows:

- $x = u\alpha\bar{\alpha}\omega_3^1 \cdots \omega_{\ell-2}^1 \bar{\beta}\beta b v, y = u\bar{\alpha}\alpha\bar{\alpha}\omega_3^2 \cdots \omega_{\ell-2}^2 \bar{\beta}\beta \bar{b} v$
 - $x = u\alpha\bar{\alpha}\omega_3^1 \cdots \omega_{\ell-2}^1 \bar{\beta}\beta b v, y = u\bar{\alpha}\alpha\bar{\alpha}\omega_3^2 \cdots \omega_{\ell-2}^2 \bar{\beta}\beta \bar{b} v$
- Similarly, if $\omega_2^1 = \omega_2^2 = \bar{\alpha}$, we can get symmetrically
- $x = u\alpha\bar{\alpha}\omega_3^1 \cdots \omega_{\ell-2}^1 \bar{\beta}\beta b v, y = u\bar{\alpha}\bar{\alpha}\omega_3^2 \cdots \omega_{\ell-2}^2 \bar{\beta}\beta \bar{b} v$
 - $x = u\alpha\bar{\alpha}\omega_3^1 \cdots \omega_{\ell-2}^1 \bar{\beta}\beta b v, y = u\bar{\alpha}\bar{\alpha}\omega_3^2 \cdots \omega_{\ell-2}^2 \bar{\beta}\beta \bar{b} v$
- The above is equivalent to $x = uas\omega^1 tbv, y = u\bar{a}\bar{s}\omega^2 \bar{t}\bar{b}v$, where

- u, v are the longest common prefix and suffix of x, y ,
- a, b, s, t are maximal alternating sequence of length at least 2.
- $\omega^1 \neq \omega^2$.

Lemma 11. Let x, y be two binary sequences of length n such that $\mathcal{D}_L(x, y) \geq 3$, if $\mathcal{D}_2(x) \cap \mathcal{D}_2(y) = 5$, then x, y must holds one of the following four structures:

- (i) $x = u\alpha\omega b v, y = u\bar{\alpha}\omega \bar{b} v$, where $\omega = u'\omega'v'$, and satisfies
 - (i) u' is a run consisting of α , ω' is alternating sequence starts with $\bar{\alpha}$ and ends with β , v' is a run consisting of $\bar{\beta}$, or (ii) u' is a run consisting of $\bar{\alpha}$, ω is alternating sequence starts with α and ends with $\bar{\beta}$, v' is a run consisting of β .
 - at least one of $|u'| \geq 2, |v'| \geq 2$ holds
 - $|\omega'| \geq 1$, and the equality is allowed to hold only if $|\bar{\alpha}| = \beta$.
- (ii) $x = u\alpha\omega b v, y = u\bar{\alpha}\omega \bar{b} v$, ω is alternating sequence and satisfies $\omega = \alpha\bar{\beta}$ or $\omega = \bar{\alpha}\beta$.
- (iii) $x = u\alpha\omega b v, y = u\bar{\alpha}\omega \bar{b} v$, $\omega = \alpha = \bar{\beta}$ or $\omega = \bar{\alpha} = \beta$.
- (iv) $x = u\alpha\bar{\alpha}\omega\beta b v, y = u\bar{\alpha}\alpha\omega\bar{\beta} \bar{b} v$, where ω is an alternating sequence starts with $\bar{\alpha}$ and ends with $\bar{\beta}$.

In (i)-(iv), u, v is the longest common prefix and suffix of x, y ; a, b is the longest alternating sequence that holds the above structure and a ends with α, b starts with β .

Proof: This lemma can be deduced from Theorem 3. ■

Lemma 8. Let x, y be two binary sequences of length n such that $\mathcal{D}_L(x, y) > 2$ and $x = u\tilde{x}v, y = u\tilde{y}v$, where u, v is the longest common prefix and suffix of x, y . Denote x starts with i , ends with j , $\mathcal{S} = \mathcal{D}_3(\tilde{x}) \cap \mathcal{D}_3(\tilde{y})$. If $\mathcal{D}_3(x) \cap \mathcal{D}_3(y) \geq 18$, then following holds:

- 1) If $|\mathcal{S}_j^i| = 6$, then $|\mathcal{S}_j^{\bar{i}}| = 4$.
- 2) If $|\mathcal{S}_j^i| = 6$, then $|\mathcal{S}_j^{\bar{i}}| = 4$.

Proof: According to Lemma 7, if $\mathcal{D}_3(\mathbf{x}) \cap \mathcal{D}_3(\mathbf{y}) \geq 18$, $\mathbf{x} = uas\omega^1tbv$, $\mathbf{y} = u\bar{a}s\omega^2\bar{t}\bar{b}v$, where

- u, v are the longest common prefix and suffix of \mathbf{x}, \mathbf{y} ,
- a, b, s, t are maximal alternating sequence of length at least 2.
- $\omega^1 \neq \omega^2$.

Suppose a ends with α , b starts with β . Since a, b are alternating sequences, then $\mathcal{S}_j^i = D_2(as\omega^1tb_1b_2 \dots b_{|b|-1}) \cap D_2(\bar{a}_2 \dots \bar{a}_{|b|}s\omega^1\bar{t}\bar{b}) = (a_1a_2 \dots a_{|a|-1}) \circ (D_2(as\omega^1t) \cap D_2(s\omega^2t\bar{\beta})) \circ (b_1b_2 \dots b_{|b|-1}), |\mathcal{S}_j^i| = D_1(a_2 \dots a_{|a|}s\omega^1tb_1 \dots b_{|b|} - 1) \cap D_3(as\omega^2tb) = (a_2 \dots a_{|a|}) \circ (D_1(s\omega^1t) \cap D_3(\bar{\alpha}s\omega^2t\bar{\beta})) \circ (b_1b_2 \dots b_{|b|-1})$.

Let $\tilde{x} = \alpha s\omega^1t\beta$, $\tilde{y} = \bar{\alpha}s\omega^2t\bar{\beta}$, $|\tilde{x}| = |\tilde{y}| = \ell$, then $|\mathcal{S}_j^i| = |D_2(\tilde{x}_1 \dots \tilde{x}_{\ell-1}) \cap D_2(\tilde{y}_2 \dots \tilde{y}_{\ell})|$, $|\mathcal{S}_j^{\bar{i}}| = |D_1(\tilde{x}_2 \dots \tilde{x}_{\ell-1}) \cap D_3(\tilde{y}_1 \dots \tilde{y}_{\ell})|$. Due to $\tilde{x}_2\tilde{x}_3$ is the first two bits of s , $\tilde{x}_{\ell-1}\tilde{x}_{\ell}$ is the last two bits of t , thus $\tilde{x}_2 = \tilde{y}_2 = \bar{x}_3 = \bar{y}_3$, $\tilde{x}_{\ell-1} = \tilde{y}_{\ell-1} = \bar{x}_{\ell} = \bar{y}_{\ell}$.

According to Lemma 3, $|\mathcal{S}_j^i| = 6$ if and only if $\tilde{x}_1 \dots \tilde{x}_{\ell-1}, \tilde{y}_2 \dots \tilde{y}_{\ell}$ satisfies the following structure in Fig.1 (1), where ψ_2, ψ_3, ψ_4 are alternating sequence of length at least 2, and only when ψ_3 is completely reversed or zero-reversed in \tilde{x} , $|\psi_3| \geq 2$, otherwise $|\psi_3| \geq 3$. Denote $|\psi_1|, |\psi_2|, |\psi_3|, |\psi_4|, |\psi_5|$ as $\ell_1, \ell_2, \ell_3, \ell_4, \ell_5$. A categorical discussion of the different values of γ, θ follows.

Case (A): $\gamma = \alpha, \theta = \beta$, refer to Fig.1(2) and $\psi_1 = \alpha, \psi_5$ is empty. (The parts marked in yellow and green are $s\omega^1t$ and $\bar{\alpha}s\omega^2t\bar{\beta}$, respectively.)

$$\begin{aligned} |\mathcal{S}_j^i| &= |D_1(\alpha\bar{\alpha}\tilde{x}_4 \dots \tilde{x}_{\ell-3}\bar{\beta}\bar{\beta}) \cap D_3(\bar{\alpha}\bar{\alpha}\tilde{y}_4 \dots \tilde{y}_{\ell-3}\bar{\beta}\bar{\beta})| \\ &= |\alpha \circ (D_1(\bar{\alpha}\tilde{x}_4 \dots \tilde{x}_{\ell-3}\beta) \cap D_1(\bar{\alpha}\tilde{y}_4 \dots \tilde{y}_{\ell-3}\bar{\beta})) \circ \beta| \\ &+ |\alpha \circ ((\bar{\alpha}\tilde{x}_4 \dots \tilde{x}_{\ell-3}) \cap D_2(\bar{\alpha}\tilde{y}_4 \dots \tilde{y}_{\ell-3}\bar{\beta})) \circ \bar{\beta}| \\ &+ |\bar{\alpha} \circ ((\tilde{x}_4 \dots \tilde{x}_{\ell-3}\bar{\beta}) \cap D_2(\alpha\bar{\alpha}\tilde{y}_4 \dots \tilde{y}_{\ell-3}\bar{\beta})) \circ \bar{\beta}| \end{aligned}$$

(A1) Since ψ_2, ψ_3 are alternating sequences, $|\alpha \circ (D_1(\bar{\alpha}\tilde{x}_4 \dots \tilde{x}_{\ell-3}\beta) \cap D_1(\bar{\alpha}\tilde{y}_4 \dots \tilde{y}_{\ell-3}\bar{\beta})) \circ \beta| = |D_1(\tilde{x}_{\ell_1+\ell_2+1} \dots \tilde{x}_{\ell_1+\ell_2+\ell_3+1}) \cap D_1(\tilde{y}_{\ell_1+\ell_2+1} \dots \tilde{y}_{\ell_1+\ell_2+\ell_3+1})|$, according Theorem 3, we have:

- (i) If $\tilde{x}_{\ell_1+\ell_2} = \tilde{x}_{\ell_1+\ell_2+1}$, $\tilde{x}_{\ell_1+\ell_2+\ell_3} = \tilde{x}_{\ell_1+\ell_2+\ell_3+1}$, then $|\psi_3| \geq 2$
- (ii) If $\tilde{x}_{\ell_1+\ell_2} = \tilde{x}_{\ell_1+\ell_2+1}$, $\tilde{x}_{\ell_1+\ell_2+\ell_3} = \bar{\tilde{x}}_{\ell_1+\ell_2+\ell_3+1}$, then $|\psi_3| \geq 3$,
- (iii) If $\tilde{x}_{\ell_1+\ell_2} = \bar{\tilde{x}}_{\ell_1+\ell_2+1}$, $\tilde{x}_{\ell_1+\ell_2+\ell_3} = \tilde{x}_{\ell_1+\ell_2+\ell_3+1}$, then $|\psi_3| \geq 3$
- (iv) If $\tilde{x}_{\ell_1+\ell_2} = \bar{\tilde{x}}_{\ell_1+\ell_2+1}$, $\tilde{x}_{\ell_1+\ell_2+\ell_3} = \bar{\tilde{x}}_{\ell_1+\ell_2+\ell_3+1}$, then $|\psi_3| \geq 2$

In case (i), $\tilde{x}_{\ell_1+\ell_2+1} = \bar{y}_{\ell_1+\ell_2+1}$, $\tilde{x}_{\ell_1+\ell_2+\ell_3+1} = \bar{y}_{\ell_1+\ell_2+\ell_3+1}$, $(\tilde{x}_{\ell_1+\ell_2+1} \dots \tilde{x}_{\ell_1+\ell_2+\ell_3+1}), (\bar{y}_{\ell_1+\ell_2+1} \dots \bar{y}_{\ell_1+\ell_2+\ell_3+1})$ are alternating sequences of length $|\psi_3| \geq 2$ and begin with different symbol, then $|\mathcal{D}_1(\tilde{x}_{\ell_1+\ell_2+1} \dots \tilde{x}_{\ell_1+\ell_2+\ell_3+1}) \cap \mathcal{D}_1(\bar{y}_{\ell_1+\ell_2+1} \dots \bar{y}_{\ell_1+\ell_2+\ell_3+1})| = |\mathcal{D}_1(\tilde{x}_{\ell_1+\ell_2+1} \dots \tilde{x}_{\ell_1+\ell_2+\ell_3}) \cap \mathcal{D}_1(\bar{y}_{\ell_1+\ell_2+1} \dots \bar{y}_{\ell_1+\ell_2+\ell_3})| = 2$.

In case (ii), $\tilde{x}_{\ell_1+\ell_2+1} = \bar{y}_{\ell_1+\ell_2+1}$, $\tilde{x}_{\ell_1+\ell_2+\ell_3+1} = \bar{y}_{\ell_1+\ell_2+\ell_3+1}$, $(\tilde{x}_{\ell_1+\ell_2+1} \dots \tilde{x}_{\ell_1+\ell_2+\ell_3+1}), (\bar{y}_{\ell_1+\ell_2+1} \dots \bar{y}_{\ell_1+\ell_2+\ell_3+1})$

are alternating sequences of length $|\psi_3| - 1 \geq 2$ and begin with different symbol, then $|\mathcal{D}_1(\tilde{x}_{\ell_1+\ell_2+1} \dots \tilde{x}_{\ell_1+\ell_2+\ell_3+1}) \cap \mathcal{D}_1(\bar{y}_{\ell_1+\ell_2+1} \dots \bar{y}_{\ell_1+\ell_2+\ell_3+1})| = 2$.

Due to the symmetry, the same result can be obtained in in case (iii), case (iv). Thus, $|\alpha \circ (D_1(\bar{\alpha}\tilde{x}_4 \dots \tilde{x}_{\ell-3}\beta) \cap D_1(\bar{\alpha}\tilde{y}_4 \dots \tilde{y}_{\ell-3}\bar{\beta})) \circ \beta| = 2$.

(A2) Since ψ_4 is alternating sequence, it implies that $\tilde{x}_{\ell_1+\ell_2+\ell_3+1} \dots \tilde{x}_{\ell-3}\bar{\beta} = \bar{y}_{\ell_1+\ell_2+\ell_3+2} \dots \bar{y}_{\ell-3}\bar{\beta}\bar{\beta}$, and $\psi_3 = \tilde{x}_{\ell_1+\ell_2+1} \dots \tilde{x}_{\ell_1+\ell_2+\ell_3} = \bar{y}_{\ell_1+\ell_2+1} \dots \bar{y}_{\ell_1+\ell_2+\ell_3}$, then $|\alpha \circ ((\bar{\alpha}\tilde{x}_4 \dots \tilde{x}_{\ell-3}) \cap D_2(\bar{\alpha}\tilde{y}'_4 \dots \tilde{y}'_{\ell-3}\bar{\beta}\bar{\beta})) \circ \beta| = |\tilde{x}_{\ell_1+\ell_2+1} \dots \tilde{x}_{\ell-3}) \cap D_2(\bar{y}'_{\ell_1+\ell_2+1} \dots \bar{y}'_{\ell-3}\bar{\beta}\bar{\beta})| = 1$.

(A3) Since ψ_2, ψ_3 is alternating sequence, it implies that $\tilde{x}_4 \dots \tilde{x}_{\ell_1+\ell_2} = \alpha\bar{\alpha}\tilde{y}_4 \dots \tilde{y}_{\ell_1+\ell_2-3}$, $\psi_3[3, \ell_3] = \tilde{x}_{\ell_1+\ell_2+3} \dots \tilde{x}_{\ell_1+\ell_2+\ell_3} = \bar{y}_{\ell_1+\ell_2+1} \dots \bar{y}_{\ell_1+\ell_2+\ell_3-2}$, and $\tilde{x}_{\ell_1+\ell_2+1} \tilde{x}_{\ell_1+\ell_2+2} \in \mathcal{D}_1(\bar{y}_{\ell_1+\ell_2+1} \bar{y}_{\ell_1+\ell_2+2} \bar{y}_{\ell_1+\ell_2+3})$, $\tilde{x}_{\ell_1+\ell_2+\ell_3+1} \in \mathcal{D}_1(\bar{y}_{\ell_1+\ell_2+\ell_3+1} \bar{y}_{\ell_1+\ell_2+\ell_3+2} \bar{y}_{\ell_1+\ell_2+\ell_3+3})$, thus $|\bar{\alpha} \circ (\tilde{x}_4 \dots \tilde{x}_{\ell-3}\bar{\beta}) \cap D_2(\alpha\bar{\alpha}\tilde{y}_4 \dots \tilde{y}_{\ell-3}\bar{\beta})) \circ \bar{\beta}| = |(\tilde{x}_4 \dots \tilde{x}_{\ell_1+\ell_2+\ell_3}) \cap D_2(\alpha\bar{\alpha}\tilde{y}_4 \dots \tilde{y}_{\ell_1+\ell_2+\ell_3})| = 1$.

The analysis of the rest three cases is similar, and here we only give the expression.

Case (B): $\gamma = \alpha, \theta = \bar{\beta}$, there exists a common suffix $\bar{\beta}$.

$$\begin{aligned} |\mathcal{S}_j^i| &= |D_1(\alpha\bar{\alpha}\tilde{x}_4 \dots \tilde{x}_{\ell-3}\bar{\beta}\bar{\beta}) \cap D_3(\bar{\alpha}\bar{\alpha}\tilde{y}_4 \dots \tilde{y}_{\ell-3}\bar{\beta}\bar{\beta})| \\ &= |D_1(\alpha\bar{\alpha}\tilde{x}_4 \dots \tilde{x}_{\ell-3}\bar{\beta}) \cap D_3(\bar{\alpha}\bar{\alpha}\tilde{y}_4 \dots \tilde{y}_{\ell-3}\bar{\beta})| \\ &= |\alpha \circ (D_1(\bar{\alpha}\tilde{x}_4 \dots \tilde{x}_{\ell-3}\bar{\beta}) \cap D_1(\bar{\alpha}\tilde{y}_4 \dots \tilde{y}_{\ell-3}\bar{\beta})) \circ \bar{\beta}| \\ &+ |\alpha \circ ((\bar{\alpha}\tilde{x}_4 \dots \tilde{x}_{\ell-4}) \cap D_2(\bar{\alpha}\tilde{y}_4 \dots \tilde{y}_{\ell-3}\bar{\beta})) \circ \bar{\beta}| \\ &+ |\bar{\alpha} \circ ((\tilde{x}_4 \dots \tilde{x}_{\ell-3}\bar{\beta}) \cap D_2(\alpha\bar{\alpha}\tilde{y}'_4 \dots \tilde{y}_{\ell-3}\bar{\beta})) \circ \bar{\beta}| \\ &= 2 + 1 + 1 = 4. \end{aligned}$$

Case (C): $\gamma = \bar{\alpha}, \theta = \beta$, there exists a common prefix $\bar{\alpha}$.

$$\begin{aligned} |\mathcal{S}_j^i| &= |D_1(\bar{\alpha}\alpha\tilde{x}_4 \dots \tilde{x}_{\ell-3}\bar{\beta}\bar{\beta}) \cap D_3(\bar{\alpha}\bar{\alpha}\tilde{y}_4 \dots \tilde{y}_{\ell-3}\bar{\beta}\bar{\beta})| \\ &= |D_1(\alpha\tilde{x}_4 \dots \tilde{x}_{\ell-3}\bar{\beta}\bar{\beta}) \cap D_3(\bar{\alpha}\alpha\tilde{y}_4 \dots \tilde{y}_{\ell-3}\bar{\beta}\bar{\beta})| \\ &= |\alpha \circ (D_1(\tilde{x}_4 \dots \tilde{x}_{\ell-3}\bar{\beta}) \cap D_1(\tilde{y}_4 \dots \tilde{y}_{\ell-3}\bar{\beta})) \circ \bar{\beta}| \\ &+ |\alpha \circ ((\tilde{x}_4 \dots \tilde{x}_{\ell-3}) \cap D_2(\bar{\alpha}\tilde{y}_4 \dots \tilde{y}_{\ell-3}\bar{\beta})) \circ \bar{\beta}| \\ &+ |\bar{\alpha} \circ ((\tilde{x}_5 \dots \tilde{x}_{\ell-3}\bar{\beta}) \cap D_2(\alpha\tilde{y}'_4 \dots \tilde{y}_{\ell-3}\bar{\beta})) \circ \bar{\beta}| \\ &= 2 + 1 + 1 = 4 \end{aligned}$$

Case (D): $\gamma = \bar{\alpha}, \theta = \bar{\beta}$, there exists a common prefix $\bar{\alpha}$ and a common suffix $\bar{\beta}$.

$$\begin{aligned} |\mathcal{S}_j^i| &= |D_1(\bar{\alpha}\alpha\tilde{x}_4 \dots \tilde{x}_{\ell-3}\bar{\beta}\bar{\beta}) \cap D_3(\bar{\alpha}\bar{\alpha}\tilde{y}_4 \dots \tilde{y}_{\ell-3}\bar{\beta}\bar{\beta})| \\ &= |D_1(\alpha\tilde{x}_4 \dots \tilde{x}_{\ell-3}\bar{\beta}) \cap D_3(\bar{\alpha}\alpha\tilde{y}_4 \dots \tilde{y}_{\ell-3}\bar{\beta})| \\ &= |\alpha \circ (D_1(\tilde{x}_4 \dots \tilde{x}_{\ell-3}) \cap D_1(\tilde{y}_4 \dots \tilde{y}_{\ell-3})) \circ \bar{\beta}| \\ &+ |\alpha \circ ((\tilde{x}_4 \dots \tilde{x}_{\ell-4}) \cap D_2(\bar{\alpha}\tilde{y}_4 \dots \tilde{y}_{\ell-3}\bar{\beta})) \circ \bar{\beta}| \\ &+ |\bar{\alpha} \circ ((\tilde{x}_5 \dots \tilde{x}_{\ell-3}) \cap D_2(\alpha\tilde{y}'_4 \dots \tilde{y}_{\ell-3}\bar{\beta})) \circ \bar{\beta}| \\ &= 2 + 1 + 1 = 4 \end{aligned}$$

Theorem 3. Let \mathbf{x}, \mathbf{y} be two binary sequences of length n such that $d_L(\mathbf{x}, \mathbf{y}) \geq 3$, $\mathcal{D}_3(\mathbf{x}) \cap \mathcal{D}_3(\mathbf{y}) = 19$ if and only they are of the form $\mathbf{x} = uas\omega^1tbv$, $\mathbf{y} = u\bar{a}s\omega^2\bar{t}\bar{b}v$, such that

- u, v are the longest common prefix and suffix of \mathbf{x}, \mathbf{y} .
- a, b are maximal alternating sequences of length ≥ 2 .

$$\begin{aligned}
(1) \quad \tilde{x} &= \overbrace{\alpha \gamma \bar{\gamma} \tilde{x}_4 \cdots \tilde{x}_{\ell_1+\ell_2-1} \tilde{x}_{\ell_1+\ell_2}}^{\psi_2} \overbrace{\tilde{x}_{\ell_1+\ell_2+1} \tilde{x}_{\ell_1+\ell_2+2} \cdots \tilde{x}_{\ell_1+\ell_2+\ell_3-1} \tilde{x}_{\ell_1+\ell_2+\ell_3}}^{\psi_3} \overbrace{\tilde{x}_{\ell_1+\ell_2+\ell_3+1} \tilde{x}_{\ell_1+\ell_2+\ell_3+2} \cdots \tilde{x}_{\ell-3} \bar{\theta} \bar{\theta} \beta}^{\psi_4} \overbrace{\beta}^{\psi_5} \\
\tilde{y} &= \overbrace{\alpha' \gamma \bar{\gamma} y_4 \cdots \tilde{y}_{1+\ell_1+\ell_2-1} \tilde{y}_{1+\ell_1+\ell_2}}^{\psi_2} \overbrace{\tilde{y}_{1+\ell_1+\ell_2+1} \tilde{y}_{1+\ell_1+\ell_2+2} \cdots \tilde{y}_{1+\ell_1+\ell_2+\ell_3-1} \tilde{y}_{1+\ell_1+\ell_2+\ell_3}}^{\psi_3} \overbrace{\tilde{y}_{1+\ell_1+\ell_2+\ell_3+1} \tilde{y}_{1+\ell_1+\ell_2+\ell_3+2} \cdots \tilde{y}_{\ell-3} \bar{\theta} \bar{\theta} \bar{\beta}}^{\psi_4} \overbrace{\bar{\beta}}^{\psi_5} \\
(2) \quad \tilde{x} &= \overbrace{\alpha \bar{\alpha} \tilde{x}_4 \cdots \tilde{x}_{\ell_1+\ell_2-1} \tilde{x}_{\ell_1+\ell_2}}^{\psi_2} \overbrace{\tilde{x}_{\ell_1+\ell_2+1} \tilde{x}_{\ell_1+\ell_2+2} \cdots \tilde{x}_{\ell_1+\ell_2+\ell_3-1} \tilde{x}_{\ell_1+\ell_2+\ell_3}}^{\psi_3} \overbrace{\tilde{x}_{\ell_1+\ell_2+\ell_3+1} \tilde{x}_{\ell_1+\ell_2+\ell_3+2} \cdots \tilde{x}_{\ell-3} \bar{\beta} \bar{\beta} \beta}^{\psi_4} \overbrace{\beta}^{\psi_5} \\
\tilde{y} &= \overbrace{\bar{\alpha} \bar{\alpha} \bar{\alpha} y_4 \cdots \tilde{y}_{1+\ell_1+\ell_2-1} \tilde{y}_{1+\ell_1+\ell_2}}^{\psi_2} \overbrace{\tilde{y}_{1+\ell_1+\ell_2+1} \tilde{y}_{1+\ell_1+\ell_2+2} \cdots \tilde{y}_{1+\ell_1+\ell_2+\ell_3-1} \tilde{y}_{1+\ell_1+\ell_2+\ell_3}}^{\psi_3} \overbrace{\tilde{y}_{1+\ell_1+\ell_2+\ell_3+1} \tilde{y}_{1+\ell_1+\ell_2+\ell_3+2} \cdots \tilde{y}_{\ell-3} \bar{\beta} \bar{\beta} \bar{\beta}}^{\psi_4} \overbrace{\bar{\beta}}^{\psi_5}
\end{aligned}$$

Fig. 1. Illustrations of \tilde{x} and \tilde{y} when $|D_2(\alpha s \omega^1 t) \cap D_2(s \omega^2 t \beta)| = 6$. (1) General case. (2) Case (A) ($\gamma = \alpha, \theta = \bar{\beta}$)

- s, t are maximal alternating sequences. If s is completely reversed or zero-reversed between a and w , then $|s| \geq 2$, otherwise $|s| \geq 3$. The same applies to t .
- $|\omega| = 3$ and ω is an alternating sequence which is neither zero-reversed nor completely reversed.

Proof: According to Lemma 7, if $\mathcal{D}_3(x) \cap \mathcal{D}_3(y) = 19$, then $x = uas\omega^1 tbv, y = \bar{u}as\omega^2 t\bar{b}\bar{v}$, where

- u, v are the longest common prefix and suffix of x, y ,
- a, b, s, t are maximal alternating sequence of length at least 2.
- $\omega^1 \neq \omega^2$.

Suppose a starts with i , ends with α , b starts with β , ends with j . Let $\tilde{x} = \alpha s \omega^1 t \beta, \tilde{y} = \bar{\alpha} s \omega^2 t \bar{\beta}, |\tilde{x}| = |\tilde{y}| = \ell$. According to Lemma 8, $\mathcal{D}_3(x) \cap \mathcal{D}_3(y) = 19$ holds if and only if:

- Type (A): $|\mathcal{S}_j^i| = 6, |\mathcal{S}_j^i| = 5, |\mathcal{S}_j^i| = 4, |\mathcal{S}_j^i| = 4$, or
- Type (B): $|\mathcal{S}_j^i| = 5, |\mathcal{S}_j^i| = 6, |\mathcal{S}_j^i| = 4, |\mathcal{S}_j^i| = 4$.

Take Type (A) for example. Consider \mathcal{S}_j^i , where the first bit i of $as\omega^1 tb$ and the last bit j of $\bar{a}s\omega^2 t\bar{b}$ must be deleted. Since a, b are alternating sequences, then we reduce to $|\mathcal{S}_j^i| = D_2(a_2 \cdots a_{|a|} s \omega^1 t b) \cap D_2(\bar{a} s \omega^2 t \bar{b}_1 \cdots \bar{b}_{|b|-1})|$

$$\begin{aligned}
&= |D_2(s \omega^1 t \beta) \cap D_2(\bar{\alpha} s \omega^2 t)| \\
&= |D_2(\tilde{x}_2 \cdots \tilde{x}_\ell) \cap D_2(\tilde{y}_1 \cdots \tilde{y}_{\ell-1})| = 6.
\end{aligned}$$

$|\mathcal{S}_j^i| = 6$ holds if and only if the structures in Fig. 2 (1) is satisfied. In Fig.2 (1), ϕ_2, ϕ_3, ϕ_4 are alternating sequence of length at least 2, and only when ϕ_3 is completely reversed or zero-reversed in $\tilde{x}, |\phi_3| \geq 2$, otherwise $|\phi_3| \geq 3$. Denote $|\phi_1|, |\phi_2|, |\phi_3|, |\phi_4|, |\phi_5|$ as $\ell_1, \ell_2, \ell_3, \ell_4, \ell_5$.

According to Lemma 11, if $|\mathcal{S}_j^i| = 5$ holds, there are four possible structures. Next we discuss the likelihood that these four structures hold in the case where $|\mathcal{S}_j^i| = 6$ holds. As before, the discussion is based on the example of $\gamma = \alpha$.

Case (A): $\tilde{x}_1 \cdots \tilde{x}_{\ell-1} = \psi_1 \psi_2 \psi_3 \psi_4 \psi_5, \tilde{y}_2 \cdots \tilde{y}_\ell = \psi_1 \bar{\psi}_2 \bar{\psi}_3 \bar{\psi}_4 \bar{\psi}_5$, where $\psi_3 = u' \psi'_3 v'$ and at least one of u', v' is a run of length at least 2. Since ϕ_2, ϕ_3, ϕ_4 are alternating sequences, it is clear that we cannot find a k such that $\tilde{x}_k = \tilde{x}_{k+1} = \tilde{x}_{k+1} = \tilde{y}_{k+2}$, a contradiction.

Case (B) $\tilde{x}_1 \cdots \tilde{x}_{\ell-1} = \psi_1 \psi_2 \psi_3 \psi_4 \psi_5, \tilde{y}_2 \cdots \tilde{y}_\ell = \psi_1 \bar{\psi}_2 \bar{\psi}_3 \bar{\psi}_4 \bar{\psi}_5$, where $|\psi_3| = 2$, and ψ_3 is neither zero-reversed nor completely reversed in \tilde{x} .

(B1) If $\tilde{x}_{1+\ell_1+\ell_2-1} \tilde{x}_{1+\ell_1+\ell_2} = \tilde{y}_{\ell_1+\ell_2+1} \tilde{y}_{\ell_1+\ell_2+2}$, then in order to satisfy the condition of ψ_3 , then there must be $\ell_3 = 2$, and $\tilde{x}_{1+\ell_1+\ell_2} = \tilde{x}_{1+\ell_1+\ell_2+1}$. Obviously, since ϕ_2, ϕ_3 are alternating sequence, $\tilde{x}_{1+\ell_1+\ell_2-1} \tilde{x}_{1+\ell_1+\ell_2} = \tilde{y}_{\ell_1+\ell_2+1} \tilde{y}_{\ell_1+\ell_2+2}$ and $\tilde{x}_{1+\ell_1+\ell_2} = \tilde{x}_{1+\ell_1+\ell_2+1}$ cannot hold at the same time, a contradiction.

(B2) If $\tilde{x}_{1+\ell_1+\ell_2-1} \tilde{x}_{1+\ell_1+\ell_2} = \tilde{y}_{\ell_1+\ell_2+1} \tilde{y}_{\ell_1+\ell_2+2}$, There are two possible structures that make the condition of ψ_3 satisfied:

- (i) $\tilde{x}_{1+\ell_1+\ell_2+3-1} \tilde{x}_{1+\ell_1+\ell_2+3} = \tilde{y}_{\ell_1+\ell_2+\ell_3+1} \tilde{y}_{\ell_1+\ell_2+\ell_3+2}$ and $\ell_3 = 2$,
- (ii) $\tilde{x}_{1+\ell_1+\ell_2+3-1} \tilde{x}_{1+\ell_1+\ell_2+3} = \tilde{y}_{\ell_1+\ell_2+\ell_3+1} \tilde{y}_{\ell_1+\ell_2+\ell_3+2}$ and $\ell_3 = 4$.

In case (i), ψ_3 corresponds to $\tilde{x}_{1+\ell_1+\ell_2+1} \tilde{x}_{1+\ell_1+\ell_2+3}$ and we have $\tilde{x}_{1+\ell_1+\ell_2} = \tilde{y}_{\ell_1+\ell_2+2} = \tilde{x}_{1+\ell_1+\ell_2+1}$, $\tilde{x}_{1+\ell_1+\ell_2+3} = \tilde{y}_{\ell_1+\ell_2+\ell_3+2} = \tilde{x}_{1+\ell_1+\ell_2+3+1}$, i.e. ω is completely reversed in \tilde{x} , a contradiction.

In case (ii), ψ_3 corresponds to $\tilde{x}_{1+\ell_1+\ell_2+1} \tilde{x}_{1+\ell_1+\ell_2+2}$. We have $\tilde{x}_{1+\ell_1+\ell_2} = \tilde{y}_{\ell_1+\ell_2+2} = \tilde{x}_{1+\ell_1+\ell_2+1}$ and $\tilde{x}_{1+\ell_1+\ell_2+2} = \tilde{x}_{1+\ell_1+\ell_2+3}$. Thus, under the condition of $|\mathcal{S}_j^i| = 6$, Case (B2)(ii) can hold simultaneously.

Case (C) $\tilde{x}_1 \cdots \tilde{x}_{\ell-1} = \psi_1 \psi_2 \psi_3 \psi_4 \psi_5, \tilde{y}_2 \cdots \tilde{y}_\ell = \psi_1 \bar{\psi}_2 \bar{\psi}_3 \bar{\psi}_4 \bar{\psi}_5$, where $|\psi_3| = 1$ and ψ_3 is neither zero-reversed nor completely reversed in \tilde{x} .

(C1) If $\tilde{x}_{1+\ell_1+\ell_2-1} \tilde{x}_{1+\ell_1+\ell_2} = \tilde{y}_{\ell_1+\ell_2+1} \tilde{y}_{\ell_1+\ell_2+2}$, it is clear that $|\psi_3| > 1$.

(C2) If $\tilde{x}_{1+\ell_1+\ell_2-1} \tilde{x}_{1+\ell_1+\ell_2} = \tilde{y}_{\ell_1+\ell_2+1} \tilde{y}_{\ell_1+\ell_2+2}$, there are only one possible structures that make the condition of ψ_3 satisfied: $\ell_3 = 3$ and $\tilde{x}_{1+\ell_1+\ell_2+3-1} \tilde{x}_{1+\ell_1+\ell_2+3} = \tilde{y}_{\ell_1+\ell_2+\ell_3+1} \tilde{y}_{\ell_1+\ell_2+\ell_3+2}$, then we have

- $\alpha \bar{\alpha} \tilde{x}_4 \cdots \tilde{x}_{1+\ell_1+\ell_2-1} = \alpha y_4 \cdots \tilde{y}_{\ell_1+\ell_2+2}$;
- $\tilde{x}_{1+\ell_1+\ell_2} = \tilde{y}_{\ell_1+\ell_2+1} = \tilde{x}_{1+\ell_1+\ell_2+1} = \tilde{y}_{\ell_1+\ell_2+3} = \tilde{y}_{\ell_1+\ell_2+3}$;
- $\tilde{x}_{1+\ell_1+\ell_2+1} = \tilde{x}_{1+\ell_1+\ell_2+3} = \tilde{y}_{\ell_1+\ell_2+4} = \tilde{y}_{\ell_1+\ell_2+3+1}$
- $\tilde{x}_{1+\ell_1+\ell_2+2} \cdots \tilde{x}_{\ell-3} = \tilde{y}_{\ell_1+\ell_2+\ell_3+2} \cdots \tilde{y}_{\ell-3} \bar{\theta} \bar{\theta}$
- either $\theta = \bar{\beta}$ or $\theta = \beta$ holds

Thus $d_L(x, y) = 2$, a contradiction.

Case (D): $\tilde{x}_1 \cdots \tilde{x}_{\ell-1} = \psi_1 \psi_2 \bar{\psi}_3 \bar{\psi}_4 \bar{\psi}_5, \tilde{y}_2 \cdots \tilde{y}_\ell = \psi_1 \bar{\psi}_2 \bar{\psi}_3 \bar{\psi}_4 \bar{\psi}_5$

(D1) If $\tilde{x}_{1+\ell_1+\ell_2-1} \tilde{x}_{1+\ell_1+\ell_2} = \tilde{y}_{\ell_1+\ell_2+1} \tilde{y}_{\ell_1+\ell_2+2}$, then $\tilde{x}_{1+\ell_1+\ell_2-1} \tilde{x}_{1+\ell_1+\ell_2}$ corresponds to $\bar{\mu} \psi_3[1]$ and

$$\begin{aligned}
(1) \quad \tilde{x} &= \alpha \underbrace{\tilde{\gamma} \tilde{\gamma} \tilde{x}_4 \cdots \tilde{x}_{1+\ell_1+\ell_2-1}}_{\tilde{\phi}_2} \underbrace{\tilde{x}_{1+\ell_1+\ell_2} \tilde{x}_{1+\ell_1+\ell_2+1} \tilde{x}_{1+\ell_1+\ell_2+2} \cdots \tilde{x}_{1+\ell_1+\ell_2+\ell_3-1}}_{\tilde{\phi}_3} \underbrace{\tilde{x}_{1+\ell_1+\ell_2+\ell_3} \tilde{x}_{1+\ell_1+\ell_2+\ell_3+1} \cdots \tilde{x}_{\ell-3}}_{\tilde{\phi}_4} \underbrace{\tilde{\theta} \tilde{\theta} \tilde{\beta}}_{\tilde{\phi}_5} \\
\tilde{y} &= \bar{\alpha} \underbrace{\bar{\gamma} \bar{\gamma} \bar{y}_4 \cdots \bar{y}_{\ell_1+\ell_2-1}}_{\bar{\phi}_2} \underbrace{\bar{y}_{\ell_1+\ell_2} \bar{y}_{\ell_1+\ell_2+1} \bar{y}_{\ell_1+\ell_2+2} \cdots \bar{y}_{\ell_1+\ell_2+\ell_3-1}}_{\bar{\phi}_3} \underbrace{\bar{y}_{\ell_1+\ell_2+\ell_3} \bar{y}_{\ell_1+\ell_2+\ell_3+1} \cdots \bar{y}_{\ell-3}}_{\bar{\phi}_4} \underbrace{\bar{\theta} \bar{\theta} \bar{\beta}}_{\bar{\phi}_5} \\
(2) \quad \tilde{x}' &= \alpha \underbrace{\tilde{\gamma} \tilde{\gamma} \tilde{x}_4 \cdots \tilde{x}_{1+\ell_1+\ell_2-1}}_{\tilde{\phi}_2} \underbrace{\tilde{x}_{1+\ell_1+\ell_2} \tilde{x}_{1+\ell_1+\ell_2+1} \tilde{x}_{1+\ell_1+\ell_2+2} \cdots \tilde{x}_{1+\ell_1+\ell_2+\ell_3-1}}_{\tilde{\phi}_3} \underbrace{\tilde{x}_{1+\ell_1+\ell_2+\ell_3} \tilde{x}_{1+\ell_1+\ell_2+\ell_3+1} \cdots \tilde{x}_{\ell-3}}_{\tilde{\phi}_4} \underbrace{\tilde{\theta} \tilde{\theta} \tilde{\beta}}_{\tilde{\phi}_5} \\
\tilde{y}' &= \bar{\alpha} \underbrace{\bar{\gamma} \bar{\gamma} \bar{y}_4 \cdots \bar{y}_{\ell_1+\ell_2-1}}_{\bar{\phi}_2} \underbrace{\bar{y}_{\ell_1+\ell_2} \bar{y}_{\ell_1+\ell_2+1} \bar{y}_{\ell_1+\ell_2+2} \cdots \bar{y}_{\ell_1+\ell_2+\ell_3-1}}_{\bar{\phi}_3} \underbrace{\bar{y}_{\ell_1+\ell_2+\ell_3} \bar{y}_{\ell_1+\ell_2+\ell_3+1} \cdots \bar{y}_{\ell-3}}_{\bar{\phi}_4} \underbrace{\bar{\theta} \bar{\theta} \bar{\beta}}_{\bar{\phi}_5} \\
(3) \quad \tilde{x}' &= \alpha \underbrace{\bar{\alpha} \bar{\alpha} \bar{x}_4 \cdots \bar{x}_{1+\ell_1+\ell_2-1}}_{\bar{\phi}_2} \underbrace{\bar{x}_{1+\ell_1+\ell_2} \bar{x}_{1+\ell_1+\ell_2+1} \bar{x}_{1+\ell_1+\ell_2+2} \cdots \bar{x}_{1+\ell_1+\ell_2+\ell_3-1}}_{\bar{\phi}_3} \underbrace{\bar{x}_{1+\ell_1+\ell_2+\ell_3} \bar{x}_{1+\ell_1+\ell_2+\ell_3+1} \cdots \bar{x}_{\ell-3}}_{\bar{\phi}_4} \underbrace{\bar{\theta} \bar{\theta} \bar{\beta}}_{\bar{\phi}_5} \\
\tilde{y}' &= \bar{\alpha} \underbrace{\bar{\alpha} \bar{\alpha} \bar{y}_4 \cdots \bar{y}_{\ell_1+\ell_2-1}}_{\bar{\phi}_2} \underbrace{\bar{y}_{\ell_1+\ell_2} \bar{y}_{\ell_1+\ell_2+1} \bar{y}_{\ell_1+\ell_2+2} \cdots \bar{y}_{\ell_1+\ell_2+\ell_3-1}}_{\bar{\phi}_3} \underbrace{\bar{y}_{\ell_1+\ell_2+\ell_3} \bar{y}_{\ell_1+\ell_2+\ell_3+1} \cdots \bar{y}_{\ell-3}}_{\bar{\phi}_4} \underbrace{\bar{\theta} \bar{\theta} \bar{\beta}}_{\bar{\phi}_5} \\
(4) \quad \tilde{x}' &= \alpha \underbrace{\bar{\alpha} \bar{\alpha} \bar{x}_4 \cdots \bar{x}_{1+\ell_1+\ell_2-1}}_{\bar{\phi}_2} \underbrace{\bar{x}_{1+\ell_1+\ell_2} \bar{x}_{1+\ell_1+\ell_2+1} \bar{x}_{1+\ell_1+\ell_2+2} \cdots \bar{x}_{1+\ell_1+\ell_2+\ell_3-1}}_{\bar{\phi}_3} \underbrace{\bar{x}_{1+\ell_1+\ell_2+\ell_3} \bar{x}_{1+\ell_1+\ell_2+\ell_3+1} \cdots \bar{x}_{\ell-3}}_{\bar{\phi}_4} \underbrace{\bar{\theta} \bar{\theta} \bar{\beta}}_{\bar{\phi}_5} \\
\tilde{y}' &= \bar{\alpha} \underbrace{\bar{\alpha} \bar{\alpha} \bar{y}_4 \cdots \bar{y}_{\ell_1+\ell_2-1}}_{\bar{\phi}_2} \underbrace{\bar{y}_{\ell_1+\ell_2} \bar{y}_{\ell_1+\ell_2+1} \bar{y}_{\ell_1+\ell_2+2} \cdots \bar{y}_{\ell_1+\ell_2+\ell_3-1}}_{\bar{\phi}_3} \underbrace{\bar{y}_{\ell_1+\ell_2+\ell_3} \bar{y}_{\ell_1+\ell_2+\ell_3+1} \cdots \bar{y}_{\ell-3}}_{\bar{\phi}_4} \underbrace{\bar{\theta} \bar{\theta} \bar{\beta}}_{\bar{\phi}_5}
\end{aligned}$$

Fig. 2. Illustrations of \tilde{x} and \tilde{y} when $|\mathcal{S}_j^i| = 6, |\mathcal{S}_j^i| = 5$. In the figure, complementary alternating segments and identical alternating segments are represented in red and blue, respectively. In (1), the three colored segments corresponds to $s, \omega^1/\omega^2, t$, correspondingly. The black entries in the middle are pending to be determined to belong to the former or the latter segment, and are analyzed as in the main texts. And $|\mathcal{S}_j^i| = 6$ if and only if $\tilde{x}_2 \cdots \tilde{x}_\ell = \phi_1 \phi_2 \phi_3 \phi_4 \phi_5, \tilde{y}_1 \cdots \tilde{y}_{\ell-1} = \phi_1 \bar{\phi}_2 \phi_3 \bar{\phi}_4 \phi_5$ as (1). Since $\phi_2, \phi_3, \phi_4, s, t$ are alternating sequences, we can obtain (2) and if $\gamma = \alpha, |\psi_1| = 0$, else $|\psi_1| = 1$. (3)(4) are the structures for the cases where $|\phi| = 4$ and γ takes different values, respectively.

$\tilde{y}_{\ell_1+\ell_2+1} \tilde{y}_{\ell_1+\ell_2+2}$ corresponds to $\bar{\mu} \bar{\psi}_3[1]$. It is clear that $\bar{\mu} \bar{\psi}_3[1] \neq \bar{\mu} \bar{\psi}_3[1]$, a contradiction.

(D2) If $\tilde{x}_{1+\ell_1+\ell_2-1} \tilde{x}_{1+\ell_1+\ell_2} = \tilde{y}_{\ell_1+\ell_2+1} \tilde{y}_{\ell_1+\ell_2+2}$. Since the same segment following the complementary segment corresponds to $\bar{\mu}$, only $\ell_3 = 3$ can satisfies that the length of the segment equals 1, then neither $\tilde{x}_{1+\ell_1+\ell_2+2} = \tilde{y}_{\ell_1+\ell_2+\ell_3+1}$ nor $\tilde{x}_{1+\ell_1+\ell_2+2} = \tilde{y}_{\ell_1+\ell_2+\ell_3+1}$ can satisfies the conditions of $\tilde{x}_1 \cdots \tilde{x}_{\ell-1} = \psi_1 \psi_2 \bar{\mu} \bar{\psi}_3 \bar{\psi}_4 \psi_5, \tilde{y}_2 \cdots \tilde{y}_\ell = \psi_1 \psi_2 \bar{\mu} \bar{\psi}_3 \bar{\psi}_4 \psi_5$.

In sum, if $|\mathcal{S}_j^i| = 6, |\mathcal{S}_j^i|$ hold simultaneously, then $\tilde{x}_1 \cdots \tilde{x}_{\ell-1} = \psi_1 \psi_2 \bar{\mu} \bar{\psi}_3 \bar{\psi}_4 \psi_5, \tilde{y}_2 \cdots \tilde{y}_\ell = \psi_1 \bar{\psi}_2 \bar{\psi}_3 \bar{\psi}_4 \psi_5$, where $\psi_1, \psi_2, \psi_3, \psi_4, \psi_5$ are alternating sequences, $|\psi_3| = 2$ and ψ_3 is neither zero reversed nor completely reversed in \tilde{x} , i.e. \tilde{x}', \tilde{y}' satisfies the structures in Fig.2 (3).

Denote $|\psi_1|, |\psi_2|, |\psi_3|, |\psi_4|, |\psi_5|$ as $\ell'_1, \ell'_2, \ell'_3, \ell'_4, \ell'_5$. It is clear that $\ell'_1 + \ell'_2 = \ell_1 + \ell_2 + 1, \ell'_3 = 2, \ell'_4 + \ell'_5 = \ell_4 + \ell_5 + 1$. Next, we analyze the structure of s, ω^1, ω^2, t when all the conditions are satisfied simultaneously.

(A) Since we have established that $\tilde{x} = \alpha s \omega^1 t \beta, \tilde{y} = \alpha s \omega^2 t \bar{\beta}$ and s, t are alternating sequence. Combined with the structure in Fig.2 (3), it is easy to see that the $\omega^1 = \tilde{x}_{1+\ell_1+\ell_2+1} \tilde{x}_{1+\ell_1+\ell_2+2} \tilde{x}_{1+\ell_1+\ell_2+3}, \omega^2 = \tilde{y}_{\ell_1+\ell_2+2} \tilde{y}_{\ell_1+\ell_2+3} \tilde{y}_{\ell_1+\ell_2+\ell_3}$, and $\omega^1 = \omega^2$, where $\omega_1^1 = s_{|s|}, \omega_3^1 = \bar{t}_1$.

(B) In Fig.2 (3), we assume that $\gamma = \bar{\alpha}$, and $|s| = \ell_1 + \ell_2 = \ell'_2 - 1$. In order to satisfy $\ell_2 \geq 2, \ell'_2 \geq 2$, then $|s| \geq 3$. In the other case $\gamma = \alpha$, we show in Fig2 (4), $|s| = \ell_2 = 1 + \ell'_2 - 1$, and in order to satisfy $\ell_2 \geq 2, \ell'_2 \geq 2$, $|s| \geq 2$. In sum, we get $|s| \geq 2$ if completely reversed in \tilde{x} and $|s| \geq 3$ if $s_1 = \bar{\alpha}, s_{|s|} = \omega_1^1$.

(C) Due to symmetry, we obtain $|t| \geq 2$ if zero-reversed in

\tilde{x} and $|t| \geq 3$ if $t_1 = \omega_1^1, t_{|t|} = \beta$.

In summary, we have obtained a necessary condition for $|\mathcal{S}_j^i| = 5, |\mathcal{S}_j^i| = 6$, i.e. $\tilde{x} = \alpha s \omega t \beta, \tilde{y} = \alpha s \bar{\omega} t \bar{\beta}$, such that

- s, t, ω are alternating sequences,
- $|\omega| = 3, \omega_1 = s_{|s|}, \omega_3 = \bar{t}_1$,
- if $|s|$ is completely reversed in $\tilde{x}, |s| \geq 2$, else $|s| \geq 3$
- if $|t|$ is zero-reversed in $\tilde{x}, |t| \geq 2$, else $|t| \geq 3$

Next we prove that the above condition is also sufficient.

Firstly, it is clear that $|\mathcal{S}_j^i| = 6$, and then $|\mathcal{S}_j^i| = 4$ according Lemma 8. Thus, we only need to prove that $|\mathcal{S}_j^i| = 5$, and $|\mathcal{S}_j^i| = 4$.

Suppose s ends with δ , according $|\omega| = 3$ and $\omega_1 = s_{|s|}, \omega_3 = \bar{t}_1$, t begins with $\bar{\delta}$, then

$$\begin{aligned}
\tilde{x} &= \alpha s_1 \cdots \delta \bar{\delta} \bar{\delta} \bar{\delta} \cdots t_{|t|} \beta \\
\tilde{y} &= \bar{\alpha} s_1 \cdots \delta \bar{\delta} \bar{\delta} \bar{\delta} \cdots t_{|t|} \bar{\beta}
\end{aligned}$$

$|\mathcal{S}_j^i| = |a_1 \cdots a_{|a|-1} \circ (\mathcal{D}_2(\alpha s \omega t) \cap \mathcal{D}_2(s \bar{\omega} t \bar{\beta})) \circ b_1 \cdots b_{|b|-1}| = |\mathcal{D}_2(\tilde{x}_2 \cdots \tilde{x}_\ell) \cap \mathcal{D}_2(\tilde{y}_1 \cdots \tilde{y}_{\ell-1})|$. Since s, ω, t are alternating sequences, then $\alpha s \omega t, s \bar{\omega} t \bar{\beta}$ can be written as $\alpha s \omega t = \psi_1 \psi_2 \bar{\psi}_3 \bar{\psi}_4 \psi_5, s \bar{\omega} t \bar{\beta} = \psi_1 \bar{\psi}_2 \bar{\psi}_3 \bar{\psi}_4 \psi_5$, where

- $\psi_1 = \alpha$ if s begins with α , else ψ_1 is empty;
- $\psi_2 = s$ if s begins with α , else $\psi_2 = \alpha s$;
- $\psi_3 = \omega_1 \omega_2$
- $\psi_4 = \delta t$ if t ends with β , else $\psi_4 = \delta t_1 \cdots t_{|t|-1}$;
- $\psi_5 = \bar{\beta}$ if t ends with $\bar{\beta}$, else ψ_5 is empty;

Denote $|\psi_1|, |\psi_2|, |\psi_3|, |\psi_4|, |\psi_5|$ as $\ell_1, \ell_2, \ell_3, \ell_4, \ell_5$

Firstly, according to Theorem 3, $|(\mathcal{D}_2(\alpha s \omega t) \cap \mathcal{D}_2(s \bar{\omega} t \bar{\beta}))| < 6$ and we can get the set

$$\begin{aligned} & \{ \\ & \quad \psi_1 \psi_2 [1, \ell_2 - 1] \psi_3 \psi_4 [1, \ell_4 - 1] \psi_5, \\ & \quad \psi_1 \psi_2 [1, \ell_2 - 1] \psi_3 \psi_4 [2, \ell_4] \psi_5, \\ & \quad \psi_1 \psi_2 [2, \ell_2] \psi_3 \psi_4 [1, \ell_4 - 1] \psi_5, \\ & \quad \psi_1 \psi_2 [2, \ell_2] \psi_3 \psi_4 [2, \ell_4] \psi_5, \\ & \quad \psi_1 \bar{\psi}_2 \psi_4 \psi_5 \\ & \} \subseteq (\mathcal{D}_2(\alpha s \omega t) \cap \mathcal{D}_2(s \bar{\omega} t \bar{\beta})). \end{aligned}$$

Thus $|\mathcal{S}_{\tilde{j}}^i| = |(\mathcal{D}_2(\alpha s \omega t) \cap \mathcal{D}_2(s \bar{\omega} t \bar{\beta}))| = 5$.

$\mathcal{S}_{\tilde{j}}^{\bar{i}} = a_2 \cdots a_{|a|} \circ (\mathcal{D}_1(s \omega t) \cap \mathcal{D}_3(\bar{\alpha}' s \bar{\omega} t \bar{\beta})) \circ b_1 \cdots b_{|b|-1}$.

Firstly $|(\mathcal{D}_1(s \omega t) \cap \mathcal{D}_3(\bar{\alpha}' s \bar{\omega} t \bar{\beta}))| \geq 4$ according to Lemma 4 and we can get the set

$$\begin{aligned} & \{ \\ & \quad s \omega_1 \omega_2 t, \\ & \quad s \omega_2 \omega_3 t, \\ & \quad \bar{\alpha} s_3 \cdots s_{|s|} \omega t, \\ & \quad s \omega t_1 \cdots t_{|t|-3} \bar{\beta}, \\ & \} \subseteq (\mathcal{D}_1(s \omega t) \cap \mathcal{D}_3(\bar{\alpha}' s \bar{\omega} t \bar{\beta})). \end{aligned}$$

Thus $|\mathcal{S}_{\tilde{j}}^{\bar{i}}| = |(\mathcal{D}_1(s \omega t) \cap \mathcal{D}_3(\bar{\alpha}' s \bar{\omega} t \bar{\beta}))| = 4$. So far, we have proved the sufficient and necessary condition condition for Type (A).

Symmetrically, we can get the sufficient and necessary condition condition for Type (B) is that $\tilde{x} = \alpha s \omega t \beta, \tilde{y} = \alpha s \bar{\omega} t \bar{\beta}$ such that

- s, t, ω are alternating sequences
- $|\omega| = 3, \omega_1 = \bar{s}_{|s|}, \omega_3 = t_1$,
- if $|s|$ is zero-reversed in $\tilde{x}, |s| \geq 2$, else $|s| \geq 3$.
- if $|t|$ is completely reversed in $\tilde{x}, |t| \geq 2$, else $|t| \geq 3$.

Combining the above two types, we obtain Theorem 3. ■