

The cost of the amount of sexual signalling in the decorated cricket

Introduction

Various species such as insects, can create acoustic signals which convey different messages for different purposes. They can be for sexual signalling, species recognition and resource defence. These acoustic signals can be powerful and can indirectly communicate an individual's condition or location (Wilkins, Seddon and Safran, 2013). The male *Gryllodes sigillatus* (decorated cricket) creates sound pulses by scissoring its forewings and drawing a scraper-like section across a toothed file on the other wing. This action of acoustic communication is used as a mode of sexual signalling to the female, signalling the quality of the males genes (Otte, 1992, Bentley and Hoy, 1974). Indeed, the act of signalling requires high energy, therefore, males of a lower condition tend to struggle to produce a high-quality and frequent signal (Tolle and Wagner, 2011, Gray and Eckhardt, 2001).

The weight change of the cricket will be different depending on diet levels. The higher the nutritional content of the diet level, the greater the increase in weight (Wagner and Hoback, 1999). Larger pronotum size will cause an increase in weight on higher percentage diets (Kell et al, 2014, Newcombe, et al 2015). Larger crickets (greater start weight and pronotum) will be more resistant to changes in weight (Kell et al, 2014). However, this body condition measure will not affect how much the crickets engage with acoustic signalling (Wagner and Hoback, 1999, Gray and Eckhardt, 2001). Even though acoustic signalling is costly, it will not change the weight of the cricket (Hunt, et al 2004).

Analysis

An ordinary least squares model was produced to investigate the energetic costs of sexual signalling in the adult male *Gryllodes sigillatus*. This was represented by the crickets' weight change over the study week in a factorial design, with diet, starting mass, amount of sexual signalling, pronotum size, and categorical predictors included. The final model also included three sets of covariates: the diet/amount of sexual signalling, the diet/pronotum size, and the amount of sexual signalling/pronotum size. The analyses and cleaning of the data took place in R version 4.2.3, using the tidyverse packages (Wickham et al., 2019) and the Janitor package (Firke, 2021). The original data set was filtered to remove incorrect values in the amount of sexual signalling, which had negative/missing values, in the pronotum and the weight change during the study week. All these changes were put into the new data set, 'filtered cricket'. After the data was properly cleaned, some data visualisations surrounding the change in weight were created to explore the spread of the data, trends, or correlations. Some other packages were used, such as patchwork (Pedersen T, 2022), It combines multiple plots to illustrate the data better; ggExtra (Attali and Baker, 2022) adds marginal histograms to ggplot2. The first model was based on past literature and data visualisations. This model included many interaction terms; the only one not included was the starting mass and the diet over the week, as this was not relevant. To make the most parsimonious model, a step-down procedure was then taken to remove the variables that were not statistically significant or had an irregular AIC from the intercept. After the final model, summary tables were completed using the packages Broom Tidy (Robinson et al., 2022) and Kable (Zhu 2020). An extra package called 'Scales' (Wickham H, Seidel D, 2022) formatted the p Values. An ANOVA was used to gain a greater depth of analysis further.

Results and Discussion

Condition of the crickets (pronotum and starting mass)

It was hypothesised that the larger the starting mass of the cricket, the more resistant the cricket would be to changes in their weight, and with the lower starting mass of the cricket, the greater the increase in their weight. Starting mass is a highly significant main effect (Linear model $F_{1,529} = 19.9$, $P < 0.001$). There was not enough evidence to support the above hypothesis. For every gram increase in the cricket's starting mass, the cricket lost -0.55296g [95% CI: -0.6227919 - -0.4831234], illustrated in Figure and Table one. The starting mass of the male is often used as a proxy of the male's quality due to females having a preferential choice of selecting larger males. Moreover, Chown and Gaston (2010) found a trade-off where the fitness advantage of a larger size is out weighted and can actually reduce the fitness of the next generation.




Figure 1: *Gryllobates sigillatus* over the first week of the study, the crickets with the higher starting masses lost more weight. This indicates that every 1-gram increase in the starting mass, causes a -0.55296g [95% CI: -0.6227919 - -0.4831234] loss in the cricket's weight. The line is the regression slope with a 95% confidence interval, and the dots are the individual data points. The marginal plots are histograms of the distribution of starting mass (top), and weight change (right).

Table 1: This shows outputs from the linear model of *Gryllobates sigillatus*. It includes the confidence intervals, high and low, p-value, t-value, standard error and coefficient. The highly significant ones are highlighted in pink.

<i>Term</i>	<i>Coefficient</i>	<i>Standard Error</i>	<i>t</i>	<i>p value</i>	<i>lower.CI</i>	<i>upper.CI</i>
(Intercept)	-0.03553	0.03241	-1.09641	0.273	-0.0991913	0.0281300
Diet_Percentage	-0.00152	0.00051	-2.98799	0.003	-0.0025269	-0.0005222
Starting_mass	-0.55296	0.03555	-15.55487	<0.001	-0.6227919	-0.4831234
Sexaul_signalling	-0.00743	0.00394	-1.88871	0.059	-0.0151657	0.0002981
Pronotum_size	0.05698	0.01434	3.97377	<0.001	0.0288108	0.0851461
Diet_Percentage:Sexaul_signalling	0.00002	0.00001	2.36772	0.018	0.0000036	0.0000387
Diet_Percentage:Pronotum_size	0.00075	0.00022	3.49713	<0.001	0.0003302	0.0011766
Sexaul_signalling:Pronotum_size	0.00358	0.00163	2.19123	0.029	0.0003700	0.0067814

It was hypothesised that the larger the pronotum, the greater the resistance to weight loss the cricket would be. Pronotum as a main interaction term is highly significant (Linear model $F_{1,529} = 19.9$, $P < 0.001$); for every pronotum millimetre increase in weight, the amount of weight gained will be 0.05698g [95% CI: 0.0288108 - 0.0851461], (seen in table 1). Due to the interactions with other variables, the main effect of the pronotum cannot be properly investigated without those interactions mentioned later. These results show that this hypothesis is true due to this evidence. As the size of the pronotum increases, so does the weight of the cricket (**Figure 2**). Indeed, Kell et al (2014) found that pronotum and weight are strongly positively correlated together.

It was hypothesised that pronotum size would not impact the amount of sexual signalling. But the amount that the crickets sexually signalled and interacted with the size of the pronotum is significant (Linear model $F_{1,529} = 4.8$, $P = 0.029$), with an increase in pronotum size in millimetres, and an increase in the amount of sexual signalling increased by a second, would cause an increase in the weight of the cricket by 0.003558g [95% CI: 0.0003700 - 0.0067814] (**Figure 2**, **Table 1**). However, there was no evidence in the literature finding this result. Most of it reads that there is no significant interaction effect between the two variables and their weight (Wagner and Hoback, 1999, Gray and Eckhardt, 2001).




Figure 2: The *Gryllodes sigillatus* weight increases with pronotum size and differs when the pronotum interacts with sexual signalling. This indicates that with an increase in pronotum size in millimetres, and an increase in the amount of sexual signalling increased by a second. This caused an increase in the weight of the cricket by 0.003558g [95% CI: 0.0003700 - 0.0067814]. The regression line has 95% confidence intervals. The dots are individual data points, and the colour indicates the amount of sexual signalling, with red being the most and purple being the least.

The effects of different nutritional diet percentages and the interactions

The hypothesis was that the higher the nutritional content of the diet was, then the greater the increase in weight (Wagner and Hoback, 1999). There was a significant overall effect of the diet percentage with the change in weight in the male crickets (Linear model $F_{1,529} = 325.2$, $P = 0.003$). Males on a higher diet percentage lost more weight, -0.00152g [95% CI: -0.0025269 - -0.0005222] (**Figure 1**), with more weight lost, for every increase in the nutritional diet content. At this time there is not enough evidence to support this hypothesis. Another study supported this by carrying out experimental reductions in nutrient quality. This typically resulted in reductions in critical weight, which could possibly be due to insulin signalling (Chown and Gaston, 2010). Yet Wagner and Hoback in 1999, and Kelly, in 2014, found evidence that diet quality affects the body mass of crickets when controlling body size.

However, the diet percentage as a main interaction term must be interpreted with caution as diet percentage also interacts with the amount of singing over the week, and the size of the pronotum. It was hypothesised that increasing the diet percentage would allow the crickets to signal additionally and gain more weight sexually. The interaction between diet and the amount of sexual signalling was significant (Linear model $F_{1,529} = 7.2$, $P = 0.018$). When the diet percentage was higher, crickets sexually signalled more, and the weight

of the crickets increased by 0.00002g [95% CI: 0.0000036 - 0.0000387], (**Figure 3, Table 1**). This result was also found by Wagner and Hoback in 1999. They found that crickets did not lose weight as they adjusted the amount of sexual signalling according to the nutritional diet that each of the crickets was fed, further backing up this hypothesis. Juxtaposing this, in a study by Hunt, et al in 2004, it was found that crickets on diets with higher percentages, would have increased sexual signals, and the more weight the crickets lost, contrasted with the crickets on the lower diet percentages. This also reduced the lifespan of the crickets (Hunt, et al 2004).




Figure 3: When the *Gryllobates sigillatus* diet percentage was higher, there was more sexual signalling, and also, their weight increased by 0.00002g [95% CI: 0.0000036 - 0.0000387]. The colours on the graph represent the different diet percentages displayed in the key at the top. The regression lines show the trends in the different diet percentages; the dots are the individual data points. A marginal density plot for the distribution of the different weights, is separated by the different diet percentages, seen to the right.

When the amount of sexual signalling is included as a main effect, it is insignificant as a predictor of weight change (Linear model $F_{1,529} = 83.7$, $P = 0.059$), yet it was still included in the model due to the above-mentioned interactions effect.

I hypothesised that the greater the size of the pronotum, the greater the increase in weight at higher diet percentages. Diet interacting with pronotum size is highly significant (Linear model $F_{1,529} = 19.9$, $P < 0.001$); with larger pronotums in millimetres and a higher diet percentage, the weight of the cricket will increase by 0.000753g [95% CI: 0.0003302 - 0.0011766] (**Figure 4, Table 1**). The diet is highly statistical in the developing stages of the crickets life. The higher diet percentage means larger pronotum, as seen in this study, yet the cricket is not an adult (Newcombe, et al 2015).




Figure 4: In *Gryllodes sigillatus* the larger the pronotum size, and the higher the diet percentage, the greater the increase in weight $0.000753g$ [95% CI: $0.0003302 - 0.0011766$]. The lines are the regression slopes, with 95% confidence intervals; the dots are individual data points. Each plot depicts a different diet percentage, with a different colour for each one.

Conclusion

It was seen that all the variables in the study had a relationship with the weight change in the male cricket. The weight of the cricket was a measure to understand the cost of sexual signalling, amongst other factors. The weight of the cricket was highly affected by the starting mass of the cricket, pronotum size, and diet/pronotum, interacting together. With the starting mass negatively affecting the weight of the diet, pronotum size increases the weight of the cricket, and diet/pronotum also increases the weight of the cricket. However, interestingly, the calling effort, or amount of sexual signalling, does not significantly affect the weight of the cricket. Yet, when the diet is included as an interaction term, it is significant, which is supported by the literature that the crickets will adjust the amount of sexual signalling (Hunt, et al 2004). The amount of sexual signalling/pronotum interaction increases the weight of the cricket. In the future, another study should include the life span of the crickets, as another potential variable, and that if it is affected by the cost of sexual signalling, and the weight of the cricket.

References

- Attali D, Baker C (2022). `__ggExtra`: Add Marginal Histograms to ‘ggplot2’, and More ‘ggplot2’ Enhancements_. R package version 0.10.0, <https://github.com/daattali/ggExtra>.
- Bentley, D. and Hoy, R.R., 1974. The neurobiology of cricket song. *Scientific American*, 231(2), pp.34-45.
- Chown, S.L. and Gaston, K.J., 2010. Body size variation in insects: a macroecological perspective. *Biological Reviews*, 85(1), pp.139-169.
- Firke S (2021) janitor: Simple Tools for Examining and Cleaning Dirty Data, URL <https://github.com/sfirke/janitor>
- Gray, D.A. and Eckhardt, G., 2001. Is cricket courtship song condition dependent?. *Animal Behaviour*, 62(5), pp.871-877.
- Hunt, J., R.Brooks, M. D.Jennions, M. J.Smith, C. L.Bentsen, and L. F.Bussière. 2004. High-quality male field crickets invest heavily in sexual display but die young. *Nature* 432:1024–1027.
- Judge, K.A., Ting, J.J. and Gwynne, D.T., 2008. Condition dependence of male life span and calling effort in a field cricket. *Evolution*, 62(4), pp.868-878.
- Kable(2021) kableExtra: Construct Complex Table with ‘kable’ and Pipe Syntax:Hao Zhu ORCID iD [aut, cre], Thomas Trivison [ctb], Timothy Tsai [ctb], Will Beasley [ctb], Yihui Xie [ctb], GuangChuang Yu [ctb], Stéphane Laurent [ctb], Rob Shepherd [ctb], Yoni Sidi [ctb], Brian Salzer [ctb], George Gui [ctb], Yeliang Fan [ctb], Duncan Murdoch [ctb], Bill Evans [ctb], URL <http://haozhu233.github.io/kableExtra/>, <https://github.com/haozhu233/kableExtra>
- Kelly, C.D., Tawes, B.R. and Worthington, A.M., 2014. Evaluating indices of body condition in two cricket species. *Ecology and Evolution*, 4(23), pp.4476-4487.
- Newcombe, D., Moore, P.J. and Moore, A.J., 2015. The role of maternal effects in adaptation to different diets. *Biological Journal of the Linnean Society*, 114(1), pp.202-211.
- Otte, D., 1992. Evolution of cricket songs. *Journal of orthoptera research*, (1), pp.25-49.
- Pedersen T (2022). `__patchwork`: The Composer of Plots_. <https://patchwork.data-imaginist.com>, <https://github.com/thomasp85/patchwork>.
- R Core Team (2022). R: A language and environment for statistical computing. R Foundation for Statistical Computing, Vienna, Austria. URL <https://www.R-project.org/>.
- RStudio Team (2020). RStudio: Integrated Development for R. RStudio, PBC, Boston, MA URL <http://www.rstudio.com/>.
- Simmons, L.W., 1995. Correlates of male quality in the field cricket, *Gryllus campestris* L.: age, size, and symmetry determine pairing success in field populations. *Behavioral Ecology*, 6(4), pp.376-381.
- Tolle, A.E. and Wagner Jr, W.E., 2011. Costly signals in a field cricket can indicate high-or low-quality direct benefits depending upon the environment. *Evolution*, 65(1), pp.283-294.
- Wagner JR, W.E. and Hoback, W.W., 1999. Nutritional effects on male calling behaviour in the variable field cricket. *Animal behaviour*, 57(1), pp.89-95.
- Wickham H, Averick M, Bryan J, Chang W, McGowan LD, François R, Golemund G, Hayes A, Henry L, Hester J, Kuhn M, Pedersen TL, Miller E, Bache SM, Müller K, Ooms J, Robinson D, Seidel DP, Spinu V, Takahashi K, Vaughan D, Wilke C, Woo K, Yutani H (2019). “Welcome to the tidyverse.” *Journal of Open Source Software*, 4(43), doi:10.21105/joss.01686.
- Wickham H, Seidel D (2022). `__scales`: Scale Functions for Visualization_. <https://scales.r-lib.org>, <https://github.com/r-lib/scales>.
- Wilke C (2021). `__ggridges`: Ridgeline Plots in ‘ggplot2’_. R package version 0.5.3, <https://wilkelab.org/ggridges/>.

Wilkins, M.R., Seddon, N. and Safran, R.J., 2013. Evolutionary divergence in acoustic signals: causes and consequences. *Trends in ecology & evolution*, 28(3), pp.156-166.