

Corner Boundary Conditions in lattice Boltzmann method

Goncalo Silva

Department of Mechanical Engineering
Instituto Superior Técnico (IST)
Lisbon, Portugal



Outline


- 1 Introduction
 - Corners in Boundary Value Problems
- 2 Motivation
 - Navier-Stokes Boundary Conditions: Corners
- 3 Lattice Boltzmann Boundary Conditions: Corners
 - Problem definition
 - Particulate dynamics
 - Using Chapman-Enskog

Outline

- 1 Introduction
 - Corners in Boundary Value Problems
- 2 Motivation
 - Navier-Stokes Boundary Conditions: Corners
- 3 Lattice Boltzmann Boundary Conditions: Corners
 - Problem definition
 - Particulate dynamics
 - Using Chapman-Enskog


Definitions

- Boundary value problem geometry:



Definitions


- Boundary value problem geometry:



- **Corner:** Place in between two boundaries, e.g. $\partial\Omega_1$ and $\partial\Omega_2$

Definitions


- Boundary value problem geometry:



- **Corner:** Place in between two boundaries, e.g. $\partial\Omega_1$ and $\partial\Omega_2$
- Particularly important when e.g. $\varphi(\partial\Omega_1) \neq \varphi(\partial\Omega_2)$

Definitions


- Boundary value problem geometry:



- **Corner:** Place in between two boundaries, e.g. $\partial\Omega_1$ and $\partial\Omega_2$
- Particularly important when e.g. $\varphi(\partial\Omega_1) \neq \varphi(\partial\Omega_2)$
- Smooth evolution of φ on $\partial\Omega$

Definitions

- Boundary value problem geometry:



- **Corner:** Place in between two boundaries, e.g. $\partial\Omega_1$ and $\partial\Omega_2$
- Particularly important when e.g. $\varphi(\partial\Omega_1) \neq \varphi(\partial\Omega_2)$
- Smooth evolution of φ on $\partial\Omega$
- Avoid singularities of φ on $\partial\Omega$

Outline

- 1 Introduction
 - Corners in Boundary Value Problems
- 2 Motivation
 - Navier-Stokes Boundary Conditions: Corners
- 3 Lattice Boltzmann Boundary Conditions: Corners
 - Problem definition
 - Particulate dynamics
 - Using Chapman-Enskog

Definitions

- Isothermal and incompressible Navier-Stokes equations

$$\begin{cases} (\mathbf{u} \cdot \nabla) \mathbf{u} = -\nabla p + \nu \Delta \mathbf{u} + \mathbf{a} \\ \nabla \cdot \mathbf{u} = 0 \end{cases} \quad \text{in } \Omega$$

... with boundary conditions setting $\{p, \mathbf{u}\}$ on $\partial\Omega$

Definitions

- Isothermal and incompressible Navier-Stokes equations

$$\begin{cases} (\mathbf{u} \cdot \nabla) \mathbf{u} = -\nabla p + \nu \Delta \mathbf{u} + \mathbf{a} \\ \nabla \cdot \mathbf{u} = 0 \end{cases} \quad \text{in } \Omega$$

... with boundary conditions setting $\{p, \mathbf{u}\}$ on $\partial\Omega$

- **Corners:** Define $\{p, \mathbf{u}\}$ from adjacent locations using interpolation or extrapolation

Definitions

- Isothermal and incompressible Navier-Stokes equations

$$\begin{cases} (\mathbf{u} \cdot \nabla) \mathbf{u} = -\nabla p + \nu \Delta \mathbf{u} + \mathbf{a} \\ \nabla \cdot \mathbf{u} = 0 \end{cases} \quad \text{in } \Omega$$


... with boundary conditions setting $\{p, \mathbf{u}\}$ on $\partial\Omega$

- **Corners:** Define $\{p, \mathbf{u}\}$ from adjacent locations using interpolation or extrapolation
- Everything else follows the same philosophy of boundary conditions

Examples

- Examples of flow domains with corners:


Backward facing step flow



Examples

- Examples of flow domains with corners:

Lid-driven cavity flow



Outline

- 1 Introduction
 - Corners in Boundary Value Problems
- 2 Motivation
 - Navier-Stokes Boundary Conditions: Corners
- 3 Lattice Boltzmann Boundary Conditions: Corners
 - Problem definition
 - Particulate dynamics
 - Using Chapman-Enskog

Introduction and motivation

- Lattice Boltzmann method (LBM)

$$f_{\alpha}(\mathbf{x} + \mathbf{c}_{\alpha}\Delta t, t + \Delta t) = f_{\alpha}(\mathbf{x}, t) - \omega(f_{\alpha} - f_{\alpha}^{(eq)})|_{(\mathbf{x}, t)} \quad \text{in } \Omega$$

Introduction and motivation

- Lattice Boltzmann method (LBM)

$$f_{\alpha}(\mathbf{x} + \mathbf{c}_{\alpha}\Delta t, t + \Delta t) = f_{\alpha}(\mathbf{x}, t) - \omega(f_{\alpha} - f_{\alpha}^{(eq)})|_{(\mathbf{x}, t)} \quad \text{in } \Omega$$

- **Corners** are subjected to the same (or even more complex) kind of problems of general plane boundary conditions in LBM

Introduction and motivation

- Lattice Boltzmann method (LBM)

$$f_{\alpha}(\mathbf{x} + \mathbf{c}_{\alpha}\Delta t, t + \Delta t) = f_{\alpha}(\mathbf{x}, t) - \omega(f_{\alpha} - f_{\alpha}^{(eq)})|_{(\mathbf{x}, t)} \quad \text{in } \Omega$$

- **Corners** are subjected to the same (or even more complex) kind of problems of general plane boundary conditions in LBM

→ Solution on $\partial\Omega$ is specified for f_{α} and **NOT** for $\{\rho, \mathbf{u}, \Pi\}$

Introduction and motivation

- Lattice Boltzmann method (LBM)

$$f_{\alpha}(\mathbf{x} + \mathbf{c}_{\alpha}\Delta t, t + \Delta t) = f_{\alpha}(\mathbf{x}, t) - \omega(f_{\alpha} - f_{\alpha}^{(eq)})|_{(\mathbf{x}, t)} \quad \text{in } \Omega$$

- **Corners** are subjected to the same (or even more complex) kind of problems of general plane boundary conditions in LBM

→ Solution on $\partial\Omega$ is specified for f_{α} and **NOT** for $\{\rho, \mathbf{u}, \Pi\}$

→ f_{α} set in a higher DoF system than $\{\rho, \mathbf{u}, \Pi\}$, hence:

- Trivial: $f_{\alpha} \longrightarrow \{\rho, \mathbf{u}, \Pi\}$
- Complex: $\{\rho, \mathbf{u}, \Pi\} \longrightarrow f_{\alpha}$

Introduction and motivation

- Lattice Boltzmann method (LBM)

$$f_{\alpha}(\mathbf{x} + \mathbf{c}_{\alpha}\Delta t, t + \Delta t) = f_{\alpha}(\mathbf{x}, t) - \omega(f_{\alpha} - f_{\alpha}^{(eq)})|_{(\mathbf{x}, t)} \quad \text{in } \Omega$$

- **Corners** are subjected to the same (or even more complex) kind of problems of general plane boundary conditions in LBM

→ Solution on $\partial\Omega$ is specified for f_{α} and **NOT** for $\{\rho, \mathbf{u}, \Pi\}$

→ f_{α} set in a higher DoF system than $\{\rho, \mathbf{u}, \Pi\}$, hence:

- Trivial: $f_{\alpha} \longrightarrow \{\rho, \mathbf{u}, \Pi\}$
- Complex: $\{\rho, \mathbf{u}, \Pi\} \longrightarrow f_{\alpha}$

→ Incorrect upscaling → Unwanted behavior, e.g. Knudsen layers

Importance of corners

- **Corners** why bothering?

Importance of corners

- **Corners** why bothering?

↑ Corner nodes are just a few nodes on $\partial\Omega$

Importance of corners

- **Corners** why bothering?

- ↑ Corner nodes are just a few nodes on $\partial\Omega$

- ↓ Inaccurate corner implementation may contaminate the solution everywhere in the domain

Importance of corners

- **Corners** why bothering?


- ↑ Corner nodes are just a few nodes on $\partial\Omega$

- ↓ Inaccurate corner implementation may contaminate the solution everywhere in the domain

- ↓ Interpolation or extrapolation may not be possible at corners

Lattice structure

...remember!



- D2Q9 model

$$c_1 = (0, 0)$$

$$c_2 = (1, 0)$$

$$c_3 = (0, 1)$$

$$c_4 = (-1, 0)$$

$$c_5 = (0, -1)$$

$$c_6 = (1, 1)$$



$$c_7 = (-1, 1)$$

$$c_8 = (-1, -1)$$

$$c_9 = (1, -1)$$

Bounceback on corner

Bottom-left Corner



Bounceback on corner

• Pros

- Mass is exactly conserved
- Stable for ω close to 2 (i.e. for high Re)
- Local
- Flexibility in handling wall, edges, corners both in 2D and 3D
- Very simple to implement from a programming viewpoint

• Cons

- Velocity accuracy may decrease from 2nd to 1st
- Pressure accuracy may decrease from 1st to 0th
- In SRT model momentum is not exactly conserved (viscosity dependent slip velocity)

Bounceback on **corner**

Question:

Apply the half-way bounceback scheme to **top-left** and **bottom-right** corners

Zou He on corner

- Boundary node and solid node coincide
- Only unknown incoming populations are modified
- Set ρ or \mathbf{u} in $f_{\alpha}^{(0)}(\rho, \mathbf{u})$
- Construct $f_{\alpha}^{(1)}$ from the symmetry requirement

Zou He on corner


- Boundary node and solid node coincide
- Only unknown incoming populations are modified
- Set ρ or \mathbf{u} in $f_{\alpha}^{(0)}(\rho, \mathbf{u})$
- Construct $f_{\alpha}^{(1)}$ from the symmetry requirement
- Additional problem: the so-called "buried links"

Zou He on Corners


- Known:

$$\rightarrow \mathbf{u} = \mathbf{0}$$

$$\rightarrow f_{\alpha} = (f_4, f_5, f_8)$$



Zou He on Corners



- Known:

- $\mathbf{u} = \mathbf{0}$


- $f_\alpha = (f_4, f_5, f_8)$

- Unknown (6 variables):

- ρ

- $f_\alpha = (f_2, f_3, f_6, f_7, f_9)$


Zou He on Corners



- Known:
 - $\rightarrow \mathbf{u} = \mathbf{0}$
 - $\rightarrow f_\alpha = (f_4, f_5, f_8)$
- Unknown (6 variables):
 - $\rightarrow \rho$
 - $\rightarrow f_\alpha = (f_2, f_3, f_6, f_7, f_9)$
- 3 Equations (2 linearly independent):
 - $\rightarrow \sum f_\alpha = \rho$
 - $\rightarrow \sum \mathbf{c}_\alpha f_\alpha = \mathbf{u}$

Zou He on corner

1) Computing $\rho...$



- Population velocity set at corner node:


$$\rightarrow C_+ = \{\mathbf{c}_3, \mathbf{c}_6, \mathbf{c}_7\}$$

$$\rightarrow C_0 = \{\mathbf{c}_1, \mathbf{c}_2, \mathbf{c}_4\}$$

$$\rightarrow C_- = \{\mathbf{c}_5, \mathbf{c}_8, \mathbf{c}_9\}$$

Zou He on corner

1) Computing ρ ...



- Population velocity set at corner node:

$$\rightarrow C_+ = \{\mathbf{c}_3, \mathbf{c}_6, \mathbf{c}_7\}$$


$$\rightarrow C_0 = \{\mathbf{c}_1, \mathbf{c}_2, \mathbf{c}_4\}$$

$$\rightarrow C_- = \{\mathbf{c}_5, \mathbf{c}_8, \mathbf{c}_9\}$$

- Not possible** to compute ρ as in boundary planes

Zou He on corner

1) Computing ρ ...



- Population velocity set at corner node:

$$\rightarrow C_+ = \{c_3, c_6, c_7\}$$

$$\rightarrow C_0 = \{c_1, c_2, c_4\}$$

$$\rightarrow C_- = \{c_5, c_8, c_9\}$$


- Not possible** to compute ρ as in boundary planes
- Inevitable **solution**: extrapolate ρ from adjacent nodes


Zou He on corner

2) Computing $\{f_2, f_3, f_6, f_7, f_9\} \dots$


- Symmetry of $f_\alpha^{(1)}$ (3 equations):

→ Bounceback of non-equilibrium populations






- Symmetry of $f_{\alpha}^{(1)}$ (3 equations):
 - Bounceback of non-equilibrium populations
- Still 2 unknowns left:
 - Populations of buried link $\{f_7, f_9\}$



- Symmetry of $f_{\alpha}^{(1)}$ (3 equations):
 - Bounceback of non-equilibrium populations
- Still 2 unknowns left:
 - Populations of buried link $\{f_7, f_9\}$
- Problem underspecified:
 - 5 eqs. and 6 unknowns

Zou He on corner

2) Computing $\{f_2, f_3, f_6, f_7, f_9\} \dots$



- Non-equilibrium bounceback:

$$f_2 - f_2^{(0)} = f_4 - f_4^{(0)}$$

$$f_3 - f_3^{(0)} = f_5 - f_5^{(0)}$$

$$f_6 - f_6^{(0)} = f_8 - f_8^{(0)}$$

Zou He on corner

2) Computing $\{f_2, f_3, f_6, f_7, f_9\}....$

- Solution for some of the unknown incoming populations (from non-equilibrium bounceback):

$$f_2 = f_4 + \frac{2}{3}u_x$$

$$f_3 = f_5 + \frac{2}{3}u_y$$

$$f_6 = f_8 + \frac{1}{6}(u_x + u_y)$$


Zou He on corner

2) Computing $\{f_2, f_3, f_6, f_7, f_9\}$

- Buried populations:


$$u_x = (f_2 + f_6 + f_9) - (f_4 + f_7 + f_8)$$

$$u_y = (f_3 + f_6 + f_7) - (f_5 + f_8 + f_9)$$



Zou He on corner

2) Computing $\{f_2, f_3, f_6, f_7, f_9\}$



- Buried populations:

$$u_x = (f_2 + f_6 + f_9) - (f_4 + f_7 + f_8)$$

$$u_y = (f_3 + f_6 + f_7) - (f_5 + f_8 + f_9)$$

- Possible solution:

$$f_7 = \frac{1}{12}(u_y - u_x)$$

$$f_9 = \frac{1}{12}(u_x - u_y)$$

Zou He on corner

2) Computing f_1

- Recall density has been extrapolated. Therefore, to ensure proper upscaling between macroscopic parameters and LB populations:

1) Set $f_1 = 0$

2) Compute $f_1 = \rho - \sum f_\alpha$

Zou He on **corner**

Question:

Apply Zou He boundary condition scheme to **top-left** and **bottom-right** corners

Exercise III

Exercise III:

Backward facing step flow