

Temporal and spatial variations of PM2 and PM10 concentrations in Mongolia

Erdenebayar Munkhtsetseg^{1,1}, Atsushi Shimizu¹,

^a*National University of Mongolia (NUM), Mongolia,*

^b*Kanazawa University, Japan,*

^c*National Institute for Environmental Studies (NIES), Japan,*

Abstract


PM2.5 and PM10 data for the 4 distinct sites of Mongolia from 2008 to 2020 is found

Keywords: particulate matters, concentrations of PM10 and PM2.5

1. Introduction

2. Data & Methods

2.1. Study area descriptions



The map demonstrates that llll

2.2. Study data and data analysis

Table ?? summarises the data. cleaned the data and filled the missing data using mtsdi

*Corresponding author

Table 1: Measured data summary

Table 1. Measured data

SITE	Location		Measured and collected data						Missing data	
	COORDINATE	ELEVATION	TOTAL ¹	WS&WD ²	VIS ³	OPC ⁴	PM2.5 ⁵	PM10 ⁵	PM2.5	PM10
Ulaanbaatar	47.92°N, 106.92°E	1350 m	76656	72603	72886	33241	67940	68777	11.4%	10.3%
Dalanzadgad	43.57°N, 104.42°E	1470 m	60336	46332	33812	-	46066	49172	23.6%	18.5%
Sainshand	44.87°N, 110.12°E	947 m	59040	50513	49720	-	47111	47313	20.2%	19.9%
Zamyn Uud	43.72°N, 111.90°E	967 m	67392	62432	63948	-	57317	58512	14.9%	13.2%

¹ Equipment height: 15 meter at urban site (Ulaanbaatar), 2 meter at Gobi sites (Dalanzadgad, Sainshand and Zamyn Uud); ² Measurement range: 0–60 m/s; 0–365 degrees. Instrument model: Wind speed and direction PGWS-100, Gill, England; ³ Range: 10–20 000 m. Visibility meter PWD10, Vaisala, Finland; ⁴ Optical Particle Counter; ⁵ Range: 0.003–100 mg/m³, Flow rate: 20 L/m, Suction rate: 2 L/ m. Measured by Kosa monitor ES-640, TDK Co. LTD, Japan;

3. Results

4. Discussion

5. Conclusions

References