

# “Calculus 3”

## Multi-Variable Calculus

Instructor: Álvaro Lozano-Robledo

### Day 6

1

### Any Reminders? Any Questions?

- Class ends at 3:15.
- Slides are being posted on GitHub!  
<https://github.com/alozanoroble/MATH-2110Q-Spring-2026>
- Videos will be posted on YouTube... but they may lag!
- Request videos!!

2

Questions?

3



*ALVARO:* Start the recording!



4

# “Calculus 3”

## Multi-Variable Calculus


Instructor: Álvaro Lozano-Robledo

## Partial Derivatives - Examples

5

## Today – Derivatives!

- Partial Derivatives
- Interpretation
- Higher Derivatives
- PDEs



6

## Partial Derivatives – The Limit Definition

If  $f$  is a function of two variables, its **partial derivatives** are the functions  $f_x$  and  $f_y$  defined by

$$f_x(x, y) = \lim_{h \rightarrow 0} \frac{f(x + h, y) - f(x, y)}{h}$$

$$f_y(x, y) = \lim_{h \rightarrow 0} \frac{f(x, y + h) - f(x, y)}{h}$$

7

## Partial Derivatives – Notation

If  $z = f(x, y)$ , we write

$$f_x(x, y) = f_x = \frac{\partial f}{\partial x} = \frac{\partial}{\partial x} f(x, y) = \frac{\partial z}{\partial x} = f_1 = D_1 f = D_x f$$

$$f_y(x, y) = f_y = \frac{\partial f}{\partial y} = \frac{\partial}{\partial y} f(x, y) = \frac{\partial z}{\partial y} = f_2 = D_2 f = D_y f$$

8

**Example:** Find the partial derivatives of  $f(x, y) = 4 - x^2 - y^2$  at  $(1, 1)$  and interpret those as slopes.

9

**Example:** Find the partial derivatives of  $f(x, y) = 4 - x^2 - y^2$  at  $(1, 1)$  and interpret those as slopes. [Extra space]

10

Example: Find the partial derivatives of  $f(x, y) = x \cdot \ln(y^2 - x)$   
at (3,2).

11

Example: Find the partial derivatives of  $f(x, y) = x \cdot \ln(y^2 - x)$   
at (3,2). [Extra space]

12

Example: Find the partial derivatives of the Body-Mass-Index formula

$$B(m, h) = m/h^2 \quad \text{at} \quad m = 64\text{kg} \quad \text{and} \quad h = 1.68\text{m}.$$

13

Example: Find the partial derivatives of the Body-Mass-Index formula

$$B(m, h) = m/h^2 \quad \text{at} \quad m = 64\text{kg} \quad \text{and} \quad h = 1.68\text{m}. \text{ [Extra]}$$

14

Example: Find the partial derivatives of  $f(x, y, z) = \sin(x^2 + y^2 + z^2)$   
at  $(1, 2, 3)$ .

15

Example: Find the partial derivatives of  $f(x, y, z) = \sin(x^2 + y^2 + z^2)$   
at  $(1, 2, 3)$ .  
[Extra space]

16



## Higher Partial Derivatives

17

Example: Find the second partial derivatives of

$$f(x, y) = 4x^2y - x^3 - y^2$$

18


Example: Find the second partial derivatives of

$$f(x, y) = 4x^2y - x^3 - y^2$$

[Extra space]

19

## Partial Differential Equations



Example: The Heat Equation

$$\frac{\partial u}{\partial t} = \alpha \left( \frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} + \frac{\partial^2 u}{\partial z^2} \right)$$

20


**Example:** Show that the function  $w(x,t) = \sin(x - a \cdot t)$  satisfies the wave equation:

$$\frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial^2 w}{\partial x^2}$$

21

**Example:** Show that the function  $w(x,t) = \sin(x - a \cdot t)$  satisfies the wave equation:

$$\frac{\partial^2 w}{\partial t^2} = a^2 \frac{\partial^2 w}{\partial x^2}$$



22

Questions?

23



*ALVARO:* Start the recording!



24

# “Calculus 3”

## Multi-Variable Calculus


Instructor: Álvaro Lozano-Robledo

### Tangent Planes

25

## Today – Tangent Planes!

- Equation
- Linear Approximations
- Differentiability
- Differentials



26

# Equation of a Tangent Plane

27

# Equation of a Tangent Plane

## 2 Equation of a Tangent Plane

Suppose  $f$  has continuous partial derivatives. An equation of the tangent plane to the surface  $z = f(x, y)$  at the point  $P(x_0, y_0, z_0)$  is

$$z - z_0 = f_x(x_0, y_0)(x - x_0) + f_y(x_0, y_0)(y - y_0)$$


Note the similarity between the equation of a tangent plane and the equation of a tangent line:

$$y - y_0 = f'(x_0)(x - x_0)$$

28

Example: Find the tangent plane at (3,2,5) to the hyperbolic paraboloid

$$z = x^2 - y^2$$



29



Example: Find the tangent plane at (3,2,5) to the hyperbolic paraboloid

$$z = x^2 - y^2$$

[Extra space]


30

## Linear Approximations



31

## Linear Approximations



32



Example: Find the linear approximation at  $(x,y)=(3,-1)$  of

$$f(x, y) = 2x^2 - xy + 3y^2$$

33

Example: Find the linear approximation at  $(x,y)=(3,-1)$  of

$$f(x, y) = 2x^2 - xy + 3y^2$$

[Extra space]

34

Example: Find the linear approximation at  $(x,y)=(2,3)$  and use it to approximate  $\sqrt{21}$ , where

$$f(x, y) = \sqrt{x^2 + 4y}$$

35

Example: Find the linear approximation at  $(x,y)=(2,3)$  and use it to approximate  $\sqrt{21}$ , where

$$f(x, y) = \sqrt{x^2 + 4y}$$

[Extra space]

$$z = \frac{3}{2} + \frac{x}{2} + \frac{y}{2}$$

$$\sqrt{21} = 4.58257569496...$$

36


# Questions?



37

# Thank you


Until next time.



38




ALVARO: Start the recording!



39

“Calculus 3”




Multi-Variable Calculus

Instructor: Álvaro Lozano-Robledo




The Chain Rule



40



# Today – The Chain Rule!

- 
- The Single Variable Case
  - Chain Rule with One Parameter
  - Chain Rule with Two Parameters

41



## The Good Ol' Chain Rule



42

Example: Find the derivative of  $f(g(t))$  with respect to  $t$  where

$$f(x) = \sin(x) \quad \text{and} \quad g(t) = t^2 + 1$$

43

Example: Find the derivative of  $f(g(t))$  with respect to  $t$  where

$$f(x) = \sin(x) \quad \text{and} \quad g(t) = t^2 + 1$$

[Extra space]

44

# The New Chain Rule – Case 1

## 1 The Chain Rule (Case 1)

Suppose that  $z = f(x, y)$  is a differentiable function of  $x$  and  $y$ , where  $x = g(t)$  and  $y = h(t)$  are both differentiable functions of  $t$ . Then  $z$  is a differentiable function of  $t$  and

$$\frac{dz}{dt} = \frac{\partial f}{\partial x} \frac{dx}{dt} + \frac{\partial f}{\partial y} \frac{dy}{dt}$$

45

Example: Find the derivative of  $f(g(t), h(t))$  with respect to  $t$  where

$$f(x, y) = x^2 + y \quad \text{and} \quad g(t) = 3t^4 + 1, \quad h(t) = 3t$$

46

Example: Find the derivative of  $f(g(t), h(t))$  with respect to  $t$  where

$$f(x, y) = x^2 + y \quad \text{and} \quad g(t) = 3t^4 + 1, \quad h(t) = 3t \quad [\text{Extra}]$$

47

## The New Chain Rule – Case 2

### 2 The Chain Rule (Case 2)

Suppose that  $z = f(x, y)$  is a differentiable function of  $x$  and  $y$ , where  $x = g(s, t)$  and  $y = h(s, t)$  are differentiable functions of  $s$  and  $t$ . Then

$$\frac{\partial z}{\partial s} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial s} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial s} \quad \frac{\partial z}{\partial t} = \frac{\partial z}{\partial x} \frac{\partial x}{\partial t} + \frac{\partial z}{\partial y} \frac{\partial y}{\partial t}$$

48



Example: Find the derivatives of  $f(g(s,t),h(s,t))$  with respect to  $s$  and  $t$  where

$$f(x,y) = x^2y^3 \quad \text{and} \quad g(s,t) = s \cdot \cos(t) , \quad h(s,t) = s \cdot e^{2t}$$

49

Example: Find the derivatives of  $f(g(s,t),h(s,t))$  with respect to  $s$  and  $t$  where

$$f(x,y) = x^2y^3 \quad \text{and} \quad g(s,t) = s \cdot \cos(t) , \quad h(s,t) = s \cdot e^{2t}$$

[Extra]

50


# Questions?



51

# Thank you

Until next time.



52