Chapter 10: Photosynthesis

Comprehensive Enhanced Master Study Guide • Bioenergetics, Photochemistry & Carbon Metabolism

Course: BSC2010C General Biology I **Institution:** University of Central Florida **Format:** Modular Master Syllabus

Module 1: Photosynthesis Overview, Autotrophs vs. Heterotrophs & Redox Dynamics

Trophic Classifications: Feeding the Biosphere

Earth's biosphere is solar-powered. Metabolic strategies are divided into two fundamental trophic modes:

- **Autotrophs ("Self-feeders"):** Organisms that synthesize complex organic molecules from inorganic raw materials (CO_2 and H_2O) without ingesting other living beings. They are the ultimate *producers* of the biosphere.
- **Photoautotrophs:** Use photon energy (sunlight) to drive organic synthesis. Includes terrestrial plants, multicellular algae, unicellular protists (e.g., *Euglena*), cyanobacteria, and purple sulfur bacteria.
- **Heterotrophs ("Other-feeders"):** Organisms unable to fix carbon; they survive by ingesting organic compounds synthesized by autotrophs (consumers, predators, and decomposers).



Figure 10.1: Solar Energy Flow Across Trophic Levels. *Sunlight powers photoautotrophic primary production (grass/plants), which supports primary consumers (herbivores), apex carnivores, and decomposers.*

Thermodynamic Profile & The Redox Nature of Photosynthesis

Photosynthesis is fundamentally a massive endergonic redox process that reverses the direction of electron flow seen in cellular respiration:




Figure 10.2: Stoichiometric Redox Tracking of Photosynthesis. *H_2O is oxidized to O_2, while CO_2 is reduced to carbohydrate ($C_6H_{12}O_6$).*

- **Thermodynamics:** Anabolic and strongly endergonic (ΔG = +686 kcal/mol; +2,870 kJ/mol). Requires external energy input provided by photons of light.
- **Water Oxidation:** Water is split; electrons and protons (H^+) are extracted, and O_2 is released as a byproduct.
- **Carbon Reduction:** Carbon dioxide is reduced as electrons and protons are added to form high-energy carbon bonds in triose sugars (G3P).

The Two Interdependent Stages of Photosynthesis

1. The Light Reactions (Thylakoids): Photo-actuated conversion of solar energy into chemical energy (ATP via photophosphorylation and NADPH via terminal electron transport). Releases O_2 from water photolysis.

2. The Calvin Cycle (Stroma): "Dark" or light-independent carbon fixation reactions that consume ATP and NADPH to reduce CO_2 into G3P sugar building blocks.

Module 2: Chloroplast Ultrastructure, Evolutionary Origin & Compartmentalization

Endosymbiosis & Subcellular Morphology

Chloroplasts originated via primary endosymbiosis when a eukaryotic ancestor engulfed an ancestral photosynthetic cyanobacterium. Chloroplasts maintain their own circular chromosome, prokaryotic 70S ribosomes, and divide by binary fission.

In terrestrial plants, chloroplasts are primarily concentrated in the **mesophyll** tissue of leaves (typically 30–40 chloroplasts per mesophyll cell). Microscopic pores called **stomata** (singular: stoma) regulate gas exchange (CO_2 entry and O_2 egress) and transpirational water loss.




Figure 10.3: Chloroplast Anatomy & Compartmental Architecture.
Double outer envelope, fluid-filled stroma, thylakoid membrane discs organized into granum stacks, and the internal thylakoid lumen.

Detailed Compartmental Division & Functions

Compartment	Biochemical Specialization & Functional Role
Outer & Inner Envelope	Selectively permeable double membrane separating organelle from cytosol; lacks photosynthetic pigments; houses TOC/TIC translocons.
Intermembrane Space	Narrow aqueous lumen (10–20 nm) between outer and inner membranes.
Stroma	Dense, gel-like fluid interior surrounding thylakoids; analogous to mitochondrial matrix. Houses circular DNA, 70S ribosomes, and soluble Calvin cycle enzymes (RuBisCO).
Thylakoid Membrane	Third internal lipid bilayer system; folded into interconnected flattened sacs. Houses photosystems (PS II & I), electron transport complexes, and ATP synthase.
Granum (pl. Grana)	Columns of tightly disc-stacked thylakoids; dramatically amplifies membrane surface area for light absorption.
Thylakoid Lumen	Continuous internal compartment inside thylakoids; acts as the proton reservoir (pH ~5) during light-driven proton pumping.

Structural Analogy: Chloroplast vs. Mitochondrion

A crucial conceptual framework tested on examinations:

- **Stroma ↔ Mitochondrial Matrix:** Both are aqueous enzymatic compartments where soluble cycles occur (Calvin Cycle ↔ Citric Acid Cycle).
- **Thylakoid Lumen ↔ Intermembrane Space:** Both serve as high-proton concentration reservoirs ($[H^+]$ high, pH acidic) that drive chemiosmotic ATP synthesis.

Module 3: The Nature of Sunlight, Electromagnetic Spectrum & Pigment Spectroscopy

Photophysics & The Electromagnetic Spectrum

Light possesses dual wave-particle properties. It propagates as rhythmic electromagnetic disturbances while delivering energy in discrete, massless packets called **photons**.

- **Wavelength (λ):** The spatial distance between consecutive wave crests. Measured in nanometers (nm) for visible light.
- **Inverse Energy Rule (Planck-Einstein):** The energy (E) of a photon is inversely proportional to its wavelength ($E = hc / \lambda$). Shorter wavelengths carry higher photon energy; longer wavelengths carry lower energy.
- **Visible Light Spectrum:** Narrow band between **380 nm and 750 nm** (ROYGBIV) that drives photosynthetic chemistry.




Figure 10.4: Wavelength of Electromagnetic Radiation. *Spatial distance from peak to peak; determines photon energy content.*

Photosynthetic Pigments & Light Absorption

Substances that absorb visible photons are called **pigments**. Wavelengths not absorbed are either transmitted or reflected:

- **Chlorophyll a:** Primary photosynthetic pigment; absorbs violet-blue (430 nm) and red (660 nm); reflects blue-green.
- **Chlorophyll b:** Accessory pigment; absorbs blue (450 nm) and red-orange (640 nm); broadens action spectrum; reflects olive-green.
- **Carotenoids:** Accessory hydrocarbons; absorb violet/blue-green (400–500 nm); reflect yellow/orange. Critical for **photoprotection** (dissipating damaging excess light).




Figure 10.5: Pigment Absorption Spectra. *Chlorophyll a (dark green), Chlorophyll b (light green), and Carotenoids (orange) absorb blue and red light while reflecting green wavelengths (500–550 nm).*

Critical Insight: Why Are Leaves Green?

Leaves appear green not because chlorophyll absorbs green light, but precisely because **chlorophyll reflects and transmits green wavelengths** (500–550 nm), while absorbing high-energy blue and red photons to power photosynthesis.

Absorption Spectrum vs. Action Spectrum

Absorption Spectrum: Graph depicting a specific pigment's light absorption versus wavelength (measured via spectrophotometer).

Action Spectrum: Graph plotting the biological effectiveness of different wavelengths in driving photosynthesis across the entire plant (oxygen output / CO_2 uptake).

Engelmann's Classic Experiment (1883): Theodor Engelmann illuminated a filamentous green alga with a prism-split spectrum and added aerobic bacteria. Bacteria clustered around blue and red illuminated zones where O_2 production was greatest.

Module 4: The Light Reactions • Architecture of Photosystems II and I

Structural Anatomy of a Photosystem

A photosystem is a multi-protein supramolecular machine embedded in the thylakoid lipid bilayer consisting of two distinct functional zones:

- **Light-Harvesting Complex (Antenna Complex):** Arrays of several hundred pigment molecules (chlorophyll a, chlorophyll b, carotenoids) bound to proteins. Functions as an antenna: absorbs a photon and transfers the excitation energy via *inductive resonance* from pigment to pigment until reaching the reaction center.
- **Reaction-Center Complex:** Centrally positioned protein complex housing a specialized pair of chlorophyll a molecules and a **primary electron acceptor**.
- **The Primary Photochemical Event:** The special pair's electrons are boosted to an excited state and captured by the primary electron acceptor via redox transfer, converting radiant energy into chemical energy.

Photosystem II (PS II • P680)

Functions first in linear electron flow. Named PS II because it was discovered second.

- **Reaction-Center Pigment: P680** (absorbs maximally at 680 nm, red light).
- **$P680^+$ Oxidizing Power:** Upon donating an electron to the primary acceptor (pheophytin), P680 becomes $P680^+$, the strongest biological oxidizing agent known (redox potential ≈ +1.25 V).
- **Water Photolysis:** $P680^+$ has such extreme electron affinity that it extracts electrons from water via the manganese-calcium oxygen-evolving complex (Mn_4CaO_5 cluster):

$$2\,H_2O \rightarrow 4\,H^+ + 4\,e^- + O_2$$

- Protons are released directly into the thylakoid lumen; molecular O_2 diffuses out of the chloroplast and leaf.

Photosystem I (PS I • P700)

Functions second in linear electron transport. Discovered first.

- **Reaction-Center Pigment: P700** (absorbs maximally at 700 nm, far-red light).
- **Photoexcitation:** Light absorbed by PS I antenna complexes excites P700 to $P700^*$, which rapidly transfers an electron to PS I's primary electron acceptor (an iron-sulfur center).
- **Filling the Electron Hole:** $P700^+$ is reduced back to P700 by an electron arriving from Photosystem II via the mobile copper protein **plastocyanin (Pc)**.
- **$NADP^+$ Reduction:** High-energy electrons from PS I's primary acceptor pass to **ferredoxin (Fd)**, then to the stromal enzyme **$NADP^+$ reductase**, which catalyzes:

$$NADP^+ + 2\,e^- + H^+ \rightarrow NADPH$$




Figure 10.6: Photosystem Architecture. *Light-harvesting antenna complex transfers excitation energy to reaction-center chlorophyll a (P680/P700).*

Summary: Comparison of PS II and PS I

Feature	Photosystem II (PS II)	Photosystem I (PS I)
Reaction Center	P680 (680 nm peak)	P700 (700 nm peak)
Primary e^- Source	Water photolysis (H_2O)	Plastocyanin (Pc) from PS II
Direct Output	O_2 released + H^+ gradient	Reduced NADPH in stroma
Location	Stacked grana thylakoids	Unstacked stroma thylakoids

Module 5: Linear vs. Cyclic Electron Flow & Photophosphorylation

Linear Electron Flow (The Z-Scheme)

The predominant pathway of electron transport; proceeds in an exergonic downhill series through an organized electron transport chain:

1. **Photon Excitation:** Light excites PS II antenna; energy reaches P680; electron transferred to primary acceptor pheophytin.
2. **Water Splitting:** Enzyme splits 2 H_2O into 4 H^+, 4 e^-, and O_2. Electrons replenish $P680^+$; protons accumulate in lumen.
3. **Intermediate ETC:** Electrons pass from primary acceptor to **Plastoquinone (Pq) → Cytochrome b_6f Complex → Plastocyanin (Pc)**.
4. **Proton Pumping:** As electrons traverse the Cytochrome b_6f complex, protons are actively pumped from the stroma into the thylakoid lumen, amplifying the proton gradient.
5. **PS I Excitation:** Photons excite P700; electron transferred to primary acceptor; hole filled by e^- from Pc.
6. **NADPH Synthesis:** Electrons pass from PS I to **Ferredoxin (Fd)**; $NADP^+$ reductase transfers 2 e^- and 1 H^+ to $NADP^+$ to make NADPH in the stroma.




Figure 10.7: Thylakoid Electron Transport & Chemiosmosis. *PS II photolysis, intermediate ETC, PS I, and ATP Synthase photophosphorylation.*

Cyclic Electron Flow • Mechanism & Purpose

Under specific physiological conditions, electrons cycle through an alternate loop involving **only PS I**:

- Electrons pass from PS I → Ferredoxin (Fd) → Cytochrome b_6f complex → Plastocyanin (Pc) → back to P700.
- **Consequences:** Protons continue to be pumped into the lumen, generating ATP via chemiosmosis, but **NO NADPH is produced and NO O_2 is evolved**.
- **Why It Exists:** The Calvin cycle consumes 9 ATP for every 6 NADPH (a 3:2 ratio). Linear flow generates roughly equal ATP and NADPH; cyclic flow supplies the additional ATP deficit to prevent carbon fixation from stalling, while protecting against photo-oxidative stress.

Photophosphorylation & Chemiosmotic Coupling

ATP synthesis in chloroplasts operates by chemiosmosis, identical in principle to mitochondria but with distinct spatial geometry:

- **Source of Proton Gradient:** Three contributing mechanisms:
  1. Water photolysis releases H^+ directly into the thylakoid lumen.
  2. Cytochrome b_6f pumps H^+ from stroma across membrane into lumen.
  3. Reduction of $NADP^+$ to NADPH in the stroma consumes stromal H^+, lowering $[H^+]$ outside.
- **Magnitude of Gradient:** The lumen reaches pH ≈ 5 while stroma stays at pH ≈ 8. This represents a **1,000-fold difference in proton concentration** ($\Delta pH = 3$)!
- **ATP Synthase (CF_0CF_1):** Protons flow exergonically down their electrochemical gradient from the thylakoid lumen back into the stroma through ATP synthase, powering rotational catalysis to produce ATP in the stroma.

Topological Orientation Comparison

Parameter	Mitochondria	Chloroplasts
H^+ Reservoir	Intermembrane Space	Thylakoid Lumen
Destination of H^+	Mitochondrial Matrix	Chloroplast Stroma
Site of ATP Synthesis	Matrix (for cellular work)	Stroma (for Calvin cycle)
Energy Source	Food (chemical bond E)	Sunlight (photon radiant E)

Module 6: The Calvin Cycle • Enzymatic Carbon Fixation, Reduction & Regeneration

Overview: Stroma-Localized Carbon Assimilation

The Calvin cycle is an anabolic metabolic pathway that uses the chemical energy of ATP and the reducing power of NADPH to convert gaseous inorganic CO_2 into high-energy triose phosphate sugars.

The true direct carbohydrate product of the Calvin cycle is not glucose, but the 3-carbon sugar **Glyceraldehyde 3-phosphate (G3P)**.

Phase 1: Carbon Fixation (RuBisCO)

Carbon fixation is catalyzed by the enzyme **RuBisCO** (Ribulose-1,5-bisphosphate carboxylase-oxygenase), the most abundant enzymatic protein on Earth:

- One molecule of CO_2 is attached to a 5-carbon sugar receptor, **Ribulose 1,5-bisphosphate (RuBP)**.
- Forms an extremely unstable 6-carbon intermediate that immediately hydrolyzes into two 3-carbon molecules of **3-phosphoglycerate (3-PGA)**.
- For 3 turns of the cycle (fixing 3 CO_2 onto 3 RuBP), a total of **six molecules of 3-PGA** are formed.

Phase 2: Reduction Phase

Converts low-energy carboxyl groups in 3-PGA into high-energy aldehyde carbonyl groups in G3P:

- Each 3-PGA is phosphorylated by **ATP** to yield 1,3-bisphosphoglycerate (consumes 6 ATP per 3 CO_2).
- Each 1,3-bisphosphoglycerate is reduced by a pair of electrons from **NADPH**, releasing an inorganic phosphate (P_i) to yield **G3P** (consumes 6 NADPH per 3 CO_2).
- Net result of 3 turns: **6 molecules of G3P** are produced. *Exactly 1 G3P exits the cycle* as net product; the other 5 G3P remain inside the cycle.

Phase 3: Regeneration of the CO_2 Acceptor (RuBP)

The remaining five molecules of G3P (15 carbons total) undergo an intricate series of enzymatic rearrangements to regenerate three molecules of RuBP (15 carbons total):

- This phase consumes **3 additional molecules of ATP**.
- RuBP is regenerated with high-energy phosphate bonds intact, ready to accept another turn of CO_2.




Figure 10.8: Calvin Cycle 3-Stage Biochemical Wheel. *Carbon fixation (RuBisCO), 3-PGA reduction to G3P, and RuBP regeneration.*

Complete Stoichiometry for 1 Net G3P Molecule

For the net synthesis of **one G3P molecule** (requiring 3 turns of the Calvin cycle):

$3\ CO_2 + 9\ ATP + 6\ NADPH \rightarrow 1\ G3P + 9\ ADP + 8\ P_i + 6\ NADP^+$

- **Fixation:** 3 CO_2 + 3 RuBP → 6 3-PGA
- **Reduction:** 6 3-PGA + 6 ATP + 6 NADPH → 6 G3P (1 Net G3P exits)
- **Regeneration:** 5 G3P + 3 ATP → 3 RuBP

Glucose Synthesis Accounting

Because glucose is a 6-carbon hexose ($C_6H_{12}O_6$), synthesizing **one molecule of glucose** requires **two G3P molecules**, representing **6 turns of the Calvin cycle**:

$6\ CO_2 + 18\ ATP + 12\ NADPH \rightarrow 1\ Glucose + 18\ ADP + 12\ NADP^+$

The exported G3P is converted in the cytosol into sucrose for transport throughout the plant, or stored in the stroma as starch.

Module 7: Photorespiration & Ecological Adaptations • C3, C4 and CAM Pathways

Photorespiration: The RuBisCO Evolutionary Dilemma

RuBisCO evolved when Earth's atmosphere had abundant CO_2 and virtually no O_2. Consequently, its active site cannot completely discriminate between CO_2 and O_2:

- **Stomatal Closure:** On hot, dry days, plants close stomata to conserve water. Transpiration halts, internal [CO_2] plummets as it is consumed, and internal [O_2] builds up from light reactions.
- **Oxygenase Activity:** RuBisCO binds O_2 instead of CO_2:

  RuBP (5C) + O_2 → 3-PGA (3C) + 2-Phosphoglycolate (2C)
- **Metabolic Cost:** 2-Phosphoglycolate is toxic and must be salvaged by a complex pathway across chloroplasts, peroxisomes, and mitochondria. It **consumes ATP and releases CO_2 without producing any sugar or ATP**!
- Can drain photosynthetic productivity in C3 crops (rice, wheat, soybeans) by up to **25–50%**.

C4 Plants: Spatial Separation of Fixation

Plants in intense sunlight and arid conditions (corn, sugarcane, crabgrass) possess anatomical and biochemical adaptations:

- **Kranz Anatomy:** Two distinct photosynthetic cell layers: outer **Mesophyll cells** and inner **Bundle-sheath cells** tightly packed around leaf veins.
- **PEP Carboxylase:** Mesophyll cells use PEP carboxylase to fix CO_2 onto PEP (3C) forming oxaloacetate (4C). PEP carboxylase has exceptionally high affinity for CO_2 and **zero affinity for O_2**.
- **Decarboxylation:** The 4C intermediate (malate) is exported via plasmodesmata to bundle-sheath cells, where it is decarboxylated to release concentrated CO_2 directly to RuBisCO, completely suppressing photorespiration.

CAM Plants: Temporal Separation of Fixation

Crassulacean Acid Metabolism (CAM) evolved in desert succulents, cacti, pineapples, and agaves to maximize water-use efficiency:

- **Night (Stomata Open):** Temperatures are cool and humidity is high. Stomata open to take up CO_2. PEP carboxylase fixes CO_2 into organic acids (principally malic acid), which are sequestered in huge central vacuoles.
- **Day (Stomata Closed):** Stomata remain sealed all day to eliminate water loss. Light reactions run, producing ATP and NADPH. Sequestered malic acid is transported out of vacuoles and decarboxylated, releasing high concentrations of CO_2 to RuBisCO in the stroma to drive the Calvin cycle.

Comparative Ecological Matrix: C3 vs. C4 vs. CAM

Feature	C3 Plants	C4 Plants	CAM Plants
Initial CO_2 Fixation	RuBisCO directly (Day)	PEP Carboxylase (Day)	PEP Carboxylase (Night)
First Product	3-PGA (3-carbon)	Oxaloacetate (4-carbon)	Malic acid (4-carbon)
Mode of Separation	None	**Spatial** (Mesophyll → Bundle-sheath)	**Temporal** (Night → Day)
Stomata Schedule	Open during day	Open during day	Open during night
Photorespiration	High in hot/dry environments	Virtually suppressed	Virtually suppressed
Typical Examples	Wheat, rice, oats, trees	Corn, sugarcane, crabgrass	Pineapple, cacti, succulents

Module 8: Comparative Bioenergetics • Respiration vs. Photosynthesis Matrix

Comparative Bioenergetic Matrix: Respiration vs. Photosynthesis

Criteria	Photosynthesis (Chloroplast)	Cellular Respiration (Mitochondrion)
Overall Chemical Nature	Anabolic (endergonic, builds complex carbohydrates from simple inorganic CO_2). ΔG = +686 kcal/mol.	Catabolic (exergonic, oxidative breakdown of organic fuels to release free energy). ΔG = -686 kcal/mol.
Redox Equation	$6\ CO_2 + 6\ H_2O + \text{Light} \rightarrow C_6H_{12}O_6 + 6\ O_2$ **CO_2 is reduced; H_2O is oxidized.**	$C_6H_{12}O_6 + 6\ O_2 \rightarrow 6\ CO_2 + 6\ H_2O + \text{Energy}$ **Glucose is oxidized; O_2 is reduced.**
Initial Electron Source	Water photolysis (H_2O split at Photosystem II).	Organic fuel molecules (Glucose, Pyruvate, Acetyl-CoA).
Final Electron Acceptor	**$NADP^+$** (reduced to NADPH in stroma).	**Molecular Oxygen (O_2)** (reduced to H_2O in matrix).
High-Energy Electron Carrier	**NADPH** (contains extra 2'-phosphate group; used exclusively for anabolic biosyntheses).	**NADH & $FADH_2$** (used for catabolic energy harvesting to drive ATP generation).
Primary Energy Driving Force	Radiant energy from absorbed photons of light.	Chemical potential energy stored in covalent C-C and C-H bonds.
Proton Reservoir Location	**Thylakoid Lumen / Space** ($[H^+]$ high, pH ≈ 5).	**Mitochondrial Intermembrane Space** ($[H^+]$ high, pH ≈ 7).
Direction of Proton Pumping	Pumps H^+ from Stroma into Thylakoid Lumen.	Pumps H^+ from Matrix into Intermembrane Space.
Direction of H^+ Chemiosmosis	Protons flow from Thylakoid Lumen → Stroma.	Protons flow from Intermembrane Space → Matrix.
Site of ATP Generation	Chloroplast Stroma (consumed by Calvin cycle).	Mitochondrial Matrix (exported for cellular work).
Major Cyclic Pathway	**Calvin Cycle:** Fixes CO_2; consumes 9 ATP & 6 NADPH per G3P; strictly anabolic.	**Citric Acid Cycle:** Oxidizes Acetyl-CoA; releases CO_2; produces NADH, $FADH_2$, ATP; catabolic.
Organismal Occurrence	Restricted to photoautotrophs (plants, algae, cyanobacteria).	All eukaryotic organisms (including plants day and night!).

Plant Bioenergetics: Day vs. Night Dynamics

Plants have both chloroplasts and mitochondria:

- **During Daylight:** Photosynthesis runs at high velocity, producing carbohydrates and O_2. A portion of synthesized sugar is immediately consumed by mitochondrial respiration; the vast remainder is stored.
- **During Darkness:** Photosynthesis stops. Plant cells rely entirely on mitochondrial cellular respiration, consuming stored sugars and O_2 to produce cellular ATP.

The Global Biospheric Carbon-Oxygen Cycle


Photosynthesis and respiration form a continuous complementary thermodynamic loop:

- The outputs of photosynthesis (O_2 + organic sugars) serve as the vital inputs for cellular respiration.
- The outputs of cellular respiration (CO_2 + H_2O) serve as the vital chemical raw materials for photosynthesis.

Module 9: Course Active Learning Practice Problems • Part I (Chloroplast Concept Map)

Chloroplast Concept Map & Energy Flow (Active Learning #3, Slide 8)

The course curriculum emphasizes the complete integration of thylakoid light reactions and stroma carbon fixation. Trace the inputs, intermediate complexes, and end products:



Component / Step	Specific Subcellular Site	Biochemical Function & In-Class Rationale
Photosystem II (PS II)	Thylakoid Membrane	Absorbs light; excites P680; extracts electrons from water via photolysis ($2\ H_2O \rightarrow 4\ H^+ + 4\ e^- + O_2$). O_2 released.
Cytochrome Complex	Thylakoid Membrane	Electron conduit between PS II and PS I; actively couples electron flow to proton pumping from the stroma into the thylakoid space.
Photosystem I (PS I)	Thylakoid Membrane	Re-excites electrons via P700 using absorbed light; transfers electrons through ferredoxin to $NADP^+$ reductase to produce NADPH in stroma.
ATP Synthase	Thylakoid Membrane	Uses the chemiosmotic proton gradient (lumen → stroma) to drive photophosphorylation, producing ATP in the stroma.
Calvin Cycle	Chloroplast Stroma	Light-independent carbon fixation: RuBisCO fixes atmospheric CO_2; consumes ATP and NADPH to produce G3P sugar building blocks.

Key Conceptual Traps from Active Learning #3

- **Trap 1: "Light reactions occur in the stroma."** False. The light reactions (PS II, ETC, PS I, ATP Synthase) are membrane-bound in the *thylakoids*. Only the Calvin cycle occurs in the *stroma*.
- **Trap 2: "Oxygen comes from carbon dioxide."** False. Oxygen produced during photosynthesis originates entirely from **water photolysis at PS II**, not from CO_2!
- **Trap 3: "ATP generated by chloroplast ATP synthase powers the whole plant cell."** False. Chloroplast ATP is synthesized directly into the stroma and is **almost entirely consumed on-site by the Calvin cycle**. Cellular work across the plant is powered by ATP produced in mitochondria!

Module 9: Course Active Learning Practice Problems • Part II (Herbicides Inhibition H-1 to H-5)

Herbicides Inhibition Diagnostic Analysis (Active Learning #3, Slide 9)

Chloroplasts are exposed to five distinct experimental herbicide compounds (H-1 through H-5). Physiological readouts are measured against untreated controls. Deducing the molecular targets represents a core examination competency:

Agent	Chlorophyll Absorption	O_2 Output	NADPH Output	ATP Synthase	G3P Output	Molecular Target & Step-by-Step Diagnostic Deduction
H-1	Normal	Normal	**Decreased**	Normal	**Decreased**	**Photosystem I (PS I):** PS II splits water normally (O_2 normal) and proton pumping drives ATP synthesis normally. However, electrons cannot reduce $NADP^+$ to NADPH. Deprived of reducing power, the Calvin cycle stalls (G3P decreases).
H-2	**Decreased**	**Decreased**	**Decreased**	**Decreased**	**Decreased**	**Chlorophyll Absorption:** Light harvesting itself is impaired. Without photon capture, no electrons are excited, water is not split, no protons are pumped, and all downstream light and dark processes collapse.
H-3	Normal	**Decreased**	**Decreased**	**Decreased**	**Decreased**	**Photosystem II (PS II):** PS II water oxidation is halted, dropping O_2. Without the source of electrons from PS II, linear electron transport stops, collapsing the proton gradient, ATP, NADPH, and G3P.
H-4	Normal	Normal	Normal	Normal	**Decreased**	**Calvin Cycle (e.g., RuBisCO):** All thylakoid light reactions run normally (absorption, O_2 evolution, ATP, and NADPH intact), but carbon fixation is selectively blocked, dropping G3P output.
H-5	Normal	Normal	Normal	**Decreased**	**Decreased**	**ATP Synthase:** Light absorption, water splitting, and NADPH synthesis proceed unaffected, but photophosphorylation is blocked. Deprived of ATP, the endergonic reduction and regeneration phases of the Calvin cycle stall, lowering G3P.

Real-World Agricultural & Chemical Correlates

- **DCMU (Diuron):** Competitively blocks the plastoquinone-binding site of Photosystem II ↔ *Matches H-3 profile* (halts O_2 evolution and linear electron transport).
- **Paraquat (Methyl Viologen):** Diverts electrons away from Photosystem I / ferredoxin ↔ *Matches H-1 profile* (depletes NADPH synthesis).
- **Atrazine:** Widely used herbicide inhibiting electron transfer within PS II ↔ *Matches H-3 profile*.

Module 10: Foundational Practice Examination • Part I (Questions 1 – 7)

1. Which of the following statements correctly characterizes the bioenergetics of photosynthesis compared to cellular respiration?

A) Photosynthesis is catabolic and exergonic ($\Delta G < 0$), oxidizing glucose to generate ATP.
B) Photosynthesis is anabolic and endergonic ($\Delta G > 0$), reducing CO_2 to carbohydrate using photon energy.
C) Photosynthesis reduces H_2O and oxidizes CO_2 in an endergonic pathway.
D) Both pathways utilize identical electron carriers (NADH and $FADH_2$) to synthesize carbohydrates.

2. During the light reactions, where do protons (H^+) accumulate to form the electrochemical gradient that drives ATP synthesis?

A) Within the stroma of the chloroplast.
B) Inside the thylakoid lumen (thylakoid space).
C) In the intermembrane space between the inner and outer chloroplast envelopes.
D) Inside the cytosol surrounding the chloroplast.

3. Why do plant leaves typically appear green to the human eye?

A) Chlorophyll pigments absorb green wavelengths (500–550 nm) more effectively than any other wavelength.
B) Chlorophyll pigments reflect and transmit green light while absorbing blue and red wavelengths.
C) Carotenoids emit green photons via fluorescence during normal photon capture.
D) The cuticle on the leaf surface selectively refracts green photons while absorbing infrared light.

4. What is the immediate source of electrons that replaces the photoexcited electrons ejected from the P680 special pair in Photosystem II?

A) NADPH oxidation via ferredoxin in the stroma.
B) Photolysis of water molecules ($H_2O \rightarrow 2\ H^+ + 2\ e^- + 1/2\ O_2$) at the manganese center.
C) Plastocyanin donating electrons from Photosystem I.
D) Carbon dioxide reduction in the Calvin cycle.

5. What is the correct sequence of electron transfer during linear electron flow in the thylakoid membrane?

A) PS I → Ferredoxin → Cytochrome complex → Plastoquinone → PS II → $NADP^+$.
B) PS II → Plastoquinone (Pq) → Cytochrome b_6f complex → Plastocyanin (Pc) → PS I → Ferredoxin (Fd) → $NADP^+$.
C) PS II → Cytochrome complex → PS I → Plastoquinone → Plastocyanin → ATP Synthase.
D) H_2O → PS I → Plastocyanin → Cytochrome complex → PS II → NADPH.

6. In what fundamental way does cyclic electron flow differ from linear electron flow?

A) Cyclic flow involves both PS II and PS I, generating both ATP and NADPH.
B) Cyclic flow involves only PS I, pumps protons to produce ATP, but does NOT produce NADPH or O_2.
C) Cyclic flow involves only PS II, splitting water without producing ATP.
D) Cyclic flow operates exclusively in the dark to fix CO_2 into G3P.

7. In chloroplast chemiosmosis, what is the orientation of proton flow through ATP synthase, and where is ATP released?

A) Protons flow from the stroma into the thylakoid lumen; ATP is released into the lumen.
B) Protons flow from the thylakoid lumen into the stroma; ATP is released into the stroma.
C) Protons flow from the intermembrane space into the matrix; ATP is released into the matrix.
D) Protons flow from the cytosol into the intermembrane space; ATP is released into the cytosol.

Module 10: Foundational Practice Examination • Part II (Questions 8 – 15)

8. What is the specific enzymatic reaction catalyzed by RuBisCO in the first step of the Calvin cycle?

A) Fixation of CO_2 onto ribulose 1,5-bisphosphate (RuBP) to yield 3-phosphoglycerate (3-PGA).

B) Oxidation of G3P to form 1,3-bisphosphoglycerate using $NADP^+$.

C) Phosphorylation of ribulose 5-phosphate using ATP to regenerate RuBP.

D) Condensation of two G3P molecules to synthesize glucose 6-phosphate.

9. How many turns of the Calvin cycle, molecules of CO_2, ATP, and NADPH are required to produce ONE net molecule of G3P?

A) 1 turn, 1 CO_2, 3 ATP, 2 NADPH.

B) 3 turns, 3 CO_2, 9 ATP, 6 NADPH.

C) 6 turns, 6 CO_2, 18 ATP, 12 NADPH.

D) 2 turns, 2 CO_2, 6 ATP, 4 NADPH.

10. Under what environmental conditions is photorespiration most pronounced, and why is it considered wasteful?

A) Cold, wet conditions; it converts excess glucose into glycogen while generating ATP.

B) Hot, dry conditions with closed stomata; RuBisCO binds O_2 instead of CO_2, consuming ATP without producing sugar.

C) Dim light at night; it causes chloroplasts to split water without photon excitation.

D) High atmospheric CO_2; it causes PEP carboxylase to overproduce malate.

11. How do C4 plants functionally prevent photorespiration in hot environments?

A) By temporally opening stomata only at night to fix CO_2 into vacuolar malic acid.

B) By spatial separation: fixing CO_2 into a 4C acid in mesophyll cells via PEP carboxylase, then delivering CO_2 to bundle-sheath cells.

C) By mutating RuBisCO so that it has zero affinity for oxygen.

D) By shutting down the Calvin cycle and relying exclusively on cyclic photophosphorylation.

12. Crassulacean Acid Metabolism (CAM) in desert succulents minimizes transpirational water loss by:

A) Opening stomata at night to fix CO_2 into organic acids, then closing stomata by day to run the Calvin cycle.

B) Conducting the light reactions at night using moonlight.

C) Eliminating the Calvin cycle entirely and storing starch in mesophyll cell walls.

D) Exporting RuBisCO into root epidermal cells.

13. A newly synthesized herbicide inhibits Photosystem I. When applied to isolated chloroplasts, what physiological profile is observed?

A) O_2 evolution stops, ATP drops, and NADPH drops.

B) O_2 evolution and ATP production remain normal, while NADPH and G3P outputs are severely decreased.

C) All light-dependent outputs are normal, but G3P synthesis is doubled.

D) Chlorophyll absorption drops to zero immediately.

14. If a chemical uncoupler makes the thylakoid membrane permeable to protons, what will be the effect on chloroplast function?

A) Electron transport stops immediately, halting O_2 and NADPH generation.

B) Protons leak across the membrane, collapsing the ΔpH gradient and abolishing ATP synthesis, while electron flow and O_2 evolution continue.

C) ATP synthesis accelerates dramatically due to rapid proton movement.

D) RuBisCO activity increases five-fold to compensate.

15. If a plant is kept in total darkness, why does the Calvin cycle eventually cease running even if abundant CO_2 is present?

A) CO_2 cannot diffuse across the leaf cuticle without light.

B) The Calvin cycle requires ATP and NADPH produced by the light reactions, which are depleted in the dark.

C) RuBisCO is degraded by proteasomes within 2 seconds of darkness.

D) Water molecules cannot enter the roots without direct sunlight.

Module 10: Comprehensive Examination Answer Key & Explanations

Q1: B • Photosynthesis Bioenergetics
Rationale: Photosynthesis is anabolic and endergonic (ΔG = +686 kcal/mol), utilizing photon energy to reduce CO_2 into carbohydrate. Cellular respiration is catabolic and exergonic.

Q2: B • Thylakoid Lumen Proton Reservoir
Rationale: Water splitting and Cytochrome b_6f pumping accumulate protons inside the thylakoid lumen (pH ≈ 5), establishing the chemiosmotic gradient across the thylakoid membrane into the stroma (pH ≈ 8).

Q3: B • Reflection of Green Wavelengths
Rationale: Pigments appear as the color of light they transmit or reflect. Chlorophyll a and b absorb violet-blue and red light while reflecting green light (500–550 nm).

Q4: B • Water Photolysis Replenishes $P680^+$
Rationale: $P680^+$ is an extraordinarily strong oxidizing agent that extracts electrons from water via the oxygen-evolving complex, releasing 4 H^+ and O_2.

Q5: B • Linear Electron Flow Pathway
Rationale: Electrons travel downhill: PS II → Pq → Cytochrome b_6f → Pc → PS I → Ferredoxin → $NADP^+$ reductase → NADPH.

Q6: B • Cyclic Electron Flow Characteristics
Rationale: Cyclic flow utilizes only PS I, ferredoxin, and the cytochrome complex to pump protons and synthesize ATP; it does NOT involve PS II, water splitting, or $NADP^+$ reduction.

Q7: B • Chemiosmosis Orientation in Chloroplast
Rationale: Protons diffuse down their electrochemical gradient from the thylakoid lumen through CF_0CF_1 ATP synthase into the stroma, releasing newly formed ATP into the stroma.

Q8: A • Carbon Fixation via RuBisCO
Rationale: RuBisCO catalyzes the addition of gaseous CO_2 to the 5-carbon acceptor RuBP, producing an unstable 6C intermediate that splits into two molecules of 3-PGA.

Q9: B • Calvin Cycle Stoichiometry (1 Net G3P)
Rationale: Synthesizing 1 net G3P requires 3 turns of the cycle (3 CO_2), consuming 6 ATP and 6 NADPH during reduction, plus 3 ATP during RuBP regeneration (total: 9 ATP, 6 NADPH).

Q10: B • Photorespiration Triggers and Costs
Rationale: When stomata close in hot/dry environments, internal [CO_2] falls and [O_2] rises. RuBisCO fixes O_2 instead of CO_2, draining ATP and releasing CO_2 without sugar yield.

Q11: B • C4 Spatial Separation Mechanism
Rationale: C4 plants fix CO_2 into a 4-carbon acid in mesophyll cells via PEP carboxylase (which cannot bind O_2), and pump the acid into bundle-sheath cells to release concentrated CO_2 to RuBisCO.

Q12: A • CAM Temporal Separation Strategy
Rationale: CAM plants open stomata at night to fix CO_2 into malic acid stored in central vacuoles, keeping stomata closed during hot daylight hours while the Calvin cycle runs.

Q13: B • Photosystem I Inhibition Profile
Rationale: Active Learning #3 (Herbicide H-1): Blocking PS I leaves PS II water splitting (O_2) and Cytochrome b_6f proton pumping (ATP) intact, but specifically abolishes NADPH synthesis and G3P output.

Q14: B • Chemical Uncoupling of Photophosphorylation
Rationale: Uncouplers dissipate the proton gradient across the thylakoid membrane; electron transport and water oxidation accelerate, but ATP synthase cannot generate ATP without a proton-motive force.

Q15: B • Light-Dark Metabolic Interdependence
Rationale: Although the Calvin cycle's enzymatic reactions do not directly absorb photons, they require continuous supplies of ATP and NADPH produced by the light reactions in thylakoids.