

Solutions to Chapter 9, Susanna Epp Discrete Math

5th Edition

<https://github.com/spamegg1>

October 24, 2023

Contents

1	Exercise Set 9.1	17
1.1	Exercise 1	17
1.2	Exercise 2	17
1.3	Exercise 3	18
1.4	Exercise 4	18
1.5	Exercise 5	18
1.6	Exercise 6	18
1.7	Exercise 7	18
1.8	Exercise 8	18
1.9	Exercise 9	19
1.10	Exercise 10	19
1.11	Exercise 11	19
	1.11.1 (a)	19
	1.11.2 (b)	19
1.12	Exercise 12	19
	1.12.1 (a)	20
	1.12.2 (b)	20
1.13	Exercise 13	20
	1.13.1 (a)	20
	1.13.2 (b)	20
1.14	Exercise 14	21
	1.14.1 (a)	21
	1.14.2 (b)	21
	1.14.3 (c)	21
1.15	Exercise 15	21
1.16	Exercise 16	21
	1.16.1 (a)	21
	1.16.2 (b)	22
	1.16.3 (c)	22
1.17	Exercise 17	22

1.17.1	(a)	22
1.17.2	(b)	22
1.18	Exercise 18	23
1.18.1	(a)	23
1.18.2	(b)	23
1.18.3	(c)	23
1.19	Exercise 19	23
1.19.1	(a)	23
1.19.2	(b)	24
1.19.3	(c)	24
1.20	Exercise 20	24
1.20.1	(a)	24
1.20.2	(b)	24
1.21	Exercise 21	25
1.21.1	(a)	25
1.21.2	(b)	25
1.21.3	(c)	25
1.22	Exercise 22	25
1.22.1	(a)	25
1.22.2	(b)	25
1.22.3	(c)	26
1.23	Exercise 23	26
1.23.1	(a)	26
1.23.2	(b)	26
1.23.3	(c)	26
1.23.4	(d)	26
1.24	Exercise 24	26
1.24.1	(a)	27
1.24.2	(b)	27
1.25	Exercise 25	27
1.25.1	(a)	27
1.25.2	(b)	27
1.26	Exercise 26	27
1.27	Exercise 27	28
1.28	Exercise 28	28
1.29	Exercise 29	28
1.30	Exercise 30	28
1.31	Exercise 31	28
1.32	Exercise 32	28
1.32.1	(a)	28
1.32.2	(b)	29
1.33	Exercise 33	29

2 Exercise Set 9.2 29

2.1	Exercise 1	29
2.2	Exercise 2	30

2.3	Exercise 3	30
2.4	Exercise 4	30
2.5	Exercise 5	30
2.6	Exercise 6	31
2.6.1	(a)	31
2.6.2	(b)	31
2.6.3	(c)	31
2.6.4	(d)	31
2.7	Exercise 7	32
2.7.1	(a)	32
2.7.2	(b)	32
2.7.3	(c)	32
2.8	Exercise 8	32
2.9	Exercise 9	32
2.9.1	(a)	33
2.9.2	(b)	33
2.9.3	(c)	33
2.10	Exercise 10	33
2.10.1	(a)	33
2.10.2	(b)	33
2.11	Exercise 11	34
2.11.1	(a)	34
2.11.2	(b)	34
2.11.3	(c)	34
2.12	Exercise 12	34
2.12.1	(a)	34
2.12.2	(b)	35
2.13	Exercise 13	35
2.13.1	(a)	35
2.13.2	(b)	35
2.13.3	(c)	36
2.14	Exercise 14	36
2.14.1	(a)	36
2.14.2	(b)	36
2.14.3	(c)	36
2.14.4	(d)	36
2.14.5	(e)	37
2.15	Exercise 15	37
2.15.1	(a)	37
2.15.2	(b)	37
2.16	Exercise 16	37
2.16.1	(a)	37
2.16.2	(b)	37
2.16.3	(c)	38
2.16.4	(d)	38
2.16.5	(e)	38

2.17	Exercise 17	38
2.17.1	(a)	38
2.17.2	(b)	38
2.17.3	(c)	39
2.17.4	(d)	39
2.17.5	(e)	39
2.18	Exercise 18	39
2.18.1	(a)	39
2.18.2	(b)	40
2.18.3	(c)	40
2.19	Exercise 19	40
2.20	Exercise 20	41
2.20.1	(a)	41
2.20.2	(b)	41
2.21	Exercise 21	41
2.21.1	(a)	41
2.21.2	(b)	42
2.21.3	(c)	42
2.22	Exercise 22	42
2.22.1	(a)	42
2.22.2	(b)	42
2.22.3	(c)	42
2.23	Exercise 23	42
2.24	Exercise 24	43
2.25	Exercise 25	43
2.26	Exercise 26	43
2.27	Exercise 27	43
2.28	Exercise 28	44
2.29	Exercise 29	44
2.30	Exercise 30	44
2.30.1	(a)	44
2.30.2	(b)	45
2.31	Exercise 31	45
2.31.1	(a)	45
2.31.2	(b)	45
2.31.3	(c)	45
2.31.4	(d)	46
2.31.5	(e)	46
2.32	Exercise 32	46
2.32.1	(a)	46
2.32.2	(b)	46
2.32.3	(c)	46
2.33	Exercise 33	47
2.33.1	(a)	47
2.33.2	(b)	47
2.33.3	(c)	47

2.34	Exercise 34	47
2.35	Exercise 35	47
2.36	Exercise 36	47
2.37	Exercise 37	48
2.37.1	(a)	48
2.37.2	(b)	48
2.37.3	(c)	48
2.37.4	(d)	48
2.38	Exercise 38	48
2.38.1	(a)	48
2.38.2	(b)	48
2.39	Exercise 39	49
2.39.1	(a)	49
2.39.2	(b)	49
2.39.3	(c)	49
2.39.4	(d)	49
2.40	Exercise 40	49
2.41	Exercise 41	49
2.42	Exercise 42	50
2.43	Exercise 43	50
2.44	Exercise 44	50
2.45	Exercise 45	51
2.46	Exercise 46	51
2.47	Exercise 47	52
2.47.1	(a)	52
2.47.2	(b)	52
2.47.3	(c)	53
2.47.4	(d)	53
3	Exercise Set 9.3	53
3.1	Exercise 1	53
3.1.1	(a)	53
3.1.2	(b)	54
3.2	Exercise 2	54
3.2.1	(a)	54
3.2.2	(b)	54
3.3	Exercise 3	54
3.3.1	(a)	54
3.3.2	(b)	55
3.3.3	(c)	55
3.4	Exercise 4	55
3.5	Exercise 5	56
3.5.1	(a)	56
3.5.2	(b)	56
3.6	Exercise 6	56
3.6.1	(a)	56

3.6.2	(b)	56
3.6.3	(c)	57
3.6.4	(d)	57
3.7	Exercise 7	57
3.7.1	(a)	57
3.7.2	(b)	58
3.7.3	(c)	58
3.7.4	(d)	58
3.8	Exercise 8	58
3.8.1	(a)	58
3.8.2	(b)	59
3.8.3	(c)	59
3.8.4	(d)	59
3.9	Exercise 9	59
3.9.1	(a)	59
3.9.2	(b)	60
3.10	Exercise 10	60
3.11	Exercise 11	61
3.11.1	(a)	61
3.11.2	(b)	61
3.11.3	(c)	61
3.12	Exercise 12	61
3.12.1	(a)	61
3.12.2	(b)	62
3.13	Exercise 13	62
3.13.1	(a)	62
3.13.2	(b)	62
3.14	Exercise 14	62
3.15	Exercise 15	63
3.16	Exercise 16	63
3.16.1	(a)	63
3.16.2	(b)	63
3.16.3	(c)	63
3.17	Exercise 17	64
3.17.1	(a)	64
3.17.2	(b)	64
3.17.3	(c)	64
3.18	Exercise 18	64
3.18.1	(a)	64
3.18.2	(b)	64
3.19	Exercise 19	65
3.20	Exercise 20	65
3.20.1	(a)	65
3.20.2	(b)	65
3.20.3	(c)	66
3.21	Exercise 21	66

3.22	Exercise 22	66
3.22.1	(a)	66
3.22.2	(b)	67
3.23	Exercise 23	67
3.23.1	(a)	67
3.23.2	(b)	67
3.23.3	(c)	68
3.24	Exercise 24	68
3.24.1	(a)	68
3.24.2	(b)	68
3.24.3	(c)	68
3.25	Exercise 25	68
3.25.1	(a)	68
3.25.2	(b)	68
3.25.3	(c)	69
3.25.4	(d)	69
3.26	Exercise 26	69
3.26.1	(a)	69
3.26.2	(b)	69
3.26.3	(c)	69
3.26.4	(d)	70
3.26.5	(e)	70
3.27	Exercise 27	70
3.27.1	(a)	71
3.27.2	(b)	71
3.28	Exercise 28	71
3.28.1	(a)	72
3.28.2	(b)	72
3.29	Exercise 29	72
3.29.1	(a)	72
3.29.2	(b)	72
3.29.3	(c)	73
3.29.4	(d)	73
3.29.5	(e)	73
3.29.6	(f)	73
3.29.7	(g)	74
3.29.8	(h)	74
3.29.9	(i)	74
3.29.10	(j)	74
3.30	Exercise 30	74
3.31	Exercise 31	75
3.31.1	(a)	75
3.31.2	(b)	75
3.31.3	(c)	75
3.31.4	(d)	75
3.31.5	(e)	76

3.32	Exercise 32	76
3.33	Exercise 33	76
3.33.1	(a)	76
3.33.2	(b)	77
3.33.3	(c)	77
3.33.4	(d)	77
3.33.5	(e)	78
3.33.6	(f)	78
3.34	Exercise 34	78
3.34.1	(a)	78
3.34.2	(b)	78
3.34.3	(c)	79
3.34.4	(d)	79
3.35	Exercise 35	79
3.36	Exercise 36	80
3.37	Exercise 37	81
3.38	Exercise 38	81
3.39	Exercise 39	81
3.40	Exercise 40	82
3.41	Exercise 41	83
3.42	Exercise 42	83
3.42.1	(a)	83
3.42.2	(b)	83
3.42.3	(c)	84
3.43	Exercise 43	84
3.43.1	(a)	84
3.43.2	(b)	84
3.43.3	(c)	84
3.44	Exercise 44	85
3.45	Exercise 45	86
3.46	Exercise 46	86
3.47	Exercise 47	87
3.48	Exercise 48	88
3.49	Exercise 49	90
3.49.1	(a)	90
3.49.2	(b)	90
4	Exercise Set 9.4	91
4.1	Exercise 1	91
4.1.1	(a)	91
4.1.2	(b)	91
4.2	Exercise 2	91
4.2.1	(a)	91
4.2.2	(b)	91
4.3	Exercise 3	91
4.4	Exercise 4	91

4.5	Exercise 5	91
4.5.1	(a)	91
4.5.2	(b)	92
4.6	Exercise 6	92
4.6.1	(a)	92
4.6.2	(b)	92
4.7	Exercise 7	92
4.8	Exercise 8	92
4.9	Exercise 9	92
4.9.1	(a)	92
4.9.2	(b)	92
4.10	Exercise 10	92
4.11	Exercise 11	92
4.12	Exercise 12	92
4.13	Exercise 13	92
4.14	Exercise 14	93
4.15	Exercise 15	93
4.16	Exercise 16	93
4.17	Exercise 17	93
4.18	Exercise 18	93
4.19	Exercise 19	93
4.20	Exercise 20	93
4.20.1	(a)	93
4.20.2	(b)	93
4.21	Exercise 21	93
4.22	Exercise 22	93
4.23	Exercise 23	93
4.24	Exercise 24	94
4.25	Exercise 25	94
4.26	Exercise 26	94
4.27	Exercise 27	94
4.28	Exercise 28	94
4.29	Exercise 29	94
4.30	Exercise 30	94
4.31	Exercise 31	94
4.32	Exercise 32	94
4.33	Exercise 33	94
4.34	Exercise 34	94
4.35	Exercise 35	94
4.36	Exercise 36	95
4.37	Exercise 37	95
4.37.1	(a)	95
4.37.2	(b)	95
4.38	Exercise 38	95
4.39	Exercise 39	95
4.40	Exercise 40	95

5	Exercise Set 9.5	95
5.1	Exercise 1	95
	5.1.1 (a)	95
	5.1.2 (b)	95
5.2	Exercise 2	95
	5.2.1 (a)	95
	5.2.2 (b)	95
5.3	Exercise 3	96
5.4	Exercise 4	96
5.5	Exercise 5	96
	5.5.1 (a)	96
	5.5.2 (b)	96
	5.5.3 (c)	96
	5.5.4 (d)	96
	5.5.5 (e)	96
5.6	Exercise 6	96
	5.6.1 (a)	96
	5.6.2 (b)	96
	5.6.3 (c)	96
	5.6.4 (d)	96
	5.6.5 (e)	97
5.7	Exercise 7	97
	5.7.1 (a)	97
	5.7.2 (b)	97
	5.7.3 (c)	97
	5.7.4 (d)	97
5.8	Exercise 8	97
	5.8.1 (a)	97
	5.8.2 (b)	97
	5.8.3 (c)	97
	5.8.4 (d)	97
5.9	Exercise 9	97
	5.9.1 (a)	97
	5.9.2 (b)	97
5.10	Exercise 10	98
5.11	Exercise 11	98
	5.11.1 (a)	98
	5.11.2 (b)	98
	5.11.3 (c)	98
	5.11.4 (d)	98
	5.11.5 (e)	98
	5.11.6 (f)	98
	5.11.7 (g)	98
	5.11.8 (h)	98
	5.11.9 (i)	98
5.12	Exercise 12	98

5.13	Exercise 13	99
5.13.1	(a)	99
5.13.2	(b)	99
5.13.3	(c)	99
5.13.4	(d)	99
5.13.5	(e)	99
5.14	Exercise 14	99
5.14.1	(a)	99
5.14.2	(b)	99
5.14.3	(c)	99
5.14.4	(d)	99
5.15	Exercise 15	99
5.15.1	(a)	99
5.15.2	(b)	99
5.15.3	(c)	100
5.15.4	(d)	100
5.16	Exercise 16	100
5.16.1	(a)	100
5.16.2	(b)	100
5.16.3	(c)	100
5.17	Exercise 17	100
5.17.1	(a)	100
5.17.2	(b)	100
5.17.3	(c)	100
5.17.4	(d)	100
5.18	Exercise 18	100
5.19	Exercise 19	100
5.19.1	(a)	100
5.19.2	(b)	101
5.19.3	(c)	101
5.20	Exercise 20	101
5.20.1	(a)	101
5.20.2	(b)	101
5.20.3	(c)	101
5.21	Exercise 21	101
5.22	Exercise 22	101
5.23	Exercise 23	101
5.24	Exercise 24	101
5.24.1	(a)	101
5.24.2	(b)	101
5.24.3	(c)	101
5.24.4	(d)	102
5.25	Exercise 25	102
5.25.1	(a)	102
5.25.2	(b)	102
5.25.3	(c)	102

5.25.4	(d)	102
5.25.5	(e)	102
5.26	Exercise 26	102
5.26.1	(a)	102
5.26.2	(b)	102
5.26.3	(c)	102
5.26.4	(d)	102
5.26.5	(e)	102
5.26.6	(f)	103
5.27	Exercise 27	103
5.27.1	(a)	103
5.27.2	(b)	103
5.27.3	(c)	103
5.27.4	(d)	103
5.28	Exercise 28	103
5.29	Exercise 29	103
5.30	Exercise 30	103
6	Exercise Set 9.6	103
6.1	Exercise 1	103
6.1.1	(a)	103
6.1.2	(b)	103
6.2	Exercise 2	104
6.2.1	(a)	104
6.2.2	(b)	104
6.3	Exercise 3	104
6.3.1	(a)	104
6.3.2	(b)	104
6.3.3	(c)	104
6.4	Exercise 4	104
6.4.1	(a)	104
6.4.2	(b)	104
6.4.3	(c)	104
6.5	Exercise 5	104
6.6	Exercise 6	104
6.7	Exercise 7	104
6.8	Exercise 8	105
6.9	Exercise 9	105
6.10	Exercise 10	105
6.11	Exercise 11	105
6.12	Exercise 12	105
6.13	Exercise 13	105
6.14	Exercise 14	105
6.15	Exercise 15	105
6.16	Exercise 16	105
6.16.1	(a)	105

6.16.2	(b)	105
6.16.3	(c)	105
6.17	Exercise 17	106
6.17.1	(a)	106
6.17.2	(b)	106
6.17.3	(c)	106
6.17.4	(d)	106
6.18	Exercise 18	106
6.18.1	(a)	106
6.18.2	(b)	106
6.18.3	(c)	106
6.18.4	(d)	106
6.19	Exercise 19	106
6.19.1	(a)	106
6.19.2	(b)	106
6.20	Exercise 20	106
6.20.1	(a)	106
6.20.2	(b)	107
6.21	Exercise 21	107
7	Exercise Set 9.7	107
7.1	Exercise 1	107
7.2	Exercise 2	107
7.3	Exercise 3	107
7.4	Exercise 4	107
7.5	Exercise 5	107
7.6	Exercise 6	107
7.7	Exercise 7	107
7.8	Exercise 8	107
7.9	Exercise 9	107
7.10	Exercise 10	108
7.10.1	(a)	108
7.10.2	(b)	108
7.10.3	(c)	108
7.11	Exercise 11	108
7.12	Exercise 12	108
7.13	Exercise 13	108
7.14	Exercise 14	108
7.15	Exercise 15	108
7.16	Exercise 16	108
7.17	Exercise 17	108
7.18	Exercise 18	108
7.19	Exercise 19	109
7.20	Exercise 20	109
7.21	Exercise 21	109
7.22	Exercise 22	109

7.23	Exercise 23	109
7.24	Exercise 24	109
7.25	Exercise 25	109
7.26	Exercise 26	109
7.27	Exercise 27	109
7.28	Exercise 28	109
7.29	Exercise 29	109
7.30	Exercise 30	109
7.31	Exercise 31	110
7.32	Exercise 32	110
7.33	Exercise 33	110
7.34	Exercise 34	110
7.35	Exercise 35	110
7.36	Exercise 36	110
7.37	Exercise 37	110
7.38	Exercise 38	110
7.39	Exercise 39	110
7.40	Exercise 40	110
7.41	Exercise 41	110
7.42	Exercise 42	110
7.43	Exercise 43	111
7.44	Exercise 44	111
7.45	Exercise 45	111
7.46	Exercise 46	111
7.47	Exercise 47	111
7.48	Exercise 48	111
7.49	Exercise 49	111
7.50	Exercise 50	111
7.51	Exercise 51	111
7.52	Exercise 52	111
7.53	Exercise 53	111
7.54	Exercise 54	111
7.55	Exercise 55	112
7.55.1	(a)	112
7.55.2	(b)	112
7.55.3	(c)	112
7.55.4	(d)	112
8	Exercise Set 9.8	112
8.1	Exercise 1	112
8.2	Exercise 2	112
8.3	Exercise 3	112
8.3.1	(a)	112
8.3.2	(b)	112
8.4	Exercise 4	112
8.5	Exercise 5	112

8.6	Exercise 6	113
8.7	Exercise 7	113
8.7.1	(a)	113
8.7.2	(b)	113
8.7.3	(c)	113
8.7.4	(d)	113
8.7.5	(e)	113
8.7.6	(f)	113
8.8	Exercise 8	113
8.9	Exercise 9	113
8.9.1	(a)	113
8.9.2	(b)	113
8.9.3	(c)	113
8.9.4	(d)	114
8.9.5	(e)	114
8.9.6	(f)	114
8.10	Exercise 10	114
8.11	Exercise 11	114
8.12	Exercise 12	114
8.13	Exercise 13	114
8.14	Exercise 14	114
8.15	Exercise 15	114
8.16	Exercise 16	114
8.17	Exercise 17	114
8.18	Exercise 18	114
8.19	Exercise 19	115
8.20	Exercise 20	115
8.21	Exercise 21	115
8.22	Exercise 22	115
8.23	Exercise 23	115
9	Exercise Set 9.9	115
9.1	Exercise 1	115
9.2	Exercise 2	115
9.3	Exercise 3	115
9.4	Exercise 4	115
9.4.1	(a)	115
9.4.2	(b)	115
9.5	Exercise 5	115
9.6	Exercise 6	116
9.6.1	(a)	116
9.6.2	(b)	116
9.6.3	(c)	116
9.7	Exercise 7	116
9.8	Exercise 8	116
9.8.1	(a)	116

9.8.2	(b)	116
9.8.3	(c)	116
9.9	Exercise 9	116
9.10	Exercise 10	116
9.11	Exercise 11	116
9.11.1	(a)	116
9.11.2	(b)	116
9.12	Exercise 12	117
9.13	Exercise 13	117
9.13.1	(a)	117
9.13.2	(b)	117
9.14	Exercise 14	117
9.14.1	(a)	117
9.14.2	(b)	117
9.15	Exercise 15	117
9.15.1	(a)	117
9.15.2	(b)	117
9.15.3	(c)	117
9.15.4	(d)	117
9.16	Exercise 16	117
9.16.1	(a)	117
9.16.2	(b)	118
9.17	Exercise 17	118
9.18	Exercise 18	118
9.19	Exercise 19	118
9.20	Exercise 20	118
9.21	Exercise 21	118
9.22	Exercise 22	118
9.23	Exercise 23	118
9.23.1	(a)	118
9.23.2	(b)	118
9.23.3	(c)	118
9.24	Exercise 24	118
9.24.1	(a)	118
9.24.2	(b)	119
9.25	Exercise 25	119
9.25.1	(a)	119
9.25.2	(b)	119
9.25.3	(c)	119
9.25.4	(d)	119
9.26	Exercise 26	119
9.27	Exercise 27	119
9.28	Exercise 28	119
9.28.1	(a)	119
9.28.2	(b)	119
9.28.3	(c)	119

9.28.4 (d)	119
9.29 Exercise 29	120
9.29.1 (a)	120
9.29.2 (b)	120
9.29.3 (c)	120
9.29.4 (d)	120
9.30 Exercise 30	120
9.30.1 (a)	120
9.30.2 (b)	120
9.30.3 (c)	120
9.30.4 (d)	120
9.31 Exercise 31	120
9.31.1 (a)	120
9.31.2 (b)	120
9.31.3 (c)	120
9.32 Exercise 32	121
9.32.1 (a)	121
9.32.2 (b)	121
9.32.3 (c)	121
9.32.4 (d)	121
9.32.5 (e)	121
9.33 Exercise 33	121
9.34 Exercise 34	121

1 Exercise Set 9.1

1.1 Exercise 1

Toss two coins 30 times and make a table showing the relative frequencies of 0, 1, and 2 heads. How do your values compare with those shown in Table 9.1.1?

<i>Proof.</i>	Event	Freq.	Rel. freq.
	2 heads obtained	7	23.33%
	1 head obtained	16	53.33%
	0 heads obtained	7	23.33%

□

1.2 Exercise 2

In the example of tossing two quarters, what is the probability that at least one head is obtained? that coin A is a head? that coins A and B are either both heads or both tails?

Proof. $3/4$, $1/2$, $1/2$

□

In 3 – 6 use the sample space given in example 9.1.1. Write each event as a set and compute its probability.

1.3 Exercise 3

The event that the chosen card is red and is not a face card.

Proof. $\{1\Diamond, 2\Diamond, 3\Diamond, 4\Diamond, 5\Diamond, 6\Diamond, 7\Diamond, 8\Diamond, 9\Diamond, 10\Diamond, 1\Heart, 2\Heart, 3\Heart, 4\Heart, 5\Heart, 6\Heart, 7\Heart, 8\Heart, 9\Heart, 10\Heart\}$, probability $= 20/52 \approx 38.5\%$ \square

1.4 Exercise 4

The event that the chosen card is black and has an even number on it.

Proof. $\{2\Spadesuit, 4\Spadesuit, 6\Spadesuit, 8\Spadesuit, 10\Spadesuit, 2\clubsuit, 4\clubsuit, 6\clubsuit, 8\clubsuit, 10\clubsuit\}$, probability $= 10/52 \approx 19.2\%$ \square

1.5 Exercise 5

The event that the denomination of the chosen card is at least 10 (counting aces high).

Proof. $\{10\Spadesuit, J\Spadesuit, Q\Spadesuit, K\Spadesuit, A\Spadesuit, 10\Diamond, J\Diamond, Q\Diamond, K\Diamond, A\Diamond, 10\Heart, J\Heart, Q\Heart, K\Heart, A\Heart, 10\clubsuit, J\clubsuit, Q\clubsuit, K\clubsuit, A\clubsuit\}$, probability $= 20/52 = 5/13 \approx 38.5\%$ \square

1.6 Exercise 6

The event that the denomination of the chosen card is at most 4 (counting aces high).

Proof. $\{2\clubsuit, 2\Heart, 2\Spadesuit, 2\Diamond, 3\clubsuit, 3\Heart, 3\Spadesuit, 3\Diamond, 4\clubsuit, 4\Heart, 4\Spadesuit, 4\Diamond\}$, probability $= 12/52 \approx 23.0\%$ \square

In 7 – 10, use the sample space given in example 9.1.2. Write each of the following events as a set and compute its probability.

1.7 Exercise 7

The event that the sum of the numbers showing face up is 8.

Proof. $\{26, 35, 44, 53, 62\}$, probability $= 5/36 \approx 13.9\%$ \square

1.8 Exercise 8

The event that the numbers showing face up are the same.

Proof. $\{11, 22, 33, 44, 55, 66\}$, probability $= 6/36 \approx 16.6\%$ \square

1.9 Exercise 9

The event that the sum of the numbers showing face up is at most 6.

Proof. $\{11, 12, 13, 14, 15, 21, 22, 23, 24, 31, 32, 33, 41, 42, 51\}$, $\text{prob.} = 15/36 \approx 41.6\%$ \square

1.10 Exercise 10

The event that the sum of the numbers showing face up is at least 9.

Proof. $\{36, 45, 46, 54, 55, 56, 63, 64, 65, 66\}$, probability $= 10/36 \approx 27.7\%$ \square

1.11 Exercise 11

Suppose that a coin is tossed three times and the side showing face up on each toss is noted. Suppose also that on each toss heads and tails are equally likely. Let HHT indicate the outcome heads on the first two tosses and tails on the third, THT the outcome tails on the first and third tosses and heads on the second, and so forth.

1.11.1 (a)

List the eight elements in the sample space whose outcomes are all the possible head-tail sequences obtained in the three tosses.

Proof. $\{HHH, HHT, HTH, HTT, THH, THT, TTH, TTT\}$ \square

1.11.2 (b)

Write each of the following events as a set and find its probability:

- (i) The event that exactly one toss results in a head.
- (ii) The event that at least two tosses result in a head.
- (iii) The event that no head is obtained.

Proof. (i) $\{HTT, THT, TTH\}$, probability $= 3/8 = 37.5\%$

(ii) $\{HHT, HTH, THH, HHH\}$, probability $= 4/8 = 50.0\%$

(iii) $\{TTT\}$, probability $= 1/8 = 12.5\%$ \square

1.12 Exercise 12

Suppose that each child born is equally likely to be a boy or a girl. Consider a family with exactly three children. Let BBG indicate that the first two children born are boys and the third child is a girl, let GBG indicate that the first and third children born are girls and the second is a boy, and so forth.

1.12.1 (a)

List the eight elements in the sample space whose outcomes are all possible genders of the three children.

Proof. $\{BBB, BBG, BGB, BGG, GBB, GBG, GGB, GGG\}$ □

1.12.2 (b)

Write each of the events in the next column as a set and find its probability.

(i) The event that exactly one child is a girl.

(ii) The event that at least two children are girls.

(iii) The event that no child is a girl.

Proof. (i) $\{GBB, BGB, BBG\}$, probability $= 3/8 = 37.5\%$

(ii) $\{GGB, GBG, BGG, GGG\}$, probability $= 4/8 = 50\%$

(iii) $\{BBB\}$, probability $= 1/8 = 12.5\%$ □

1.13 Exercise 13

Suppose that on a true/false exam you have no idea at all about the answers to three questions. You choose answers randomly and therefore have a 50–50 chance of being correct on any one question. Let CCW indicate that you were correct on the first two questions and wrong on the third, let WCW indicate that you were wrong on the first and third questions and correct on the second, and so forth.

1.13.1 (a)

List the elements in the sample space whose outcomes are all possible sequences of correct and incorrect responses on your part.

Proof. $\{CCC, CCW, CWC, CWW, WCC, WCW, WWC, WWW\}$ □

1.13.2 (b)

Write each of the following events as a set and find its probability:

(i) The event that exactly one answer is correct.

(ii) The event that at least two answers are correct.

(iii) The event that no answer is correct.

Proof. (i) $\{CWW, WCW, WWC\}$, probability $= 3/8 = 37.5\%$

(ii) $\{CCW, CWC, WCC, CCC\}$, probability $= 4/8 = 50\%$

(iii) $\{WWW\}$, probability $= 1/8 = 12.5\%$ □

1.14 Exercise 14

Three people have been exposed to a certain illness. Once exposed, a person has a 50-50 chance of actually becoming ill.

1.14.1 (a)

What is the probability that exactly one of the people becomes ill?

Proof. probability = $3/8 = 37.5\%$

□

1.14.2 (b)

What is the probability that at least two of the people become ill?

Proof. probability = $4/8 = 50\%$

□

1.14.3 (c)

What is the probability that none of the three people becomes ill?

Proof. probability = $1/8 = 12.5\%$

□

1.15 Exercise 15

When discussing counting and probability, we often consider situations that may appear frivolous or of little practical value, such as tossing coins, choosing cards, or rolling dice. The reason is that these relatively simple examples serve as models for a wide variety of more complex situations in the real world. In light of this remark, comment on the relationship between your answer to exercise 11 and your answers to exercises 12 – 14.

Proof. The answers to exercises 11, 12, 13, 14 are the same, because all 4 situations are modeled exactly the same way. They each consist of 3 instances of the same event, each of which has 2 possible outcomes with 50% probability (heads or tails, boy or girl, correct or incorrect, ill or not ill). They each have a sample space with 8 elements following the same pattern.

□

1.16 Exercise 16

Two faces of a six-sided die are painted red, two are painted blue, and two are painted yellow. The die is rolled three times, and the colors that appear face up on the first, second, and third rolls are recorded.

1.16.1 (a)

Let BBR denote the outcome where the color appearing face up on the first and second rolls is blue and the color appearing face up on the third roll is red. Because there are

as many faces of one color as of any other, the outcomes of this experiment are equally likely. List all 27 possible outcomes.

Proof. $\{RRR, RRB, RRY, RBR, RBB, RBY, RYR, RYB, RYY,$
 $BBR, BRB, BRY, BBR, BBB, BBY, BYR, BYB, BYY,$
 $YRR, YRB, YRY, YBR, YBB, YBY, YYR, YYB, YYY\}$ \square

1.16.2 (b)

Consider the event that all three rolls produce different colors. One outcome in this event is RBY and another RYB . List all outcomes in the event. What is the probability of the event?

Proof. $\{RBY, RYB, YBR, BRY, BYR, YRB\}$, probability $= 6/27 = 2/9 \approx 22.2\%$ \square

1.16.3 (c)

Consider the event that two of the colors that appear face up are the same. One outcome in this event is RRB and another is RBR . List all outcomes in the event. What is the probability of the event?

Proof. $\{RRB, RBR, BRR, RRY, RYR, YRR, BBR, BRB, RBB, BBY, BYB, YBB,$
 $YYR, YRY, RYY, YYB, YBY, BYY\}$, probability $= 18/27 = 2/3 \approx 66.6\%$ \square

1.17 Exercise 17

Consider the situation described in exercise 16.

1.17.1 (a)

Find the probability of the event that exactly one of the colors that appears face up is red.

Proof. $\{RBB, RBY, RYB, RYY, BRB, BRY, BBR, BYR, YRB, YRY, YBR, YYR\}$,
probability $= 12/27 = 4/9 \approx 44.4\%$ \square

1.17.2 (b)

Find the probability of the event that at least one of the colors that appears face up is red.

Proof. $\{RRR, RRB, RRY, RBR, RBB, RBY, RYR, RYB, RYY, BRR, BRB, BRY,$
 $BBR, BYR, YRR, YRB, YRY, YBR, YYR\}$, probability $= 20/27 \approx 74\%$ \square

1.18 Exercise 18

An urn contains two blue balls (denoted B_1 and B_2) and one white ball (denoted W). One ball is drawn, its color is recorded, and it is replaced in the urn. Then another ball is drawn, and its color is recorded.

1.18.1 (a)

Let B_1W denote the outcome that the first ball drawn is B_1 and the second ball drawn is W . Because the first ball is replaced before the second ball is drawn, the outcomes of the experiment are equally likely. List all nine possible outcomes of the experiment.

Proof. $\{B_1B_1, B_1B_2, B_1W, B_2B_1, B_2B_2, B_2W, WB_1, WB_2, WW\}$ □

1.18.2 (b)

Consider the event that the two balls that are drawn are both blue. List all outcomes in the event. What is the probability of the event?

Proof. $\{B_1B_1, B_1B_2, B_2B_1, B_2B_2\}$, probability $= 4/9 \approx 44.4\%$ □

1.18.3 (c)

Consider the event that the two balls that are drawn are of different colors. List all outcomes in the event. What is the probability of the event?

Proof. $\{B_1W, B_2W, WB_1, WB_2\}$, probability $= 4/9 \approx 44.4\%$ □

1.19 Exercise 19

An urn contains two blue balls (denoted B_1 and B_2) and three white balls (denoted W_1, W_2 , and W_3). One ball is drawn, its color is recorded, and it is replaced in the urn. Then another ball is drawn and its color is recorded.

1.19.1 (a)

Let B_1W_2 denote the outcome that the first ball drawn is B_1 and the second ball drawn is W_2 . Because the first ball is replaced before the second ball is drawn, the outcomes of the experiment are equally likely. List all 25 possible outcomes of the experiment.

Proof. $\{B_1B_1, B_1B_2, B_1W_1, B_1W_2, B_1W_3,$
 $B_2B_1, B_2B_2, B_2W_1, B_2W_2, B_2W_3,$
 $W_1B_1, W_1B_2, W_1W_1, W_1W_2, W_1W_3,$
 $W_2B_1, W_2B_2, W_2W_1, W_2W_2, W_2W_3,$
 $W_3B_1, W_3B_2, W_3W_1, W_3W_2, W_3W_3\}$ □

1.19.2 (b)

Consider the event that the first ball that is drawn is blue. List all outcomes in the event. What is the probability of the event?

Proof. $\{B_1B_1, B_1B_2, B_1W_1, B_1W_2, B_1W_3, B_2B_1, B_2B_2, B_2W_1, B_2W_2, B_2W_3\}$,

probability = $10/25 = 40\%$

□

1.19.3 (c)

Consider the event that only white balls are drawn. List all outcomes in the event. What is the probability of the event?

Proof. $\{W_1W_1, W_1W_2, W_1W_3, W_2W_1, W_2W_2, W_2W_3, W_3W_1, W_3W_2, W_3W_3\}$

probability = $9/25 = 36\%$

□

1.20 Exercise 20

Refer to Example 9.1.3. Suppose you are appearing on a game show with a prize behind one of five closed doors: A, B, C, D, and E. If you pick the correct door, you win the prize. You pick door A. The game show host then opens one of the other doors and reveals that there is no prize behind it. Then the host gives you the option of staying with your original choice of door A or switching to one of the other doors that is still closed.

1.20.1 (a)

If you stick with your original choice, what is the probability that you will win the prize?

Proof. There are 5 cases.

In Case 1, the prize is behind door A. In this case if I stick with door A, I would win.

In the remaining 4 cases, the prize is behind one of the doors B,C,D,E. In these cases if I stick with door A, I would lose.

Each case has 20% probability. If I stick with door A, I can only win in Case 1. Therefore the probability that I win the prize is 20%.

□

1.20.2 (b)

If you switch to another door, what is the probability that you will win the prize?

Proof. Like above, if I switch to another door, I lose in Case 1, and I have a $1/4$ chance of winning in each of the other 4 cases. Therefore the probability that I win the prize is 25%.

□

1.21 Exercise 21

1.21.1 (a)

How many positive two-digit integers are multiples of 3?

Proof. Between 10 and 99 (inclusive), the multiples of 3 are: 12, 15, 18, \dots , 93, 96, 99. Notice that $12 = 3 \cdot 4$ and $99 = 3 \cdot 33$. So there are as many positive two-digit integers that are multiples of 3 as there are integers from 4 to 33 inclusive. By Theorem 9.1.1 there are $33 - 4 + 1 = 30$ such integers. \square

1.21.2 (b)

What is the probability that a randomly chosen positive two-digit integer is a multiple of 3?

Proof. There are $99 - 10 + 1 = 90$ positive two-digit integers in all, and by part (a), 30 of these are multiples of 3. So the probability that a randomly chosen positive two-digit integer is a multiple of 3 is $30/90 = 1/3 \approx 33.3\%$. \square

1.21.3 (c)

What is the probability that a randomly chosen positive two-digit integer is a multiple of 4?

Proof. Of the integers from 10 through 99 that are multiples of 4, the smallest is 12 ($= 4 \cdot 3$) and the largest is 96 ($= 4 \cdot 24$). Thus there are $24 - 3 + 1 = 22$ two-digit integers that are multiples of 4. Hence the probability that a randomly chosen two-digit integer is a multiple of 4 is $22/90 \approx 36.6\%$. \square

1.22 Exercise 22

1.22.1 (a)

How many positive three-digit integers are multiples of 6?

Proof. They are 102, 108, 114, \dots , 984, 990, 996. Notice $102 = 6 \cdot 17$ and $996 = 6 \cdot 166$. So there are $166 - 17 + 1 = 150$ such integers. \square

1.22.2 (b)

What is the probability that a randomly chosen positive three-digit integer is a multiple of 6?

Proof. There are $999 - 100 + 1 = 900$ positive three-digit integers. So the probability is, by part (a), $150/900 = 1/6 \approx 16.6\%$. \square

1.22.3 (c)

What is the probability that a randomly chosen positive three-digit integer is a multiple of 7?

Proof. Multiples of 7 are $7 \cdot 15 = 105, 7 \cdot 16 = 112, \dots, 7 \cdot 141 = 987, 7 \cdot 142 = 994$. So there are $142 - 15 + 1 = 128$ such integers, then the probability is $128/900 = 32/225 \approx 14.22\%$. \square

1.23 Exercise 23

Suppose $A[1], A[2], A[3], \dots, A[n]$ is a one-dimensional array and $n > 50$.

1.23.1 (a)

How many elements are in the array?

Proof. n elements. \square

1.23.2 (b)

How many elements are in the subarray $A[4], A[5], \dots, A[39]$?

Proof. $39 - 4 + 1 = 36$ elements. \square

1.23.3 (c)

If $3 \leq m \leq n$, what is the probability that a randomly chosen array element is in the subarray $A[3], A[4], \dots, A[m]$?

Proof. There are $m - 3 + 1 = m - 2$ elements in the subarray. There are n elements in the array. So the probability is $\frac{m-2}{n}$. \square

1.23.4 (d)

What is the probability that a randomly chosen array element is in the subarray shown below if $n = 39$?

$$A[\lfloor n/2 \rfloor], A[\lfloor n/2 \rfloor + 1], \dots, A[n]$$

Proof. $\lfloor 39/2 \rfloor = \lfloor 19.5 \rfloor = 19$, therefore there are $39 - 19 + 1 = 21$ elements in the subarray, and there are 39 elements in the array, so the probability is $21/39 \approx 53.84\%$. \square

1.24 Exercise 24

Suppose $A[1], A[2], \dots, A[n]$ is a one-dimensional array and $n \geq 2$. Consider the subarray $A[1], A[2], \dots, A[\lfloor n/2 \rfloor]$.

1.24.1 (a)

How many elements are in the sub-array (i) if n is even? and (ii) if n is odd?

Proof. (i) There are $\lfloor \frac{n}{2} \rfloor = \frac{n}{2}$ elements in the sub-array.

(ii) There are $\lfloor \frac{n}{2} \rfloor = \frac{n-1}{2}$ elements in the sub-array. \square

1.24.2 (b)

What is the probability that a randomly chosen array element is in the sub-array (i) if n is even? and (ii) if n is odd?

Proof. There are n elements in the array, so

(i) The probability that an element is in the given sub-array is $\frac{n/2}{n} = \frac{1}{2}$,

(i) The probability that an element is in the given sub-array is $\frac{(n-1)/2}{n} = \frac{n-1}{2n}$. \square

1.25 Exercise 25

Suppose $A[1], A[2], \dots, A[n]$ is a one-dimensional array and $n \geq 2$. Consider the sub-array $A[\lfloor n/2 \rfloor], A[\lfloor n/2 \rfloor + 1], \dots, A[n]$.

1.25.1 (a)

How many elements are in the sub-array (i) if n is even? and (ii) if n is odd?

Proof. (i) There are $n - \lfloor \frac{n}{2} \rfloor + 1 = n - \frac{n}{2} + 1 = \frac{n+2}{2}$ elements in the sub-array.

(ii) There are $n - \lfloor \frac{n}{2} \rfloor + 1 = n - \frac{n-1}{2} + 1 = \frac{n+3}{2}$ elements in the sub-array. \square

1.25.2 (b)

What is the probability that a randomly chosen array element is in the sub-array (i) if n is even? and (ii) if n is odd?

Proof. There are n elements in the array, so

(i) The probability that an element is in the given sub-array is $\frac{(n+2)/2}{n} = \frac{n+2}{2n}$,

(i) The probability that an element is in the given sub-array is $\frac{(n+3)/2}{n} = \frac{n+3}{2n}$. \square

1.26 Exercise 26

What is the 27th element in the one-dimensional array $A[42], A[43], \dots, A[100]$?

Proof. Let k be the 27th element in the array. By Theorem 9.1.1, $k - 42 + 1 = 27$, and so $k = 42 + 27 - 1 = 68$. Thus the 27th element in the array is $A[68]$. \square

1.27 Exercise 27

What is the 62nd element in the one-dimensional array $B[29], B[30], \dots, B[100]$?

Proof. Let k be the 62nd element in the array. By Theorem 9.1.1, $k - 29 + 1 = 62$, and so $k = 29 + 62 - 1 = 90$. Thus the 62th element in the array is $A[90]$. \square

1.28 Exercise 28

If the largest of 56 consecutive integers is 279, what is the smallest?

Proof. Let m be the smallest of the integers. By Theorem 9.1.1, $279 - m + 1 = 56$, and so $m = 279 - 56 + 1 = 224$. Thus the smallest of the integers is 224. \square

1.29 Exercise 29

If the largest of 87 consecutive integers is 326, what is the smallest?

Proof. Let m be the smallest of the integers. By Theorem 9.1.1, $326 - m + 1 = 87$, and so $m = 326 - 87 + 1 = 240$. Thus the smallest of the integers is 240. \square

1.30 Exercise 30

How many even integers are between 1 and 1,001?

Proof. They are $2 = 2 \cdot 1, 4 = 2 \cdot 2, \dots, 998 = 2 \cdot 499, 1000 = 2 \cdot 500$. So there are 500 of them. \square

1.31 Exercise 31

How many integers that are multiples of 3 are between 1 and 1,001?

Proof. They are $3 = 3 \cdot 1, 6 = 3 \cdot 2, \dots, 996 = 3 \cdot 332, 999 = 3 \cdot 333$. So there are 333 of them. \square

1.32 Exercise 32

A certain non-leap year has 365 days, and January 1 occurs on a Monday.

1.32.1 (a)

How many Sundays are in the year?

Proof. Sundays occur on days 7, 14, 21, ..., 364 of the year. Since $7 = 7 \cdot 1$ and $364 = 7 \cdot 52$, there are 52 Sundays in the year. \square

1.32.2 (b)

How many Mondays are in the year?

Proof. For each Sunday, there is a Monday in the same week. However there is also the 365th day, which comes directly after the last Sunday, which is the 364th day. Therefore there are $52 + 1 = 53$ Mondays in the year. \square

1.33 Exercise 33

Prove Theorem 9.1.1. (Let m be any integer and prove the theorem by mathematical induction on n .)

Proof. Let m be any integer and let $P(n)$ be the statement “if $m \leq n$, then there are $n - m + 1$ integers from m to n inclusive.” The base case is $n = m$.

Show $P(m)$ is true: There is only one integer from m to m inclusive, namely m itself. And $n - m + 1 = m - m + 1 = 1$, so $P(m)$ is true.


Show that for any integer $k \geq m$ if $P(k)$ is true then $P(k + 1)$ is true: Assume $k \geq m$ and assume there are $k - m + 1$ integers from m to k inclusive. Then there is one more integer, namely $k + 1$, from m to $k + 1$ inclusive, thus there are $(k - m + 1) + 1 = (k + 1) - m + 1$ integers from m to $k + 1$ inclusive. So $P(k + 1)$ is true. \square

2 Exercise Set 9.2

In 1 – 4, use the fact that in baseball’s World Series, the first team to win four games wins the series.

2.1 Exercise 1


Suppose team A wins the first three games. How many ways can the World Series be completed? (Draw a tree.)



Proof. There are five ways to complete the series: A , $B - A$, $B - B - A$, $B - B - B - A$, and $B - B - B - B$. \square

2.2 Exercise 2

Suppose team A wins the first two games. How many ways can the World Series be completed? (Draw a tree.)



Proof. There are 15 ways: AA, ABA, ABBA, ABBBA, ABBBB, BAA, BABA, BABBA, BABBB, BBAA, BBABA, BBABB, BBBAA, BBBAB, BBBB. \square

2.3 Exercise 3

How many ways can a World Series be played if team A wins four games in a row?

Proof. Four ways: $A - A - A - A$, $B - A - A - A - A$, $B - B - A - A - A - A$, and $B - B - B - A - A - A - A$. \square


2.4 Exercise 4

How many ways can a World Series be played if no team wins two games in a row?

Proof. Two ways: $A - B - A - B - A - B - A$ and $B - A - B - A - B - A - B$. \square

2.5 Exercise 5

In a competition between players X and Y , the first player to win three games in a row or a total of four games wins. How many ways can the competition be played if X wins the first game and Y wins the second and third games? (Draw a tree.)




Proof. There are seven ways: $XXX, XXYX, XXYY, XYXX, XYXY, XYY, Y$. □

2.6 Exercise 6

One urn contains two black balls (labeled B_1 and B_2) and one white ball. A second urn contains one black ball and two white balls (labeled W_1 and W_2). Suppose the following experiment is performed: One of the two urns is chosen at random. Next a ball is randomly chosen from the urn. Then a second ball is chosen at random from the same urn without replacing the first ball.

2.6.1 (a)

Construct the possibility tree showing all possible outcomes of this experiment.



Proof. □

2.6.2 (b)

What is the total number of outcomes of this experiment?

Proof. 12 □

2.6.3 (c)

What is the probability that two black balls are chosen?

Proof. $2/12 = 1/6 \approx 16.6\%$ □

2.6.4 (d)

What is the probability that two balls of opposite color are chosen?


Proof. $8/12 = 2/3 \approx 66.6\%$ □

2.7 Exercise 7

One urn contains one blue ball (labeled B_1) and three red balls (labeled R_1, R_2 , and R_3). A second urn contains two red balls (R_4 and R_5) and two blue balls (B_2 and B_3). An experiment is performed in which one of the two urns is chosen at random and then two balls are randomly chosen from it, one after the other without replacement.

2.7.1 (a)

Construct the possibility tree showing all possible outcomes of this experiment.



Proof.

□

2.7.2 (b)

What is the total number of outcomes of this experiment?

Proof. 24

□

2.7.3 (c)

What is the probability that two red balls are chosen?

Proof. $8/24 = 1/3 \approx 33.3\%$

□

2.8 Exercise 8

A person buying a personal computer system is offered a choice of three models of the basic unit, two models of keyboard, and two models of printer. How many distinct systems can be purchased?

Proof. By the multiplication rule, the answer is $3 \cdot 2 \cdot 2 = 12$.

□

2.9 Exercise 9

Suppose there are three roads from city A to city B and five roads from city B to city C .

2.9.1 (a)

How many ways is it possible to travel from city A to city C via city B ?

Proof. In going from city A to city B , one may take any of the 3 roads. In going from city B to city C , one may take any of the 5 roads. So, by the multiplication rule, there are $3 \cdot 5 = 15$ ways to travel from city A to city C via city B . \square

2.9.2 (b)

How many different round-trip routes are there from city A to B to C to B and back to A ?

Proof. A round-trip journey can be thought of as a four-step operation: **Step 1:** Go from A to B . **Step 2:** Go from B to C . **Step 3:** Go from C to B . **Step 4:** Go from B to A .

Since there are 3 ways to perform step 1, 5 ways to perform step 2, 5 ways to perform step 3, and 3 ways to perform step 4, by the multiplication rule, there are $3 \cdot 5 \cdot 5 \cdot 3 = 225$ round-trip routes. \square

2.9.3 (c)

How many different routes are there from cities A to B to C to B and back to A in which no road is traversed twice?

Proof. In this case the steps for making a round-trip journey are the same as in part (b), but since no route segment may be repeated, there are only 4 ways to perform step 3 and only 2 ways to perform step 4. So, by the multiplication rule, there are $3 \cdot 5 \cdot 4 \cdot 2 = 120$ round-trip routes in which no road is traversed twice. \square

2.10 Exercise 10

Suppose there are three routes from North Point to Boulder Creek, two routes from Boulder Creek to Beaver Dam, two routes from Beaver Dam to Star Lake, and four routes directly from Boulder Creek to Star Lake. (Draw a sketch.)

2.10.1 (a)

How many routes from North Point to Star Lake pass through Beaver Dam?

Proof. By the multiplication rule $3 \cdot 2 \cdot 2 = 12$ routes. \square

2.10.2 (b)

How many routes from North Point to Star Lake bypass Beaver Dam?

Proof. By the multiplication rule $3 \cdot 4 = 12$ routes. \square

2.11 Exercise 11

2.11.1 (a)

A bit string is a finite sequence of 0's and 1's. How many bit strings have length 8?

Proof. Imagine constructing a bit string of length 8 as an eight-step process:

Step 1: Choose either a 0 or a 1 for the left-most position,

Step 2: Choose either a 0 or a 1 for the next position to the right.

⋮

Step 8: Choose either a 0 or a 1 for the right-most position.

Since there are 2 ways to perform each step, the total number of ways to accomplish the entire operation, which is the number of different bit strings of length 8, is $2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 \cdot 2 = 2^8 = 256$. \square

2.11.2 (b)

How many bit strings of length 8 begin with three 0's?

Proof. Imagine that there are three 0's in the three left-most positions, and imagine filling in the remaining 5 positions as a five-step process, where step i is to fill in the $(i + 3)$ rd position. Since there are 2 ways to perform each of the 5 steps, there are 2^5 ways to perform the entire operation. So there are 2^5 , or 32, 8-bit strings that begin with three 0's. \square

2.11.3 (c)

How many bit strings of length 8 begin and end with a 1?

Proof. $2^6 = 64$ \square

2.12 Exercise 12

Hexadecimal numbers are made using the sixteen hexadecimal digits 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, A, B, C, D, E, F and are denoted using the subscript 16. For example, 9A2D₁₆ and BC54₁₆ are hexadecimal numbers.

2.12.1 (a)

How many hexadecimal numbers begin with one of the digits 3 through B, end with one of the digits 5 through F, and are 5 digits long?

Proof. Think of creating a hexadecimal number that satisfies the given requirements as a five-step process.

Step 1: Choose the left-most hexadecimal digits. It can be any of the 9 hexadecimal digits from 3 through B.

Steps 2-4: Choose the three hexadecimal digits for the middle three positions. Each can be any of the 16 hexadecimal digits.

Step 5: Choose the right-most hexadecimal digit. It can be any of the 11 hexadecimal digits from 5 through F.

There are 9 ways to perform step 1, 16 ways to perform each of steps 2 through 4, and 11 ways to perform step 5. Thus, the total number of specified hexadecimal numbers is $9 \cdot 16 \cdot 16 \cdot 16 \cdot 11 = 405,504$. \square

2.12.2 (b)

How many hexadecimal numbers begin with one of the digits 4 through D, end with one of the digits 2 through E, and are 6 digits long?

Proof. There are 10 choices for the first digit: 4, 5, 6, 7, 8, 9, A, B, C, D.

There are 14 choices for the last digit: 2, 3, 4, 5, 6, 7, 8, 9, A, B, C, D, E.

There are 16 choices for each one of the 4 middle digits.

So: $10 \cdot 16 \cdot 16 \cdot 16 \cdot 16 \cdot 14 = 9175040$ \square

2.13 Exercise 13

A coin is tossed four times. Each time the result H for heads or T for tails is recorded. An outcome of HHTT means that heads were obtained on the first two tosses and tails on the second two. Assume that heads and tails are equally likely on each toss.

2.13.1 (a)

How many distinct outcomes are possible?

Proof. In each of the four tosses there are two possible results: Either a head (H) or a tail (T) is obtained. Thus, by the multiplication rule, the number of outcomes is $2 \cdot 2 \cdot 2 \cdot 2 = 2^4 = 16$. \square

2.13.2 (b)

What is the probability that exactly two heads occur?

Proof. There are six outcomes with two heads: HHTT, HTHT, HTTH, THHT, THTH, TTHH. Thus the probability of obtaining exactly two heads is $6/16 = 3/8$. \square

2.13.3 (c)

What is the probability that exactly one head occurs?

Proof. There are four outcomes with exactly one head: HTTT, THTT, TTHT, TTTH. Thus the probability of obtaining exactly two heads is $4/16 = 1/4$. \square

2.14 Exercise 14

Suppose that in a certain state, all automobile license plates have four uppercase letters followed by three digits.

2.14.1 (a)

How many different license plates are possible?

Proof. Think of creating license plates that satisfy the given conditions as the following seven-step process: In steps 1–4 choose the letters to put in positions 1–4, and in steps 5–7, choose the digits to put in positions 5–7. Since there are 26 letters and 10 digits and since repetition is allowed, there are 26 ways to perform each of steps 1–4 and 10 ways to perform each of steps 5–7. Thus the number of license plates is $26 \cdot 26 \cdot 26 \cdot 26 \cdot 10 \cdot 10 \cdot 10 = 456,976,000$. \square

2.14.2 (b)

How many license plates could begin with A and end in 0?

Proof. In this case there is only one way to perform step 1 (because the first letter must be an A) and only one way to perform step 7 (because the last digit must be a 0). Therefore, the number of license plates is $26 \cdot 26 \cdot 26 \cdot 10 \cdot 10 = 1,757,600$. \square

2.14.3 (c)

How many license plates could begin with TGIF?

Proof. There are 3 digits left to be chosen. Each has 10 possibilities. So by the multiplication rule $10^3 = 1000$ can begin with TGIF. \square

2.14.4 (d)

How many license plates are possible in which all the letters and digits are distinct?

Proof. In this case there are 26 ways to perform step 1, 25 ways to perform step 2, 24 ways to perform step 3, 23 ways to perform step 4, 10 ways to perform step 5, 9 ways to perform step 6, and 8 ways to perform step 7, so the number of license plates is $26 \cdot 25 \cdot 24 \cdot 23 \cdot 10 \cdot 9 \cdot 8 = 258,336,000$. \square

2.14.5 (e)

How many license plates could begin with AB and have all letters and digits distinct?

Proof. 24 choices for the third letter, 23 for the fourth letter. Then 10, 9, 8 choices for the 1st, 2nd, 3rd digits. So: $24 \cdot 23 \cdot 10 \cdot 9 \cdot 8 = 397440$. \square

2.15 Exercise 15

A combination lock requires three selections of numbers, each from 1 through 30.

2.15.1 (a)

How many different combinations are possible?

Proof. $30^3 = 27000$ \square

2.15.2 (b)

Suppose the locks are constructed in such a way that no number may be used twice. How many different combinations are possible?

Proof. $30 \cdot 29 \cdot 28 = 24360$ \square

2.16 Exercise 16

2.16.1 (a)

How many integers are there from 10 through 99?

Proof. Two solutions:

(i) By the multiplication rule, the number of integers from 10 through 99 = (the number of ways to pick the first digit) \times (the number of ways to pick the second digit) = $9 \cdot 10 = 90$.

(ii) By Theorem 9.1.1, the number of integers from 10 through 99 = $90 - 10 + 1 = 90$. \square

2.16.2 (b)

How many odd integers are there from 10 through 99?

Proof. Because odd integers end in 1, 3, 5, 7, or 9, the number of odd integers from 10 through 99 = (the number of ways to pick the first digit) \times (the number of ways to pick the second digit) = $9 \cdot 5 = 45$.

An alternative solution uses the listing method shown in the solution for Example 9.1.4. \square

2.16.3 (c)

How many integers from 10 through 99 have distinct digits?

Proof. (the number of ways to pick the first digit) \times (the number of ways to pick the second digit) $= 9 \cdot 9 = 81$.

Another solution is to start with the solution to part (a), which is 90, and subtract the number of integers that have the same digit repeated, of which there are 9: 11, 22, 33, 44, 55, 66, 77, 88, 99. \square

2.16.4 (d)

How many odd integers from 10 through 99 have distinct digits?

Proof. (the number of ways to pick the second digit) \times (the number of ways to pick the first digit) $= 5 \cdot 8 = 40$. Here the second digit can be 1,3,5,7,9; and the first digit cannot be 0 or the same as the first digit. \square

2.16.5 (e)

What is the probability that a randomly chosen two-digit integer has distinct digits? has distinct digits and is odd?

Proof. $81/90 = 9/10$, $40/90 = 4/9$ \square

2.17 Exercise 17

2.17.1 (a)

How many integers are there from 1000 through 9999?

Proof. By Theorem 9.1.1, $9999 - 1000 + 1 = 9000$. Another solution: there are 9 choices for the first digit (cannot be 0), and 10 choices for the other 3 digits, so $9 \cdot 10 \cdot 10 \cdot 10 = 9000$. \square

2.17.2 (b)

How many odd integers are there from 1000 through 9999?

Proof. 4500.

One solution is that it's half of the answer to part (a), since half the integers are even, and the other odd.

Another solution: there are 9 choices for the first digit (cannot be 0), 10 choices for the second and third digits, and 5 choices for the last (1,3,5,7,9), therefore $9 \cdot 10 \cdot 10 \cdot 5 = 4500$. \square

2.17.3 (c)

How many integers from 1000 through 9999 have distinct digits?

Proof. There are 9 choices for the first digit (cannot be 0). There are 9 choices for the second digit (cannot be the same as the first digit, but can be 0). Then 8 and 7 choices for the third and fourth digits. So there are $9 \cdot 9 \cdot 8 \cdot 7 = 4536$ such integers. \square

2.17.4 (d)

How many odd integers from 1000 through 9999 have distinct digits?

Proof. There are 5 choices for the last digit (1, 3, 5, 7, 9). There are 8 choices for the first digit (cannot be the same as the last digit, and cannot be 0). Then 8 choices for the second digit (cannot be the same as the first or the last, but can be 0) and 7 choices for the third digit. So there are $5 \cdot 8 \cdot 8 \cdot 7 = 2240$ such integers. \square


2.17.5 (e)

What is the probability that a randomly chosen four-digit integer has distinct digits?
has distinct digits and is odd?

Proof. $4536/9000$, $2240/9000$ \square

2.18 Exercise 18

The following diagram shows the keypad for an automatic teller machine. As you can see, the same sequence of keys represents a variety of different PINs. For instance, 2133, AZDE, and BQ3F are all keyed in exactly the same way.



2.18.1 (a)

How many different PINs are represented by the same sequence of keys as 2133?

Proof. Let step 1 be to choose either the number 2 or one of the letters corresponding to the number 2 on the keypad, let step 2 be to choose either the number 1 or one of

the letters corresponding to the number 1 on the keypad, and let steps 3 and 4 be to choose either the number 3 or one of the letters corresponding to the number 3 on the keypad. There are 4 ways to perform step 1, 3 ways to perform step 2, and 4 ways to perform each of steps 3 and 4. So by the multiplication rule, there are $4 \cdot 3 \cdot 4 \cdot 4 = 192$ ways to perform the entire operation. Thus there are 192 different PINs that are keyed the same as 2133. Note that on a computer keyboard, these PINs would not be keyed the same way. \square

2.18.2 (b)

How many different PINs are represented by the same sequence of keys as 5031?

Proof. $4 \cdot 1 \cdot 4 \cdot 3 = 48$ \square

2.18.3 (c)


How many different numeric sequences on the machine contain no repeated digit?

Proof. $10 \cdot 9 \cdot 8 \cdot 7 = 5040$ \square

2.19 Exercise 19

Three officers - a president, a treasurer, and a secretary - are to be chosen from among four people: Ann, Bob, Cyd, and Dan. Suppose that Bob is not qualified to be treasurer and Cyd's other commitments make it impossible for her to be secretary. How many ways can the officers be chosen? Can the multiplication rule be used to solve this problem?

Proof. There are 14 different paths from “root” to “leaf” of this possibility tree, and so there are 14 ways the officers can be chosen. Because $14 = 2 \cdot 7$, reordering the steps will not make it possible to use the multiplication rule alone to solve this problem.



2.20 Exercise 20

Modify Example 9.2.2 by supposing that a PIN must not begin with any of the letters A-M and must end with a digit. Continue to assume that no symbol may be used more than once and that the total number of PINs is to be determined.

2.20.1 (a)

Find the error in the following “solution.” “Constructing a PIN is a four-step process.

Step 1: Choose the left-most symbol.

Step 2: Choose the second symbol from the left.

Step 3: Choose the third symbol from the left.

Step 4: Choose the right-most symbol.

Because none of the thirteen letters from A through M may be chosen in step 1, there are $36 - 13 = 23$ ways to perform step 1. There are 35 ways to perform step 2 and 34 ways to perform step 3 because previously used symbols may not be used. Since the symbol chosen in step 4 must be a previously unused digit, there are $10 - 3 = 7$ ways to perform step 4. Thus there are $23 \cdot 35 \cdot 34 \cdot 7 = 191,590$ different PINs that satisfy the given conditions.”

Proof. The number of ways to perform step 4 is not constant; it depends on how the previous steps were performed. For instance, if 3 digits had been chosen in steps 1-3, then there would be $10 - 3 = 7$ ways to perform step 4, but if 3 letters had been chosen in steps 1-3, then there would be 10 ways to perform step 4. \square

2.20.2 (b)

Reorder steps 1-4 in part (a) as follows: **Step 1:** Choose the right-most symbol. **Step 2:** Choose the left-most symbol. **Step 3:** Choose the second symbol from the left. **Step 4:** Choose the third symbol from the left. Use the multiplication rule to find the number of PINs that satisfy the given conditions.

Proof. $10 \cdot 22 \cdot 34 \cdot 33 = 246840$ \square

2.21 Exercise 21

Suppose A is a set with m elements and B is a set with n elements.

2.21.1 (a)

How many relations are there from A to B ? Explain.

Proof. 2^{mn} . There are mn pairs in $A \times B$. A relation is simply any subset of $A \times B$. So the number of relations from A to B is the same as the size of the power set of $A \times B$, which is 2^{mn} . \square

2.21.2 (b)

How many functions are there from A to B ? Explain.

Proof. n^m . For each input $a \in A$ there are n possible outputs $b \in B$. Since there are m inputs, by the multiplication rule there are $n \cdot n \cdots n = n^m$ functions. \square

2.21.3 (c)

What fraction of the relations from A to B are functions?

Proof. $n^m/2^{mn}$ \square

2.22 Exercise 22

2.22.1 (a)

How many functions are there from a set with three elements to a set with four elements?

Proof. The answer is $4 \cdot 4 \cdot 4 = 4^3 = 64$. Imagine creating a function from a 3-element set to a 4-element set as a three-step process: Step 1 is to send the first element of the 3-element set to an element of the 4-element set (there are four ways to perform this step); step 2 is to send the second element of the 3-element set to an element of the 4-element set (there are also four ways to perform this step); and step 3 is to send the third element of the 3-element set to an element of the 4-element set (there are four ways to perform this step). Thus the entire process can be performed in $4 \cdot 4 \cdot 4$ different ways. \square

2.22.2 (b)

How many functions are there from a set with five elements to a set with two elements?

Proof. $2^5 = 32$ \square

2.22.3 (c)

How many functions are there from a set with m elements to a set with n elements, where m and n are positive integers?

Proof. n^m . \square

2.23 Exercise 23

In Section 2.5 we showed how integers can be represented by strings of 0's and 1's inside a digital computer. In fact, through various coding schemes, strings of 0's and 1's can be used to represent all kinds of symbols. One commonly used code is the Extended Binary-Coded Decimal Interchange Code (EBCDIC) in which each symbol has an 8-bit representation. How many distinct symbols can be represented by this code?

Proof. $2^8 = 256$

□

In each of 24 – 28, determine how many times the innermost loop will be iterated when the algorithm segment is implemented and run. (assume that m, n, p, a, b, c , and d are all positive integers.)

2.24 Exercise 24

```
for  $i := 1$  to 30
  for  $j := 1$  to 15
    [Statements in body of inner loop.]
    [None contain branching statements that lead outside the loop.]
  next  $j$ 
next  $i$ 
```

Proof. The outer loop is iterated 30 times, and during each iteration of the outer loop there are 15 iterations of the inner loop. Hence, by the multiplication rule, the total number of iterations of the inner loop is $30 \cdot 15 = 450$. □

2.25 Exercise 25

```
for  $j := 1$  to  $m$ 
  for  $k := 1$  to  $n$ 
    [Statements in body of inner loop.]
    [None contain branching statements that lead outside the loop.]
  next  $k$ 
next  $j$ 
```

Proof. mn

□

2.26 Exercise 26

```
for  $i := 1$  to  $m$ 
  for  $j := 1$  to  $n$ 
    for  $k := 1$  to  $p$ 
      [Statements in body of inner loop.]
      [None contain branching statements that lead outside the loop.]
    next  $k$ 
  next  $j$ 
next  $i$ 
```

Proof. mnp

□

2.27 Exercise 27

```
for  $i := 5$  to 50
```

```

for  $j := 10$  to 20
    [Statements in body of inner loop.]
    [None contain branching statements that lead outside the loop.]
next  $j$ 
next  $i$ 

```

Proof. The outer loop is iterated $50 - 5 + 1 = 46$ times, and during each iteration of the outer loop there are $20 - 10 + 1 = 11$ iterations of the inner loop. Hence, by the multiplication rule, the total number of iterations of the inner loop is $46 \cdot 11 = 506$. \square

2.28 Exercise 28

Assume $a \leq b$ and $c \leq d$.

```

for  $i := a$  to  $b$ 
    for  $j := c$  to  $d$ 
        [Statements in body of inner loop.]
        [None contain branching statements that lead outside the loop.]
    next  $j$ 
next  $i$ 

```

Proof. $(b - a + 1)(d - c + 1)$ \square

2.29 Exercise 29

Consider the numbers 1 through 99,999 in their ordinary decimal representations. How many contain exactly one of each of the digits 2, 3, 4, and 5?

Hint: An efficient solution is to add leading zeros as needed to make each number five digits long. For instance, write 1 as 00001. Then, instead of choosing digits for the positions, choose positions for the digits. The answer is 720.

Proof. Following the Hint, there are 5 possible positions to place 2, then 4 positions to place 3, 3 positions to place 4, 2 positions to place 5, and 6 choices for the last position (1, 6, 7, 8, 9, 0). Therefore $5 \cdot 4 \cdot 3 \cdot 2 \cdot 6 = 720$. (Notice that there is no issue with the digit 0, since it can be placed in the first position.) \square

2.30 Exercise 30

Let $n = p_1^{k_1} p_2^{k_2} \cdots p_m^{k_m}$ where p_1, p_2, \dots, p_m are distinct prime numbers and k_1, k_2, \dots, k_m are positive integers. How many ways can n be written as a product of two positive integers that have no common factors, assuming the following?

2.30.1 (a)

Order matters (that is, $8 \cdot 15$ and $15 \cdot 8$ are regarded as different).

Proof. To split n into two positive integers a, b that have no common factors, we need to split the set of all the prime powers $\{p_1^{k_1}, \dots, p_m^{k_m}\}$ into two disjoint subsets A, B . Then a is the product of the prime powers in A , and b is the product of those in B .

When one of the subsets is chosen, the other is automatically forced to be what's left. For example, if the set is $p_1^{k_1}, p_2^{k_2}, p_3^{k_3}$ and we choose A to be $\{p_1^{k_1}, p_3^{k_3}\}$ then B has to be $\{p_2^{k_2}\}$.

However, since order matters, choosing A to be $\{p_1^{k_1}, p_3^{k_3}\}$ and B to be $\{p_2^{k_2}\}$ is considered different than choosing A to be $\{p_2^{k_2}\}$ and B to be $\{p_1^{k_1}, p_3^{k_3}\}$.

This means that the number of ways to write n as the product of two positive integers with no common prime factors is equal to the number of ways we can choose A to be a subset of $\{p_1^{k_1}, \dots, p_m^{k_m}\}$. This is a set with m elements, so it has 2^m subsets. So there are 2^m ways to choose A , which is the answer. \square

2.30.2 (b)

Order does not matter (that is, $8 \cdot 15$ and $15 \cdot 8$ are regarded as the same).

Proof. The solution is the same as in part (a) except the choices where A and B are swapped do not count. For example, choosing A to be $\{p_1^{k_1}, p_3^{k_3}\}$ and B to be $\{p_2^{k_2}\}$ is considered the same as choosing A to be $\{p_2^{k_2}\}$ and B to be $\{p_1^{k_1}, p_3^{k_3}\}$.

So, in this case the answer is half the answer to part (a), namely, $2^m/2 = 2^{m-1}$. \square

2.31 Exercise 31

2.31.1 (a)

If p is a prime number and a is a positive integer, how many distinct positive divisors does p^a have?

Proof. There are $a + 1$ divisors: $1, p, p^2, \dots, p^a$. \square

2.31.2 (b)

If p and q are distinct prime numbers and a and b are positive integers, how many distinct positive divisors does $p^a q^b$ have?

Proof. A divisor is a product of any one of the $a + 1$ numbers listed in part (a) times any one of the $b + 1$ numbers $1, q, q^2, \dots, q^b$. So, by the multiplication rule, there are $(a + 1)(b + 1)$ divisors in all. \square

2.31.3 (c)

If p, q , and r are distinct prime numbers and a, b , and c are positive integers, how many distinct positive divisors does $p^a q^b r^c$ have?

Proof. $(a + 1)(b + 1)(c + 1)$ \square

2.31.4 (d)

If p_1, p_2, \dots, p_m are distinct prime numbers and a_1, a_2, \dots, a_m are positive integers, how many distinct positive divisors does $p_1^{a_1} p_2^{a_2} \cdots p_m^{a_m}$ have?

Proof. $(a_1 + 1)(a_2 + 1) \cdots (a_m + 1)$ □

2.31.5 (e)

What is the smallest positive integer with exactly 12 divisors?

Proof. Assume p_1, p_2, \dots, p_m are distinct prime numbers and a_1, a_2, \dots, a_m are positive integers, and $n = p_1^{a_1} p_2^{a_2} \cdots p_m^{a_m}$ is the integer we are looking for. Assume n has exactly 12 divisors.

By part (d), $12 = 2 \cdot 2 \cdot 3$ so $m = 3$ and $2 \cdot 2 \cdot 3 = (a_1 + 1)(a_2 + 1)(a_3 + 1)$ so it follows $a_1 = 1, a_2 = 1, a_3 = 2$.

To make n as small as possible, we can choose the smallest first three primes 2, 3, 5, and since $a_3 = 2$ we can let $p_1 = 3, p_2 = 5, p_3 = 2$. (This way the highest power 2 goes on top of the smallest base prime, 2.)

Therefore $n = p_1^{a_1} p_2^{a_2} p_3^{a_3} = 3^1 \cdot 5^1 \cdot 2^2 = 60$ is the smallest positive integer with exactly 12 divisors: 1, 2, 3, 4, 5, 6, 10, 12, 15, 20, 30, 60. □

2.32 Exercise 32

2.32.1 (a)

How many ways can the letters of the word ALGORITHM be arranged in a row?

Proof. Since the nine letters of the word ALGORITHM are all distinct, there are as many arrangements of these letters in a row as there are permutations of a set with nine elements: $9! = 362,880$. □

2.32.2 (b)

How many ways can the letters of the word ALGORITHM be arranged in a row if A and L must remain together (in order) as a unit?

Proof. In this case there are effectively eight symbols to be permuted (because AL may be regarded as a single symbol). So the number of arrangements is $8! = 40,320$. □

2.32.3 (c)

How many ways can the letters of the word ALGORITHM be arranged in a row if the letters GOR must remain together (in order) as a unit?

Proof. In this case there are effectively seven symbols to be permuted (because GOR may be regarded as a single symbol). So the number of arrangements is $7! = 5040$. □

2.33 Exercise 33

Six people attend the theater together and sit in a row with exactly six seats.

2.33.1 (a)

How many ways can they be seated together in the row?

Proof. $6! = 720$ □

2.33.2 (b)

Suppose one of the six is a doctor who must sit on the aisle in case she is paged. How many ways can the people be seated together in the row with the doctor in an aisle seat?

Proof. Excluding the doctor who must sit on the aisle seat, $5! = 120$ ways. (However the question is a bit ambiguous, are there 2 aisle seats, one on each side of the row? In that case the answer would be 240.) □

2.33.3 (c)

Suppose the six people consist of three married couples and each couple wants to sit together with the older partner on the left. How many ways can the six be seated together in the row?

Proof. The couples can be treated as one person, so $3! = 6$. □

2.34 Exercise 34

Five people are to be seated around a circular table. Two seatings are considered the same if one is a rotation of the other. How many different seatings are possible?

Proof. The same reasoning as in Example 9.2.9 gives an answer of $4! = 24$. □

2.35 Exercise 35

Write all the 2-permutations of $\{W, X, Y, Z\}$.

Proof. $WX, WY, WZ, XW, XY, XZ, YW, YX, YZ, ZW, ZX, ZY$ □

2.36 Exercise 36

Write all the 3-permutations of $\{s, t, u, v\}$.

Proof. There are $P(4, 3) = \frac{4!}{(4-3)!} = 24$ 3-permutations of a 4-element set. So:

$stu, stv, suv, sut, svt, svu, tsu, tsv, tuv, tus, tvs, tvu,$

$ust, usv, utv, uts, uvs, uvt, vst, vsu, vtu, vts, vus, vut.$ □

2.37 Exercise 37

Evaluate the following quantities.

2.37.1 (a)

$$P(6, 4)$$

$$\text{Proof. } P(6, 4) = \frac{6!}{(6-4)!} = \frac{6!}{2!} = \frac{6 \cdot 5 \cdot 4 \cdot 3 \cdot \cancel{2 \cdot 1}}{\cancel{2 \cdot 1}} = 360$$

□

2.37.2 (b)

$$P(6, 6)$$

$$\text{Proof. } P(6, 6) = \frac{6!}{(6-6)!} = \frac{6!}{0!} = 720/1 = 720$$

□

2.37.3 (c)

$$P(6, 3)$$

$$\text{Proof. } P(6, 3) = \frac{6!}{(6-3)!} = \frac{6!}{3!} = \frac{6 \cdot 5 \cdot 4 \cdot \cancel{3 \cdot 2 \cdot 1}}{\cancel{3 \cdot 2 \cdot 1}} = 120$$

□

2.37.4 (d)

$$P(6, 1)$$

$$\text{Proof. } P(6, 1) = \frac{6!}{(6-1)!} = \frac{6!}{5!} = \frac{\cancel{6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}}{\cancel{5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}} = 6$$

□

2.38 Exercise 38

2.38.1 (a)

How many 3-permutations are there of a set of five objects?

$$\text{Proof. } P(5, 3) = \frac{5!}{(5-3)!} = \frac{5!}{2!} = \frac{5 \cdot 4 \cdot 3 \cdot \cancel{2 \cdot 1}}{\cancel{2 \cdot 1}} = 60$$

□

2.38.2 (b)

How many 2-permutations are there of a set of eight objects?

$$\text{Proof. } P(8, 2) = \frac{8!}{(8-2)!} = \frac{8!}{6!} = \frac{\cancel{8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}}{\cancel{6 \cdot 5 \cdot 4 \cdot 3 \cdot 2 \cdot 1}} = 56$$

□

2.39 Exercise 39

2.39.1 (a)

How many ways can three of the letters of the word ALGORITHM be selected and written in a row?

$$\text{Proof. } P(9, 3) = \frac{9!}{(9-3)!} = \frac{9!}{6!} = \frac{9 \cdot 8 \cdot 7 \cdot \cancel{6!}}{\cancel{6!}} = 504 \quad \square$$

2.39.2 (b)

How many ways can six of the letters of the word ALGORITHM be selected and written in a row?

$$\text{Proof. } P(9, 6) = \frac{9!}{(9-6)!} = \frac{9!}{3!} = \frac{9 \cdot 8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot \cancel{3!}}{\cancel{3!}} = 60480 \quad \square$$

2.39.3 (c)

How many ways can six of the letters of the word ALGORITHM be selected and written in a row if the first letter must be A?

$$\text{Proof. } P(8, 5) = \frac{8!}{(8-5)!} = \frac{8!}{3!} = \frac{8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 \cdot \cancel{3!}}{\cancel{3!}} = 6720 \quad \square$$

2.39.4 (d)

How many ways can six of the letters of the word ALGORITHM be selected and written in a row if the first two letters must be OR?

$$\text{Proof. } P(7, 4) = \frac{7!}{(7-4)!} = \frac{7!}{3!} = \frac{7 \cdot 6 \cdot 5 \cdot 4 \cdot \cancel{3!}}{\cancel{3!}} = 840 \quad \square$$

2.40 Exercise 40

Prove that for every integer $n \geq 2$, $P(n+1, 3) = n^3 - n$.

$$\begin{aligned} \text{Proof. } P(n+1, 3) &= \frac{(n+1)!}{(n+1-3)!} = \frac{(n+1)!}{(n-2)!} = \frac{(n+1) \cdot n \cdot (n-1) \cdot \cancel{(n-2)!}}{\cancel{(n-2)!}} \\ &= (n+1)n(n-1) = (n^2-1)n = n^3 - n \quad \square \end{aligned}$$

2.41 Exercise 41

Prove that for every integer $n \geq 2$, $P(n+1, 2) - P(n, 2) = 2P(n, 1)$.

$$\text{Proof. } P(n+1, 2) = \frac{(n+1)!}{(n+1-2)!} = \frac{(n+1)!}{(n-1)!} = \frac{(n+1) \cdot n!}{(n-1)!} = (n+1)P(n, 1)$$

$$P(n, 2) = \frac{n!}{(n-2)!} = \frac{(n-1) \cdot n!}{(n-1) \cdot (n-2)!} = (n-1) \frac{n!}{(n-1)!} = (n-1)P(n, 1)$$

$$\text{So } P(n+1, 2) - P(n, 2) = (n+1)P(n, 1) - (n-1)P(n, 1) = 2P(n, 1). \quad \square$$

2.42 Exercise 42

Prove that for every integer $n \geq 3$, $P(n+1, 3) - P(n, 3) = 3P(n, 2)$.

$$\text{Proof. } P(n+1, 3) = \frac{(n+1)!}{(n+1-3)!} = \frac{(n+1)!}{(n-2)!} = \frac{(n+1) \cdot n!}{(n-2)!} = (n+1)P(n, 2)$$

$$P(n, 3) = \frac{n!}{(n-3)!} = \frac{(n-2) \cdot n!}{(n-2) \cdot (n-3)!} = (n-2) \frac{n!}{(n-2)!} = (n-2)P(n, 2)$$

$$\text{So } P(n+1, 3) - P(n, 3) = (n+1)P(n, 2) - (n-2)P(n, 2) = 3P(n, 2). \quad \square$$

2.43 Exercise 43

Prove that for every integer $n \geq 2$, $P(n, n) = P(n, n-1)$.

$$\text{Proof. } P(n, n) = \frac{n!}{(n-n)!} = \frac{n!}{0!} = \frac{n!}{1} = \frac{n!}{1!} = \frac{n!}{(n-(n-1))!} = P(n, n-1) \quad \square$$

2.44 Exercise 44

Prove Theorem 9.2.1 by mathematical induction.

Proof. Let $P(k)$ be the statement: “If an operation consists of k steps and the first step can be performed in n_1 ways, the second step can be performed in n_2 ways [regardless of how the first step was performed], ..., the k th step can be performed in n_k ways [regardless of how the preceding steps were performed], then the entire operation can be performed in $n_1 n_2 \cdots n_k$ ways.”

Show that $P(1)$ is true: Assume an operation consists of 1 step, and the first step can be performed in n_1 ways. Then the whole operation can be performed in n_1 ways, so $P(1)$ is true.

Show that for any integer $k \geq 1$ if $P(k)$ is true then $P(k+1)$ is true: Assume an operation consists of $k+1$ steps, and assume steps $1, 2, 3, \dots, k+1$ can be performed in n_1, n_2, \dots, n_{k+1} ways, respectively, regardless of how preceding steps are performed. Assume that the first k steps can be performed in $n_1 n_2 \cdots n_k$ ways. [This is the inductive hypothesis.]

Since the $k+1$ st step can be performed in n_{k+1} ways regardless of how the preceding k steps were performed, for each way of performing the preceding k steps, there are n_{k+1} ways to perform step $k+1$. Thus there are

$$\underbrace{n_{k+1} + n_{k+1} + \cdots + n_{k+1}}_{n_1 n_2 \cdots n_k \text{ times}}$$

ways to perform the whole task, where the sum has $n_1 n_2 \cdots n_k$ terms. And the sum equals

$$n_{k+1} \cdot (n_1 n_2 \cdots n_k) = n_1 n_2 \cdots n_k n_{k+1}$$

[as was to be shown.] □

2.45 Exercise 45

Prove Theorem 9.2.2 by mathematical induction.


Proof. Let $P(n)$ be the statement “For any integer n with $n \geq 1$, the number of permutations of a set with n elements is $n!$.”

Show that $P(1)$ is true: There is only 1 permutation of a set with 1 element, and $1! = 1$, so $P(1)$ is true.

Show that for any integer $k \geq 1$ if $P(k)$ is true then $P(k+1)$ is true: Assume $k \geq 1$ and assume that the number of permutations of any set with k elements is $k!$. [This is the inductive hypothesis.] Assume $A = \{a_1, \dots, a_{k+1}\}$ is a set with $k+1$ elements.

Consider the subset $A' = \{a_1, \dots, a_k\}$ of A with k elements. By the inductive hypothesis A' has $k!$ permutations. Now we need to find a way to write the permutations of A in terms of the permutations of A' .

Every permutation of A can be thought of as taking a permutation of A' and then inserting a_{k+1} into a position in that permutation. For example, if $a_3, a_1, a_{k-2}, \dots, a_k, a_2, a_{k-1}$ is a permutation of A' , then there are $k+1$ positions in which a_{k+1} can be inserted to obtain a permutation of A :



So for each permutation of A' , there are $k+1$ permutations of A . Therefore the number of permutations of A is $k! \cdot (k+1) = (k+1)!$. □

2.46 Exercise 46

Prove Theorem 9.2.3 by mathematical induction.

Proof. Let $Q(n)$ be the statement: “for every integer r with $1 \leq r \leq n$, $P(n, r) = n(n-1)(n-2) \cdots (n-r+1)$.”

Show that $Q(1)$ is true: There is only one 1-permutation of a set of 1 element, and when $n = r = 1$ the formula $n(n-1)(n-2) \cdots (n-r+1)$ is equal to 1. Therefore $Q(1)$ is true.

Show that for any integer $k \geq 1$ if $Q(k)$ is true then $Q(k+1)$ is true: Assume $k \geq 1$ and assume for every integer r with $1 \leq r \leq k$, $P(k, r) = k(k-1)(k-2) \cdots (k-r+1)$. [This is the inductive hypothesis.]

We want to show that for every integer r with $1 \leq r \leq k+1$, $P(k+1, r) = (k+1)k(k-1) \cdots (k-r+2)$.

When $r = 1$, there are $k+1$ ways to choose 1 element from a set of $k+1$ elements. So $P(k+1, 1) = k+1$, and the formula $(k+1)k(k-1) \cdots (k-r+2)$ for $r = 1$ becomes $(k+1)$, therefore the formula holds.

Now assume $2 \leq r \leq k+1$. We can think of choosing r elements from a set of $k+1$ elements as follows: first choosing 1 element out of $k+1$ elements, then choosing $r-1$ elements from the remaining k elements.

There are $k+1$ ways to choose the first element. Then by the inductive hypothesis there are $P(k, r-1) = k(k-1)(k-2) \cdots (k-(r-1)+1) = k(k-1)(k-2) \cdots (k-r+2)$ ways to choose $r-1$ elements from the remaining k elements. Then by the multiplication rule, $P(k+1, r) = (k+1) \cdot k(k-1)(k-2) \cdots (k-r+2)$, [as was to be shown.] \square

2.47 Exercise 47

A permutation on a set can be regarded as a function from the set to itself. For instance, one permutation of $\{1, 2, 3, 4\}$ is 2341. It can be identified with the function that sends each position number to the number occupying that position. Since position 1 is occupied by 2, 1 is sent to 2 or $1 \rightarrow 2$; since position 2 is occupied by 3, 2 is sent to 3 or $2 \rightarrow 3$; and so forth. The entire permutation can be written using arrows as follows:

$$\begin{array}{cccc} 1 & 2 & 3 & 4 \\ \downarrow & \downarrow & \downarrow & \downarrow \\ 2 & 3 & 4 & 1 \end{array}$$

2.47.1 (a)

Use arrows to write each of the six permutations of $\{1, 2, 3\}$.

$$\text{Proof.} \quad \begin{array}{ccc|ccc|ccc|ccc|ccc|ccc} 1 & 2 & 3 & 1 & 2 & 3 & 1 & 2 & 3 & 1 & 2 & 3 & 1 & 2 & 3 & 1 & 2 & 3 \\ \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\ 1 & 2 & 3 & 1 & 3 & 2 & 2 & 1 & 3 & 2 & 3 & 1 & 3 & 1 & 2 & 3 & 2 & 1 \end{array}$$

\square

2.47.2 (b)

Use arrows to write each of the permutations of $\{1, 2, 3, 4\}$ that keep 2 and 4 fixed.

$$\text{Proof.} \quad \begin{array}{cccc|cccc} 1 & 2 & 3 & 4 & 1 & 2 & 3 & 4 \\ \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\ 1 & 2 & 3 & 4 & 3 & 2 & 1 & 4 \end{array}$$

\square

2.47.3 (c)

Which permutations of $\{1, 2, 3\}$ keep no elements fixed?

$$\begin{array}{c} \text{Proof.} \end{array} \begin{array}{ccc|ccc} 1 & 2 & 3 & 1 & 2 & 3 \\ \downarrow & \downarrow & \downarrow & \downarrow & \downarrow & \downarrow \\ 2 & 3 & 1 & 3 & 1 & 2 \end{array}$$

□

2.47.4 (d)

Use arrows to write all permutations of $\{1, 2, 3, 4\}$ that keep no elements fixed.

$$\begin{array}{c} \text{Proof.} \end{array} \begin{array}{cccc|cccc|cccc|cccc|cccc|cccc} 1 & 2 & 3 & 4 & 1 & 2 & 3 & 4 & 1 & 2 & 3 & 4 & 1 & 2 & 3 & 4 & 1 & 2 & 3 & 4 & 1 & 2 & 3 & 4 & 1 & 2 & 3 & 4 \\ \downarrow & \downarrow \\ 2 & 1 & 4 & 3 & 2 & 3 & 4 & 1 & 2 & 4 & 1 & 3 & 3 & 1 & 4 & 2 & 3 & 4 & 1 & 2 & 3 & 4 & 2 & 1 & 3 & 4 & 2 & 1 \\ \\ 1 & 2 & 3 & 4 & 1 & 2 & 3 & 4 & 1 & 2 & 3 & 4 & 1 & 2 & 3 & 4 & 1 & 2 & 3 & 4 & 1 & 2 & 3 & 4 & 1 & 2 & 3 & 4 \\ \downarrow & \downarrow \\ 4 & 1 & 2 & 3 & 4 & 3 & 1 & 2 & 4 & 3 & 2 & 1 & 4 & 3 & 2 & 1 & 4 & 3 & 2 & 1 & 4 & 3 & 2 & 1 & 4 & 3 & 2 & 1 \end{array}$$

□


3 Exercise Set 9.3

3.1 Exercise 1

3.1.1 (a)

How many bit strings consist of from one through four digits? (Strings of different lengths are considered distinct. Thus 10 and 0010 are distinct strings.)

Proof. Think of creating a bit string with n bits as an n -step process where a general step k is to place either a 0 or a 1 in the k th position. Since there are two ways to do this for each position, by the multiplication rule, the number of bit strings of length k is 2^k . Now the set of all bit strings consisting of from 1 through 4 bits can be broken into four disjoint subsets:



Applying the addition rule to the figure shows that there are $2 + 2^2 + 2^3 + 2^4 = 30$ bit strings consisting of from one through four bits. \square

3.1.2 (b)

How many bit strings consist of from five through eight digits?

Proof. By reasoning similar to that of part (a), there are $2^5 + 2^6 + 2^7 + 2^8 = 480$ bit strings of from five through eight bits. \square

3.2 Exercise 2

3.2.1 (a)

How many strings of hexadecimal digits consist of from one through three digits? (Recall that hexadecimal numbers are constructed using the 16 digits 0, 1, 2, 3, 4, 5, 6, 7, 8, 9, A, B, C, D, E, F.)

Proof. $16^1 + 16^2 + 16^3 = 16 + 256 + 4096 = 4368$ \square

3.2.2 (b)

How many strings of hexadecimal digits consist of from two through five digits?

Proof. $16^2 + 16^3 + 16^4 + 16^5 = 256 + 4096 + 65536 + 1048576 = 1118464$ \square

3.3 Exercise 3

3.3.1 (a)

How many integers from 1 through 999 do not have any repeated digits?

Proof. The set of integers from 1 through 999 with no repeated digit can be broken into three disjoint subsets: those from 1 through 9, those from 1 through 99, and those from 100 through 999. Now constructing an integer from 100 through 999 with no repeated digit can be thought of as a three-step process.

Step 1: Choose a digit for the left-most position (where there are 9 choices because 0 cannot be chosen).

Step 2: Choose a digit for the middle position (where there are also 9 choices because the digit in the left-most position cannot be reused but 0 can be used).

Step 3: Choose a digit for the right-most position (where there are 8 choices because neither of the other two digits can be reused).

Thus there are $9 \cdot 9 \cdot 8$ integers from 100 through 999 with no repeated digit. Similar reasoning shows that there are $9 \cdot 9$ integers from 10 through 99 with no repeated digit. Finally, there are clearly 9 integers from 1 through 9 with no repeated digit. Hence, by

the addition rule, the number of integers from 1 through 999 with no repeated digit is $9 + 9 \cdot 9 + 9 \cdot 9 \cdot 8 = 738$. \square

3.3.2 (b)

How many integers from 1 through 999 have at least one repeated digit?

Proof. Let

x = number of integers from 1 through 999 with at least one repeated digit,

y = total number of integers from 1 through 999,

z = number of integers from 1 through 999 with no repeated digits.

Then $x = y - z = 999 - 738 = 261$ \square

3.3.3 (c)


What is the probability that an integer chosen at random from 1 through 999 has at least one repeated digit?

Proof. The probability that an integer chosen at random has at least one repeated digit is $261/999 \approx 26.1\%$. \square

3.4 Exercise 4

How many arrangements in a row of no more than three letters can be formed using the letters of the word NETWORK (with no repetitions allowed)?

Proof. Use the multiplication rule to count the elements in each of the three sets containing 1, 2, and 3 letters, respectively. Then, because these sets are disjoint, use the addition rule to compute the total number of elements in the three sets taken together.



Applying the addition rule to the figure above shows that there are $7 + 7 \cdot 6 + 7 \cdot 6 \cdot 5 = 259$ arrangements of three letters of the word NETWORK if repetition of letters is not permitted. \square

3.5 Exercise 5

3.5.1 (a)

How many five-digit integers (integers from 10,000 through 99,999) are divisible by 5?

Proof. Let

x = number of integers from 10,000 through 99,999 that are divisible by 5,

y = number of integers from 1 through 99,999 that are divisible by 5 = $100,000/5 - 1 = 20,000 - 1 = 19,999$,

z = number of integers from 1 through 9999 that are divisible by 5 = $10,000/5 - 1 = 2000 - 1 = 1999$.

Then $x = y - z = 19,999 - 1999 = 18,000$. □

3.5.2 (b)

What is the probability that a five-digit integer chosen at random is divisible by 5?

Proof. There are $99,999 - 10,000 + 1 = 90,000$ five-digit integers. So: $18,000/90,000 = 1/5 = 20\%$. □

3.6 Exercise 6

In a certain state, all license plates consist of from four to six symbols chosen from the 26 uppercase letters of the Roman alphabet together with the ten digits 0–9.

3.6.1 (a)

How many license plates are possible if repetition of symbols is allowed?

Proof. In this exercise the 26 letters in the alphabet plus the 10 digits give a total of 36 symbols that can be used on a license plate.

Imagine constructing a license plate with 4 symbols as a four-step process: [step 1](#) is to fill in the first symbol, [step 2](#) is to fill in the second symbol, [step 3](#) is to fill in the third symbol, and [step 4](#) is to fill in the fourth symbol. Because any one of the 36 symbols can be used in each step, by the multiplication rule, the number of license plates that use four symbols is 36^4 . Similarly, the number that use 5 symbols is 36^5 , and the number that use six symbols is 36^6 . Thus because license plates can have anywhere from 4 to 6 symbols, the total number of plates with repeated symbols allowed is $36^4 + 36^5 + 36^6 = 2,238,928,128$. □

3.6.2 (b)

How many license plates do not contain any repeated symbols?

Proof. When repetition is not allowed, the number of license plates that use four symbols is $36 \cdot 35 \cdot 34 \cdot 33$. The reason is that in the second step the symbol used in the first step cannot be used, so there are only 35 choices for the second step. In the third step, neither of the symbols used in the first two steps can be used, and so there are only 34 choices for the third step. And in the fourth step, none of the symbols used in the first three steps can be used, and so there are only 33 choices for the fourth step. Similarly, the number of license plates that use 5 symbols is $36 \cdot 35 \cdot 34 \cdot 33 \cdot 32$, and the number that use six symbols is $36 \cdot 35 \cdot 34 \cdot 33 \cdot 32 \cdot 31$. Thus the total number of license plates is $36 \cdot 35 \cdot 34 \cdot 33 + 36 \cdot 35 \cdot 34 \cdot 33 \cdot 32 + 36 \cdot 35 \cdot 34 \cdot 33 \cdot 32 \cdot 31 = 1,449,063,000$. \square

3.6.3 (c)

How many license plates have at least one repeated symbol?

Proof. Consider two sets: the set of plates with repetition not allowed and the set of plates that have a repeated symbol. Note that these two sets have no elements in common, and that since every license plate either has a repeated symbol or does not have a repeated symbol, every license plate considered in part (a) is in one of the two sets. In other words, the set of all license plates with repetition allowed is composed of two disjoint subsets: the set of plates with repetition not allowed and the set of plates that have a repeated symbol. Thus, by the difference rule, the number of license plates with a repeated symbol is the difference between the number of plates with repetition allowed minus the number of plates with repetition not allowed: $2,238,928,128 - 1,449,063,000 = 789,865,128$. \square

3.6.4 (d)

What is the probability that a license plate chosen at random has at least one repeated symbol?

Proof. The probability that a license plate chosen at random has at least one repeated symbol is $789,865,128 / 2,238,928,128 \approx 35.3\%$. \square

3.7 Exercise 7

At a certain company, passwords must be from 3–5 symbols long and composed from the 26 uppercase letters of the Roman alphabet, the ten digits 0–9, and the 14 symbols !, @, #, \$, %, ^, &, *, (,), -, +, {, and }.

3.7.1 (a)

How many passwords are possible if repetition of symbols is allowed?

Proof. The 26 letters in the alphabet plus the 10 digits plus the 14 special characters give a total of 50 symbols that can be used. By the multiplication rule, the number of passwords with 3, 4, and 5 symbols is 50^3 , 50^4 , and 50^5 . Since the sets consisting of these passwords are disjoint, by the addition rule, the number of passwords is

$$50^3 + 50^4 + 50^5 = 318,875,000. \quad \square$$

3.7.2 (b)

How many passwords contain no repeated symbols?

Proof. There are $50 \cdot 49 \cdot 48 = 117600$ passwords of length 3 with no repeated symbols.

There are $50 \cdot 49 \cdot 48 \cdot 47 = 5527200$ passwords of length 4 with no repeated symbols.

There are $50 \cdot 49 \cdot 48 \cdot 47 \cdot 46 = 254251200$ passwords of length 5 with no repeated symbols.

So by the addition rule, the total is $254251200 + 5527200 + 117600 = 259,896,000. \quad \square$

3.7.3 (c)

How many passwords have at least one repeated symbol?

Proof. $318,875,000 - 259,896,000 = 58,979,000 \quad \square$

3.7.4 (d)

What is the probability that a password chosen at random has at least one repeated symbol?

Proof. $58,979,000/318,875,000 \approx 18.5\% \quad \square$

3.8 Exercise 8

In a certain country license plates consist of zero or one digit followed by four or five uppercase letters from the Roman alphabet.

3.8.1 (a)

How many different license plates can the country produce?

Hint: One approach is to divide the license plates into four groups depending on the number of digits and letters they contain. Another approach is to consider creating a license plate as a two-step process: step 1: either choose one digit or do not choose a digit; and step 2: choose 4 or 5 letters.

Proof. There are four groups: 1. No digit, 4 letters, 2. One digit, 4 letters, 3. No digit, 5 letters, 4. One digit, 5 letters.

First group has 26^4 possibilities. Second group has $10 \cdot 26^4$ possibilities.

Third group has 26^5 possibilities. Fourth group has $10 \cdot 26^5$ possibilities.

In total: $26^4 + 10 \cdot 26^4 + 26^5 + 10 \cdot 26^5 = 11(26^4 + 26^5) = 11 \cdot 26^4(1 + 26) = 11 \cdot 26^4 \cdot 27 = 135,721,872 \quad \square$

3.8.2 (b)

How many license plates have no repeated letter?

Proof. Choosing 4 non-repeated letters, no digit: $26 \cdot 25 \cdot 24 \cdot 23 = 358,800$, with one digit: $10 \cdot 358,800 = 3,588,000$

Choosing 5 non-repeated letters, no digit: $26 \cdot 25 \cdot 24 \cdot 23 \cdot 22 = 7,893,600$, with one digit: $10 \cdot 7,893,600 = 78,936,000$

Total: $358,800 + 3,588,000 + 7,893,600 + 78,936,000 = 90,776,400$ □

3.8.3 (c)

How many license plates have at least one repeated letter?

Proof. $135,721,872 - 90,776,400 = 44,945,472$ □

3.8.4 (d)

What is the probability that a license plate has a repeated letter?

Proof. $44,945,472/135,721,872 \approx 33.11\%$ □

3.9 Exercise 9

3.9.1 (a)

Consider the following algorithm segment:

```

for  $i := 1$  to 4
  for  $j := 1$  to  $i$ 
    [Statements in body of inner loop.]
    [None contain branching statements that lead outside the loop.]
  next  $j$ 
next  $i$ 

```

How many times will the inner loop be iterated when the algorithm is implemented and run?

Proof. Each column of the table below corresponds to a pair of values of i and j for which the inner loop will be iterated.

i	1	2	→	3	→	→	4	→	→	→
j	1	1	2	1	2	3	1	2	3	4
	1		2		3			4		

Since there are $1 + 2 + 3 + 4 = 10$ columns, the inner loop will be iterated ten times. □

3.9.2 (b)

Let n be a positive integer, and consider the following algorithm segment:

```
for  $i := 1$  to  $n$ 
  for  $j := 1$  to  $i$ 
    [Statements in body of inner loop.]
    [None contain branching statements that lead outside the loop.]
  next  $j$ 
next  $i$ 
```

How many times will the inner loop be iterated when the algorithm is implemented and run?

Proof. $1 + 2 + \cdots + n = n(n + 1)/2$

□

3.10 Exercise 10

A calculator has an eight-digit display and a decimal point that is located at the extreme right of the number displayed, or at the extreme left, or between any pair of digits. The calculator can also display a minus sign at the extreme left of the number. How many distinct numbers can the calculator display? (Note that certain numbers are equal, such as 1.9, 1.90, and 01.900, and should, therefore, not be counted twice.)

Proof. Let's solve this problem for a 1-digit display, then 2-digit, and so on. Also, let's first solve the problem only for nonnegative numbers.

1-digit display: First let's consider nonnegative numbers. A number x can be in the range $0 \leq x < 1$ (if the decimal point is on the left) or in the range $1 \leq x < 10$ (if the decimal point is on the right).

For the first range there are 10 possibilities: .0, .1, .2, .3, .4, .5, .6, .7, .8, .9;

for the second range we cannot allow the digit to be 0, otherwise $x < 1$, so there are only 9 possibilities: 1., 2., 3., 4., 5., 6., 7., 8., 9.;

so we have $10^1 + (10^1 - 1) = 19$ nonnegative numbers.

2-digit display: First let's consider nonnegative numbers. A number x can be in the range $0 \leq x < 1$ (if the decimal point is on the left) or in the range $1 \leq x < 10$ (if the decimal point is in the middle) or in the range $10 \leq x < 100$ (if the decimal point is on the right).

For the first range there are 10^2 possibilities: .00, .01, ..., .09, .10, .11, ..., .99;

for the second range we cannot allow the left-most digit to be 0, otherwise $x < 1$, so there are only 90 possibilities: 1.0, 1.1, 1.2, ..., 9.9;

for the third range there are also 90 possibilities: 10., 11., 12., ..., 99. because, similarly, the first digit cannot be zero; so we have $10^2 + 10^1 \cdot (10^1 - 1) + 10^1 \cdot (10^1 - 1) = 100 + 90 + 90 = 280$ nonnegative numbers.

3-digit display: A number x can be in one of the ranges $0 \leq x < 1$ or $1 \leq x < 10$ or $10 \leq x < 100$ or $100 \leq x < 1000$.

Similar to the above, we have $10^3 + 10^2 \cdot (10^1 - 1) + 10^2 \cdot (10^1 - 1) + 10^2 \cdot (10^1 - 1) = 1000 + 900 + 900 + 900 = 3700$ nonnegative numbers.

So the general formula for n digits is: $10^n + n \cdot 10^{n-1} \cdot 9$. For an $n = 8$ -digit display, there are $10^8 + 8 \cdot 10^7 \cdot 9 = 100,000,000 + 10,000,000 \cdot 72 = 820,000,000$ unique nonnegative numbers.

Conclusion: 820,000,000 nonnegative and 820,000,000 nonpositive numbers (obtained by putting the negative sign in front of them) gives 1,640,000,000 numbers. However 0 and -0 are the same, so it's 1,639,999,999 numbers. \square

3.11 Exercise 11

3.11.1 (a)

How many ways can the letters of the word QUICK be arranged in a row?

Proof. The answer is the number of permutations of the five letters in QUICK, which equals $5! = 120$. \square

3.11.2 (b)

How many ways can the letters of the word QUICK be arranged in a row if the Q and the U must remain next to each other in the order QU?

Proof. Because \boxed{QU} (in order) is to be considered as a single unit, the answer is the number of permutations of the four symbols \boxed{QU} , I, C, K. This is $4! = 24$. \square

3.11.3 (c)

How many ways can the letters of the word QUICK be arranged in a row if the letters QU must remain together but may be in either the order QU or the order UQ?

Proof. By part (b), there are $4!$ arrangements of \boxed{QU} , I, C, K. Similarly, there are $4!$ arrangements of \boxed{UQ} , I, C, K. Therefore, by the addition rule, there are $4! + 4! = 48$ arrangements in all. \square

3.12 Exercise 12

3.12.1 (a)

How many ways can the letters of the word THEORY be arranged in a row?

Proof. $6! = 720$ \square

3.12.2 (b)

How many ways can the letters of the word THEORY be arranged in a row if T and H must remain next to each other as either TH or HT?

Proof. $5! + 5! = 240$ □

3.13 Exercise 13

A group of eight people are attending the movies together.

3.13.1 (a)

Two of the eight insist on sitting side-by-side. In how many ways can the eight be seated together in a row?

Proof. Let

x = the number of ways to place eight people in a row keeping A and B together,

y = the number of ways to arrange \boxed{AB} , C, D, E, F, G, H,

z = the number of ways to arrange \boxed{BA} , C, D, E, F, G, H.

Then $x = y + z = 7! + 7! = 5040 + 5040 = 10080$. □

3.13.2 (b)

Two of the people do not like each other and do not want to sit side-by-side. Now how many ways can the eight be seated together in a row?

Proof. Let

x = the number of ways to place eight people in a row keeping A and B apart,

y = the number of ways to place eight people in a row,

z = the number of ways to place eight people in a row keeping A and B together.

By part (a) $z = 10080$. So $x = y - z = 8! - 10080 = 40320 - 10080 = 30240$. □

3.14 Exercise 14

An early compiler recognized variable names according to the following rules: Numeric variable names had to begin with a letter, and then the letter could be followed by another letter or a digit or by nothing at all. String variable names had to begin with the symbol \$ followed by a letter, which could then be followed by another letter or a digit or by nothing at all. How many distinct variable names were recognized by this compiler?

Proof. the number of variable names = the number of numeric variable names + the number of string variable names = $(26 + 26 \cdot 36) + (26 + 26 \cdot 36) = 1924$. □

3.15 Exercise 15

Identifiers in a certain database language must begin with a letter, and then the letter may be followed by other characters, which can be letters, digits, or underscores (_). However, 82 keywords (all consisting of 15 or fewer characters) are reserved and cannot be used as identifiers. How many identifiers with 30 or fewer characters are possible? (Write the answer using summation notation and evaluate it using a formula from Section 5.2.)

Proof. For the first character there are 26 choices (letters).

For other characters there are 37 choices (letters, digits, or underscore).

So, there are 26 length 1 identifiers, $26 \cdot 37$ length-2 identifiers, ..., and $26 \cdot 37^{29}$ length-30 identifiers. So the number of identifiers with 30 or fewer characters is:

$$26 + 26 \cdot 37 + \cdots + 26 \cdot 37^{29} = 26(1 + 37 + 37^2 + \cdots + 37^{29}) = 26 \cdot \frac{37^{30} - 1}{37 - 1}$$

□

3.16 Exercise 16

3.16.1 (a)

If any seven digits could be used to form a telephone number, how many seven-digit telephone numbers would not have any repeated digits?

Proof. $10 \cdot 9 \cdot 8 \cdot 7 \cdot 6 \cdot 5 \cdot 4 = 604,800$

□

3.16.2 (b)

How many seven-digit telephone numbers would have at least one repeated digit?

Proof. Let x = the number of phone numbers with at least one repeated digit,

y = the total number of phone numbers,

z = the number of phone numbers with no repeated digits,

then $x = y - z = 10^7 - 604,800 = 9,395,200$.

□

3.16.3 (c)

What is the probability that a randomly chosen seven-digit telephone number would have at least one repeated digit?

Proof. $9,395,200/10,000,000 \approx 93.95\%$

□

3.17 Exercise 17

3.17.1 (a)

How many strings of four hexadecimal digits do not have any repeated digits?

Proof. $16 \cdot 15 \cdot 14 \cdot 13 = 43680$

□

3.17.2 (b)

How many strings of four hexadecimal digits have at least one repeated digit?

Proof. $16^4 - 43680 = 65536 - 43680 = 21856$

□

3.17.3 (c)

What is the probability that a randomly chosen string of four hexadecimal digits has at least one repeated digit?

Proof. $21856/65536 \approx 33.34$

□

3.18 Exercise 18

Just as the difference rule gives rise to a formula for the probability of the complement of an event, so the addition and inclusion/exclusion rules give rise to formulas for the probability of the union of mutually disjoint events and for a general union of (not necessarily mutually exclusive) events.

3.18.1 (a)

Prove that for mutually disjoint events A and B , $P(A \cup B) = P(A) + P(B)$.

Proof. Let A and B be mutually disjoint events in a sample space S . By the addition rule, $N(A \cup B) = N(A) + N(B)$. Therefore, by the equally likely probability formula,

$$P(A \cup B) = \frac{N(A \cup B)}{N(S)} = \frac{N(A) + N(B)}{N(S)} = \frac{N(A)}{N(S)} + \frac{N(B)}{N(S)} = P(A) + P(B).$$

□

3.18.2 (b)

Prove that for any events A and B , $P(A \cup B) = P(A) + P(B) - P(A \cap B)$.

Proof. Let A and B be any events in a sample space S . By the inclusion / exclusion rule, $N(A \cup B) = N(A) + N(B) - N(A \cap B)$. Therefore, by the equally likely probability formula,

$$P(A \cup B) = \frac{N(A \cup B)}{N(S)} = \frac{N(A) + N(B) - N(A \cap B)}{N(S)} = \frac{N(A)}{N(S)} + \frac{N(B)}{N(S)} - \frac{N(A \cap B)}{N(S)}$$

$$= P(A) + P(B) - P(A \cap B).$$

□

3.19 Exercise 19

A combination lock requires three selections of numbers, each from 1 through 39. Suppose the lock is constructed in such a way that no number may be used twice in a row but the same number may occur both first and third. For example, 20 13 20 would be acceptable, but 20 20 13 would not. How many different combinations are possible?

Proof. There are 39 choices for the first selection. Then 38 choices for the second selection. Now there are 38 choices for the third selection: it cannot be the same as the second selection, but it can be the same as the first. So by the multiplication rule $39 \cdot 38 \cdot 38$ possible combinations. □

3.20 Exercise 20

3.20.1 (a)

How many integers from 1 through 100,000 contain the digit 6 exactly once?

Proof. Use strings of five digits to represent integers from 1 to 100,000 that contain the digit 6 exactly once. For example, use 00306 to represent 306. Strings of six digits are not needed because 100,000 does not contain a 6. Imagine constructing a five-digit string that contains exactly one 6 as a five-step operation to fill in five positions with five digits:

$\frac{\quad}{1} \quad \frac{\quad}{2} \quad \frac{\quad}{3} \quad \frac{\quad}{4} \quad \frac{\quad}{5} .$

Step 1: Choose one of the five positions for the 6.

Step 2: Choose a digit for the left-most remaining position.

Step 3: Choose a digit for the next remaining position to the right.

Step 4: Choose a digit for the next remaining position to the right.

Step 5: Choose a digit for the right-most position.

Since there are 5 choices for step 1 (any one of the five positions) and 9 choices for each of steps 2–5 (any digit except 6), by the multiplication rule, the number of ways to perform this operation is $5 \cdot 9 \cdot 9 \cdot 9 \cdot 9 = 32,805$. Hence there are 32,805 integers from 1 to 100,000 that contain the digit 6 exactly once. □

3.20.2 (b)

How many integers from 1 through 100,000 contain the digit 6 at least once?

Proof. Let x = the number of integers from 1 through 100,000,

y = the number of integers from 1 through 100,000, that do not contain the digit 6,

z = the number of integers from 1 through 100,000, that contain the digit 6 at least once.

Then $z = x - y = 100,000 - (9^5 + 1) = 40950$. □

3.20.3 (c)

If an integer is chosen at random from 1 through 100,000, what is the probability that it contains two or more occurrences of the digit 6?

Proof. Let x = the number of integers from 1 through 100,000, that contain the digit 6 at least once,

y = the number of integers from 1 through 100,000, that contain the digit 6 exactly once,

z = the number of integers from 1 through 100,000, that contain the digit 6 two or more times.

Then $z = x - y = 40950 - 32805 = 8145$. Then the probability is $8145/100000 = 8.145\%$ □

3.21 Exercise 21

Six new employees, two of whom are married to each other, are to be assigned six desks that are lined up in a row. If the assignment of employees to desks is made randomly, what is the probability that the married couple will have nonadjacent desks? (Hint: The event that the couple have nonadjacent desks is the complement of the event that they have adjacent desks.)

Proof. There are $6! = 720$ possible assignments. The number of assignments where the married couple are adjacent is $2 \cdot 5! = 240$ since we can think of the adjacent married couple as one, and with the remaining 4 employees, there are 5 people to assign desks, but we can also switch the married couple's positions. So $720 - 240 = 480$ assignments have the married couple in nonadjacent desks. The probability is $480/720 = 2/3$. □

3.22 Exercise 22

3.22.1 (a)

Consider strings of length n over the set $\{a, b, c, d\}$. How many such strings contain at least one pair of adjacent characters that are the same?

Proof. Let x = the total number of all strings $= 4^n$,

y = the number of strings that have no pairs of adjacent characters that are the same $= 4 \cdot \underbrace{3 \cdot \dots \cdot 3}_{n-1 \text{ times}} = 4 \cdot 3^{n-1}$,

z = the number of strings contain at least one pair of adjacent characters that are the same $= x - y = 4^n - 4 \cdot 3^{n-1}$. □

3.22.2 (b)

If a string of length ten over $\{a, b, c, d\}$ is chosen at random, what is the probability that it contains at least one pair of adjacent characters that are the same?

Proof. When $n = 10$, the number of all strings is $4^{10} = 1048576$ and the number of strings that contain at least one pair of adjacent characters that are the same is $4^{10} - 4 \cdot 3^9 = 1048576 - 4 \cdot 19683 = 969844$ so the probability is $969844/1048576 \approx 92.49\%$. \square

3.23 Exercise 23

3.23.1 (a)

How many integers from 1 through 1,000 are multiples of 4 or multiples of 7?

Proof. Let A = the set of integers that are multiples of 4 and B = the set of integers that are multiples of 7. Then $A \cap B$ = the set of integers that are multiples of 28.

Now $N(A) = 250$ because

$$\begin{array}{cccccccccccc} 1 & 2 & 3 & 4 & 5 & 6 & 7 & 8 & \dots & 999 & 1000 \\ & & & \downarrow & & & & \downarrow & & & \downarrow \\ & & & 4 \cdot 1 & & & & 4 \cdot 2 & \dots & & 4 \cdot 250 \end{array}$$

or equivalently, because $1000 = 4 \cdot 250$. Also $N(B) = 142$ because

$$\begin{array}{cccccccccccccccc} 1 & 2 & 3 & 4 & 5 & 6 & 7 & \dots & 14 & \dots & 994 & 995 & \dots & 1000 \\ & & & & & & \downarrow & & \downarrow & & \downarrow & & & \\ & & & & & & 7 \cdot 1 & & 7 \cdot 2 & & 7 \cdot 142 & & & \end{array}$$

or equivalently because $1000 = 7 \cdot 142 + 6$. And $N(A \cap B) = 35$ because

$$\begin{array}{cccccccccccc} 1 & 2 & 3 & \dots & 28 & \dots & 56 & \dots & 980 & \dots & 1000 \\ & & & & \downarrow & & \downarrow & & \downarrow & & \\ & & & & 28 \cdot 1 & & 28 \cdot 2 & & 28 \cdot 35 & & \end{array}$$

or equivalently because $1000 = 28 \cdot 35 + 20$. So $N(A \cup B) = 250 + 142 - 35 = 357$. \square

3.23.2 (b)

Suppose an integer from 1 through 1,000 is chosen at random. Use the result of part (a) to find the probability that the integer is a multiple of 4 or a multiple of 7.

Proof. $357/1000 = 35.7\%$ \square

3.23.3 (c)

How many integers from 1 through 1,000 are neither multiples of 4 nor multiples of 7?

Proof. $1000 - 357 = 643$ □

3.24 Exercise 24

3.24.1 (a)

How many integers from 1 through 1,000 are multiples of 2 or multiples of 9?

Proof. $\lfloor 1000/2 \rfloor = 500$ multiples of 2, $\lfloor 1000/9 \rfloor = 111$ multiples of 9, and $\lfloor 1000/18 \rfloor = 55$ multiples of 18 (which are double-counted). So: $500 + 111 - 55 = 556$. □

3.24.2 (b)

Suppose an integer from 1 through 1,000 is chosen at random. Use the result of part (a) to find the probability that the integer is a multiple of 2 or a multiple of 9.

Proof. $556/1000 = 55.6\%$ □

3.24.3 (c)

How many integers from 1 through 1,000 are neither multiples of 2 nor multiples of 9?

Proof. $1000 - 556 = 444$ □

3.25 Exercise 25

Counting Strings:

3.25.1 (a)

Make a list of all bit strings of lengths 0, 1, 2, 3, and 4 that do not contain the bit pattern 111.

Proof. $\lambda, 0, 1, 00, 01, 10, 11, 000, 001, 010, 100, 011, 101, 110, 0000, 0001, 0010, 0011, 0100, 0101, 0110, 1000, 1001, 1010, 1011, 1100, 1101$ □

3.25.2 (b)

For each integer $n \geq 0$, let d_n = the number of bit strings of length n that do not contain the bit pattern 111. Find d_0, d_1, d_2, d_3 , and d_4 .

Proof. $d_0 = 1, d_1 = 2, d_2 = 4, d_3 = 7, d_4 = 13$ □

3.25.3 (c)

Find a recurrence relation for d_0, d_1, d_2, \dots

Proof. Let k be an integer with $k \geq 3$. Any string of length k that does not contain the bit pattern 111 starts either with a 0 or with a 1. If it starts with a 0, this can be followed by any string of $k - 1$ bits that does not contain the pattern 111. There are d_{k-1} of these. If the string starts with a 1, then the first two bits are 10 or 11. If the first two bits are 10, then these can be followed by any string of $k - 2$ bits that does not contain the pattern 111. There are d_{k-2} of these. If the string starts with a 11, then the third bit must be 0 (because the string does not contain 111), and these three bits can be followed by any string of $k - 3$ bits that does not contain the pattern 111. There are d_{k-3} of these. Therefore, for every integer $k \geq 3$, $d_k = d_{k-1} + d_{k-2} + d_{k-3}$. \square

3.25.4 (d)

Use the results of parts (b) and (c) to find the number of bit strings of length 5 that do not contain the pattern 111.

Proof. $d_5 = d_4 + d_3 + d_2 = 13 + 7 + 4 = 24$ \square

3.26 Exercise 26

Counting Strings: Consider the set of all strings of a 's, b 's, and c 's.

3.26.1 (a)

Make a list of all of these strings of lengths 0, 1, 2, and 3 that do not contain the pattern aa .

Proof. Length 0: λ , Length 1: a, b, c , Length 2: $ab, ac, ba, bb, bc, ca, cb, cc$,

Length 3: $abc, acb, bac, bca, bcb, bcc, cab, cba, cbb, cbc, cca, ccb, ccc$. \square

3.26.2 (b)

For each integer $n \geq 0$, let s_n = the number of strings of a 's, b 's, and c 's of length n that do not contain the pattern aa . Find s_0, s_1, s_2 , and s_3 .

Proof. $s_0 = 1, s_1 = 3, s_2 = 8, s_3 = 22$ \square

3.26.3 (c)

Find a recurrence relation for s_0, s_1, s_2, \dots

Proof. Let s be a string of length k that does not contain the pattern aa .

If x starts with b or c then it can be followed by any string of length $k - 1$ that does not contain the pattern aa . There are s_{k-1} of those, so there are $2s_{k-1}$ such strings which x can be: s_{k-1} of them starting with b and s_{k-1} of them starting with c .

If x starts with a then it can be followed by a b or a c and then a string of length $k - 2$ that does not contain the pattern aa . There are s_{k-2} of those, so there are $2s_{k-2}$ such strings that x can be: s_{k-2} of them starting with ab and s_{k-2} of them starting with ac .

So $s_k = 2s_{k-1} + 2s_{k-2}$. □

3.26.4 (d)

Use the results of parts (b) and (c) to find the number of strings of a 's, b 's, and c 's of length four that do not contain the pattern aa .

Proof. $s_4 = 2s_3 + 2s_2 = 2(22 + 8) = 60$. □

3.26.5 (e)

Use the technique described in Section 5.8 to find an explicit formula for s_0, s_1, s_2, \dots

Proof. $s_k = 2s_{k-1} + 2s_{k-2}$ so the characteristic equation is $t^2 - 2t - 2 = 0$ which has the solutions

$$\frac{2 \pm \sqrt{(-2)^2 - 4(1)(-2)}}{2} = \frac{2 \pm \sqrt{4 + 8}}{2} = \frac{2 \pm 2\sqrt{3}}{2} = 1 \pm \sqrt{3}$$

The general solution has the form $s_k = A(1 + \sqrt{3})^k + B(1 - \sqrt{3})^k$. When $k = 0$ we have

$$s_0 = 1 = A(1 + \sqrt{3})^0 + B(1 - \sqrt{3})^0 = A + B$$

which gives $B = 1 - A$; and when $k = 1$ we have

$$s_1 = 3 = A(1 + \sqrt{3})^1 + B(1 - \sqrt{3})^1 = A + B + \sqrt{3}(A - B)$$

so we get $3 = 1 + \sqrt{3}(A - (1 - A)) = 1 + \sqrt{3}(2A - 1)$, solving we get $\frac{2}{\sqrt{3}} = 2A - 1$ so $\frac{1}{\sqrt{3}} + \frac{1}{2} = A$. Then $B = \frac{1}{2} - \frac{1}{\sqrt{3}}$. So

$$s_n = \left(\frac{1}{2} + \frac{1}{\sqrt{3}}\right) (1 + \sqrt{3})^n + \left(\frac{1}{2} - \frac{1}{\sqrt{3}}\right) (1 - \sqrt{3})^n$$

□

3.27 Exercise 27

For each integer $n \geq 0$, let a_k be the number of bit strings of length n that do not contain the pattern 101.

3.27.1 (a)

Show that $a_k = a_{k-1} + a_{k-3} + a_{k-4} + \cdots + a_0 + 2$, for every integer $k \geq 3$.

Proof. Assume $k \geq 3$ and s is a string of length k that does not contain the pattern 101.

If s begins with a 0 then it can be followed by a string of length $k - 1$ that does not contain the pattern 101. There are a_{k-1} of these.

If s begins with a 1 then consider the following. If it begins with:

100 then it can be followed by a string of length $k - 3$ that does not contain the pattern 101, and there are a_{k-3} of them,

1100 then it can be followed by a string of length $k - 4$ that does not contain the pattern 101, and there are a_{k-4} of them,

11100 then it can be followed by a string of length $k - 5$ that does not contain the pattern 101, and there are a_{k-5} of them,

$\underbrace{11 \dots 1}_{k-3}00$, then it can be followed by a string of length $k - (k - 1) = 1$ that does not contain the pattern 101, and there are a_1 of them,

$\underbrace{11 \dots 1}_{k-2}00$, then it can be followed by a string of length $k - k = 0$ that does not contain the pattern 101, and there are a_0 of them,

$\underbrace{11 \dots 1}_{k-1}$, then it can be followed by 1 or 0, and there are 2 of them.

So $a_k = a_{k-1} + a_{k-3} + a_{k-4} + \cdots + a_0 + 2$. □

3.27.2 (b)

Use the result of part (a) to show that if $k \geq 3$, then $a_k = 2a_{k-1} - a_{k-2} + a_{k-3}$.

Proof. By part (a) we have $a_{k-1} = a_{k-2} + a_{k-4} + a_{k-5} + \cdots + a_0 + 2$.

So $a_{k-1} + a_{k-3} = a_{k-2} + a_{k-3} + a_{k-4} + a_{k-5} + \cdots + a_0 + 2$.

So $a_{k-1} - a_{k-2} + a_{k-3} = a_{k-3} + a_{k-4} + a_{k-5} + \cdots + a_0 + 2$.

So $2a_{k-1} - a_{k-2} + a_{k-3} = a_{k-1} + a_{k-3} + a_{k-4} + a_{k-5} + \cdots + a_0 + 2$.

Notice the right-hand side is equal to a_k by part (a). □

3.28 Exercise 28

For each integer $n \geq 2$ let a_n be the number of permutations of $\{1, 2, 3, \dots, n\}$ in which no number is more than one place removed from its “natural” position. Thus $a_1 = 1$ since the one permutation of $\{1\}$, namely, 1, does not move 1 from its natural position. Also, $a_2 = 2$ since neither of the two permutations of $\{1, 2\}$, namely, 12 and 21, moves either number more than one place from its natural position.

3.28.1 (a)

Find a_3 .

Proof. $a_3 = 3$ (The three permutations that do not move more than one place from their “natural” positions are 213, 132, and 123.) \square

3.28.2 (b)

Find a recurrence relation for a_1, a_2, a_3, \dots

Proof. Let's try to see a relationship between $a_2 = 2$ and $a_3 = 3$. We need to add the new element 3. Since we cannot move 3 more than one from its natural position, it can be in position 2 or 3. If it's in position 3, like $_3$, then both of the permutations from a_2 work: 123, 213. If it's in position 2, then 2 must be in position 3: $_23$, so there is only one possibility left: 123.

Let's find a_4 . There are $4! = 24$ permutations of $\{1, 2, 3, 4\}$, but there are only 5 which do not move more than one place: 1234, 2134, 1243, 2143, 1324. So $a_4 = 5$.

But how is it obtained from a_3 ? There are 3 permutations in a_3 : 123, 132, 213. Now we need to add the element 4 to them. 4 can be either in position 3 or position 4. If it's in position 4, like $___4$, then all the permutations from a_3 work: 1234, 2134, 1324. If 4 is in position 3, then 3 must be in position 4: $__43$, so now there are a_2 ways to fill the two spots: 1243, 2143.

So we can see that the general form of the recurrence is: $a_n = a_{n-1} + a_{n-2}$. \square

3.29 Exercise 29

Refer to Example 9.3.5.

3.29.1 (a)

Write the following IP address in dotted decimal form: 11001010 00111000 01101011 11101110

Proof. $11001010_2 = 2 + 2^3 + 2^6 + 2^7 = 202$

$00111000_2 = 2^3 + 2^4 + 2^5 = 56$

$01101011_2 = 1 + 2 + 2^3 + 2^5 + 2^6 = 107$

$11101110_2 = 2 + 2^2 + 2^3 + 2^5 + 2^6 + 2^7 = 238$

So the answer is 202.56.107.238. \square

3.29.2 (b)

How many Class A networks can there be?

Proof. The network ID for a Class A network consists of 8 bits and begins with 0. If all possible combinations of eight 0's and 1's that start with a 0 were allowed, there would be 2 choices (0 or 1) for each of the 7 positions from the second through the eighth. This would give $2^7 = 128$ possible ID's. But because neither 00000000 nor 01111111 is allowed, the total is reduced by 2, so there are 126 possible Class A networks. \square

3.29.3 (c)

What is the dotted decimal form of the IP address for a computer in a Class A network?

Proof. Let $w.x.y.z$ be the dotted decimal form of the IP address for a computer in a Class A network. Because the network IDs for a Class A network go from 00000001 ($= 1$) through 01111110 ($= 126$), w can be any integer from 1 through 126. In addition, each of x , y , and z can be any integer from 0 ($= 00000000$) through 255 ($= 11111111$), except that x , y , and z cannot all be 0 simultaneously and cannot all be 255 simultaneously. \square

3.29.4 (d)

How many host IDs can there be for a Class A network?

Proof. Twenty-four positions are allocated for the host ID in a Class A network. If each could be either 0 or 1, there would be $2^{24} = 16,777,216$ possible host IDs. But neither all 0's nor all 1's is allowed, which reduces the total by 2. Thus there are 16,777,214 possible host IDs in a Class A network. \square

3.29.5 (e)

How many Class C networks can there be?

Proof. The network ID for a Class C network consists of 24 bits and begins with 110. There are 2 choices (0 or 1) for each of the 21 positions from the fourth through the twenty-fourth. This gives 2^{21} possible ID's. \square

3.29.6 (f)

What is the dotted decimal form of the IP address for a computer in a Class C network?

Proof. Let $w.x.y.z$ be the dotted decimal form of the IP address for a computer in a Class C network. The network IDs for a Class C network go from 110000000000000000000000 through 110111111111111111111111.

If we break these down into chunks of 8 bits and convert them to binary, we get:

$$11000000_2 = 192_{10}, 00000000_2 = 0_{10}, 00000000_2 = 0_{10},$$

$$11011111_2 = 223_{10}, 11111111_2 = 255_{10}, 11111111_2 = 255_{10}.$$

As dotted decimals they range from 192.0 to 223.255. So $192 \leq w \leq 223$, $0 \leq x \leq 255$, $0 \leq y \leq 255$, $0 \leq z \leq 255$. \square

3.29.7 (g)

How many host IDs can there be for a Class C network?

Proof. There are only 8 bits for host IDs in a Class C network, but a host ID may not consist of either all 0's or all 1's. Therefore $2^8 - 2 = 254$ host IDs. \square

3.29.8 (h)

How can you tell, by looking at the first of the four numbers in the dotted decimal form of an IP address, what kind of network the address is from? Explain.

Proof. For Class A networks $0 \leq w \leq 126$, for Class B networks $128 \leq w \leq 191$, for Class C networks $192 \leq w \leq 223$. So we can tell from this info what kind of network an address is from. \square

3.29.9 (i)

An IP address is 140.192.32.136. What class of network does it come from?

Proof. Observe that $140 = 128 + 8 + 4 = 10001100_2$, which begins with 10. Thus the IP address comes from a Class B network. An alternative solution uses the result of Example 9.3.5: Network IDs for Class B networks range from 128 through 191. Thus, since $128 \leq 140 \leq 191$, the given IP address is from a Class B network. \square

3.29.10 (j)

An IP address is 202.56.107.238. What class of network does it come from?

Proof. For Class C networks we have $192 \leq w \leq 223$, and 202 is in this range, so it the address belongs to a Class C network. \square

3.30 Exercise 30

A row in a classroom has n seats. Let s_n be the number of ways nonempty sets of students can sit in the row so that no student is seated directly adjacent to any other student. (For instance, a row of three seats could contain a single student in any of the seats or a pair of students in the two outer seats. Thus $s_3 = 4$.) Find a recurrence relation for s_1, s_2, s_3, \dots

Proof. Let us denote students with a * and empty seats with a . .

$n = 1$: The seats . can be filled as *, so $s_1 = 1$.

$n = 2$: The seats .. can be filled as *. or .*, so $s_2 = 2$.

$n = 3$: The seats ... can be filled as *.., .*, ..*, or *.* , so $s_3 = 4$.

$n = 4$: The seats can be filled as *..., .*..., ..*, ...*, *.*., *..*, or *.*., so $s_4 = 7$.

Proof. The probability that at least two of the four people share the same birth month is $8,856/20,736 \approx 42.7\%$. \square

3.31.5 (e)

How large must n be so that in any group of n people, the probability that two or more share the same birth month is at least 50%?

Proof. When there are five people, the probability that at least two share the same birth month is $\frac{12^5 - 12 \cdot 11 \cdot 10 \cdot 9 \cdot 8}{12^5} \approx 61.8\%$, and when there are more than five people, the probability is even greater. Thus, since the probability for four people is less than 50%, the group must contain five or more people for the probability to be at least 50% that two or more share the same birth month. \square

3.32 Exercise 32

Assuming that all years have 365 days and all birthdays occur with equal probability, how large must n be so that in any randomly chosen group of n people, the probability that two or more have the same birthday is at least $1/2$? (This is called the birthday problem. Many people find the answer surprising.)

Proof. The probability that two out of n people share the same birthday is

$$\frac{365^n - 365 \cdot 364 \cdot \dots \cdot (365 - n + 1)}{365^n}$$

We want this to be $\geq 50\%$. By trying values on the calculator I found that for $n = 23$ it's $\approx 50.73\%$ and for $n = 22$ it's $\approx 47.57\%$ so $n = 23$ is the smallest value. \square

3.33 Exercise 33

A college conducted a survey to explore the academic interests and achievements of its students. It asked students to place checks beside the numbers of all the statements that were true of them. Statement #1 was “I was on the Dean’s list last term,” statement #2 was “I belong to an academic club, such as the math club or the Spanish club,” and statement #3 was “I am majoring in at least two subjects.” Out of a sample of 100 students, 28 checked #1, 26 checked #2, and 14 checked #3, 8 checked both #1 and #2, 4 checked both #1 and #3, 3 checked both #2 and #3, and 2 checked all three statements.

3.33.1 (a)

How many students checked at least one of the statements?


Proof. The number of students who checked at least one of the statements is $N(H) + N(C) + N(D) - N(H \cap C) - N(H \cap D) - N(C \cap D) + N(H \cap C \cap D) = 28 + 26 + 14 - 8 - 4 - 3 + 2 = 55$. \square

3.33.2 (b)


How many students checked none of the statements?

Proof. By the difference rule, the number of students who checked none of the statements is the total number of students minus the number who checked at least one statement. This is $100 - 55 = 45$. \square

3.33.3 (c)



Let H be the set of students who checked #1, C the set of students who checked #2, and D the set of students who checked #3. Fill in the numbers for all eight regions of the diagram.



Proof.

\square

3.33.4 (d)

How many students checked #1 and #2 but not #3?

Proof. The number of students who checked #1 and #2 but not #3 is $N(H \cap D) - N(H \cap C \cap D) = 8 - 2 = 6$. \square

3.33.5 (e)

How many students checked #2 and #3 but not #1?

Proof. The number of students who checked #2 and #3 but not #1 is $N(C \cap D) - N(H \cap C \cap D) = 3 - 2 = 1$. \square

3.33.6 (f)

How many students checked #2 but neither of the other two?

Proof. The number of students who checked #2 but not #1 or #3 is $N(C) - N(C \cap (H \cup D)) = 17 - (6 + 2 + 1) = 8$. \square

3.34 Exercise 34

A study was done to determine the efficacy of three different drugs: A, B, and C, in relieving headache pain. Over the period covered by the study, 50 subjects were given the chance to use all three drugs. The following results were obtained:

21 reported relief from drug A

21 reported relief from drug B

31 reported relief from drug C

9 reported relief from both drugs A and B

14 reported relief from both drugs A and C

15 reported relief from both drugs B and C

41 reported relief from at least one of the drugs.

Note that some of the 21 subjects who reported relief from drug A may also have reported relief from drugs B or C. A similar occurrence may be true for the other data.

3.34.1 (a)

How many people got relief from none of the drugs?

Proof. $50 - 41 = 9$ \square

3.34.2 (b)

How many people got relief from all three drugs?

Proof. By the Inclusion-Exclusion Principle,


$$N(A \cup B \cup C) = N(A) + N(B) + N(C) - N(A \cap B) - N(A \cap C) - N(B \cap C) + N(A \cap B \cap C).$$

By using the values given in the problem, we get


$$41 = 21 + 21 + 31 - 9 - 14 - 15 + N(A \cap B \cap C).$$

Solving we get $N(A \cap B \cap C) = 6$. □

3.34.3 (c)



Let A be the set of all subjects who got relief from drug A, B the set of all subjects who got relief from drug B, and C the set of all subjects who got relief from drug C. Fill in the numbers for all eight regions of the diagram.



Proof. □

3.34.4 (d)

How many subjects got relief from A only?

Proof. 4 □

3.35 Exercise 35

An interesting use of the inclusion/exclusion rule is to check survey numbers for consistency. For example, suppose a public opinion polltaker reports that out of a national sample of 1,200 adults, 675 are married, 682 are from 20 to 30 years old, 684 are female, 195 are married and are from 20 to 30 years old, 467 are married females, 318 are females

from 20 to 30 years old, and 165 are married females from 20 to 30 years old. Are the polltaker's figures consistent? Could they have occurred as a result of an actual sample survey?

Proof. Let M = the set of married people in the sample,

Y = the set of people between 20 and 30 in the sample, and

F = the set of females in the sample.

Then the number of people in the set $M \cup Y \cup F$ is less than or equal to the size of the sample. And so

$$\begin{aligned} 1,200 &\geq N(M \cup Y \cup F) \\ &= N(M) + N(Y) + N(F) - N(M \cap Y) \\ &\quad - N(M \cap F) - N(Y \cap F) + N(M \cap Y \cap F) \\ &= 675 + 682 + 684 - 195 - 467 - 318 + 165 \\ &= 1,226. \end{aligned}$$

This is impossible since $1,200 < 1,226$, so the polltaker's figures are inconsistent. They could not have occurred as a result of an actual sample survey. \square

3.36 Exercise 36

Fill in the reasons for each step below. If A and B are sets in a finite universe U , then

$$\begin{aligned} N(A \cap B) &= N(U) - N((A \cap B)^c) && \text{(a) } \underline{\hspace{2cm}} \\ &= N(U) - N(A^c \cup B^c) && \text{(b) } \underline{\hspace{2cm}} \\ &= N(U) - (N(A^c) + N(B^c) - N(A^c \cap B^c)) && \text{(c) } \underline{\hspace{2cm}} \end{aligned}$$

Proof. (a) because

$$\begin{aligned} A \cap B &= (A \cap B) \cap U && \text{by identity law} \\ &= U \cap (A \cap B) && \text{by commutative law} \\ &= U \cap ((A \cap B)^c)^c && \text{by double complement law} \\ &= U - (A \cap B)^c && \text{by set difference law} \end{aligned}$$

and because $N(U - (A \cap B)^c) = N(U) - N((A \cap B)^c)$.

(b) by De Morgan laws (c) by inclusion-exclusion principle for 2 sets \square

For each of exercises 37 – 39, the number of elements in a certain set can be found by computing the number in a larger universe that are not in the set and subtracting this from the total in the larger universe. In each of these, as was the case for the solution to example 9.3.6(b), De Morgan's laws and the inclusion / exclusion rule can be used.

3.37 Exercise 37

How many positive integers less than 1,000 have no common factors with 1,000?

Proof. Let A be the set of all positive integers less than 1,000 that are not multiples of 2, and let B be the set of all positive integers less than 1,000 that are not multiples of 5. Since the only prime factors of 1,000 are 2 and 5, the number of positive integers that have no common factors with 1,000 is $N(A \cap B)$. Let the universe U be the set of all positive integers less than 1,000. Then A^c is the set of positive integers less than 1,000 that are multiples of 2, B^c is the set of positive integers less than 1,000 that are multiples of 5, and $A^c \cap B^c$ is the set of positive integers less than 1,000 that are multiples of 10. By one of the procedures discussed in Section 9.1 or 9.2, it is easily found that $N(A^c) = 499$, $N(B^c) = 199$, and $N(A^c \cap B^c) = 99$. Thus, by the inclusion/exclusion rule, $N(A^c \cup B^c) = N(A^c) + N(B^c) - N(A^c \cap B^c) = 499 + 199 - 99 = 599$. But by De Morgan's law, $N(A^c \cup B^c) = N((A \cap B)^c)$, and so (*) $N((A \cap B)^c) = 599$. Now since $(A \cap B)^c = U - (A \cap B)$, by the difference rule we have (**) $N((A \cap B)^c) = N(U) - N(A \cap B)$. Equating the right-hand sides of (*) and (**) gives $N(U) - N(A \cap B) = 599$. And because $N(U) = 999$, we conclude that $999 - N(A \cap B) = 599$, or, equivalently, $N(A \cap B) = 999 - 599 = 400$. So there are 400 positive integers less than 1,000 that have no common factor with 1,000. \square

3.38 Exercise 38

How many permutations of $abcde$ are there in which the first character is a, b , or c and the last character is c, d , or e ?

Proof. **Case 1: first character is a .** For the last character there are 3 choices: c, d, e . Then the middle 3 characters can be permuted in $3!$ ways. So $3 \cdot 3! = 3 \cdot 6 = 18$ possibilities.

Case 2: first character is b . For the last character there are 3 choices: c, d, e . Then the middle 3 characters can be permuted in $3!$ ways. So $3 \cdot 3! = 3 \cdot 6 = 18$ possibilities.

Case 3: first character is c . For the last character there are 2 choices: d, e . Then the middle 3 characters can be permuted in $3!$ ways. So $2 \cdot 3! = 2 \cdot 6 = 12$ possibilities.

In total, there are $18 + 18 + 12 = 48$ permutations in which the first character is a, b , or c and the last character is c, d , or e . \square

3.39 Exercise 39

How many integers from 1 through 999,999 contain each of the digits 1, 2, and 3 at least once? (*Hint:* For each $i = 1, 2$, and 3, let A_i be the set of all integers from 1 through 999,999 that do not contain the digit i .)

Proof. Let A_1, A_2, A_3 be the sets of all integers from 1 through 999,999 that do not contain the digits 1, 2, 3, respectively.

$N(A_1) = 9^6 - 1$ because, excluding the digit 1, there are 9 choices for each digit, for a total of 9^6 , but one of them, namely 000,000 corresponds to 0 which is out of the range 1 through 999,999.

Similarly $N(A_2) = N(A_3) = 9^6 - 1$.

$N(A_1 \cap A_2) = 8^6 - 1$ because, excluding the digits 1 and 2, there are 8 choices for each digit, for a total of 8^6 , but one of them, namely 000,000 corresponds to 0 which is out of the range 1 through 999,999.

Similarly $N(A_1 \cap A_3) = N(A_2 \cap A_3) = 8^6 - 1$.

Finally $N(A_1 \cap A_2 \cap A_3) = 7^6 - 1$ by a similar argument to the above arguments.

Now by the inclusion / exclusion principle, $N(A_1 \cup A_2 \cup A_3) = N(A_1) + N(A_2) + N(A_3) - N(A_1 \cap A_2) - N(A_1 \cap A_3) - N(A_2 \cap A_3) + N(A_1 \cap A_2 \cap A_3) = 3(9^6 - 1) - 3(8^6 - 1) + 7^6 - 1 = 925,539$

So the number of integers that do not contain at least one of the digits 1,2,3 is 925,539. Then $999,999 - 925,539 = 74460$ integers contain all three of the digits 1,2,3.

We can verify this with a computer:

```
scala> var count = 0
var count: Int = 0

scala> for i <- 1 to 999999 do
  |   if "123".forall(c => i.toString.contains(c)) then
  |     count += 1
  |
scala> count
val res0: Int = 74460
```

□

For 40 and 41, use the definition of the Euler phi function ϕ from Section 7.1, exercises 51 – 53.

3.40 Exercise 40

Use the inclusion/exclusion principle to prove the following: If $n = pq$, where p and q are distinct prime numbers, then $\phi(n) = (p - 1)(q - 1)$.

Hint: Let A and B be the sets of all positive integers less than or equal to n that are divisible by p and q , respectively. Then $\phi(n) = n - N(A \cup B)$.

Proof. (following the Hint)

Notice $N(A) = q$ since there are q multiples of p from 1 to n : $p, 2p, 3p, \dots, (q - 1)p, qp = n$. Similarly $N(B) = p$.

Notice $N(A \cap B) = 1$ since $n = pq$ is the only number from 1 through n that is divisible by both p and q .

By the inclusion / exclusion principle, $N(A \cup B) = N(A) + N(B) - N(A \cap B) = q + p - 1$. So $\phi(n) = n - (q + p - 1) = pq - q - p + 1 = q(p - 1) - (p - 1) = (p - 1)(q - 1)$. \square

3.41 Exercise 41

Use the inclusion/exclusion principle to prove the following: If $n = pqr$, where p, q , and r are distinct prime numbers, then $\phi(n) = (p - 1)(q - 1)(r - 1)$.

Proof. We can use an approach similar to Exercise 40. Let P, Q, R be the sets of positive integers less than or equal to n that are divisible by p, q, r respectively.

Then $N(P) = qr, N(Q) = pr, N(R) = pq$ by similar arguments.

Now $N(P \cap Q) = r$ because: if an integer is divisible by both p and q then it is divisible by pq since p and q are both prime; and there are r multiples of pq from 1 through n : $pq, 2pq, 3pq, \dots, (r - 1)pq, rpq = n$.

By similar arguments $N(P \cap R) = q$ and $N(Q \cap R) = p$. Finally $N(P \cap Q \cap R) = 1$ because $n = pqr$ itself is the only integer divisible by all 3 primes.

By the inclusion / exclusion principle, $N(P \cup Q \cup R) = N(P) + N(Q) + N(R) - N(P \cap Q) - N(P \cap R) - N(Q \cap R) + N(P \cap Q \cap R) = qr + pr + pq - r - q - p + 1$.

$N(P \cup Q \cup R)$ is the number of integers that are divisible by at least one of p, q, r , thus $\phi(n) = n - N(P \cup Q \cup R) = pqr - qr + pr + pq - r - q - p + 1$, which, after factoring, equals $(p - 1)(q - 1)(r - 1)$. \square

3.42 Exercise 42

A gambler decides to play successive games of blackjack until he loses three times in a row. (Thus the gambler could play five games by losing the first, winning the second, and losing the final three or by winning the first two and losing the final three. These possibilities can be symbolized as LWLLL and WWLLL.) Let g_n be the number of ways the gambler can play n games.

3.42.1 (a)

Find g_3, g_4 , and g_5 .

Proof. 3 games: LLL so $g_3 = 1$. 4 games: WLLL so $g_4 = 1$.

5 games: LWLLL, WWLLL so $g_5 = 2$. \square

3.42.2 (b)

Find g_6 .

Proof. 6 games: WWWLLL, LWLLL, WLWLLL, LLWLLL so $g_6 = 4$. \square

3.42.3 (c)

Find a recurrence relation for g_3, g_4, g_5, \dots

Hint: If $k \geq 6$, any sequence of k games must begin with W, LW, or LLW.

Proof. By the Hint we can see that for $k \geq 6$, $g_k = g_{k-1} + g_{k-2} + g_{k-3}$. □

3.43 Exercise 43

A derangement of the set $\{1, 2, \dots, n\}$ is a permutation that moves every element of the set away from its “natural” position. Thus 21 is a derangement of $\{1, 2\}$, and 231 and 312 are derangements of $\{1, 2, 3\}$. For each positive integer n , let d_n be the number of derangements of the set $\{1, 2, \dots, n\}$.

3.43.1 (a)

Find d_1, d_2 , and d_3 .

Proof. $d_1 = 0$ because the only permutation of the set $\{1\}$ is 1, which cannot move 1 from its natural position.

$d_2 = 1$ because the only two permutations of the set $\{1, 2\}$ are 12 and 21, and only 21 moves both 1 and 2 away from their natural positions.

$d_3 = 2$ because there are 6 permutations of the set $\{1, 2, 3\}$: 123, 132, 213, 231, 312, 321; and only 231 and 312 move every element away from their natural positions. □

3.43.2 (b)

Find d_4 .

Proof. There are 24 permutations: 1234, 1243, 1324, 1342, 1423, 1432, 2134, 2143, 2314, 2341, 2413, 2431, 3124, 3142, 3214, 3241, 3412, 3421, 4123, 4132, 4213, 4231, 4312, 4321.

The only ones that move every element away from their natural positions are: 2143, 2341, 2413, 3142, 3412, 3421, 4123, 4312, 4321.

So $d_4 = 9$. □

3.43.3 (c)

Find a recurrence relation for d_1, d_2, d_3, \dots

Hint: Divide the set of all derangements into two subsets: one subset consists of all derangements in which the number 1 changes places with another number, and the other subset consists of all derangements in which the number 1 goes to position $i \neq 1$ but i does not go to position 1. The answer is $d_k = (k-1)d_{k-1} + (k-1)d_{k-2}$. Can you justify it?

Proof. (following the Hint)

Consider a derangement in the first subset. Then 1 switches places with another number i . There are $k - 1$ other numbers that 1 can switch places with. For each one of them, then the remaining $k - 2$ numbers have to also derange, and there are d_{k-2} ways to do that. Thus the number of derangements in the first subset is $(k - 1)d_{k-2}$.

Consider a derangement in the second subset. Then 1 moves to position i for some number i . There are $k - 1$ choices for this i . For each one of these choices, there are d_{k-1} ways to derange the remaining $k - 1$ numbers. Why?

Because number i cannot go to position 1, but it can go to any one of the remaining $k - 2$ positions, and similarly any other number j ($j \neq 1, j \neq i$) can go to $k - 2$ positions (any position except j and i), so this situation is the same as derangements of $k - 1$ elements.

Thus the number of derangements in the second subset is $(k - 1)d_{k-1}$. So the total is: $d_k = (k - 1)d_{k-1} + (k - 1)d_{k-2}$. \square

3.44 Exercise 44

Note that a product $x_1x_2x_3$ may be parenthesized in two different ways: $(x_1x_2)x_3$ and $x_1(x_2x_3)$. Similarly, there are several different ways to parenthesize $x_1x_2x_3x_4$. Two such ways are $(x_1x_2)(x_3x_4)$ and $x_1((x_2x_3)x_4)$. Let P_n be the number of different ways to parenthesize the product $x_1x_2 \dots x_n$. Show that if $P_1 = 1$, then

$$P_n = \sum_{k=1}^{n-1} P_k P_{n-k} \text{ for every integer } n \geq 2.$$

(It turns out that the sequence P_1, P_2, P_3, \dots is the same as the sequence of Catalan numbers: $P_n = C_{n-1}$ for every integer $n \geq 1$. See Example 5.6.4.)

Proof. Consider the product $x_1x_2 \dots x_n$. All the different ways to parenthesize it can be classified as follows: split it into two products x_1 and $x_2 \dots x_n$, or into x_1x_2 and $x_3 \dots x_n$, or into $x_1x_2x_3$ and $x_4 \dots x_n$, ..., or into $x_1x_2 \dots x_{n-1}$ and x_n .

By the addition rule, P_n is the sum of the number of ways each one of these product pairs can be parenthesized.

For each of the product pairs, by the product rule, the number of ways to parenthesize the product pair is the product of the numbers of ways to parenthesize each product.

There are P_1 ways to parenthesize x_1 and P_{n-1} ways to parenthesize $x_2 \dots x_n$, so there are P_1P_{n-1} ways to parenthesize their product.

Similarly there are P_2 ways to parenthesize x_1x_2 and P_{n-2} ways to parenthesize $x_3 \dots x_n$, so there are P_2P_{n-2} ways to parenthesize their product.

And so on. Therefore $P_n = \sum_{k=1}^{n-1} P_k P_{n-k}$ for every $n \geq 2$. \square

3.45 Exercise 45

Use mathematical induction to prove Theorem 9.3.1: “Suppose a finite set A equals the union of k distinct mutually disjoint subsets A_1, A_2, \dots, A_k . Then $N(A) = N(A_1) + N(A_2) + \dots + N(A_k)$.”

Proof. Let $P(n)$ be the statement “if a finite set A equals the union of n distinct mutually disjoint subsets A_1, A_2, \dots, A_n then $N(A) = N(A_1) + N(A_2) + \dots + N(A_n)$.”

Show that $P(1)$ is true: In this case $A = A_1$ therefore $N(A) = N(A_1)$, so $P(1)$ is true.

Show that for any integer $k \geq 1$ if $P(k)$ is true then $P(k+1)$ is true: Assume $P(k)$ is true and assume A is a finite set that equals the union of $k+1$ mutually disjoint subsets A_1, \dots, A_k, A_{k+1} . [We want to show $N(A) = N(A_1) + N(A_2) + \dots + N(A_{k+1})$].

Let $A' = A_1 \cup \dots \cup A_k$. Then A' is a finite set that equals the union of k mutually disjoint subsets A_1, \dots, A_k . By the inductive hypothesis $N(A') = N(A_1) + N(A_2) + \dots + N(A_k)$.

Notice that since A_1, \dots, A_{k+1} are mutually disjoint, $A' = A_1 \cup \dots \cup A_k$ and A_{k+1} are also disjoint. So $N(A) = N(A' \cup A_{k+1}) = N(A') + N(A_{k+1}) = N(A_1) + N(A_2) + \dots + N(A_k) + N(A_{k+1})$, [as was to be shown.] \square

3.46 Exercise 46

Prove the inclusion/exclusion rule for two sets A and B by showing that $A \cup B$ can be partitioned into $A \cap B$, $A - (A \cap B)$, and $B - (A \cap B)$, and then using the addition and difference rules. (See the hint for exercise 39 in Section 6.2.)

Proof. **Claim:** $A \cup B = (A \cap B) \cup [A - (A \cap B)] \cup [B - (A \cap B)]$ and $(A \cap B)$, $A - (A \cap B)$ and $B - (A \cap B)$ are mutually disjoint sets.

Proof of Claim.

First to prove the equality:

1. Assume $x \in A \cup B$.
2. By definition of union, $x \in A$ or $x \in B$.
3. **Case 1:** $x \in A$.
 - 3.1 **Case 1.1:** $x \in B$. Then by definition of intersection, $x \in A \cap B$. So by definition of union $x \in (A \cap B) \cup [A - (A \cap B)] \cup [B - (A \cap B)]$.
 - 3.2 **Case 1.2:** $x \notin B$. Then by definition of intersection, $x \notin A \cap B$. So by definition of difference $x \in A - (A \cap B)$. So by definition of union, $x \in (A \cap B) \cup [A - (A \cap B)] \cup [B - (A \cap B)]$.
4. **Case 2:** $x \in B$. Similar to Case 1. We conclude $x \in (A \cap B) \cup [A - (A \cap B)] \cup [B - (A \cap B)]$.
5. By 3 and 4, $x \in (A \cap B) \cup [A - (A \cap B)] \cup [B - (A \cap B)]$.

6. By 1, 5 and definition of subset, $A \cup B \subseteq (A \cap B) \cup [A - (A \cap B)] \cup [B - (A \cap B)]$.
7. Assume $x \in (A \cap B) \cup [A - (A \cap B)] \cup [B - (A \cap B)]$.
8. By definition of union, $x \in A \cap B$ or $x \in A - (A \cap B)$ or $x \in B - (A \cap B)$.
9. **Case 1:** $x \in A \cap B$. By definition of intersection $x \in A$ and $x \in B$. So by definition of union, $x \in A \cup B$.
10. **Case 2:** $x \in A - (A \cap B)$. By definition of difference, $x \in A$ and $x \notin A \cap B$. So by definition of union, $x \in A \cup B$.
11. **Case 3:** $x \in B - (A \cap B)$. By definition of difference, $x \in B$ and $x \notin A \cap B$. So by definition of union, $x \in A \cup B$.
12. By 9, 10 and 11, $x \in A \cup B$.
13. By 7, 12 and definition of subset, $(A \cap B) \cup [A - (A \cap B)] \cup [B - (A \cap B)] \subseteq A \cup B$.
14. By 6, 13 and definition of set equality $(A \cap B) \cup [A - (A \cap B)] \cup [B - (A \cap B)] = A \cup B$.

Now to prove that the three sets are mutually disjoint:

Assume $x \in A \cap B$. We want to show $x \notin A - (A \cap B)$ and $x \notin B - (A \cap B)$.

Argue by contradiction and assume $x \in A - (A \cap B)$. Then by definition of difference $x \in A$ and $x \notin A \cap B$, contradicting $x \in A \cap B$. Similarly we can prove $x \notin B - (A \cap B)$.

Assume $x \in A - (A \cap B)$. We want to show $x \notin A \cap B$ and $x \notin B - (A \cap B)$.

By definition of difference, $x \in A$ and $x \notin A \cap B$. Then by definition of intersection either $x \notin A$ or $x \notin B$. The first case $x \notin A$ contradicts the fact that $x \in A$, so $x \notin B$. So by definition of intersection $x \notin A \cap B$ (otherwise $x \in B$, contradiction) and by definition of difference $x \notin B - (A \cap B)$ either (otherwise $x \in B$, contradiction).

Assume $x \in B - (A \cap B)$. We want to show $x \notin A \cap B$ and $x \notin A - (A \cap B)$. This proof is very similar to the case above.

[end of proof of the Claim.]

By the claim $N(A \cup B) = N((A \cap B) \cup [A - (A \cap B)] \cup [B - (A \cap B)])$.

Since these sets are disjoint by the claim, by the addition rule $N(A \cup B) = N(A \cap B) + N(A - (A \cap B)) + N(B - (A \cap B))$.

By the difference rule $N(A - (A \cap B)) = N(A) - N(A \cap B)$ and $N(B - (A \cap B)) = N(B) - N(A \cap B)$. Substituting these into the equation above, we get

$$N(A \cup B) = N(A \cap B) + N(A) - N(A \cap B) + N(B) - N(A \cap B) = N(A) + N(B) - N(A \cap B)$$

[which proves the Inclusion / Exclusion principle for two sets.] □

3.47 Exercise 47

Proof. □

3.48 Exercise 48

Use mathematical induction to prove the general inclusion/exclusion rule: If A_1, \dots, A_n are finite sets, then $N(A_1 \cup A_2 \cup \dots \cup A_n)$

$$= \sum_{1 \leq i \leq n} N(A_i) - \sum_{1 \leq i < j \leq n} N(A_i \cap A_j) + \sum_{1 \leq i < j < k \leq n} N(A_i \cap A_j \cap A_k) \\ - \dots + (-1)^{n+1} N(A_1 \cap A_2 \cap \dots \cap A_n)$$

(The notation $\sum_{1 \leq i < j \leq n} N(A_i \cap A_j)$ means that quantities of the form $N(A_i \cap A_j)$ are to be added together for all integers i and j with $1 \leq i < j \leq n$.)

Hint: Use the associative law for sets from Theorem 6.2.2 and the generalized distributive law for sets from exercise 40, Section 6.2.

Proof. Let $P(n)$ be the statement: “if A_1, \dots, A_n are finite sets then the equation above is true.”

Show that $P(1)$ is true: When $n = 1$ the equation becomes $N(A_1) = \sum_{1 \leq i \leq 1} N(A_i)$, with all the other sums becoming empty sums, since $1 \leq i < j \leq 1$ is impossible for integers i, j . Both sides are equal to $N(A_1)$ therefore $P(1)$ is true.

Show that for any integer $n \geq 1$ if $P(n)$ is true then $P(n+1)$ is true: Assume $n \geq 1$ and assume $P(n)$ is true. Assume A_1, \dots, A_{n+1} are finite sets. [We want to prove $P(n+1)$.]

Let $B = \bigcup_{i=1}^n A_i$. Notice $\bigcup_{i=1}^{n+1} A_i = B \cup A_{n+1}$. Then by the inclusion / exclusion principle for two sets,

$$N\left(\bigcup_{i=1}^{n+1} A_i\right) = N(B \cup A_{n+1}) = N(B) + N(A_{n+1}) - N(B \cap A_{n+1}) \quad (1)$$

By the inductive hypothesis

$$N(B) = \sum_{1 \leq i \leq n} N(A_i) - \sum_{1 \leq i < j \leq n} N(A_i \cap A_j) + \dots + (-1)^{n+1} N(A_1 \cap A_2 \cap \dots \cap A_n) \quad (2)$$

By the generalized distributive law,

$$B \cap A_{n+1} = (A_1 \cap A_{n+1}) \cup (A_2 \cap A_{n+1}) \cup \dots \cup (A_n \cap A_{n+1}) = \bigcup_{i=1}^n (A_i \cap A_{n+1})$$

By the inductive hypothesis again, $N(B \cap A_{n+1}) = N\left(\bigcup_{i=1}^n (A_i \cap A_{n+1})\right)$

$$= \sum_{1 \leq i \leq n} N(A_i \cap A_{n+1}) - \sum_{1 \leq i < j \leq n} N((A_i \cap A_{n+1}) \cap (A_j \cap A_{n+1})) \\ + \dots + (-1)^{n+1} N\left(\bigcap_{i=1}^n (A_i \cap A_{n+1})\right) \quad (3)$$

Now let's try to substitute and combine all the terms.

The term $N(B)$ in (1) will be replaced by the right hand side of (2), and the term $N(B \cap A_{n+1})$ in (1) will be replaced by the right hand side of (3). So let's imagine we've done that already, and start combining terms in (1).

The term $\sum_{1 \leq i \leq n} N(A_i)$ on the right hand side of (2), and the term $N(A_{n+1})$ in (1), can be combined into: $\sum_{1 \leq i \leq n+1} N(A_i)$.

The term $\sum_{1 \leq i < j \leq n} N(A_i \cap A_j)$ in (2) and the term $\sum_{1 \leq i \leq n} N(A_i \cap A_{n+1})$ on the right side of (3) can be combined into $\sum_{1 \leq i < j \leq n+1} N(A_i \cap A_j)$. Why? The first sum has all the two-way intersections between all the A_i from 1 to n , and the second sum has all the two-way intersections of the A_i from 1 to n with A_{n+1} . Putting them together we end up with all the two-way intersections between all the A_i from 1 to $n+1$. Both of these sums have negative signs (the second sum gets a negative sign from $N(B \cap A_{n+1})$).

Notice that the next term in (3) is, by the associative law and the idempotent law for intersection:

$$\sum_{1 \leq i < j \leq n} N((A_i \cap A_{n+1}) \cap (A_j \cap A_{n+1})) = \sum_{1 \leq i < j \leq n} N(A_i \cap A_j \cap A_{n+1})$$

So, similarly the term $\sum_{1 \leq i < j < k \leq n} N(A_i \cap A_j \cap A_k)$ in (2) and the term $\sum_{1 \leq i < j \leq n} N(A_i \cap A_j \cap A_{n+1})$ in (3) can be combined into $\sum_{1 \leq i < j < k \leq n+1} N(A_i \cap A_j \cap A_k)$ because the first sum has all the three-way intersections between A_i from 1 to n , and the second sum has all the three-way intersections of A_{n+1} with the A_i from 1 to n . Both sums have a positive sign (the second sum has a double negative sign, one from $N(B \cap A_{n+1})$ in (1) and another from itself).

All the other i -way intersection term sums from (2) and (3) can be combined in a similar way. So we end up with: $N\left(\bigcup_{i=1}^{n+1} A_i\right)$

$$\begin{aligned} &= \sum_{1 \leq i \leq n+1} N(A_i) - \sum_{1 \leq i < j \leq n+1} N(A_i \cap A_j) + \sum_{1 \leq i < j < k \leq n+1} N(A_i \cap A_j \cap A_k) \\ &\quad - \cdots + (-1)^{n+2} N(A_1 \cap A_2 \cap \cdots \cap A_{n+1}) \end{aligned}$$

which proves $P(n+1)$, [as was to be shown.] □

3.49 Exercise 49

A circular disk is cut into n distinct sectors, each shaped like a piece of pie and all meeting at the center point of the disk. Each sector is to be painted red, green, yellow, or blue in such a way that no two adjacent sectors are painted the same color. Let S_n be the number of ways to paint the disk.

3.49.1 (a)

Find a recurrence relation for S_k in terms of S_{k-1} and S_{k-2} for each integer $k \geq 4$.

Hint: Use the solution method described in Section 5.8. The answer is $S_k = 2S_{k-1} + 3S_{k-2}$ for every integer $k \geq 4$.

Proof. NOTE: I have to assume, without any extra information, that the disk is FIXED, in other words, rotation, flipping etc. are considered different ways to paint.

For example, for $n = 3$ consider the clockwise paintings starting from the sector that occupies 12 o'clock:

RGB and RBG: in both paintings, R is adjacent to G and B, G is adjacent to R and B, B is adjacent to R and G. In another way, they are equivalent if we traverse the second disk counter-clockwise.

YBG and BGY: These would be equivalent if rotation is allowed, because it amounts to choosing a different starting point.

On to the problem:

Consider a disk cut into k sectors and painted as described in the problem. Number the sectors as s_1, s_2, \dots, s_k . So s_k and s_1 are adjacent, therefore of different colors. There are two cases: either s_1 and s_{k-1} are painted the same color, or not.

Case 1: s_1 and s_{k-1} are painted the same color. In this case s_k can be painted one of the other 3 colors. Now remove s_k from the disk. The remaining $k - 1$ sectors can be considered a disk of $k - 2$ sectors, by merging s_1 and s_{k-1} (since they are painted the same color). There are S_{k-2} ways to color this remaining disk. Therefore there are $3S_{k-2}$ ways to color the original k -sector disk.

Case 2: s_1 and s_{k-1} are painted different colors. In this case s_k can be painted with one of the remaining 2 colors. Now remove s_k from the disk. The remaining $k - 1$ sectors form a valid painting by making s_1 and s_{k-1} adjacent (since they are painted with different colors). There are S_{k-1} ways to paint this remaining disk. Therefore there are $2S_{k-1}$ ways to color the original k -sector disk.

By the addition rule $S_k = 2S_{k-1} + 3S_{k-2}$. □

3.49.2 (b)

Find an explicit formula for S_n for $n \geq 2$.

Proof. $S_2 = 12$ because there are 4 colors to paint s_1 , 3 colors to paint s_2 .

$S_3 = 24$ because there are 4 colors to paint s_1 , 3 colors to paint s_2 , and, since s_3 is adjacent to both s_1 and s_2 , there are 2 colors to paint s_3 .

The characteristic equation is $t^2 - 2t - 3 = 0$ so $(t - 3)(t + 1) = 0$ which gives $t = -1, 3$. So the general form of the solution is $S_n = A \cdot (-1)^n + B \cdot 3^n$.

Solving $S_2 = 12 = A(-1)^2 + B \cdot 3^2$ we get $12 = A + 9B$. Solving $S_3 = 24 = A(-1)^3 + B \cdot 3^3$ we get $24 = -A + 27B$. Adding these two equations together gives $36 = 36B$ so $B = 1$ and $A = 3$.

Thus $S_n = 3 \cdot (-1)^n + 3^n$ for all $n \geq 2$. □

4 Exercise Set 9.4

4.1 Exercise 1

4.1.1 (a)

Proof. □

4.1.2 (b)

Proof. □

4.2 Exercise 2

4.2.1 (a)

Proof. □

4.2.2 (b)

Proof. □

4.3 Exercise 3

Proof. □

4.4 Exercise 4

Proof. □

4.5 Exercise 5

4.5.1 (a)

Proof. □

4.5.2 (b)

Proof.



4.6 Exercise 6

4.6.1 (a)

Proof.



4.6.2 (b)

Proof.



4.7 Exercise 7

Proof.



4.8 Exercise 8

Proof.



4.9 Exercise 9

4.9.1 (a)

Proof.



4.9.2 (b)

Proof.



4.10 Exercise 10

Proof.



4.11 Exercise 11

Proof.



4.12 Exercise 12

Proof.



4.13 Exercise 13

Proof.



4.14 Exercise 14

Proof.



4.15 Exercise 15

Proof.



4.16 Exercise 16

Proof.



4.17 Exercise 17

Proof.



4.18 Exercise 18

Proof.



4.19 Exercise 19

Proof.



4.20 Exercise 20

4.20.1 (a)

Proof.



4.20.2 (b)

Proof.



4.21 Exercise 21

Proof.



4.22 Exercise 22

Proof.



4.23 Exercise 23

Proof.



4.24 Exercise 24

Proof.



4.25 Exercise 25

Proof.



4.26 Exercise 26

Proof.



4.27 Exercise 27

Proof.



4.28 Exercise 28

Proof.



4.29 Exercise 29

Proof.



4.30 Exercise 30

Proof.



4.31 Exercise 31

Proof.



4.32 Exercise 32

Proof.



4.33 Exercise 33

Proof.



4.34 Exercise 34

Proof.



4.35 Exercise 35

Proof.



4.36 Exercise 36

Proof.



4.37 Exercise 37

4.37.1 (a)

Proof.



4.37.2 (b)

Proof.



4.38 Exercise 38

Proof.



4.39 Exercise 39

Proof.



4.40 Exercise 40

Proof.



5 Exercise Set 9.5

5.1 Exercise 1

5.1.1 (a)

Proof.



5.1.2 (b)

Proof.



5.2 Exercise 2

5.2.1 (a)

Proof.



5.2.2 (b)

Proof.



5.3 Exercise 3

Proof.



5.4 Exercise 4

Proof.



5.5 Exercise 5

5.5.1 (a)

Proof.



5.5.2 (b)

Proof.



5.5.3 (c)

Proof.



5.5.4 (d)

Proof.



5.5.5 (e)

Proof.



5.6 Exercise 6

5.6.1 (a)

Proof.



5.6.2 (b)

Proof.



5.6.3 (c)

Proof.



5.6.4 (d)

Proof.



5.6.5 (e)

Proof.



5.7 Exercise 7

5.7.1 (a)

Proof.



5.7.2 (b)

Proof.



5.7.3 (c)

Proof.



5.7.4 (d)

Proof.



5.8 Exercise 8

5.8.1 (a)

Proof.



5.8.2 (b)

Proof.



5.8.3 (c)

Proof.



5.8.4 (d)

Proof.



5.9 Exercise 9

5.9.1 (a)

Proof.



5.9.2 (b)

Proof.



5.10 Exercise 10

Proof.



5.11 Exercise 11

5.11.1 (a)

Proof.



5.11.2 (b)

Proof.



5.11.3 (c)

Proof.



5.11.4 (d)

Proof.



5.11.5 (e)

Proof.



5.11.6 (f)

Proof.



5.11.7 (g)

Proof.



5.11.8 (h)

Proof.



5.11.9 (i)

Proof.



5.12 Exercise 12

Proof.



5.13 Exercise 13

5.13.1 (a)

Proof.



5.13.2 (b)

Proof.



5.13.3 (c)

Proof.



5.13.4 (d)

Proof.



5.13.5 (e)

Proof.



5.14 Exercise 14

5.14.1 (a)

Proof.



5.14.2 (b)

Proof.



5.14.3 (c)

Proof.



5.14.4 (d)

Proof.



5.15 Exercise 15

5.15.1 (a)

Proof.



5.15.2 (b)

Proof.



5.15.3 (c)

Proof.



5.15.4 (d)

Proof.



5.16 Exercise 16

5.16.1 (a)

Proof.



5.16.2 (b)

Proof.



5.16.3 (c)

Proof.



5.17 Exercise 17

5.17.1 (a)

Proof.



5.17.2 (b)

Proof.



5.17.3 (c)

Proof.



5.17.4 (d)

Proof.



5.18 Exercise 18

Proof.



5.19 Exercise 19

5.19.1 (a)

Proof.



5.19.2 (b)

Proof.



5.19.3 (c)

Proof.



5.20 Exercise 20

5.20.1 (a)

Proof.



5.20.2 (b)

Proof.



5.20.3 (c)

Proof.



5.21 Exercise 21

Proof.



5.22 Exercise 22

Proof.



5.23 Exercise 23

Proof.



5.24 Exercise 24

5.24.1 (a)

Proof.



5.24.2 (b)

Proof.



5.24.3 (c)

Proof.



5.24.4 (d)

Proof.



5.25 Exercise 25

5.25.1 (a)

Proof.



5.25.2 (b)

Proof.



5.25.3 (c)

Proof.



5.25.4 (d)

Proof.



5.25.5 (e)

Proof.



5.26 Exercise 26

5.26.1 (a)

Proof.



5.26.2 (b)

Proof.



5.26.3 (c)

Proof.



5.26.4 (d)

Proof.



5.26.5 (e)

Proof.



5.26.6 (f)

Proof.



5.27 Exercise 27

5.27.1 (a)

Proof.



5.27.2 (b)

Proof.



5.27.3 (c)

Proof.



5.27.4 (d)

Proof.



5.28 Exercise 28

Proof.



5.29 Exercise 29

Proof.



5.30 Exercise 30

Proof.



6 Exercise Set 9.6

6.1 Exercise 1

6.1.1 (a)

Proof.



6.1.2 (b)

Proof.



6.2 Exercise 2

6.2.1 (a)

Proof.



6.2.2 (b)

Proof.



6.3 Exercise 3

6.3.1 (a)

Proof.



6.3.2 (b)

Proof.



6.3.3 (c)

Proof.



6.4 Exercise 4

6.4.1 (a)

Proof.



6.4.2 (b)

Proof.



6.4.3 (c)

Proof.



6.5 Exercise 5

Proof.



6.6 Exercise 6

Proof.



6.7 Exercise 7

Proof.



6.8 Exercise 8

Proof.



6.9 Exercise 9

Proof.



6.10 Exercise 10

Proof.



6.11 Exercise 11

Proof.



6.12 Exercise 12

Proof.



6.13 Exercise 13

Proof.



6.14 Exercise 14

Proof.



6.15 Exercise 15

Proof.



6.16 Exercise 16

6.16.1 (a)

Proof.



6.16.2 (b)

Proof.



6.16.3 (c)

Proof.



6.17 Exercise 17

6.17.1 (a)

Proof.



6.17.2 (b)

Proof.



6.17.3 (c)

Proof.



6.17.4 (d)

Proof.



6.18 Exercise 18

6.18.1 (a)

Proof.



6.18.2 (b)

Proof.



6.18.3 (c)

Proof.



6.18.4 (d)

Proof.



6.19 Exercise 19

6.19.1 (a)

Proof.



6.19.2 (b)

Proof.



6.20 Exercise 20

6.20.1 (a)

Proof.



6.20.2 (b)

Proof.



6.21 Exercise 21

Proof.



7 Exercise Set 9.7

7.1 Exercise 1

Proof.



7.2 Exercise 2

Proof.



7.3 Exercise 3

Proof.



7.4 Exercise 4

Proof.



7.5 Exercise 5

Proof.



7.6 Exercise 6

Proof.



7.7 Exercise 7

Proof.



7.8 Exercise 8

Proof.



7.9 Exercise 9

Proof.



7.10 Exercise 10

7.10.1 (a)

Proof.



7.10.2 (b)

Proof.



7.10.3 (c)

Proof.



7.11 Exercise 11

Proof.



7.12 Exercise 12

Proof.



7.13 Exercise 13

Proof.



7.14 Exercise 14

Proof.



7.15 Exercise 15

Proof.



7.16 Exercise 16

Proof.



7.17 Exercise 17

Proof.



7.18 Exercise 18

Proof.



7.19 Exercise 19

Proof.



7.20 Exercise 20

Proof.



7.21 Exercise 21

Proof.



7.22 Exercise 22

Proof.



7.23 Exercise 23

Proof.



7.24 Exercise 24

Proof.



7.25 Exercise 25

Proof.



7.26 Exercise 26

Proof.



7.27 Exercise 27

Proof.



7.28 Exercise 28

Proof.



7.29 Exercise 29

Proof.



7.30 Exercise 30

Proof.



7.31 Exercise 31

Proof.



7.32 Exercise 32

Proof.



7.33 Exercise 33

Proof.



7.34 Exercise 34

Proof.



7.35 Exercise 35

Proof.



7.36 Exercise 36

Proof.



7.37 Exercise 37

Proof.



7.38 Exercise 38

Proof.



7.39 Exercise 39

Proof.



7.40 Exercise 40

Proof.



7.41 Exercise 41

Proof.



7.42 Exercise 42

Proof.



7.43 Exercise 43

Proof.



7.44 Exercise 44

Proof.



7.45 Exercise 45

Proof.



7.46 Exercise 46

Proof.



7.47 Exercise 47

Proof.



7.48 Exercise 48

Proof.



7.49 Exercise 49

Proof.



7.50 Exercise 50

Proof.



7.51 Exercise 51

Proof.



7.52 Exercise 52

Proof.



7.53 Exercise 53

Proof.



7.54 Exercise 54

Proof.



7.55 Exercise 55

7.55.1 (a)

Proof.



7.55.2 (b)

Proof.



7.55.3 (c)

Proof.



7.55.4 (d)

Proof.



8 Exercise Set 9.8

8.1 Exercise 1

Proof.



8.2 Exercise 2

Proof.



8.3 Exercise 3

8.3.1 (a)

Proof.



8.3.2 (b)

Proof.



8.4 Exercise 4

Proof.



8.5 Exercise 5

Proof.



8.6 Exercise 6

Proof.

☐

8.7 Exercise 7

8.7.1 (a)

Proof.

☐

8.7.2 (b)

Proof.

☐

8.7.3 (c)

Proof.

☐

8.7.4 (d)

Proof.

☐

8.7.5 (e)

Proof.

☐

8.7.6 (f)

Proof.

☐

8.8 Exercise 8

Proof.

☐

8.9 Exercise 9

8.9.1 (a)

Proof.

☐

8.9.2 (b)

Proof.

☐

8.9.3 (c)

Proof.

☐

8.9.4 (d)

Proof.



8.9.5 (e)

Proof.



8.9.6 (f)

Proof.



8.10 Exercise 10

Proof.



8.11 Exercise 11

Proof.



8.12 Exercise 12

Proof.



8.13 Exercise 13

Proof.



8.14 Exercise 14

Proof.



8.15 Exercise 15

Proof.



8.16 Exercise 16

Proof.



8.17 Exercise 17

Proof.



8.18 Exercise 18

Proof.



8.19 Exercise 19

Proof.



8.20 Exercise 20

Proof.



8.21 Exercise 21

Proof.



8.22 Exercise 22

Proof.



8.23 Exercise 23

Proof.



9 Exercise Set 9.9

9.1 Exercise 1

Proof.



9.2 Exercise 2

Proof.



9.3 Exercise 3

Proof.



9.4 Exercise 4

9.4.1 (a)

Proof.



9.4.2 (b)

Proof.



9.5 Exercise 5

Proof.



9.6 Exercise 6

9.6.1 (a)

Proof.



9.6.2 (b)

Proof.



9.6.3 (c)

Proof.



9.7 Exercise 7

Proof.



9.8 Exercise 8

9.8.1 (a)

Proof.



9.8.2 (b)

Proof.



9.8.3 (c)

Proof.



9.9 Exercise 9

Proof.



9.10 Exercise 10

Proof.



9.11 Exercise 11

9.11.1 (a)

Proof.



9.11.2 (b)

Proof.



9.12 Exercise 12

Proof.



9.13 Exercise 13

9.13.1 (a)

Proof.



9.13.2 (b)

Proof.



9.14 Exercise 14

9.14.1 (a)

Proof.



9.14.2 (b)

Proof.



9.15 Exercise 15

9.15.1 (a)

Proof.



9.15.2 (b)

Proof.



9.15.3 (c)

Proof.



9.15.4 (d)

Proof.



9.16 Exercise 16

9.16.1 (a)

Proof.



9.16.2 (b)

Proof.



9.17 Exercise 17

Proof.



9.18 Exercise 18

Proof.



9.19 Exercise 19

Proof.



9.20 Exercise 20

Proof.



9.21 Exercise 21

Proof.



9.22 Exercise 22

Proof.



9.23 Exercise 23

9.23.1 (a)

Proof.



9.23.2 (b)

Proof.



9.23.3 (c)

Proof.



9.24 Exercise 24

9.24.1 (a)

Proof.



9.24.2 (b)

Proof.



9.25 Exercise 25

9.25.1 (a)

Proof.



9.25.2 (b)

Proof.



9.25.3 (c)

Proof.



9.25.4 (d)

Proof.



9.26 Exercise 26

Proof.



9.27 Exercise 27

Proof.



9.28 Exercise 28

9.28.1 (a)

Proof.



9.28.2 (b)

Proof.



9.28.3 (c)

Proof.



9.28.4 (d)

Proof.



9.29 Exercise 29

9.29.1 (a)

Proof.



9.29.2 (b)

Proof.



9.29.3 (c)

Proof.



9.29.4 (d)

Proof.



9.30 Exercise 30

9.30.1 (a)

Proof.



9.30.2 (b)

Proof.



9.30.3 (c)

Proof.



9.30.4 (d)

Proof.



9.31 Exercise 31

9.31.1 (a)

Proof.



9.31.2 (b)

Proof.



9.31.3 (c)

Proof.



9.32 Exercise 32

9.32.1 (a)

Proof.



9.32.2 (b)

Proof.



9.32.3 (c)

Proof.



9.32.4 (d)

Proof.



9.32.5 (e)

Proof.



9.33 Exercise 33

Proof.



9.34 Exercise 34

Proof.

