Study tables

Table 1 Sociodemographic characteristics

Forensic inpatients n = 66; psychiatric outpatients n = 25; healthy controls n = 24.

Characteristic	Forensic	Outpatient	Control	Test	p
Age, mean ± SD	39.32 ± 10.68	39.72 ± 8.32	37.00 ± 7.27	F(2, 112) = 0.626	.537
Education				$\chi^2(6) = 13.720$	.033
Primary school	9 (13.6)	0 (0.0)	0 (0.0)		
Middle school	33 (50.0)	10 (40.0)	9 (37.5)		
High school	18 (27.3)	13 (52.0)	14 (58.3)		
College/university	6 (9.1)	2 (8.0)	1 (4.2)		
Marital status				$\chi^2(4) = 30.889$	<.001
Single	44 (66.7)	13 (52.0)	3 (12.5)		
Married	15 (22.7)	10 (40.0)	21 (87.5)		
Widowed/divorced	7 (10.6)	2 (8.0)	0 (0.0)		
Employment				$\chi^2(2) = 15.238$	<.001
Employed	41 (62.1)	13 (52.0)	24 (100.0)		
Not employed	25 (37.9)	12 (48.0)	0 (0.0)		
Social security coverage				$\chi^2(2) = 17.188$	<.001
No coverage	31 (47.0)	10 (40.0)	0 (0.0)		
Coverage	35 (53.0)	15 (60.0)	24 (100.0)		

Values are n (%) unless stated otherwise. Employment refers to the pre-admission period for forensic inpatients. SD, standard deviation. Pearson chi-square results for education and marital status require caution because of sparse expected cells.

Table 2 Diagnostic distribution

Diagnosis	Forensic inpatients, n (%)	Outpatients, n (%)
Paranoid schizophrenia	28 (42.4)	9 (36.0)
Other schizophrenia subtypes and atypical psychosis	29 (43.9)	13 (52.0)
Bipolar affective disorder	9 (13.6)	3 (12.0)

Values are n (%), calculated within each patient group. Diagnostic labels follow the study team’s coding; healthy controls are not included in these diagnostic denominators. Rounding may prevent percentages from summing to exactly 100%.

Table 3 Affective temperament scores

Dimension	Forensic mean ± SD	Outpatient mean ± SD	Control mean ± SD	F(2,112)	Overall p
Depressive	10.09 ± 4.27	12.68 ± 4.29	3.79 ± 3.02	32.126	<.001
Cyclothymic	11.27 ± 5.34	11.28 ± 4.95	4.71 ± 3.92	16.418	<.001
Hyperthymic	12.52 ± 4.76	10.24 ± 5.40	10.96 ± 3.20	2.568	.081
Irritable	11.58 ± 4.78	8.12 ± 4.00	2.29 ± 1.83	44.248	<.001
Anxious	10.12 ± 4.83	12.32 ± 5.12	3.38 ± 3.90	24.966	<.001

Forensic n = 66; outpatient n = 25; control n = 24. Groups were compared using one-way ANOVA followed by Tukey tests. Comparisons between forensic inpatients and outpatients were significant for depressive (p = .020; SE of difference = 0.952) and irritable scores (p = .002; SE = 0.978). Outpatients had higher depressive scores, and forensic inpatients had higher irritable scores. The other three comparisons were not significant. Both patient groups scored higher than controls in depressive, cyclothymic, irritable and anxious temperament (each Tukey p < .001). No dominant-temperament thresholds were applied. SD, standard deviation; SE, standard error.