

Mathematical Analysis Study Notes

Emrys

Contents

1	Numerical Sequences and Series	2
1.1	Series	2
1.2	Positive Series	3
1.3	The Root and Ratio Tests	7

1 Numerical Sequences and Series

1.1 Series

Definition 1.1.1 Given a sequence $\{a_n\}$, we use the notation

$$\sum_{n=p}^q a_n \quad (p \leq q)$$

to denote the sum $a_p + a_{p+1} + \cdots + a_q$. And we call the symbolic expression

$$\sum_{n=1}^{\infty} a_n = a_1 + a_2 + \cdots + a_n + \cdots$$

an infinite series, or just a series.

With $\{a_n\}$ we associate a sequence $\{S_n\}$, where

$$S_n = \sum_{k=1}^n a_k.$$

The numbers S_n are called the partial sums of the series.

If $\{S_n\}$ converges to s , we say that the series converges, and write

$$\sum_{n=1}^{\infty} a_n = s.$$

The number s is called the sum of series; If $\{S_n\}$ diverges, the series is said to diverge.

Remark 1.1.1 The above s is called the sum of the series; but it should be clearly understood that s is the limit of a sequence of sums, and is not obtained simply by addition.

Remark 1.1.2 It is clear that every theorem about sequences can be state in series. The Cauchy criterion can be restated in the following form.

Remark 1.1.3 For convenience of notation, we shall simply write $\sum a_n$ in place of $\sum_{n=1}^{\infty} a_n$.

Theorem 1.1.1 (*Cauchy criterion*) $\sum a_n$ converges \iff for every $\varepsilon > 0$ there

exists an integer N such that if $m \geq n \geq M$

$$\left| \sum_{k=n}^m a_k \right| \leq \varepsilon \quad (1.1)$$

It can be proved by the Cauchy criterion of the limit of sequences. And in particular, by taking $m = n$, (1.1) becomes

$$|a_n| \leq \varepsilon \quad (n \geq N).$$

So we can get the following vanishing condition.

Proposition 1.1.1 (*Absolute Value Property*) $\sum a_n$ converges if $\sum |a_n|$ converges.

Proof: Since

$$|a_{m+1}| + \cdots + |a_n| \leq ||a_{m+1}| + \cdots + |a_n||,$$

we can easily prove the original proposition by Cauchy criterion.

Theorem 1.1.2 (*Vanishing condition*) If $\sum a_n$ converges, then $\lim_{n \rightarrow \infty} a_n = 0$.

Remark 1.1.4 The condition $a_n \rightarrow 0$ is not, however, sufficient to ensure convergence of $\sum a_n$. For instance, the series

$$\sum_{n=1}^{\infty} \frac{1}{n}$$

diverges; and we will prove it in the *Example 1.1.2*.

Example 1.1.1 let $q \in \mathbb{R}$, discuss the convergence and divergence of the geometric series $\sum q^n$.

Solution: We know that the partial sum of it is $S_n = \frac{q(1-q^n)}{1-q}$, so we have the series converges when $|q| < 1$ and diverges when $|q| \geq 1$.

Example 1.1.2 Determine the convergence and divergence of the harmonic series $\sum \frac{1}{n}$.

Solution: When $n \geq 1$, we have

$$\sum_{k=n+1}^{2n} \frac{1}{k} \geq \sum_{k=n+1}^{2n} \frac{1}{2n} = \frac{1}{2},$$

so by Cauchy criterion, we know the original series diverges.

1.2 Positive Series

Introduction

When studying a subject, we usually proceed from the specific to the general. Therefore, when researching series, we first focus on a specific type of series. It is called the positive series, which, as the name suggests, are series where each term is a positive number. But next, we will also research the non-negative series meanwhile.

Theorem 1.2.1 (*Fundamental Test*) The positive series converges $\iff \{S_n\}$ is bounded above.

Proof: Because $a_n > 0$, We conclude that this partial sum S_n is monotonically increasing with respect to n . It follows from the fact that a monotonically increasing and bounded above sequence must converge that $\{S_n\}$ converges.

Remark 1.2.1 The above test also holds for $a_n \geq 0$.

Example 1.2.1 If $\sum a_n$ is a positive series, and $\frac{a_{n+1}}{a_n} < \frac{n}{n+\alpha}$, where $\alpha > 1$, prove that $\sum a_n$ converges.

Proof: Since

$$\frac{a_{n+1}}{a_n} < \frac{n}{n+\alpha},$$

it follows that

$$a_n n + 1 < \frac{1}{\alpha - 1(na_n - (n+1)a_{n+1})},$$

Thus, we can conclude

$$S_n < a_1 + \frac{1}{\alpha - 1} \sum_{k=1}^{n-1} (ka_k - (k+1)a_{k+1}) = a_1 + \frac{1}{\alpha - 1} (a_1 - na_n) < \frac{\alpha}{\alpha - 1} a_1$$

Then we have the series is bounded above, and by the above theorem, it follows that the series converges.

Theorem 1.2.2 (*Cauchy Condensation Test*) Suppose $a_1 \geq a_2 \geq a_3 \geq \dots \geq 0$. Then the series $\sum_{n=1}^{\infty} a_n$ converges if and only if the series $\sum_{k=0}^{\infty} 2^k a_{2^k}$ converges.

Proof: By the *Theorem 1.2.1*, it suffices to consider boundness of the partial sums. let

$$S_n = \sum_{k=1}^n a_k = a_1 + a_2 + \dots + a_n,$$

$$T_m = \sum_{k=0}^m 2^k a_{2^k} = a_1 + 2a_2 + \dots + 2^m a_{2^m}$$

When $2^m \leq n \leq 2^{m+1}$, we have

$$\begin{aligned} S_n &\leq a_1 + (a_2 + a_3) + \cdots + (a_{2^m} + \cdots + a_{2^{m+1}-1}) \\ &\leq a_1 + 2a_2 + \cdots + 2^m a_{2^m} = T_m \end{aligned}$$

and

$$\begin{aligned} S_n &\geq a_1 + a_2 + (a_3 + a_4) + \cdots + (a_{2^{m-1}+1} + \cdots + a_{2^m}) \\ &\geq \frac{1}{2}a_1 + a_2 + 2a_4 + \cdots + 2^{m-1}a_{2^m} = \frac{1}{2}T_m \end{aligned}$$

Thus, we have S_n is bounded above $\iff T_m$ is bounded above. This completes the proof.

Example 1.2.2 $\sum \frac{1}{n^p}$ converges if $p > 1$ and diverges if $p \leq 1$.

Proof: If $p \leq 0$, divergence follows from the vanishing condition of series.

If $p > 0$, *Theorem 1.2.2* is applicable, we are led to the series

$$\sum_{k=0}^{\infty} 2^k \cdot \frac{1}{2^{kp}} = \sum_{k=0}^{\infty} 2^{(1-p)k} = \sum_{k=0}^{\infty} (2^{1-p})^k.$$

Now, $|2^{1-p}| < 1 \iff p > 1$, and the proof follows by the first example of the previous subsection.

Example 1.2.3 $\sum_{n=2}^{\infty} \frac{1}{n(\ln n)^p}$ converges if $p > 1$ and diverges if $p \leq 1$.

Proof: We consider the series

$$\sum_{k=1}^{\infty} 2^k \cdot \frac{1}{2^k (\ln 2^k)^p} = (\ln 2)^{-p} \sum_{k=1}^{\infty} \frac{1}{k^p}$$

By *Example 1.2.2*, we can easily prove the original proposition.

Corollary 1.2.1 The above procedure may evidently be continued. In other words, we can also conclude

$$\sum_{n=3}^{\infty} \frac{1}{n \ln n (\ln(\ln n))^p}$$

converges if and only if $p > 1$, and so forth. For instance,

$$\sum_{n=3}^{\infty} \frac{1}{n \ln n (\ln(\ln n))} \tag{1.2}$$

diverges, whereas

$$\sum_{n=3}^{\infty} \frac{1}{n \ln n (\ln(\ln n))^2} \quad (1.3)$$

converges.

Introduction

From the above corollary, we may now observe that the series (1.2) differ very little from those of (1.3). Still, one diverges, and the other converges. If we continue the process which led us from *Example 1.2.1* to *Example 1.2.2*, and then to (1.2) and (1.3), we also get pairs of convergent and divergent series whose terms differ even less than those of (1.2) and (1.3).

Thus, one might be led to the conjecture that there is a limiting situation of some sort, i.e. a "boundary" with all convergent series on one side, all divergent series on the other side. Or when n is sufficiently large, Is there a largest convergent series?

In the following, We shall show that this conjecture is false.

Proposition 1.2.1 Suppose positive series $\sum a_n$ converges, then there must exist positive series $\sum b_n$ converges and satisfies $\lim_{n \rightarrow \infty} \frac{b_n}{a_n} = +\infty$.

Proof: Let

$$S_n = a_1 + \cdots + a_n,$$

$$T_n = a_n + a_{n+1} + \cdots = \sum_{k=n}^{\infty} a_k.$$

We initially prove that S_n converges $\iff T_n$ converges to 0. We can easily prove that T_n converges iff T_n converges to 0. Hence, next we use the Cauchy criterion to prove

$$S_n \text{ converges} \iff T_n \text{ converges}.$$

Since for $m > n$,

$$T_m - T_n = a_n + \cdots + a_{m-1} = S_{m-1} - S_{n-1}$$

So we have for every $\varepsilon > 0$, there exists $N_1 > 0$, such that for $m > n > N$, $|T_m - T_n| < \varepsilon$, if and only if for every $\varepsilon > 0$, there exists $N_1 > 0$, such that for $m > n > N$, $|S_m - S_n| < \varepsilon$. In other words, the above proposition holds.

By

$$a_n = T_n - T_{n+1} = (\sqrt{T_n} - \sqrt{T_{n+1}})(\sqrt{T_n} + \sqrt{T_{n+1}}),$$

we set $b_n = \sqrt{T_n} - \sqrt{T_{n+1}}$, and $B_n = \sum b_n = \sqrt{T_1} - \sqrt{T_n + 1}$.

Hence, Σb_n converges owing to the convergence of T_n , and we have

$$\frac{b_n}{a_n} = \frac{1}{\sqrt{T_n} + \sqrt{T_{n+1}}} \rightarrow \infty, \quad (n \rightarrow \infty)$$

Introduction

From the proof of *Theorem 1.2.2*, we can know that for the non-negative series $\Sigma a_n, \Sigma b_n$, if $A_n \leq B_n$ and Σb_n converges, then Σa_n converges.

This is rooted from the *Theorem 1.2.1*, we can expand this to more general case.

Theorem 1.2.3 (*Comparison Test*) 1. If $|a_n| \leq c_n$ for $n \geq N$, where N is some fixed integer, and if Σc_n converges, then Σa_n converges.

2. If $a_n \geq d_n \geq 0$ or $a_n \leq d_n \leq 0$ for $n \geq N$, and if Σd_n diverges, then Σa_n diverges.

Remark 1.2.2 It is also possible to multiply c_n and d_n by a constant $C > 0$.

Remark 1.2.3 Whether analyzing a single series or comparing two series, we actually don't need to consider the finite initial terms. That is to say, we can only consider the case when n is sufficiently large, and we can even remove the finite terms from one series to compare it with another series.

1.3 The Root and Ratio Tests

Introduction

The above comparison test could generate other useful tests if we let b be some series whose convergence or divergence is known.

Sometimes if we cannot find some series which have some relationship with the original series, we can try some typical series like Σq^n .

Theorem 1.3.1 (*Root Test*) Given Σa_n , put $\alpha = \overline{\lim}_{n \rightarrow \infty} \sqrt[n]{|a_n|} > 0$. Then

- if $\alpha < 1$, Σa_n converges;
- if $\alpha > 1$, Σa_n diverges;
- if $\alpha = 1$, the test gives no information.

Proof: By

$$\alpha = \overline{\lim}_{n \rightarrow \infty} \sqrt[n]{|a_n|},$$

we can know

$$\forall \varepsilon > 0, \exists N > 0, \text{ s.t. } \forall n > N, |\sup\{\sqrt[k]{|a_k|} \mid k \geq n\} - \alpha| < \varepsilon,$$

i.e.

$$\alpha - \varepsilon < \sup\{\sqrt[k]{|a_k|} < \alpha + \varepsilon,$$

if $\alpha < 1$, it follows that

$$\forall k \geq n \geq N, \sqrt[k]{|a_k|} < \alpha + \varepsilon,$$

Because $0 < \alpha < 1$, we can set ε such that $\alpha + \varepsilon = q < 1$,

i.e.

$$\forall k \geq n \geq N, |a_k| < (\alpha + \varepsilon)^k = q^k, (0 < q < 1).$$

Since Σq^n converges if $|q| < 1$, $\Sigma |a_n|$ also converges.

Then it follows that Σa_n converges by absolute value proposition of series.

If $\alpha > 1$, it follows that

$$\forall n \geq N, \exists k \geq n, \text{ s.t. } \sqrt[k]{|a_k|} > \alpha - \varepsilon,$$

Because $\alpha > 1$, we can set ε such that $\alpha - \varepsilon = q > 1$,

i.e.

$$\forall N > 0, \exists k \geq N, |a_k| > (\alpha - \varepsilon)^k = q^k > 1.$$

Thus, we can conclude that $|a_n|$ fails to converge to 0, and then it follows that a_n also fails to converge to 0.

By vanishing condition, Σa_n diverges.

To prove the case $\alpha = 1$, we need to consider the series

$$\Sigma \frac{1}{n}, \quad \Sigma \frac{1}{n^2}.$$

For each of these series $\alpha = 1$, but the first diverges and the second converges.

Remark 1.3.1 The above theorem also applies to the case where the limit exists, and we can also write it in the form for sufficiently large n .

Introduction

when we utilize the comparison test, we just want to find a constant $C > 0$ and a positive series Σb_n such that $\frac{|a_n|}{b_n} < C$ when n is sufficiently large. And we can find that if when n is sufficiently large, $\{\frac{|a_n|}{b_n}\}$ is monotonically decreasing, then the previous equation holds.

In other words, $|\frac{a_{n+1}}{a_n}| \leq \frac{b_{n+1}}{b_n}$ for $n > N$ where N is a fixed integer. And next

we can let b_n be some typical series like Σq^n .

Theorem 1.3.2 (*Ratio Test*) The series Σa_n

- converges if $\limsup_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| < 1$,
- diverges if $\liminf_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| > 1$.
- cannot be determined to be convergent or divergent in other cases.

Proof: If $\limsup_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| < 1$, we can find $0 < q < 1$, and an integer N_1 , such that

$$\left| \frac{a_{n+1}}{a_n} \right| < q = \frac{q^{n+1}}{q^n}$$

for $n > N_1$, so it follows that $\left\{ \left| \frac{a_n}{q^n} \right| \right\}$ is monotonically decreasing when n is sufficiently large. Then the original proposition holds.

If $\liminf_{n \rightarrow \infty} \left| \frac{a_{n+1}}{a_n} \right| < 1$, we can find $r > 1$, and an integer N_2 , such that

$$\left| \frac{a_{n+1}}{a_n} \right| > r > 1$$

for $n > N_2$, and it is easily seen that the vanishing condition $a_n \rightarrow 0$ does not hold and the second proposition follows.

Remark 1.3.2 The above two theorems can also be written into the form of sufficient n . And this form can usually include the case when the ratio equal 1. However, in the next, we will introduce a more refined test to tackle the case when the previous theorem can not determine, i.e. the ratio $\frac{a_n}{a_{n+1}}$ is close to 1, but may fluctuate slightly around 1 or have higher-order corrections although n is sufficiently large.

Theorem 1.3.3 (*Guass Test*) Given a positive series Σa_n , Suppose that for sufficiently large n , the ratio of successive terms admits an asymptotic expansion:

$$\frac{a_n}{a_{n+1}} = 1 + \frac{h}{n} + \frac{p_n}{n^\alpha},$$

where $\alpha > 1$ and p_n is bounded, then Σa_n satisfies:

- if $h > 1$, Σa_n converges.
- if $h < 1$, Σa_n diverges.
- if $h = 1$, Σa_n is inconclusive.

Corollary 1.3.1 the above theorem have another form: Suppose that for sufficiently large n , the ratio of successive terms admits an expansion:

$$\frac{a_n}{a_{n+1}} = 1 + \frac{h}{n} + o\left(\frac{1}{n \ln n}\right),$$

then Σa_n satisfies:

- if $h > 1$, Σa_n converges.
- if $h \leq 1$, Σa_n diverges.

Theorem 1.3.4 (*Raabe Test*) If we turn the Guass test into the limit form, then we can obtain the Rabbe test: Given a positive series Σa_n , set

$$\lim_{n \rightarrow \infty} n\left(\frac{a_n}{a_{n+1}} - 1\right) = \alpha,$$

- if $\alpha > 1$, Σa_n converges;
- if $\alpha < 1$, Σa_n diverges;
- if $\alpha = 1$, the test gives no information.

Remark 1.3.3 When we research the convergence or divergence of series Σa_n , we can all initially determine $\Sigma |a_n|$ which is a non-positive series and even a positive series. $\Sigma |a_n|$ is more easily than Σa_n to analysis and can be applied by more tests. In the above root or ratio tests, actually, we all utilize this thought, just first analysis $\Sigma |a_n|$ and then think how to connect the original series Σa_n .