

Relational Geometry: An Architecture for Difference, Novelty, and Closure

Nature of the Program, Current Results, and the Navier–Stokes Laboratory

Le Matt Ansatz Di Ego

September 2026

Abstract

This document presents, in a common language intended for both academic readers and technically curious non-specialists, a research program that attempts to order several notions usually treated as primitive—difference, cardinality, dimension, identity, time, and scale—according to a more austere logical dependence. Its central thread is

unit $\rightarrow$ reflection $\rightarrow$ difference $\rightarrow$ relation $\rightarrow$ comparison $\rightarrow$ residual $\rightarrow$ recursion $\rightarrow$ closure $\rightarrow$ identity.

The proposal is not presented as a complete physical theory. Its strongest present form is that of a *research architecture*: it separates definitions, mathematical realizations, physical correspondences, exact results, hypotheses, and NO-GO statements. Current results include a formal distinction between novelty and closure; several conditional realizations of order four; a precise identification of where π enters when a discrete cycle is continuously parametrized; a separation between oscillatory and parabolic scale-rate laws; and, in Navier–Stokes, an exact material decomposition that turns the idea of relational “attunement” into a measurable dynamical question. The numerical evidence currently identifies mechanisms compatible with this reading, but it is not a proof of regularity or blowup.

For a first reading. The main thesis is not that “everything is relation.” It is more specific: if a theory aims to explain how structure and identity arise, it should avoid using as premises the very notions it seeks to derive. The program attempts to turn that conceptual discipline into mathematical obligations and, eventually, into falsifiable physical tests.

1 Nature and scope

The starting question is deliberately prior to any particular physics: *what is the minimal structure required for a difference to exist, to be preserved, and eventually to become part of an identity?*

The initial unit is not imagined as a particle, a zero-size point, or a spatial vacuum. It is a logical limit with no accredited internal differences and no available external reference. To speak at that stage of distance, direction, time, energy, or number would introduce structures that the program is trying to explain later.

The first proposed operation is minimal self-reference: the unit can act as the reference for an expression of itself. If reference and expression cease to be indistinguishable, a difference appears. Difference is therefore placed before cardinality: first there must be

something that can be distinguished; only afterward can an architecture preserve several records and count them.

In this sense, “information” in the program does not begin as a bit or as entropy. More austere, it begins as a difference whose removal would make indistinguishable states that the architecture must keep distinct.

Epistemic rule. The program distinguishes between [Definition], [Exact], [Program derivation], [Correspondence], [Hypothesis], [Open], and **NO-GO**. The purpose of this classification is not rhetorical caution. It is to prevent a philosophical intuition from being presented as though it were already a mathematical or physical consequence.

2 From difference to residual

A difference can only be sustained relative to a reference. Relation therefore appears before richer physical notions such as force or interaction. A relation allows comparison, and a comparison can be represented schematically as

$$\mathcal{C}(a, b) = (q, r),$$

where q is the part that the current architecture can resolve and r is what remains unrepresented without loss of relevant information.

The residual is not matter, energy, or noise by definition. It is a *representational debt*. If $r \neq 0$ and discarding it would change the distinguishable state, the architecture must preserve it somehow. The result of a comparison can then become a new reference. Recursion appears there, together with a logical depth that should not yet be confused with physical time.

This makes it possible to formulate novelty more strictly than mere surprise: something is *new* when it cannot be reduced to the currently available closure without losing an accredited difference.

3 The central conceptual result: novelty is not closure

One of the most important advances of the program has been to separate concepts that are often conflated. Return, consistency, and identity closure are not the same thing.

A cycle may return to its starting point without producing identity. A new relation may increase the rank of a structure without closing anything. A residual may disappear in a local comparison without the architecture becoming able to reconstruct itself.

The working definition is stronger:

Relational closure. An image closes when the architecture can contain it and reconstruct its belonging through internal relations, without requiring a new external reference in order to recognize it.

This leads to a stronger conclusion about identity. A new identity cannot be pure repetition, because then it would not be new; nor can it be absolute novelty without reference, because there would be no criterion by which to recognize it as belonging to the system.

Particularization therefore requires, at least conceptually,

$$\boxed{\text{inheritance} + \text{novelty} + \text{closure}.}$$

Identity is no longer treated as an immobile substance, but as an architecture capable of preserving its criterion of recognition while changing. In the current terminology of the program, this dynamic stability is better described as *attunement*: transformations remain compatible with the relations that make it possible to recognize “the same” through change.

4 What has been learned about four and π

The number four appears through several routes in the program, but the mature conclusion is precisely that these routes should not be treated as though they constituted a single proof.

In one algebraic realization, if there exists an operator J such that

$$J^2 = -I,$$

then

$$I \rightarrow J \rightarrow -I \rightarrow -J \rightarrow I, \quad J^4 = I,$$

and a discrete cycle of order four is obtained. This is exact once $J^2 = -I$ is assumed. What remains unproved is that the foundational architecture must necessarily select an operator of this kind.

Other routes—a projective/quaternionic construction and a binary realization in which the first combination of inheritance and novelty appears at $\varepsilon = 4$ —again point toward four, but under different assumptions. The scientific problem is no longer to observe that “four appears several times,” but to prove or falsify that these appearances are realizations of one underlying abstract structure.

A similar distinction applies to π . The discrete cycle above does not require π . If a continuous interpolation is chosen,

$$e^{\theta J} = \cos \theta I + \sin \theta J,$$

then periodicity 2π appears exactly. The conclusion is not that π has been derived “from nothing,” but something more precise: *π enters when a discrete orientation of order four is represented as a continuous periodic trajectory.*

Methodological result. Separating “why four” from “why π ” removes an important circularity. In the present realizations, the first is a problem of discrete closure; the second appears when continuous and metric structure is added.

5 Time, rate, and scale

Recursion introduces logical order, but logical order is not yet duration. A disciplined way to introduce time is to compare rates of transformation. Before choosing a clock, the

elementary object may be a dimensionless ratio

$$\frac{\varpi_i}{\varpi_j}.$$

Choosing one process as reference turns that comparison into an operational measure of duration.

In a quadratic spectral realization,

$$G = C^\dagger C, \quad \ddot{f} + \Gamma G f = 0,$$

an eigenmode $G u_i = \lambda_i u_i$ satisfies

$$\varpi_i^2 = \Gamma \lambda_i.$$

If $R_i = \lambda_i^{-1/2}$, then

$$R_i |\varpi_i| = \sqrt{\Gamma}.$$

But this relation is not universal. Navier–Stokes has parabolic scaling; there a natural combination is of the form $R^2 \varpi$. This difference produced an important NO-GO result: a linear scale–rate law cannot be exported to every dynamics merely because it is elegant in an oscillatory realization.

The conceptual consequence matters: the relation between scale and rate depends on the dynamical class. The program must derive the appropriate law from each operator rather than impose one temporal geometry on every system.

6 Navier–Stokes as a physical laboratory

The program’s most concrete contact with a mature physical theory is currently in vorticity dynamics. For $q = |\omega| > 0$ and $\xi = \omega/q$,

$$\omega = q\xi, \quad \frac{D_t \omega}{q} = a\xi + b,$$

with

$$a = D_t \log q, \quad b = D_t \xi, \quad b \perp \xi.$$

The identity separates exactly two aspects of material change: logarithmic growth along the current direction and turning of that direction.

If

$$\frac{D_t \omega}{q} = S\xi + r, \quad S\xi = \alpha\xi + \tau, \quad r = r_{\parallel}\xi + r_{\perp},$$

then

$$b = \tau + r_{\perp}.$$

For $a > 0$, define the diagnostic quantities

$$D_{\perp} = \frac{|b|}{|\tau| + |r_{\perp}|}, \quad \Lambda_{\perp} = \frac{|\tau| + |r_{\perp}|}{a},$$

so that

$$\frac{|b|}{a} = D_{\perp} \Lambda_{\perp}.$$

$D_{\perp} \ll 1$ identifies cancellation between transverse contributions; $\Lambda_{\perp} \ll 1$ identifies transverse scarcity. This factorization is diagnostic: it classifies a mechanism, but does not by itself impose a dynamical law.

The strongest step in the present stage appears when accumulated growth is used as the coordinate

$$dG = a dt, \quad G = \log \frac{q(t_2)}{q(t_1)}.$$

Defining

$$V = \frac{D_t \omega}{q} = a\xi + b, \quad \mathcal{Y} = D_t V,$$

the transverse component of the evolution of b satisfies exactly

$$\boxed{P_{\xi}^{\perp} \frac{db}{dG} = \frac{\mathcal{Y}_{\perp}}{a} - b, \quad a > 0.}$$

Dynamical reading. For each unit of logarithmic growth of the reference, the existing transverse difference carries a homogeneous term $-b$. Preserving or increasing that difference requires a transverse contribution from $\mathcal{Y}/a$. In the language of the program, transverse novelty is not preserved merely because it has appeared: it must be continually *re-accredited* by the dynamics.

This reading does not turn the equation into an ontological proof. Its importance is different: it provides an exact, measurable realization of an idea that was only conceptual at the foundational level.

7 What the current data say

The associated technical manuscript analyzes the forced JHTDB dataset `isotropic1024coarse`. In a screen of 100,000 points, the low-material-turning tail remained present within the sampled top 0.1% of vorticity. Two states were audited with fourth-order finite differences; one is more sensitive to stencil width than the other. Short episodes showed growth of q while the direction remained nearly parallel, and one representative case exhibited substantial, nearly opposite transverse contributions, consistent with a cancellation-dominated mechanism.

ID	q (FD4)	$\rho_{\perp}$ (7 pt)	$\rho_{\perp}$ (5 pt)	approx. G
961	146.364	0.0404	0.0411	0.0685
995	223.441	0.0715	0.0465	0.0514

Table 1: Two representative trajectories from the diagnostic. ID 995 illustrates the methodological sensitivity that prevents the classification from being presented as a universal robust law.

The most defensible result is not that turbulence “prefers” a particular state. It is more concrete: *small material turning can arise from cancellation between large transverse terms, not only because those terms are themselves small*. This changes the physical question: it is not enough to measure how much the direction turns; one must identify which terms

sustain or destroy the cancellation.

The Burgers vortex also provides an exact NO-GO: zero transverse turning can coexist with positive amplification without implying a singularity. Therefore, “low turning” cannot be used in isolation as a danger criterion. Any useful criterion must also incorporate scale, amplitude, domain/energy assumptions, or an additional dynamical discriminant.

The base manuscript states the boundary explicitly: the present data classify a mechanism, but do not provide a PDE bound or a proof of regularity or blowup for unforced three-dimensional Navier–Stokes.¹

8 Academic positioning

The program does not arise in an intellectual vacuum. It has clear neighbors, but should not be conflated with them.

Neighboring area	Genuine overlap	Difference that must be preserved
Relational quantum mechanics	priority of relations over absolute states	it begins from quantum formalism; here the attempt is to order concepts before choosing a physical theory
Operational reconstructions	deriving formalisms from more transparent principles	they often assume systems, operations, and probabilities that are already defined
Information geometry	distinguishability can acquire geometric structure	it does not derive the existence of difference or identify that geometry with physical space
Fixed points and self-reference	they provide rigorous languages for closure and self-reference	a fixed point alone is insufficient to produce novelty or identity
Vorticity geometry	the direction of ω has dynamical and regularity consequences	the proposed contribution must go beyond “direction matters” and yield a new discriminant

This comparison suggests a simple academic requirement: the program becomes scientifically substantive if one of its concepts produces something more than a new description—a necessity theorem, an invariant, a bound, a predictive classification, or a falsification not already built into the definitions.

9 An interpretive extension: mathematical discovery and generative systems

The same architecture can be used, cautiously, to think about machine-assisted mathematical discovery. Not as a theory of the internal operation of an LLM, but as a language for the process of representation.

¹The associated technical paper states this boundary exactly: the material decomposition and factorization identify the cancellation mechanism; the dynamics of persistence and accumulated growth remain open.

A mathematical difficulty may contain constraints that the available representation cannot hold simultaneously. In that case, the “residual” would not be an error, but a sign that the current architecture cannot yet express all relevant relations. A reorganization may turn that residual into a new reference and allow a closure that was previously inaccessible.

Under this reading, discovery does not always mean receiving more information. Sometimes it may mean constructing a different relational space in which information already available can finally coexist. This is an interpretive extension of the program, not evidence about the internal mechanics of the models. Its value will depend on whether it can eventually be translated into observable comparisons between successful and failed reasoning trajectories.

10 What can be sustained today

Status	Claim
[Definition]	Difference, residual, closure, identity, and attunement have an explicit conceptual order and are not identified with one another.
[Conditionally exact]	$J^2 = -I \Rightarrow J^4 = I$, and the interpolation $e^{\theta J}$ has period 2π .
[Exact]	The material vorticity decomposition, the diagnostic factorization, and the transverse law in growth coordinates.
[Finite evidence]	Forced high-vorticity episodes with low material turning and, in selected cases, significant transverse cancellation.
[NO-GO]	Novelty does not imply closure; independence does not imply orthogonality; low turning does not imply singularity; $R\varpi = \text{const}$ is not a universal law.
[Open]	A theorem unifying the closure realizations, a necessary derivation of order four, and a relational invariant imposing a genuinely new physical restriction.

11 Conclusions and decisive tests

The program has advanced far enough to leave two extremes behind. It is no longer only a philosophical narrative, because it contains exact identities, algebraic realizations, NO-GO results, and computational protocols. But it is not yet a closed physical theory, because its most ambitious bridges remain identifiable hypotheses.

The strongest conceptual conclusion can be stated as follows: *a difference does not become identity merely by existing or by repeating, but when an architecture can preserve novelty while at the same time reconstructing its belonging*. This distinction between difference, novelty, and closure organizes the entire program.

The cleanest mathematical conclusion is that several suggestive structures—order four, transversality, periodicity, and scale–rate relations—can appear without being postulated as primitives, but only within realizations whose assumptions must remain visible. The next major result will not be finding another “four,” but proving why a minimal class of closures must produce it, or proving that it need not.

The most promising physical conclusion currently comes from Navier–Stokes: directional

attunement can be studied as a dynamical competition between growth of the reference and re-accreditation of transverse difference. The equation

$$P_{\xi}^{\perp} \frac{db}{dG} = \frac{\mathcal{Y}_{\perp}}{a} - b$$

turns that sentence into an exact identity. The question now is whether pressure, viscosity, and forcing terms allow one to derive from it a new restriction that survives adversarial benchmarks.

The decisive test of the program is not rhetorical. It must produce at least one of the following: a nontrivial closure theorem; a demonstrable necessity of order four; an operational construction of time that recovers known limits; or a dynamical invariant that imposes a new quantitative physical restriction. If none of these routes succeeds, the program itself will have delimited its scope.

Perhaps this is its most useful feature in the current state: metaphysics is not presented as a conclusion, but as a source of questions that must end in reproducible mathematics. The aim is not to force the world to resemble the initial image. It is to subject that image to enough relations to find out which part of it can close without contradiction.

Selected references

- [1] C. Rovelli, “Relational Quantum Mechanics,” *International Journal of Theoretical Physics* 35 (1996), 1637–1678.
- [2] J. B. Barbour and B. Bertotti, “Mach’s Principle and the Structure of Dynamical Theories,” *Proceedings of the Royal Society A* 382 (1982), 295–306.
- [3] L. Hardy, “Quantum Theory From Five Reasonable Axioms,” arXiv:quant-ph/0101012 (2001).
- [4] G. Chiribella, G. M. D’Ariano and P. Perinotti, “Informational derivation of quantum theory,” *Physical Review A* 84 (2011), 012311.
- [5] F. W. Lawvere, “Diagonal arguments and cartesian closed categories,” *Lecture Notes in Mathematics* 92, Springer (1969).
- [6] F. Nielsen, “An Elementary Introduction to Information Geometry,” *Entropy* 22 (2020), 1100.
- [7] P. Constantin and C. Fefferman, “Direction of vorticity and the problem of global regularity for the Navier–Stokes equations,” *Indiana University Mathematics Journal* 42 (1993), 775–789.
- [8] H. Beirão da Veiga and L. C. Berselli, “On the regularizing effect of the vorticity direction in incompressible viscous flows,” *Differential and Integral Equations* 15 (2002), 345–356.
- [9] J. D. Gibbon, D. D. Holm, R. M. Kerr and I. Roulstone, “Quaternions and particle dynamics in the Euler fluid equations,” *Nonlinearity* 19 (2006), 1969–1983.
- [10] M. Holzner et al., “Viscous tilting and production of vorticity in homogeneous turbulence,” *Physics of Fluids* 22 (2010), 061701.
- [11] D. Buaria, J. M. Lawson and M. Wilczek, “Twisting vortex lines regularize Navier–Stokes turbulence,” *Science Advances* 10 (2024), eado1969.
- [12] Johns Hopkins Turbulence Databases, “Forced Isotropic Turbulence,” dataset and documentation.
- [13] Le Matt Ansatz Di Ego, “Transient Transverse Cancellation in Material Vorticity Amplification: Exact identities, forced-turbulence evidence, and a relational closure interpretation,” manuscript, September 2026.