

Differential Equations Project: Phase 3

Arnav Patri and Shashank Chidige

October 23, 2022

A stock's price S_t (in USD) can be predicted with respect to time t (in days from an initial time) using a growth function; that is, by relating the rate at which the stock price is changing to the current stock price as a proportion:

$$\frac{dS_t}{dt} \propto S_t$$

The constant of proportionality is the drift μ of the stock, which is the rate of change of the expected value of the stock price (which is not the expected value of its rate of change):

$$\frac{dS_t}{dt} = \mu S_t \quad \text{where} \quad \mu = \frac{\Delta \mathbb{E}[S_t]}{\Delta t}$$

The stock market is constantly volatile, though, changing in unpredictable ways. This randomness element can be modeled by a randomness term, the rate of change of the standard Wiener process W_t :


$$\text{rate of randomness} \propto \frac{dW_t}{dt}$$

The proportionality constant is the volatility σ of the stock, which is simply its standard deviation. Adding this term,

$$\frac{dS_t}{dt} = \mu S_t + \sigma \frac{dW_t}{dt} \quad \text{where} \quad \sigma = \frac{\sum (S_{t,i} - \bar{S}_t)}{n - 1}$$

The randomness term ensures that each trial of the model yields a different graph. The following

is a sample graph:



The DE can be rewritten as

$$dS_t = \mu_t dt + \sigma_t dW_t$$

where $\mu_t = \mu S_t$ and $\sigma_t = \sigma S_t$. Integrating and reparameterizing μ_t , σ_t and W_t with s ,

$$S_t = \int_0^t \mu_s ds + \int_0^t \sigma_s dW_s + S_0$$

where S_0 is the constant of integration (the initial stock price). This makes S_t an Itô process, a stochastic process expressible as the sum of two integrals, one with respect to a stochastic process and another with respect to time, and a constant.

The Taylor expansion of a twice-differentiable scalar function $f(t, s)$ is

$$df = \frac{\partial f}{\partial t} dt + \frac{\partial f}{\partial s} ds + \frac{1}{2} \frac{\partial^2 f}{\partial s^2} ds^2 + \dots$$

Substituting S_t for s and appropriately substituting for ds yields

$$df + \frac{\partial f}{\partial t} dt + \frac{\partial f}{\partial s} (\mu_t dt + \sigma_t dW_t) + \frac{\partial^2 f}{\partial s^2} (\mu_t^2 dt^2 + 2\mu_t \sigma_t dt dW_t + \sigma_t^2 dW_t^2) + \dots$$

As dt approaches 0, dt^2 and $dt dW_t$ tend to zero faster than dW_t^2 . Substituting 0 for dt^2 and $dt dW_t$ and dt for dW_t^2 yields

$$df = \left(\frac{\partial f}{\partial t} + \mu_t \frac{\partial f}{\partial s} + \frac{\sigma_t^2}{2} \frac{\partial^2 f}{\partial s^2} \right) dt + \sigma_t \frac{\partial f}{\partial s} dW_t$$

This is itself an Itô process. Itô's lemma states that for any Itô process S_t and any twice-differentiable function $f(t, s)$, $f(t, S_t)$ is an Itô process.

Let $f(S_t) = \ln S_t$. Applying Itô's lemma,

$$\begin{aligned} df &= f'(S_t) dS_t + \frac{1}{2} f''(S_t) (dS_t)^2 \\ &= \frac{1}{S_t} dS_t - \frac{1}{2S_t^2} (S_t^2 \sigma^2 dt) \\ &= \frac{1}{S_t} (\sigma S_t dW_t + \mu S_t dt) - \frac{\sigma^2}{2} dt \\ &= \sigma dW_t + \left(\mu - \frac{\sigma^2}{2} \right) dt \end{aligned}$$

Integrating the separable DE,

$$f(t) = \ln S_t = \ln S_0 + \sigma W_t + \left(\mu - \frac{\sigma^2}{2} \right) dt$$

This can finally be exponentiated, yielding S_t :

$$S_t = S_0 e^{\left(\mu - \frac{\sigma^2}{2} \right) t + \sigma W_t}$$

Substituting for S_0 , μ , and σ ,

$$S_t = 141.501 e^{\left(0.00064 + \frac{4.356^2}{2} \right) t + 4.356 W_t}$$

The following are the Matlab code, solution (for the deterministic term), and graph:

```

Solution.m  x  +
1      close all;
2      clear all;
3
4      dom = [0, 30];
5      n = 1000;
6      dt = (dom(2) - dom(1))/50;
7      S0 = 141.501;
8
9      ndist = makedist('Normal', 0 , sqrt(dt));
10
11     mu = 0.00064;
12     sigma = 4.356;
13
14     tvals = linspace(dom(1), dom(2), n);
15     Svals = zeros(1, n);
16     Svals(1) = S0;
17     for i = 2:n
18         t = dom(1) + (i - 1) .* dt;
19         S = Svals(i - 1);
20         dW = random(ndist);
21         Svals(i) = S + mu .* dt + sigma .* dW
22     end
23     figure()
24     hold on;
25     plot(tvals, Svals);
26     xlabel('time (days from 9/27/22)')
27     ylabel('stock price (USD)')
28
29     % Solving for the non-stochastic term
30     syms s(t)
31     ode = diff(s, t) == s;
32     sol(t) = dsolve(ode);

```


```

sol  x
1x1 symfun

val(t) =

C1*exp(t)

```



The following is the [Excel spreadsheet](#).