Discrete Math

Arnav Patri

September 3, 2022

Contents

4	Nui	mber Theory and Cryptography
	4.2	Integer Representations and Algorithms
		4.2.2 Representations of Integers
	4.3	Primes and Greatest Common Divisors
		4.3.2 Primes
		4.3.3 Trial Division
	4.1	Divisibility and Modular Arithmetic
6	Coı	ınting
	6.1	The Basics of Counting
	6.3	Permutations and Combinations
	6.4	Binomial Coefficients
	6.5	Generalized Permutations and Combinations

Chapter 4

Number Theory and Cryptography

4.2 Integer Representations and Algorithms

Definition of a Number A number is dependent on a given base and its place value and digits.

4.2.2 Representations of Integers

A base b has b-1 digits. The first digit from the right is multiplied by b^0 , the second by b^1 , and so on. The number itself is the sum of each digit multiplied by b raised to the power of its respective place value.

0 is a member of every base (except sometimes base 1).

Let b be an integer greater than 1. If b is an integer greater than 1 and n is positive, then n can be expressed uniquely in the form

$$n = a_k b^k + a_{k-1} b^{k-1} + \dots + a_1 b + a_0$$

A number in base b is denoted by $(n)_b$.

A number is a linear combination of its digits and their place values.

Constructing Base b Expansions Given an integer n to be represented in base b,

```
q:=n
k:=0
while q \neq 0
a:= a \mod b
q:= q \operatorname{div} b
k:= k+1
return (a_{k-1}, \ldots, a_1, a_0) \{(a_{k-1} \ldots a_1 a_0)_b \text{ is the base } b \text{ expansion of } n\}
```

A number in its own base is always represented as 10.

Addition and multiplication in base b follows the same conventions as that of base 10.

To add two numbers a and b in base 2, their rightmost bits a_0 and b_0 can be added such that

$$a_0 + b_0 = 2c_0 + s_0$$

where s_0 is s_0 is the rightmost bit of the binary expansion of the sum and c_0 is the **carry**, being either 0 or 1. This process can be repeated.

$$c_0 = \frac{a_0 + b_0 - s_0}{2}$$

4.3 Primes and Greatest Common Divisors

4.3.2 Primes

A **prime number** is a whole number whose only factors are 1 and itself. By definition, it does not appear on the multiplication table. A nonprime positive integer is called **composite**

The Fundamental Theorem of Arithmetic Every integer greater greater than 1 can be written uniquely as the product of one or more primes.

Two numbers are relatively prime or coprime if their greatest common factor (GCF) is 1. If n is divisible by a and b, then it is also divisible by $a \times b$.

4.3.3 Trial Division

4.1 Divisibility and Modular Arithmetic

Divisibility Rules

If the difference between a 2 times a number's last digit and the rest of the number is divisible by 7 or 0, the number is as well. If the difference between a number's last digit multiplied by 5 and the rest of the numbers is divisible by 17 or 0, the number is divisible by 17.

If the sum of 2 times the last digit of a number and the rest of the digits is divisible by 19, the number is divisible by 19.

If the sum of 7 times the last digit of a number and the rest of the number is divisible by 23, then so is the number.

If the difference between 3 times the last digit of a number and the rest of the number is divisible by 31, then so is the number.

Ends in 1 : not divisible by 2.

1 + 1 + 1 = 3, $3 \mod 3 = 0$: divisible by 3.

101,001/3 = 33,667 Ends in 1 : not divisible by 5.

Ends in 1: not divisible by 2.

1+1+1+1+1+1=6, $6 \mod 3=0$: divisible by 3.

Ends in 1 : not divisible by 5. 111 - 111 = 0 : divisible by 7. 33,667/7 = 5,291.

Chapter 6

Counting

- 6.1 The Basics of Counting
- 6.3 Permutations and Combinations
- 6.4 Binomial Coefficients
- 6.5 Generalized Permutations and Combinations