

whole adaptive diffusion

Adaptive diffusion tree built from the full allGO feature matrix, using the full matrix rather than a decomposed cosine subspace.

Method: whole adaptive diffusion. Values shown on each tree page are: GO-BIC/gene, where lower is better within comparable non-degenerate trees; coherent, the fraction of clusters passing the GO-enrichment rule; within TF-IDF, the weighted mean within-cluster GO-vector cosine; largest, the fraction of genes in the largest cluster; and singletons, the singleton-gene fraction.

Best tree in this method-specific PDF:

whole_adaptive_diffusion




GO-BIC/gene: 1475.90

clusters: 20

coherent clusters: 15/20 (0.750)

largest cluster fraction: 0.568

singleton-gene fraction: 0.000



```
whole_adaptive_diffusion

display rank: 1
raw GO-IC rank: 15
quality tier: quality_plausible

method family: whole_adaptive_diffusion
weighting: whole
block: whole matrix
modes: None-None

GO-BIC active: 888491.19
GO-BIC active/gene: 1475.90
GO active params: 22978
GO logLik: -270141.06

clusters: 20
largest cluster: 342 (0.568)
singletons: 0 (0.000 of clusters)
median cluster size: 11.5

coherent clusters: 15 (0.750)
median significant GO terms: 140.5
min cluster q: 1.04e-39
median within TF-IDF cosine: 0.3445
weighted within TF-IDF cosine: 0.1433
```

