

MPF objective function for an Ising model

Jascha Sohl-Dickstein

August 31, 2011

This document derives the MPF objective function for the case of an Ising model, and connections to all states which differ by a single bit flip. It is written quickly, and not well proofread!! Typos and missing steps are likely...

For the Ising model, the energy function is

$$E = \mathbf{x}^T \mathbf{J} \mathbf{x} \quad (1)$$

where $\mathbf{x} \in \{0, 1\}^N$, $\mathbf{J} \in \mathcal{R}^{N \times N}$, and \mathbf{J} is symmetric ($\mathbf{J} = \mathbf{J}^T$). The MPF objective function is

$$K = \sum_{\mathbf{x} \in \mathcal{D}} \sum_n \exp \left(\frac{1}{2} [E(\mathbf{x}) - E(\mathbf{x} + \mathbf{d}(\mathbf{x}, n))] \right) \quad (2)$$

where the sum over n indicates a sum over all data dimensions, and the function $\mathbf{d}(\mathbf{x}, n) \in \{-1, 0, 1\}^N$ is

$$\mathbf{d}(\mathbf{x}, n)_i = \begin{cases} 0 & i \neq n \\ -(2x_i - 1) & i = n \end{cases} \quad (3)$$

For the Ising model, the MPF objective function becomes

$$K = \sum_{\mathbf{x} \in \mathcal{D}} \sum_n \exp \left(\frac{1}{2} [\mathbf{x}^T \mathbf{J} \mathbf{x} - (\mathbf{x} + \mathbf{d}(\mathbf{x}, n))^T \mathbf{J} (\mathbf{x} + \mathbf{d}(\mathbf{x}, n))] \right) \quad (4)$$

$$= \sum_{\mathbf{x} \in \mathcal{D}} \sum_n \exp \left(\frac{1}{2} [\mathbf{x}^T \mathbf{J} \mathbf{x} - (\mathbf{x}^T \mathbf{J} \mathbf{x} + 2\mathbf{x}^T \mathbf{J} \mathbf{d}(\mathbf{x}, n) + \mathbf{d}(\mathbf{x}, n)^T \mathbf{J} \mathbf{d}(\mathbf{x}, n))] \right) \quad (5)$$

$$= \sum_{\mathbf{x} \in \mathcal{D}} \sum_n \exp \left(-\frac{1}{2} [2\mathbf{x}^T \mathbf{J} \mathbf{d}(\mathbf{x}, n) + \mathbf{d}(\mathbf{x}, n)^T \mathbf{J} \mathbf{d}(\mathbf{x}, n)] \right) \quad (6)$$

$$= \sum_{\mathbf{x} \in \mathcal{D}} \sum_n \exp \left(-\frac{1}{2} \left[2 \sum_i x_i J_{in} (1 - 2x_n) + J_{nn} \right] \right) \quad (7)$$

$$= \sum_{\mathbf{x} \in \mathcal{D}} \sum_n \exp \left(\left[(2x_n - 1) \sum_i x_i J_{in} - \frac{1}{2} J_{nn} \right] \right). \quad (8)$$

Assume the symmetry constraint on \mathbf{J} is enforced by writing it in terms of a another possibly asymmetric matrix $\mathbf{J}' \in \mathcal{R}^{N \times N}$,

$$\mathbf{J} = \frac{1}{2}\mathbf{J}' + \frac{1}{2}\mathbf{J}'^T. \quad (9)$$

The derivative of the MPF objective function with respect to \mathbf{J}' is

$$\begin{aligned} \frac{\partial K}{\partial J'_{lm}} = & \frac{1}{2} \sum_{\mathbf{x} \in \mathcal{D}} \exp \left(\left[(2x_m - 1) \sum_i x_i J_{im} - \frac{1}{2} J_{mm} \right] \right) \left[(2x_m - 1) x_l - \delta_{lm} \frac{1}{2} \right] \\ & + \frac{1}{2} \sum_{\mathbf{x} \in \mathcal{D}} \exp \left(\left[(2x_l - 1) \sum_i x_i J_{il} - \frac{1}{2} J_{ll} \right] \right) \left[(2x_l - 1) x_m - \delta_{ml} \frac{1}{2} \right], \end{aligned} \quad (10)$$

where the second term is simply the first term with indices l and m reversed.

Note that both the objective function and gradient can be calculated using matrix operations (no for loops). See the code.