

# A Lattice Boltzmann Method for immiscible multiphase flow simulations using the Level Set Method

Lorenz Hufnagel, Daniel Zint

BGCE Student Project

August 20th, 2015

# Multiphase flow

## Examples


- Liquid-liquid mixtures (e.g. oil in water)
- Gas-liquid mixtures (e.g. bubble dynamics)



# Multiphase flow

## Examples

- Liquid-liquid mixtures (e.g. oil in water)
- Gas-liquid mixtures (e.g. bubble dynamics)



# Macroscopic fluid mechanics

- $N$  immiscible fluids.
- Each has own  $\rho_i, \nu_i$
- Hydrodynamics described by (incompressible) NSE

# Macroscopic fluid mechanics

- $N$  immiscible fluids.
- Each has own  $\rho_i, \nu_i$
- Hydrodynamics described by (incompressible) NSE

# Macroscopic fluid mechanics

- $N$  immiscible fluids.
- Each has own  $\rho_i, \nu_i$
- Hydrodynamics described by (incompressible) NSE

$$\nabla \cdot \vec{v}_i = 0 \text{ in } \Omega_i$$

$$\frac{\partial \vec{v}_i}{\partial t} + (\vec{v}_i \cdot \nabla) \vec{v}_i = -\frac{1}{\rho_i} \nabla p_i + \nu_i \nabla^2 \vec{v}_i \text{ in } \Omega_i$$

# Mathematic foundation

## Interface conditions





Figure : Two fluid domains  $\Omega_i$  and interface  $\Gamma$  inbetween

- Velocity across interface is  $C_0$ -continuous

$$[v] = \lim_{\epsilon \rightarrow 0} (\vec{v}(x + \epsilon \vec{n}) - \vec{v}(x - \epsilon \vec{n})) = 0$$

# Mathematic foundation

## Interface conditions cont'd




- Normal stress is balanced by surface tension  $\rightarrow$  pressure jump

$$[T] \cdot \vec{n} = \lim_{\epsilon \rightarrow 0} (\mathbf{T}_2(x + \epsilon \vec{n}) - \mathbf{T}_1(x - \epsilon \vec{n})) \cdot \vec{n} = 2\sigma \kappa \vec{n}$$



# Mathematic foundation

## Interface conditions cont'd



- Normal stress is balanced by surface tension  $\rightarrow$  pressure jump

$$[T] \cdot \vec{n} = \lim_{\epsilon \rightarrow 0} (\mathbf{T}_2(x + \epsilon \vec{n}) - \mathbf{T}_1(x - \epsilon \vec{n})) \cdot \vec{n} = 2\sigma \kappa \vec{n}$$

where  $\mathbf{T}_i$  is the stress tensor  $\mathbf{T}_i = 2\nu_i \rho_i \mathbf{S}_i - p \mathbb{I}$  and  $\kappa$  is the curvature of the interface  $\nabla \cdot \vec{n}$ .  $\mathbf{S}_{ij} = \frac{1}{2}(\partial_i v_j + \partial_j v_i)$ .

# Mathematic foundation

## Solvers for the two-phase problem

- Colour gradient method of Gunstensen and Rothman
  - only for small density and viscosity differences
- Method of Shan and Chen
  - models miscible fluids
  - immiscible flows only approximatively
- Free surface methods
  - only for big density and viscosity differences
- Method of Thömmes and Becker
  - using Level Set Method for interface motion
  - small and big density/viscosity differences possible
  - mass loss in Level Set Method

# Mathematic foundation

## Solvers for the two-phase problem

- Colour gradient method of Gunstensen and Rothman
  - only for small density and viscosity differences
- Method of Shan and Chen
  - models miscible fluids
  - immiscible flows only approximatly
- Free surface methods
  - only for big density and viscosity differences
- Method of Thoenmes and Becker
  - using Level Set Method for interface motion
  - small and big density/viscosity differences possible
  - mass loss in Level Set Method

# Mathematic foundation

## Solvers for the two-phase problem

- Colour gradient method of Gunstensen and Rothman
  - only for small density and viscosity differences
- Method of Shan and Chen
  - models miscible fluids
  - immiscible flows only approximatively
- Free surface methods
  - only for big density and viscosity differences
- Method of Thömmes and Becker
  - using Level Set Method for interface motion
  - small and big density/viscosity differences possible
  - mass loss in Level Set Method

# Mathematic foundation

## Solvers for the two-phase problem

- Colour gradient method of Gunstensen and Rothman
  - only for small density and viscosity differences
- Method of Shan and Chen
  - models miscible fluids
  - immiscible flows only approximatively
- Free surface methods
  - only for big density and viscosity differences
- Method of Thömmes and Becker
  - using Level Set Method for interface motion
  - small and big density/viscosity differences possible
  - mass loss in Level Set Method

# Mathematic foundation

## Ingredients for solver with Level Set Method

To solve the two-phase problem we need to:

- solve the flow equations  $\rightarrow$  LBM
- compute the motion of the interface  $\rightarrow$  Level Set Method
- couple the two algorithms  $\rightarrow$  according BC's

# Mathematic foundation

## Ingredients for solver with Level Set Method

To solve the two-phase problem we need to:

- solve the flow equations  $\rightarrow$  LBM
- compute the motion of the interface  $\rightarrow$  Level Set Method
- couple the two algorithms  $\rightarrow$  according BC's

# Mathematic foundation

## Ingredients for solver with Level Set Method

To solve the two-phase problem we need to:

- solve the flow equations  $\rightarrow$  LBM
- compute the motion of the interface  $\rightarrow$  Level Set Method
- couple the two algorithms  $\rightarrow$  according BC's



# Mathematic foundation

## LBM

### ■ Stream

$$f_i(x + c_i, t + 1) = f_i(x, t) - \frac{1}{\tau}(f_i - f_i^{eq}) + G_i$$

### ■ Collide (Bhatnagar-Gross-Krook approximation)

$$f_i^{eq}(\rho, v) = f_i^*(\rho + 3c_i \cdot v + \frac{9}{2}(c_i \cdot v)^2 - \frac{3}{2}v^2)$$

# Mathematic foundation

## LBM

### ■ Stream

$$f_i(x + c_i, t + 1) = f_i(x, t) - \frac{1}{\tau}(f_i - f_i^{eq}) + G_i$$


### ■ Collide (Bhatnagar-Gross-Krook approximation)

$$f_i^{eq}(\rho, v) = f_i^*(\rho + 3c_i \cdot v + \frac{9}{2}(c_i \cdot v)^2 - \frac{3}{2}v^2)$$

# Mathematic foundation

## LBM

the corresponding D2Q9 weight factors



$$f_i^* = \begin{cases} \frac{4}{9} & i = C \\ \frac{1}{9} & i = N, E, S, W \\ \frac{1}{36} & i = NE, SE, SW, NW \end{cases}$$

# Mathematic foundation

## LBM

we can compute for

- Density

$$\rho(x, t) = \sum f_i(x, t)$$

- Velocity

$$u = \sum f_i(x, t) c_i$$

# Mathematic foundation

## Interface capturing

Interface between fluid phases is captured by level set method

# Mathematic foundation

## Interface capturing

Interface between fluid phases is captured by level set method

- Level set function:  $\varphi := \varphi(x, t) \rightarrow \mathbb{R}$   
Interface is the zero-isosurface of this function
- Level set equation:

$$\frac{\partial \varphi}{\partial t} + \vec{v} \cdot \nabla \varphi = 0$$

where  $\vec{v}$  is the velocity at the interface, given by NSE.

# Mathematic foundation

## Interface capturing

Interface between fluid phases is captured by level set method

- Level set function:  $\varphi := \varphi(x, t) \rightarrow \mathbb{R}$   
Interface is the zero-isosurface of this function
- Level set equation:

$$\frac{\partial \varphi}{\partial t} + \vec{v} \cdot \nabla \varphi = 0$$

where  $\vec{v}$  is the velocity at the interface, given by NSE.

# Mathematic foundation

## Interface capturing

### Benefits of Level-Set Method:

- Implicit surface description eases bubble coalescence/breakup in code
- High density and viscosity ratios ( $> 10^3$ ) possible for simulation



# Mathematic foundation

## Interface capturing

### Benefits of Level-Set Method:

- Implicit surface description eases bubble coalescence/breakup in code
- High density and viscosity ratios ( $> 10^3$ ) possible for simulation

# Mathematic foundation

## Interface capturing

### Benefits of Level-Set Method:

- Implicit surface description eases bubble coalescence/breakup in code
- High density and viscosity ratios ( $> 10^3$ ) possible for simulation

# Mathematic foundation

## Interface implementation (Methods Coupling)


Hydrodynamics are solved by LBM (Stream-Collide, D2Q9, SRT for our first test)

- Interface becomes a new boundary condition for LBM

$$f_i(x, t + 1) = f_{i*}^+(x, t) + 6\Delta x f_i^* c_i \cdot \tilde{v} + R_i$$

- $\tilde{v}$  is the velocity on the interface along the direction  $c_i$
- calculated by linear interpolation:

$$\tilde{v} = qv(x_2, t) + (1 - q)v(x_1, t)$$



# Mathematic foundation

## Interface implementation (Methods Coupling)


Hydrodynamics are solved by LBM (Stream-Collide, D2Q9, SRT for our first test)

- Interface becomes a new boundary condition for LBM

$$f_i(x, t + 1) = f_{i*}^+(x, t) + 6\Delta x f_i^* c_i \cdot \tilde{v} + R_i$$

- $\tilde{v}$  is the velocity on the interface along the direction  $c_i$
- calculated by linear interpolation:

$$\tilde{v} = qv(x_2, t) + (1 - q)v(x_1, t)$$



# Mathematic foundation

## Interface implementation (Methods Coupling)


Hydrodynamics are solved by LBM (Stream-Collide, D2Q9, SRT for our first test)

- Interface becomes a new boundary condition for LBM

$$f_i(x, t + 1) = f_{i*}^+(x, t) + 6\Delta x f_i^* c_i \cdot \tilde{v} + R_i$$

- $\tilde{v}$  is the velocity on the interface along the direction  $c_i$
- calculated by linear interpolation:

$$\tilde{v} = qv(x_2, t) + (1 - q)v(x_1, t)$$



# Mathematic foundation

## Interface implementation (Methods Coupling)

$R_i$  ensures the jump conditions of the normal stress and corrects the error terms resulting from the bounce back treatment.

- $R_i = 6\Delta x^2 f_i^* \Lambda_i : A$ ,  
with  $A : B = \sum_{i,j} A_{ij} B_{ij}$ .
- $\Lambda_i = c_i \otimes c_i - \frac{1}{3} |c_i|^2 \mathbb{I}$  ,  
with  $c_i \otimes c_i = c_i c_i^T$ .
- $A = -q(1-q)[S] - (q-1/2)S^{(k)}$ 
  - $S^{(k)}$  velocity gradient at  $x_k$
  - $[S]$  jump of velocity gradient at the interface. Depends on  $\vec{n}$ ,  $\vec{t}$  and  $\kappa$ .

# Mathematic foundation

## Interface implementation (Methods Coupling)

$R_i$  ensures the jump conditions of the normal stress and corrects the error terms resulting from the bounce back treatment.

- $R_i = 6\Delta x^2 f_i^* \Lambda_i : A$ ,  
with  $A : B = \sum_{i,j} A_{ij} B_{ij}$ .
- $\Lambda_i = c_i \otimes c_i - \frac{1}{3} |c_i|^2 \mathbb{I}$  ,  
with  $c_i \otimes c_i = c_i c_i^T$ .
- $A = -q(1-q)[S] - (q-1/2)S^{(k)}$ 
  - $S^{(k)}$  velocity gradient at  $x_k$
  - $[S]$  jump of velocity gradient at the interface. Depends on  $\vec{n}$ ,  $\vec{t}$  and  $\kappa$ .

# Mathematic foundation

## Interface implementation (Methods Coupling)

$R_i$  ensures the jump conditions of the normal stress and corrects the error terms resulting from the bounce back treatment.

- $R_i = 6\Delta x^2 f_i^* \Lambda_i : A$ ,  
with  $A : B = \sum_{i,j} A_{ij} B_{ij}$ .
- $\Lambda_i = c_i \otimes c_i - \frac{1}{3} |c_i|^2 \mathbb{I}$  ,  
with  $c_i \otimes c_i = c_i c_i^T$ .
- $A = -q(1 - q)[S] - (q - 1/2)S^{(k)}$ 
  - $S^{(k)}$  velocity gradient at  $x_k$
  - $[S]$  jump of velocity gradient at the interface. Depends on  $\vec{n}$ ,  $\vec{t}$  and  $\kappa$ .



# Mathematic foundation

## Interface implementation (Methods Coupling)

$R_i$  ensures the jump conditions of the normal stress and corrects the error terms resulting from the bounce back treatment.

- $R_i = 6\Delta x^2 f_i^* \Lambda_i : A$ ,  
with  $A : B = \sum_{i,j} A_{ij} B_{ij}$ .
- $\Lambda_i = c_i \otimes c_i - \frac{1}{3} |c_i|^2 \mathbb{I}$  ,  
with  $c_i \otimes c_i = c_i c_i^T$ .
- $A = -q(1 - q)[S] - (q - 1/2)S^{(k)}$ 
  - $S^{(k)}$  velocity gradient at  $x_k$
  - $[S]$  jump of velocity gradient at the interface. Depends on  $\vec{n}$ ,  $\vec{t}$  and  $\kappa$ .

# Mathematic foundation

## Algorithm for LBM with level set

- Create initial interface
- Run level set method to calculate surface description
- Run LBM for a prescribed number of steps
- Run level set method for the same time interval
- Repeat until end of simulation

# Mathematic foundation

## Algorithm for LBM with level set

- Create initial interface
- Run level set method to calculate surface description
- Run LBM for a prescribed number of steps
- Run level set method for the same time interval
- Repeat until end of simulation

# Mathematic foundation

## Algorithm for LBM with level set

- Create initial interface
- Run level set method to calculate surface description
- Run LBM for a prescribed number of steps
- Run level set method for the same time interval
- Repeat until end of simulation

# Mathematic foundation

## Algorithm for LBM with level set

- Create initial interface
- Run level set method to calculate surface description
- Run LBM for a prescribed number of steps
- Run level set method for the same time interval
- Repeat until end of simulation

# Mathematic foundation

## Algorithm for LBM with level set

- Create initial interface
- Run level set method to calculate surface description
- Run LBM for a prescribed number of steps
- Run level set method for the same time interval
- Repeat until end of simulation

# Results

## Implementation

- Implementation in Matlab
  - Own LBM-solver
  - Level set method solved by Toolbox of Ian Mitchell

# Results

## Implementation

- Implementation in Matlab
- Own LBM-solver
- Level set method solved by Toolbox of Ian Mitchell



# Results

## Implementation

- Implementation in Matlab
- Own LBM-solver
- Level set method solved by Toolbox of Ian Mitchell

# Results

## Setup

- Transient couette flow with immersed drop
- Fluids have the same  $\rho_i, \nu_i \rightarrow [S]$  vanishes
- Goal: See interactions at the interface

# Results

## Setup

- Transient couette flow with immersed drop
- Fluids have the same  $\rho_i, \nu_i \rightarrow [S]$  vanishes
- Goal: See interactions at the interface



# Results

## Setup

- Transient couette flow with immersed drop
- Fluids have the same  $\rho_i, \nu_i \rightarrow [S]$  vanishes
- Goal: See interactions at the interface



# Results

## Simulation





# Results

## Simulation



# Results

## Simulation



# Outlook

*...so far intermediate results*

## Outlook:

- Create more examples
- Add correction term to prevent mass loss
- Reduce computational effort: Store Level-Set function only in narrow band around interface, Adalsteinsson and Sethian



# Outlook

*...so far intermediate results*

Outlook:

- Create more examples
- Add correction term to prevent mass loss
- Reduce computational effort: Store Level-Set function only in narrow band around interface, Adalsteinsson and Sethian

# Outlook

*...so far intermediate results*

Outlook:

- Create more examples
- Add correction term to prevent mass loss
- Reduce computational effort: Store Level-Set function only in narrow band around interface, Adalsteinsson and Sethian

# References



Thömmes, Guido, et al. "A lattice Boltzmann method for immiscible multiphase flow simulations using the level set method." *Journal of Computational Physics* 228.4 (2009): 1139-1156



D. Adalsteinsson, and J. A. Sethian. A fast level set method for propagating interfaces. *J. Comput. Phys.*, 118(2):269-277, 1995



Mitchell, Ian, *A Toolbox of Level Set Methods*