

Table of Laplace Transforms

Remember that we consider all functions (signals) as defined only on $t \geq 0$.

General

$$f(t)$$

$$F(s) = \int_0^{\infty} f(t) e^{-st} dt$$

$$f + g$$

$$F + G$$

$$\alpha f \quad (\alpha \in \mathbf{R})$$

$$\alpha F$$

$$\frac{df}{dt}$$

$$sF(s) - f(0)$$

$$\frac{d^k f}{dt^k}$$

$$s^k F(s) - s^{k-1} f(0) - s^{k-2} \frac{df}{dt}(0) - \dots - \frac{d^{k-1} f}{dt^{k-1}}(0)$$

$$g(t) = \int_0^t f(\tau) d\tau$$

$$G(s) = \frac{F(s)}{s}$$

$$f(\alpha t), \alpha > 0$$

$$\frac{1}{\alpha} F(s/\alpha)$$

$$e^{at} f(t)$$

$$F(s - a)$$

$$t f(t)$$

$$-\frac{dF}{ds}$$

$$t^k f(t)$$

$$(-1)^k \frac{d^k F(s)}{ds^k}$$

$$\frac{f(t)}{t}$$

$$\int_s^{\infty} F(s) ds$$

$$g(t) = \begin{cases} 0 & 0 \leq t < T \\ f(t - T) & t \geq T \end{cases}, \quad T \geq 0 \quad G(s) = e^{-sT} F(s)$$

Specific

1	$\frac{1}{s}$
δ	1
$\delta^{(k)}$	s^k
t	$\frac{1}{s^2}$
$\frac{t^k}{k!}, k \geq 0$	$\frac{1}{s^{k+1}}$
e^{at}	$\frac{1}{s-a}$
$\cos \omega t$	$\frac{s}{s^2 + \omega^2} = \frac{1/2}{s - j\omega} + \frac{1/2}{s + j\omega}$
$\sin \omega t$	$\frac{\omega}{s^2 + \omega^2} = \frac{1/2j}{s - j\omega} - \frac{1/2j}{s + j\omega}$
$\cos(\omega t + \phi)$	$\frac{s \cos \phi - \omega \sin \phi}{s^2 + \omega^2}$
$e^{-at} \cos \omega t$	$\frac{s + a}{(s + a)^2 + \omega^2}$
$e^{-at} \sin \omega t$	$\frac{\omega}{(s + a)^2 + \omega^2}$

Notes on the derivative formula at $t = 0$

The formula $\mathcal{L}(f') = sF(s) - f(0_-)$ must be interpreted very carefully when f has a discontinuity at $t = 0$. We'll give two examples of the correct interpretation.

First, suppose that f is the constant 1, and has no discontinuity at $t = 0$. In other words, f is the constant function with value 1. Then we have $f' = 0$, and $f(0_-) = 1$ (since there is no jump in f at $t = 0$). Now let's apply the derivative formula above. We have $F(s) = 1/s$, so the formula reads

$$\mathcal{L}(f') = 0 = sF(s) - 1$$

which is correct.

Now, let's suppose that g is a unit step function, *i.e.*, $g(t) = 1$ for $t > 0$, and $g(0) = 0$. In contrast to f above, g has a jump at $t = 0$. In this case, $g' = \delta$, and $g(0_-) = 0$. Now let's apply the derivative formula above. We have $G(s) = 1/s$ (exactly the same as F !), so the formula reads

$$\mathcal{L}(g') = 1 = sG(s) - 0$$

which again is correct.

In these two examples the functions f and g are the same except at $t = 0$, so they have the same Laplace transform. In the first case, f has no jump at $t = 0$, while in the second case g does. As a result, f' has no impulsive term at $t = 0$, whereas g does. As long as you keep track of whether your function has, or doesn't have, a jump at $t = 0$, and apply the formula consistently, everything will work out.