

# Real Analysis

Yunwei Ren

## Contents

<b>3</b>	<b>Lebesgue Measure</b>	<b>2</b>
3.1	Introduction . . . . .	2

## 3 Lebesgue Measure

### 3.1 Introduction

1.

*Proof.* Since  $\mathfrak{M}$  is an  $\sigma$ -algebra,  $B \setminus A \in \mathfrak{M}$  as long as  $A, B \in \mathfrak{M}$ . Since  $B \setminus A$  and  $A$  are disjoint,  $mB = mA + m(B \setminus A) \geq mA$  since  $m$  is nonnegative.  $\square$

2.

*Proof.* Let  $A_0 = E_0$  and  $E_k = A_k \setminus A_{k-1}$  for  $k \geq 1$ . Clear that  $E_i$  and  $E_j$  are disjoint for distinct  $i$  and  $j$ ,  $\bigcup A_n = \bigcup E_n$  and  $A_i \subset E_i$  for every  $i$ . Hence,

$$m\left(\bigcup E_n\right) = m\left(\bigcup A_n\right) = \sum mA_n \leq \sum mE_n,$$

where the last inequality comes from Exercise 1.  $\square$

3.

*Proof.* Suppose that  $mA < \infty$ . Then  $mA = m(A \cup \emptyset) = mA + m\emptyset$ , implying that  $m\emptyset = 0$ .  $\square$