

Convex Optimization

Yunwei Ren

Contents

2	Convex Sets	2
---	-------------	---

2 Convex Sets

2.1

Proof. For $k = 2$, $\theta_1 x_1 + \theta_2 x_2 \in C$ holds by definition. We argue by induction on k and assume that the inclusion holds for $k < m$. When $k = m$, denoting $\sum_{i=1}^{m-1} \theta_i$ by s ,

$$\sum_{i=1}^m \theta_i x_i = s \sum_{i=1}^{m-1} \frac{\theta_i x_i}{s} + \theta_m x_m.$$

Since $\sum_{i=1}^{m-1} \theta_i/s = 1$, by the induction hypothesis, $\sum_{i=1}^{m-1} \theta_i x_i/s \in C$. Meanwhile, as $s + \theta_m = 1$, $\sum_{i=1}^m \theta_i x_i \in C$, completing the proof. \square

2.2

Proof. Clear that the intersection of two convex sets is still convex. Hence, the intersection of $C \subset \mathbb{R}^n$ and any line is convex as long as C is convex.

Now we suppose that the intersection of C and any line is convex. For any $x_1, x_2 \in C$, $C_l = C \cap \{\theta x_1 + (1 - \theta)x_2 : \theta \in \mathbb{R}\}$ is convex and therefore $\theta x_1 + (1 - \theta)x_2 \in C_l \subset C$ for every $0 \leq \theta \leq 1$. Thus, C is convex. \square

2.8

Proof.

(a) It is trivial when a_1 and a_2 are linearly dependent, so we assume that a_1 and a_2 are linearly independent. We first tackle the problem for orthonormal a_1 and a_2 and then reduce the general situation to it.

Suppose that a_1 and a_2 are orthonormal. Let $S_0 = \text{span}(a_1, a_2)$ and (b_1, \dots, b_{n-2}) a basis of S_0^\perp . Then

$$x \in S_0 \Leftrightarrow \begin{bmatrix} b_1^T \\ \vdots \\ b_{n-2}^T \end{bmatrix} x = Bx = 0.$$

For $y = y_1 a_1 + y_2 a_2 \in S_0$, $y_1 \leq 1$ iff $a_1^T y \leq 1$ as (a_1, a_2) is an orthonormal basis of S_0 . Hence,

$$-1 \leq y_1, y_2 \leq 1 \Leftrightarrow \begin{bmatrix} a_1^T \\ a_2^T \\ -a_1^T \\ -a_2^T \end{bmatrix} y = Ay \preceq \mathbf{1}.$$

Thus, for orthonormal a_1 and a_2 , $S = \{x : Bx = 0, Ax \preceq \mathbf{1}\}$, a polyhedron.

Now we only assume the linear independence of a_1 and a_2 . We know that there exists some invertible n -by- n matrix¹ R such that $[\tilde{a}_1, \tilde{a}_2] = R[a_1, a_2]$ and \tilde{a}_1 and \tilde{a}_2 are orthonormal. Denoting the set described in the problem with respect to u_1 and u_2 by $S(u_1, u_2)$, $x \in S(a_1, a_2)$ iff $Rx \in S(\tilde{a}_1, \tilde{a}_2)$ iff $Rx \in \{x : \tilde{B}x = 0, \tilde{A}x \preceq \mathbf{1}\}$ where the meaning of \tilde{A} and \tilde{B} are described in the previous passage. Hence,

$$S(a_1, a_2) = \{x : \tilde{B}Rx = 0, \tilde{A}Rx \preceq \mathbf{1}\}.$$

¹We can use QR factorization to construct the matrix explicitly

- (b) Yes, and the provided form has already satisfied the requirement.
(c) No. Note that $\langle x, y \rangle_2 \leq 1$ for all y with 2-norm 1 implies

$$\|x\|_2 = \langle x, x/\|x\| \rangle_2 \leq 1.$$

And by the Cauchy-Schwarz inequality, for every $\|x\| \leq 1$, $\langle x, y \rangle_2$ holds for every $\|y\|_2 = 1$. Hence, S is the intersection of the unit ball and $\{x : x \succeq 0\}$, which is not a polyhedron.

(d) Yes. Let $\tilde{S} = \{x \in \mathbb{R}^n : x \succeq 0, \|x\|_\infty \leq 1\}$, which is clearly a polyhedron since when $x \succeq 0$, $\|x\|_\infty \leq 1$ is equivalent to $[e_1, \dots, e_n]x \preceq \mathbf{1}$ where e_i is the i -th vector in the standard basis of \mathbb{R}^n .

Now we show that $S = \tilde{S}$. Suppose that $x \succeq 0$. If $\langle x, y \rangle_2 \leq 1$ for all y with 1-norm 1, then $x_i = \langle x, e_i \rangle_2 \leq 1$. Namely, $\|x\|_\infty \leq 1$. Meanwhile, if $\|x\|_\infty \leq 1$,

$$\langle x, y \rangle \leq \sum_{i=1}^n x_i |y_i| \leq 1$$

as it is just the weighted average of x_1, \dots, x_n . Hence, $S = \tilde{S}$, completing the proof. \square

2.9

Proof.

(a) By the definition,

$$\begin{aligned} x \in V &\Leftrightarrow \|x - x_0\|_2^2 - \|x - x_i\|_2^2 \leq 0 \\ &\Leftrightarrow 2\langle x, x_i - x_0 \rangle \leq \langle x_i, x_i \rangle - \langle x_0, x_0 \rangle \quad \text{for } i = 1, \dots, K \\ &\Leftrightarrow 2 \begin{bmatrix} \langle x, x_1 - x_0 \rangle \\ \vdots \\ \langle x, x_K - x_0 \rangle \end{bmatrix} \preceq \begin{bmatrix} \|x_1\|_2^2 - \|x_0\|_2^2 \\ \vdots \\ \|x_K\|_2^2 - \|x_0\|_2^2 \end{bmatrix} \\ &\Leftrightarrow 2 \begin{bmatrix} (x_1 - x_0)^T \\ \vdots \\ (x_K - x_0)^T \end{bmatrix} x \preceq \begin{bmatrix} \|x_1\|_2^2 - \|x_0\|_2^2 \\ \vdots \\ \|x_K\|_2^2 - \|x_0\|_2^2 \end{bmatrix} \end{aligned}$$

Hence, V is a polyhedron. Intuitively, the border of a Voronoi set are the lines with the same distances to x_0 and x_i .

(b) Suppose that $P = \{x : \alpha_k^T x \leq b_k, k = 1, \dots, K\}$. Let x_0 be any point of P and we construct the other points by reflection. For each k , let \tilde{x}_k be any point of $\{x : \alpha_k^T x = b_k\}$, $U_k = I - 2\alpha_k \alpha_k^T / \|\alpha_k\|_2^2$, the Householder matrix, and

$$R_k(x) = U_k(x - \tilde{x}_k) + \tilde{x}_k = x + 2 \frac{\alpha_k}{\|\alpha_k\|_2^2} (b_k - \alpha_k^T x).$$

It is easy to verified that P is the Voronoi region of x_0 with respect to $R_1(x_0), \dots, R_K(x_0)$. \square

2.16

Proof. For every $(a, b_1 + b_2), (c, d_1 + d_2) \in S$ and $0 \leq \theta \leq 1$, let

$$z_\theta = \theta(a, b_1 + b_2) + (1 - \theta)(c, d_1 + d_2) = (x, y_1 + y_2)$$

where

$$x = \theta a + (1 - \theta)c, \quad y_i = \theta b_i + (1 - \theta)d_i \quad \text{for } i = 1, 2.$$

Since S_i is convex and $(a, b_i), (c, d_i) \in S_i$,

$$(x, y_i) = \theta(a, b_i) + (1 - \theta)(c, d_i) \in S_i.$$

Hence, S is convex. □