

# Solutions to *Topology*

Yunwei Ren

## Contents

<b>2</b>	<b>Topological Spaces and Continuous Functions</b>	<b>2</b>
13	Basis for a Topology . . . . .	2

## 2 Topological Spaces and Continuous Functions

### 13 Basis for a Topology

1.

*Proof.* Let  $\mathcal{T}$  be the topology of  $X$ . Since  $\mathcal{T}$  is a basis for itself and the hypothesis implies that  $A$  is a set in the topology generated by  $\mathcal{T}$ ,  $A \in \mathcal{T}$ , i.e.,  $A$  is open.  $\square$

4.

*Proof.*

(a) Put  $\mathcal{T} = \bigcap_{\alpha} \mathcal{T}_{\alpha}$ . Since  $\emptyset$  and  $X$  are contained in all  $\mathcal{T}_{\alpha}$ , they are also contained in  $\mathcal{T}$ . Let  $\{U_{\beta}\}_{\beta \in J}$  be an indexed family of elements of  $\mathcal{T}$  and put  $U = \bigcup_{\beta \in J} U_{\beta}$ . For every  $\beta$ , since  $U_{\beta}$  is open with respect to each  $\mathcal{T}_{\alpha}$ , by definition, so is  $\bigcup_{\beta \in J} U_{\beta}$ . Similarly, we can show that  $\mathcal{T}$  is closed under finite intersection. Thus,  $\mathcal{T}$  is a topology.

The union  $\bigcup \mathcal{T}_{\alpha}$ , however, may not be a topology. Take  $X = \{a, b, c\}$  for example.  $\mathcal{T}_a = \{\emptyset, a, X\}$  and  $\mathcal{T}_b = \{\emptyset, b, X\}$  are two topologies, but their union is not.

(b) Let  $\mathcal{T}$  be the intersection of all topologies containing all  $\mathcal{T}_{\alpha}$ . By (a),  $\mathcal{T}$  is a topology and clear that it is the unique smallest one. Now, let  $\mathcal{T}' = \bigcap \mathcal{T}_{\alpha}$ , which is again a topology and is contained in all  $\mathcal{T}_{\alpha}$ . It can be verified that  $\mathcal{T}'$  is the unique largest one.

(c)  $\{\emptyset, X, \{a\}, \{b\}, \{a, b\}, \{b, c\}\}; \{\emptyset, X, \{a\}\}$ .  $\square$

5.

*Proof.* Let  $\mathcal{A}$  be a basis,  $\mathcal{T}$  the topology generated by  $\mathcal{A}$ ,  $\{\mathcal{T}_{\alpha}\}$  the collection of all topologies containing  $\mathcal{A}$  and  $\mathcal{T}' = \bigcap \mathcal{T}_{\alpha}$ . For every union  $U$  of elements of  $\mathcal{A}$ , since, for every  $\alpha$ ,  $\mathcal{A} \subset \mathcal{T}_{\alpha}$  and  $\mathcal{T}_{\alpha}$  is closed under arbitrary union,  $U \in \mathcal{T}_{\alpha}$ . Hence,  $\mathcal{T} \subset \mathcal{T}'$ . Consequently,  $\mathcal{T}'$  is also the intersection of all topologies containing  $\mathcal{T}$ . Since  $\mathcal{T}$  contains itself as a subset,  $\mathcal{T}' \subset \mathcal{T}$ . Thus,  $\mathcal{T} = \mathcal{T}'$ .

Consider the collection of all finite intersections of  $\mathcal{A}$ , which is a basis, and apply the previous result to complete the proof.  $\square$

6.

*Proof.* Let  $\mathcal{T}_l$  and  $\mathcal{T}_K$  be the topology of  $\mathbb{R}_l$  and  $\mathbb{R}_K$  respectively.  $B = (-1, 1) - K$  is a basis element of  $\mathcal{T}_K$  and  $0 \in B$ . However, no half-open interval containing 0 is in  $B$ . Hence,  $\mathcal{T}_l$  is no finer than  $\mathcal{T}_K$ . Conversely,  $C = [1, 2)$  is a basis element of  $\mathcal{T}_l$  and  $1 \in C$ , but as  $1 \in K$ , there is no basis element of  $\mathcal{T}_K$  containing 1. Hence,  $\mathcal{T}_K$  is no finer than  $\mathcal{T}_l$ . Thus, they are not comparable.  $\square$

8.

*Proof.*

(a) First clear that  $\mathcal{B} \subset \mathcal{T}$ . For every  $U \in \mathcal{T}$  and  $x \in U$ , since  $U$  is open, there exists some  $\delta > 0$  such that  $(x - \delta, x + \delta) \subset U$ . Hence, there exists some rational  $a$  and  $b$  such that  $x - \delta < a < x < b < x + \delta$ . Thus, by Lemma 13.2,  $\mathcal{B}$  generates the standard topology on  $\mathbb{R}$ .

(b) Since  $x \in [[x], [x] + 1) \in \mathcal{C}$  for every  $x \in \mathbb{R}$ , the first condition for a basis is satisfied. Meanwhile, for every  $B_1 = [a, b)$  and  $B_2 = [c, d)$  in  $\mathcal{C}$ , if they are not disjoint,  $[c, b) = B_1 \cap B_2$  is also in  $\mathcal{C}$ . Hence, the second condition is satisfied. Thus,  $\mathcal{C}$  is a basis.

Since  $[\sqrt{2}, 2)$  can not be represented by union of elements in  $\mathcal{C}$ ,  $\mathcal{C}$  does not generate the lower limit topology.  $\square$