GROUND RING OF TWO DIMENSIONAL STRING THEORY

EDWARD WITTEN*

School of Natural Sciences
Institute for Advanced Study
Olden Lane
Princeton, NJ 08540

ABSTRACT

String theories with two dimensional space-time target spaces are characterized by the existence of a "ground ring" of operators of spin (0,0). By understanding this ring, one can understand the symmetries of the theory and illuminate the relation of the critical string theory to matrix models. The symmetry groups that arise are, roughly, the area preserving diffeomorphisms of a two dimensional phase space that preserve the fermi surface (of the matrix model) and the volume preserving diffeomorphisms of a three dimensional cone. The three dimensions in question are the matrix eigenvalue, its canonical momentum, and the time of the matrix model.

[★] Research supported in part by NSF Grant PHY86-20266.

1. Introduction

Critical string theory with two dimensional target space time has been the subject of much recent interest. The simplest target space is a flat space-time with a linear dilaton field, and perhaps an additional "Liouville" term. The corresponding world-sheet Lagrangian is

$$L = \frac{1}{4\pi\alpha'} \int d^2x \sqrt{h} \left(h^{ij} \partial_i X \partial_j X + h^{ij} \partial_i \phi \partial_j \phi \right) - \frac{1}{2\pi\sqrt{\alpha'}} \int d^2x \sqrt{h} \cdot \phi R^{(2)}$$

$$+ \mu \int d^2x \sqrt{h} \exp(-2\phi/\sqrt{\alpha'}).$$
(1.1)

(h is the world-sheet metric, $R^{(2)}$ is the Ricci scalar, and μ is the cosmological constant.) The "matter" field X is decoupled from the "Liouville" field ϕ (and from the ghosts that appear upon quantization), so some vertex operators can be constructed as products of X operators and ϕ operators. The obvious conformal fields constructed from X are standard vertex operators e^{ipX} . These depend on the continuously variable parameter p. In addition, at particular values of the momentum, additional "discrete" primary fields appear. The first of these, at zero momentum, is the current ∂X (or, when left and right movers are combined, the gravitional vertex operator $\bar{\partial}X\partial X$). These discrete states have long been known [1]. In the context of two dimensional string theory they were discussed at an early stage by N. Seiberg (unpublished) and have showed up in matrix model calculations [2]. It has been argued that they should be interpreted in terms of a "stringy" target space topological field theory [3]; this indicates that understanding these modes is essential for extracting the real physical lessons of two dimensional string theory. These modes may be important in understanding two dimensional black holes [4,5]. They have been the subject of several recent studies [6,7,8].

In addition to the discrete vertex operators that have usually been considered, which occur (if one considers (1,1) operators) at ghost number zero, there are also discrete vertex operators which occur at the same values of the momenta at adjacent values of the ghost number. This is obvious in the BRST formalism. If

one has a family of states $|\alpha(p)\rangle$, of unit norm, parametrized by a momentum variable p, and BRST invariant only at, say, $p = p_0$, then one has

$$Q|\alpha(p)\rangle = f(p)|\beta(p)\rangle,$$
 (1.2)

where $|\beta(p)\rangle$ is of unit norm and f(p) vanishes precisely at $p = p_0$. Then $|\alpha(p_0)\rangle$ and $|\beta(p_0)\rangle$ are a pair of discrete states at adjacent values of the ghost number.

The whole pattern of discrete states of various ghost number has been thoroughly described by Lian and Zuckerman [9]. However, the physical consequences of the existence of discrete states of non-standard ghost number have not yet been considered. This is the problem that we will consider in the present paper.

2. The Discrete States

To begin with, we consider the holomorphic part of the current algebra. Thus, X is a free field with

$$X(z)X(w) \sim -\ln(z-w). \tag{2.1}$$

The stress tensor is $T_{zz} = -(\partial_z X)^2/2$. The tachyon operators $V_p = e^{ipX}$ are conformal fields of dimension $p^2/2$.

For reasons that are standard and will become clear, it is convenient to first consider the theory compactified at the SU(2) radius and with zero cosmological constant. At the SU(2) radius, the allowed values of the momenta are $p = n/\sqrt{2}$, with $n \in \mathbb{Z}$. There is an SU(2) current algebra with (T_+, T_3, T_-) represented by $(e^{iX\sqrt{2}}, i\sqrt{2}\partial X, e^{-iX\sqrt{2}})$. For $s \geq 0$, there is no singularity in the short distance product

$$e^{iX\sqrt{2}}(z) \cdot e^{isX/\sqrt{2}}(w) \tag{2.2}$$

for $z \to w$, so the operator $e^{isX/\sqrt{2}}$ transforms as the highest weight of an SU(2) representation. In view of the J_3 value, this is a representation of spin s. By

repeatedly acting with the lowering operators J_- , we fill out the SU(2) multiplets with operators $V_{s,n}$, $n \in \{s, s-1, \ldots, -s\}$, such that $V_{s,s} = e^{isX/\sqrt{2}}$, $V_{s,-s} = e^{-isX/\sqrt{2}}$. The $V_{s,n}$ are all Virasoro primaries, since the $V_{s,s}$ are standard primary fields and the SU(2) current algebra intertwines with Virasoro. For |n| = s, the $V_{s,n}$ are standard tachyon operators at particular momenta, but for |n| < s, they are new objects, "discrete primaries." The conformal dimension of $V_{s,n}$ is $s^2/4$. More detailed investigation [1,9] shows that regardless of the radius of compactification of the X field, the only primary fields of the X system are the tachyon vertex operators and the $V_{s,n}$. At an arbitrary radius, the discrete primaries are whatever combinations of $V_{s,n}$'s with anti-holomorphic counterparts are permitted (by the standard momentum restrictions) at that particular radius.

Coupling To Other Degrees Of Freedom

Now, let us couple the $V_{s,n}$'s to the rest of the system. First, we consider the Liouville field ϕ . Its stress tensor is $T_{zz} = -(1/2)(\partial \phi)^2 + \sqrt{2}\phi$. The operator $e^{iw\phi}$ has conformal dimension $w^2/2 + iw\sqrt{2}$. Looking for a spin one primary field of the form $V_{s,n}e^{iw\phi}$, we see that there are two possibilities:

$$W_{s,n}^{\pm} = V_{s,n} \cdot e^{\sqrt{2}\phi \mp s\phi\sqrt{2}}.$$
 (2.3)

Note that $W_{s,n}^{\pm}$ has momentum $(n, i(-1 \pm s)) \cdot \sqrt{2}$. The importance of the spin one condition is that (1,1) vertex operators corresponding to infinitesimal deformations of the theory can be constructed in the form

$$W_{s,n}^{\pm}(z) \cdot \overline{W}_{s,n'}^{\pm}(z) \tag{2.4}$$

where $\overline{W}_{s,n}$ are the antiholomorphic operators analogous to the $W_{s,n}$. Notice that as the Liouville field is noncompact, the Liouville momenta must be matched

^{*} The latter holds since a rotation about the J_1 axis gives $X \to -X$ and reverses the sign of J_3 .

between left and right movers, which is the reason that the left-movers and right-movers in (2.4) share the same value of s and of the sign \pm . At the SU(2) radius, there is no similar matching of n. Note that $W_{0,0}^+ = W_{0,0}^-$ is the cosmological constant operator $e^{\phi\sqrt{2}}$.

Now we wish to include the ghosts, which are fermi fields b and c of spins 2 and -1, with $T_{zz} = -2b_{zz}\partial_z c^z - \partial_z b_{zz} \cdot c^z$. One of the important merits of the ghosts is that in their presence, the physical state condition can be incorporated by considering the cohomology of the BRST operator Q. In the standard fashion, the spin one fields $W_{s,n}^{\pm}$ are related to spin 0, BRST invariant fields

$$Y_{s,n}^{\pm} = cW_{s,n}^{\pm} \tag{2.5}$$

of ghost number 1.

2.1. Additional States At Other Values Of The Ghost Number

This cannot be the whole story, for a reason indicated in the introduction. The $Y_{s,n}^{\pm}$ of |n|=s are not truly discrete (but are special tachyon states). However, the ghost number one discrete operators $Y_{s,n}^{\pm}$ of |n| < s must have "partners" at an adjoining value of the ghost number, namely 0 or 2. In fact, the $Y_{s,n}^{+}$ have partners at ghost number 0, and the $Y_{s,n}^{-}$ have partners at ghost number 2 [9]. The ghost number 0 partner of $Y_{s,n}^{+}$ will be called in this paper $\mathcal{O}_{u,n}$, with u=s-1. The momentum of $\mathcal{O}_{u,n}$ is thus $(n,iu) \cdot \sqrt{2}$. Since |n| < s, we have $|n| \le u$, so the allowed values of u and n are $u=0,1/2,1,\ldots$ and $n=u,u-1,\ldots,-u$.

For instance, the first case of a $Y_{s,n}^+$ with |n| < s is $Y_{1,0}^+ = c\partial X$, a discrete primary field of ghost number 1 and momentum (0,0). The corresponding discrete primary field of ghost number 0 and momentum (0,0) is the identity operator, $\mathcal{O}_{0,0} = 1$. At the next level, there are two discrete primaries, $Y_{3/2,1/2}^+$ and $Y_{3/2,-1/2}^+$, of spin 0 and ghost number 1. The corresponding spin 0, ghost number 0, operators

are

$$x = \mathcal{O}_{1/2,1/2} = \left(cb + \frac{i}{\sqrt{2}}(\partial X - i\partial\phi)\right) \cdot e^{i(X+i\phi)/\sqrt{2}}$$

$$y = \mathcal{O}_{1/2,-1/2} = \left(cb - \frac{i}{\sqrt{2}}(\partial X + i\partial\phi)\right) \cdot e^{-i(X-i\phi)/\sqrt{2}}.$$
(2.6)

Some details about their construction can be found in appendix 1.

2.2. The Ground Ring

There are two immediate reasons that the existence of spin zero, ghost number zero operators is of vital importance:

- (1) It leads to the existence of symmetries of the conformal field theory under study.
- (2) It leads to the existence of a ring, which we will call the "ground ring," which largely controls the properties of the theory, as we will see.

To understand the first point, note that we already know of the existence of infinitely many spin (1,0) and (0,1) operators of zero ghost number, namely $W_{s,n}^{\pm}$ and $\overline{W}_{s,n}^{\pm}$. Because of the Liouville momentum that they carry, which must be matched between left and right movers, to construct true quantum field operators as opposed to the holomorphic or antiholomorphic chiral vertex operators of the above discussion, W (or \overline{W}) should not be considered by itself but must be paired with antiholomorphic (or holomorphic) fields of the same Liouville momentum. One way to do this is to pair $W \cdot \overline{W}$, as in (2.4), to make fields of spin (1,1), corresponding to infinitesimal moduli. To make quantum field operators of spin (1,0), however, we must combine a W with an antiholomorphic spin zero field of the same Liouville momentum, and conversely for \overline{W} . The spin (1,0) and spin (0,1) currents of the theory are consequently

$$J_{s,n,n'} = W_{s,n}^{+} \overline{\mathcal{O}}_{s-1,n'}$$

$$\overline{J}_{s,n,n'} = \mathcal{O}_{s-1,n} \overline{W}_{s,n'}^{+}$$
(2.7)

On general grounds, these currents generate a Lie algebra of symmetries – which

we will presently determined to be the Lie algebra of volume preserving diffeomorphisms of a certain three manifold.

The spin 0 BRST invariant operators generate a commutative, associative ring for the following reason. Let \mathcal{O} and \mathcal{O}' be two such operators. The operator product $\mathcal{O}(z)\mathcal{O}'(0)$ is BRST invariant, so all the terms in its short distance expansion for $z \to 0$ are BRST invariant. Negative powers of z may arise in this short distance expansion, but the operators multiplying the negative powers of z are negative dimension operators which must be BRST commutators (as there is no BRST cohomology at negative dimension in this theory). Therefore, modulo the BRST commutators, the short distance limit of $\mathcal{O}(z)\mathcal{O}'(0)$ is some BRST invariant spin zero operator $\mathcal{O}''(0)$ (which may vanish):

$$\mathcal{O}(z)\mathcal{O}'(0) \sim \mathcal{O}''(0) + \{Q, \ldots\}. \tag{2.8}$$

This is the desired multiplication law, $\mathcal{O} \cdot \mathcal{O}' = \mathcal{O}''$. This procedure obviously defines a commutative, associative ring.

By combining left and right movers in the usual way we can form spin (0,0) quantum field operators

$$\mathcal{V}_{u,n,n'} = \mathcal{O}_{u,n} \overline{\mathcal{O}}_{u,n'} \tag{2.9}$$

which can be multiplied in just the same way. We will call the ring of the \mathcal{V} 's the ground ring of the theory, and we will call the ring of the chiral \mathcal{O} 's the chiral ground ring.

Since the ground ring is naturally associated with the conformal field theory, any symmetries of the theory must be symmetries of the ground ring. This will make it easy to determine the symmetry algebra. In fact, if J is a spin one current generating a symmetry, then the action of J on a spin (0,0) operator \mathcal{O} is defined by

$$J(\mathcal{O}(P)) = \frac{1}{2\pi i} \oint_{\mathcal{C}} J(z) \cdot \mathcal{O}(P)$$
 (2.10)

(modulo BRST commutators) where C is any contour surrounding the point P. Standard contour manipulations show that if J and J' are two currents and [J, J'] is their commutator then

$$[J, J'](\mathcal{O}(P)) = J(J'(\mathcal{O}(P))) - J'(J(\mathcal{O}(P))),$$
 (2.11)

so that the $\mathcal{O}(P)$'s form a representation of the symmetry algebra. What is more unusual is that since a contour enclosing two points P and Q is homologous to the sum of a contour surrounding P and a contour surrounding Q, we get

$$J(\mathcal{O}(P)\mathcal{O}'(Q)) = J(\mathcal{O}(P)) \cdot \mathcal{O}(Q) + \mathcal{O}(P) \cdot J(\mathcal{O}'(Q)), \tag{2.12}$$

which in the limit $P \to Q$ gives

$$J(\mathcal{O} \cdot \mathcal{O}') = J(\mathcal{O}) \cdot \mathcal{O}' + \mathcal{O} \cdot J(\mathcal{O}'). \tag{2.13}$$

Thus, the J's act as derivations of the ring of \mathcal{O} 's. These considerations can be applied in two different ways: the chiral currents $W_{s,n}^{\pm}$ act as derivations of the chiral ground ring, and the quantum currents J and \overline{J} of (2.7) are derivations of the full quantum ground ring.

2.3. Determination Of The Ground Ring

The ground ring is easily determined. First, we consider the chiral ground ring. We note that the product $x^n y^m$ has the quantum numbers of $\mathcal{O}_{(n+m)/2,(n-m)/2}$, so $\mathcal{O}_{u,n}$ is a multiple of $x^{u+n}y^{u-n}$ if the latter is not zero. We will prove that $x^n y^m \neq 0$ for all n, m in the next subsection. Accepting this for the moment, the chiral ground ring of the $\mathcal{O}_{u,n}$'s is just the ring of polynomial functions in x and y.

Similarly, the anti-chiral ground ring (of BRST invariant antiholomorphic spin (0,0) operators) consists of polynomial functions in $x' = \overline{\mathcal{O}}_{1/2,1/2}$ and $y' = \overline{\mathcal{O}}_{1/2,-1/2}$.

Now let us combine left and right movers to make the full quantum ground ring. It is generated by operators $x^n y^m x'^n y'^m$ with n + m = n' + m' (to balance the Liouville momenta). Such operators are multiplied in the obvious way, by multiplying the left and right moving parts separately. At n + m = 0 there is only the identity operator. At n + m = 1 there are four operators,

$$a_1 = xx'$$

$$a_2 = yy'$$

$$a_3 = xy'$$

$$a_4 = yx'.$$

$$(2.14)$$

The ground ring is generated by the a_i since any monomial $x^n y^m x'^n y'^m$ with n + m = n' + m' can obviously be written as a monomial in the a_i . The a_i obey the one obvious relation

$$a_1 a_2 - a_3 a_4 = 0 (2.15)$$

and are otherwise independent. (In algebraic geometry, one says that the left hand side of (2.15) is a homogeneous polynomial whose zeros form a quadric in \mathbf{P}^3 , and the equations (2.14) are an isomorphism of this quadric with $\mathbf{P}^1 \times \mathbf{P}^1$.)

Thus, the ground ring of the theory, at the SU(2) point, is the ring of polynomial functions on the three dimensional quadric cone Q defined by $a_1a_2 - a_3a_4 = 0$. In comparing with matrix models, we will learn that the three dimensions of this cone correspond to the matrix eigenvalue, its canonical momentum, and the "time" of the matrix model Schodinger equation. At the SU(2) point, there is a complete symmetry among these variables, which is partly lost upon decompactification.

2.4. Determination Of The Chiral Symmetry Algebra

Now we will determine the symmetry algebra of the theory, by using the fact that it acts as an algebra of derivations of the ground ring. First we consider the chiral currents.

To determine how a chiral current J acts on the chiral ground ring, it is enough, from (2.13), to know what it does to x and y. Indeed if

$$J(x) = f(x, y), \text{ and } J(y) = g(x, y),$$
 (2.16)

then it follows from (2.13) that for an arbitrary element w(x, y) of the ground ring, we get

$$J(w) = \left(f\frac{\partial}{\partial x} + g\frac{\partial}{\partial y}\right) \cdot w. \tag{2.17}$$

In particular, J can be identified with the vector field $f\partial_x + g\partial_y$ on the (x,y) plane, and so the chiral currents generate an algebra of diffeomorphisms of the (x,y) plane. (And similarly, the quantum currents which combine left and right movers act by diffeomorphisms of the quadric cone Q.)

Let us determine the vector fields corresponding to some particular $W_{s,n}^{\pm}$. First of all, the $W_{s,n}^{-}$ are trivially disposed of. They have momenta $(n,i(-1-s))\sqrt{2}$, and would map x or y to spin (0,0) operators of ghost number 0 and Liouville momentum $i(-1/2-s)\sqrt{2}$. As there is no BRST cohomology with those quantum numbers, the $W_{s,n}^{-}$ annihilate x and y and therefore the whole ground ring. Similarly, the cosmological constant operator $W_{0,0}^{+}$ (which in any case is the same as $W_{0,0}^{-}$) annihilates x and y. Now we consider the $W_{s,n}^{+}$. The first non-trivial case is $W_{1/2,\pm 1/2}^{+}$. These are tachyon operators,

$$W_{1/2,\pm 1/2}^{+} = e^{(\pm iX + \phi)/\sqrt{2}}.$$
 (2.18)

Recalling the definition of y, we have

$$W_{1/2,1/2}^{+}(z)\cdot y(w) = e^{(iX+\phi)/\sqrt{2}}(z)\cdot \left(cb - \frac{i}{\sqrt{2}}\partial(X+i\phi)\right)e^{-(iX+\phi)/\sqrt{2}}(w) \sim \frac{1}{z-w}.$$
(2.19)

The action of $W_{1/2,1/2}^+$ on y is given by the residue of the pole, which is 1, so

$$W_{1/2,1/2}^{+}(y) = 1.$$
 (2.20)

On the other hand, $W^+_{1/2,1/2}$ annihilates x (there is no short distance singularity in the operator product $W^+_{1/2,1/2} \cdot x$). Hence we have determined that $W^+_{1/2,1/2}$ acts on the chiral ground ring as

$$W_{1/2,1/2}^{+} = \frac{\partial}{\partial y}.$$
 (2.21)

Similarly, $W_{1/2,-1/2}^+$ acts as

$$W_{1/2,-1/2}^{+} = \frac{\partial}{\partial x}.$$
 (2.22)

At this point, we can fill a gap in our determination of the chiral ground ring by proving that $x^n y^m \neq 0$. Indeed, an *n*-fold repeated commutator of $x^n y^m$ with $W_{1/2,-1/2}^+$ followed by an *m*-fold repeated commutator with $W_{1/2,1/2}^+$ gives 1, which does not vanish, so $x^n y^m$ must not be zero.

From (2.21) and (2.22), we see that $W_{1/2,1/2}^+$ and $W_{1/2,-1/2}^+$ commute in their action on the x-y plane. One might be inclined to think that they commute altogether. In fact

$$W_{1/2,1/2}^+(z)W_{1/2,-1/2}^+(w) = e^{(iX+\phi)/\sqrt{2}}(z)e^{(-iX+\phi)/\sqrt{2}}(w) \sim \frac{1}{z-w}e^{\phi\sqrt{2}}(w) = \frac{1}{z-w}W_{0,0}^+(w).$$
(2.23)

The object $W_{0,0}^+(w)$ is "central" in the sense that it commutes with all of the currents, as we will see, and acts trivially on the x-y plane. (2.23) is the beginning of the discovery that what is going on in the problem is quantum mechanics on the x-y plane, as the commutator of (the operators representing) ∂_x and ∂_y is not zero but is a central object.

Now, let us determine the rest of the algebra. The operators $W_{1,n}^+$ are simply the SU(2) currents. x and y have $T_3 = \pm 1/2$, respectively (as we see from their

momentum values), and as they are the only cohomology classes with their values of the ghost number and Liouville momentum, they form an SU(2) representation which must have spin 1/2. Thus, without further ado we have

$$W_{1,1}^{+} = x \frac{\partial}{\partial y}$$

$$W_{1,0}^{+} = \frac{1}{2} \left(x \frac{\partial}{\partial x} - y \frac{\partial}{\partial y} \right)$$

$$W_{1,-1}^{+} = y \frac{\partial}{\partial x}.$$
(2.24)

(These formulas have been written as if x and y might be real variables, with symmetry group $SL(2,\mathbb{R})$ rather than the SU(2) that is natural in the Euclidean theory. Upon complexification, (2.24) is equivalent to the standard SU(2) generators.)

Now we consider the $W_{s,n}^+$ with s > 1. These operators transform under SU(2) with spin s. By considering the Liouville momenta, we see that they map x or y to polynomials in x and y that are homogeneous of degree 2s - 1. Such a polynomial has spin s - 1/2. On the other hand, we know that x and y have spin 1/2. In combining spin s with spin 1/2 to make spin s - 1/2, there is only one invariant coupling. If these couplings are not zero (which we will prove presently), they can be set to any desired non-zero value by scaling the $W_{s,n}^+$ by factors depending only on s. Therefore, if we can find any Lie algebra in which these couplings are all non-zero, and in which the $W_{s,n}$ of $s \le 1$ act correctly, it is the one we want.

In view of suggestions about the possible relation of area-preserving diffeomorphisms to high spin fields in two dimensions [10,11], and the above observation that $[\partial_x, \partial_y]$ is central, there is an obvious Lie algebra with the right properties. Let ω be the area form

$$\omega = dx \ dy \tag{2.25}$$

on the x-y plane. Every polynomial h(x,y) determines a vector field

$$\frac{\partial h}{\partial x}\frac{\partial}{\partial y} - \frac{\partial h}{\partial y}\frac{\partial}{\partial x} \tag{2.26}$$

that generates an infinitesimal area-preserving transformation of the x-y plane. If we identify $W_{s,n}^+$ with the transformation determined by the vector field that corresponds to $h(x,y) = x^{s+n}y^{s-n}$, then all of the conditions are obeyed since (i) these W's are non-zero; (ii) this assignment is SU(2) (or $SL(2,\mathbb{R})$) covariant; (iii) these W's obviously generate a Lie algebra. The area preserving diffeomorphisms of the x-y plane are therefore the Lie algebra that we are looking for.

It remains to show that for any s, the $W_{s,n}^+$ act non-trivially on x and y. First let us note that for any s, by counting the Liouville momenta, the commutator $[W_{1/2,n}^+, W_{s,n'}^+]$ is a multiple of $W_{s-1/2,n+n'}^+$. Moreover, for given s, these commutators are not all zero. Indeed, for n=-1/2 and n'=s one has $W_{1/2,-1/2}^+=e^{(-iX+\phi)/\sqrt{2}}$ and $W_{s,s}^+=e^{(isX+(1-s)\phi)\sqrt{2}}$. As

$$e^{(-iX+\phi)/\sqrt{2}}(z) \cdot e^{(isX+(1-s)\phi)\sqrt{2}}(w) \sim \frac{1}{z-w} e^{(i(s-1/2)X+(3/2-s)\phi)\sqrt{2}}(w) = \frac{1}{z-w} W_{s-1/2,s-1/2}(w),$$
(2.27)

one has

$$[W_{1/2,-1/2}^+, W_{s,s}^+] = W_{s-1/2,s-1/2}^+. (2.28)$$

Inductively, if $W_{s-1/2,s-1/2}^+$ does not annihilate x and y, then $W_{s,s}^+$ does not either. As we know that the $W_{s,n}^+$ do not all annihilate x and y for sufficiently small s, this is true for all s, as assumed above.

It is interesting to note that the "energy" operator of the theory, that is, the operator that generates translations of X, is ∂X which is none other than $W_{1,0}^+$ in our present notation. According to the above, this operator generates on the x-y plane the motion derived from the Hamiltonian function h(x,y)=xy. With p=x+y, q=x-y, this is none other than the inverted harmonic oscillator Hamiltonian $h=p^2-q^2$ familiar in the c=1 matrix model. We will make a more extensive comparison with the matrix model in §3, after combining left and right movers. The conclusion will be much the same.

The Other Operators

Now we want to extend this discussion to determine the commutators of the $W_{s,n}^-$.

As $W_{s,n}^-$ has Liouville momentum $-i\sqrt{2}(s+1)$, a commutator $[W_{s,n}^-, W_{s',n'}^-]$ must be a multiple of $W_{s+s'+1,n+n'}^-$. But $W_{s,n}^-$ transforms under SU(2) with spin s. Since spin s+s'+1 does not appear in the tensor product of spin s and spin s', we must have

$$[W_{s,n}^-, W_{s',n'}^-] = 0$$
, for all s, n, s', n' . (2.29)

Next we consider a commutator $[W^+_{s,n},W^-_{s',n'}]$. As $W^+_{s,n}$ has Liouville momentum $-i\sqrt{2}(-s+1)$, the Liouville momentum of the commutator is $-i\sqrt{2}(s'-s+2)$. If $s \geq 1+s'$, we must write this as $-i\sqrt{2}(-s''+1)$ where s''=s-s'-1, and then $[W^+_{s,n},W^-_{s',n'}]$ is a multiple of $W^+_{s'',n+n'}$. As the latter transforms under SU(2) with spin s''=s-s'-1, and spin s-s'-1 does not appear in the tensor product of spin s with spin s', $[W^+_{s,n},W^-_{s',n'}]$ necessarily vanishes for such s and s'. It remains to consider the case s<1+s'. Then we must have

$$[W_{s,n}^+, W_{s',n'}^-] = \text{const} \cdot W_{s'',n+n'}^-, \tag{2.30}$$

with s'' = s' + 1 - s. Therefore the problem is just to determine how the W^- transform under the algebra of the W^+ .

I claim that the answer is that (after scaling the $W_{s,n}^-$ by suitable factors that depend only on s), $W_{s,n}^-$ transforms like

$$W_{s,n}^{-} \sim \frac{\partial^{(s-n)/2}}{\partial x^{(s-n)/2}} \frac{\partial^{(s+n)/2}}{\partial y^{(s+n)/2}} \delta^{2}(x,y). \tag{2.31}$$

I will just briefly indicate how this can be proved. The proposed formula is obviously SU(2) covariant, so the $W_{1,n}^+$ act correctly. For s=1/2, the commutators $[W_{1/2,1/2}^+, W_{s,s}^-]$ are all non-zero. This is proved by explicitly evaluating them; the

calculation is just analogous to the one above for $[W^+_{1/2,-1/2},W^+_{s,s}]$. The commutators $[W^+_{1/2,n},W^-_{s,n'}] \sim W^-_{s+1/2,n+n'}$ are all determined by SU(2) in terms on one unknown s dependent coefficient, which is nonzero for all s in view of what was just said. This being so, the $W^-_{s,n}$ can be uniquely normalized so as to transform under $W^+_{1/2,n}$ as claimed in (2.31). The derivatives of $\delta^2(x,y)$ appearing in (2.31) give a representation of the algebra of polynomial area-preserving vector fields, so to prove that the $W^-_{s,n}$ transform as claimed, it is enough to show that there is only one possible representation in which the $W^+_{s,n}$ for $s \leq 1$ act as claimed. This is an immediate consequence of the formula $[W^+_{1/2,n},W^+_{s,n'}] \sim W^+_{s-1/2,n+n'}$ which combined with SU(2) invariance uniquely determines the action of the $W^+_{s,n}$ once the action of the $W^+_{s-1/2,n}$ is known.

The $W_{s,n}^-$ act trivially on the x-y plane; this can be demonstrated by considering the Liouville momentum and SU(2) transformations, as above.

2.5. The Quantum Symmetry Algebra

Now we will combine the left and right movers and determine the quantum symmetry algebra, that is the algebra of the operators introduced earlier in (2.7):

$$J_{s,n,n'} = W_{s,n}^{+} \overline{\mathcal{O}}_{s-1,n'}$$

$$\overline{J}_{s,n,n'} = \mathcal{O}_{s-1,n} \overline{W}_{s,n'}^{+}$$
(2.32)

I will again proceed indirectly, using the fact that the J's and \overline{J} 's must correspond to vector fields on the quadric surface Q defined by

$$a_1 a_2 - a_3 a_4 = 0. (2.33)$$

A hypersurface H in affine space, with defining equation f = 0, has a natural volume form

$$\Theta = -\frac{da_1 da_2 da_3}{\partial f / \partial a_4}. (2.34)$$

It can be described more invariantly by saying that Θ is the unique three form on

H such that on H

$$\Lambda = -df \cdot \Theta \tag{2.35}$$

where $\Lambda = da_1 da_2 da_3 da_4$ is (up to a constant multiple) the unique translation and SL(4) invariant volume form on the ambient affine space. The latter description makes it possible to define Θ without singling out one of the coordinates. For the quadric Q, $f = a_1 a_2 - a_3 a_4$, so

$$\Theta = \frac{da_1 da_2 da_3}{a_3}. (2.36)$$

If we write

$$a_3 = \rho e^{\psi}, \quad a_4 = \rho e^{-\psi},$$
 (2.37)

so that the equation of Q is $\rho^2 = a_1 a_2$, then

$$\Theta = d\psi da_1 da_2. \tag{2.38}$$

The polynomial vector fields on Q transform under SO(2,2), whose complexification is the same as that of $SU(2) \times SU(2)$, as

$$\bigoplus_{n=0,1/2,1,\dots} ((n+1,n) \oplus (n,n+1) \oplus (n+1,n+1)). \tag{2.39}$$

The volume preserving polynomial vector fields transform as

$$\bigoplus_{n=0,1/2,1,\dots} ((n+1,n) \oplus (n,n+1)). \tag{2.40}$$

These facts will be demonstrated in appendix 2. I claim that the Lie algebra of the J's and \overline{J} 's is the algebra of volume preserving polynomial vector fields on Q, and in fact that the J's and \overline{J} 's correspond respectively to the pieces of spin (n+1,n) and (n,n+1). To demonstrate this, it is enough to note that $J_{s,n,n'}$ and $\overline{J}_{s,n,n'}$

have spin (s, s-1) and (s-1, s), for $s=1, 3/2, 2, \ldots$ Moreover, the vector fields corresponding to the $J_{s,n,n'}$ and $\overline{J}_{s,n,n'}$ are nonzero because of arguments similar to those that we gave in the chiral case. The $SU(2) \times SU(2)$ content therefore ensures that the J's and \overline{J} 's must exactly fill up the volume preserving polynomial vector fields on Q.

2.6. Departures From The SU(2) Point

Now we will briefly discuss how some of the structures change in the compactified theory under a small departure from the SU(2) point. Some of the most interesting points will be speculative.

The infinitesimal moduli of the theory, for small departure from the SU(2) point, are

$$Z_{s,n,n'}^{\pm} = W_{s,n}^{\pm} \overline{W}_{s,n'}^{\pm}.$$
 (2.41)

I will focus on deformations in which only the Z^+ are excited. (However, the Z^- moduli are also important, for instance in two dimensional black holes.) The Z^+ moduli transform under $SU(2) \times SU(2)$ as

$$\bigoplus_{n=0,1/2,1,\dots} (n,n). \tag{2.42}$$

As we will see in appendix (2), this is precisely the $SU(2) \times SU(2)$ content of a general polynomial function ϕ on the hypersurface Q. Thus, the Z^+ moduli naturally combine into such a function.

Now let us try to guess the geometric meaning of the function ϕ . Such a function gives precisely the data we need to deform the quadric Q to a general nearby hypersurface described by an equation

$$a_1 a_2 - a_3 a_4 = \phi(a_i). \tag{2.43}$$

I conjecture that the first order Z^+ corrections to the operator algebra deform the relation $a_1a_2 - a_3a_4 = 0$ into (2.43). In particular, the cosmological constant corresponds to the case $\phi = \mu$ (a constant). Thus, according to our ansatz, the ground ring at $\mu \neq 0$ but still at the SU(2) radius is the ring of functions on the smooth quadric

$$a_1 a_2 - a_3 a_4 = \mu. (2.44)$$

Thus, conjecturally, to first order, the hypersurface Q can be deformed to an arbitrary nearby affine hypersurface in the ambient four-space. However, the general deformation will be obstructed in second order by a non-zero beta function. This can be computed as follows [12]. Consider a general theory with left and right moving currents J_a and \overline{J}_a , with structure constants $f_{ab}{}^c$ and $\overline{f}_{ab}{}^c$. For an infinitesimal perturbation of the Lagrangian

$$L \to L + \sum w^{aa'} J_a \overline{J}_{a'}, \tag{2.45}$$

the quadratic beta function corresponds to the counterterm

$$\beta = w^{aa'} w^{bb'} f_{ab}{}^c f_{a'b'}{}^{c'} J_c \overline{J}_{c'}. \tag{2.46}$$

We can easily implement this prescription in the present situation. Since the structure constants for left and right movers are just the Poisson brackets in x - y and x' - y', respectively, the beta function is

$$\beta = \epsilon^{ij} \epsilon^{i'j'} \frac{\partial^2 \phi}{\partial x^i \partial x'^{i'}} \frac{\partial^2 \phi}{\partial x^j \partial x'^{j'}}.$$
 (2.47)

After a relatively long but straightforward calculation, one finds that this can be written in terms of the a_i as follows. Let Z be the scaling operator $Z = \sum_i a_i \cdot \partial/\partial a_i$. Let η_{ij} be the metric such that $a_1a_2 - a_3a_4 = \eta_{ij}a^ia^j/2$. Let $\Delta = \eta^{ij}\partial_i\partial_j$ be the corresponding Laplacian. Then

$$\beta = \frac{1}{2} \left(\eta^{ij} (\partial_i Z\phi) (\partial_j Z\phi) - (Z^2 \phi) \cdot \Delta \phi \right). \tag{2.48}$$

An important check on this formula is that if ϕ is a solution of the equation $\beta = 0$, then so is $\phi' = \phi + (a_1a_2 - a_3a_4)f$ for any f; indeed we wish to identify ϕ

and ϕ' , as they coincide on Q. As will be explained in the appendix, there is always a unique choice of f such that $\Delta \phi' = 0$. It is easy to see that if ϕ is a homogenous polynomial (so $Z\phi = \lambda \phi$ for some λ) such that $\Delta \phi = 0$ and ϕ is a solution of $\beta = 0$, then $w(\phi)$ is also a solution of $\beta = 0$ for any function w. The simplest nonconstant function obeying the stated conditions is $\phi = (a_1 a_2 + a_3 a_4)/2$. Consequently (since in fact $\phi = a_1 a_2$ on Q)

$$\overline{\phi} = -\sum_{n=0}^{\infty} \epsilon_n (a_1 a_2)^n \tag{2.49}$$

is a perturbation with vanishing second order beta function, for any values of the constants ϵ_n .

According to our ansatz for the meaning of ϕ , the marginal operator (2.49) corresponds to deforming the quadric Q to a more general hypersurface

$$a_3 a_4 = a_1 a_2 + \sum_{n=0}^{\infty} \epsilon_n (a_1 a_2)^n.$$
 (2.50)

In comparing to matrix models, as we will see, the eigenvalue phase space will correspond to the $a_1 - a_2$ plane, and the curve $a_3a_4 = 0$ in the $a_1 - a_2$ plane will correspond to the fermi surface. The function a_1a_2 will correspond to the standard inverted harmonic oscillator Hamiltonian $H = p^2 - q^2$, and the function on the right hand side of (2.50) is the Hamiltonian of a perturbed matrix model.

The problem can probably be analyzed more completely using a criterion of Chaudhuri and J. A. Schwarz, who claim [12] that a perturbation of the type (2.45) is exactly marginal if and only if the $w^{aa'}$ are non-zero only for a and a' in some abelian subalgebra of the full current algebra.

Actually, under a generic perturbation of the compactified theory near the SU(2) point, much of the structure we have found disappears. For instance, at a generic value of the radius of the X field, it is necessary to match the left and right moving momenta, and so to discard most of the states. This instability of the structure seems not to have an analog in the uncompactified theory. Its proper interpretation is unclear.

3. The Uncompactified Theory; Comparison To The Matrix Model

We now want to consider the uncompactified string theory. Henceforth, therefore, X will be real valued, rather than circle valued. Of course, this is the case that is best understood from the point of view of matrix models.

Of the operators that we have found in §2, the ones that survive in the uncompactified theory are the ones that have $p_L - p_R = 0$, where p_L and p_R are the left and right moving momenta. Recalling the definitions

$$a_1 = xx'$$

$$a_2 = yy'$$

$$a_3 = xy'$$

$$a_4 = yx'$$

$$(3.1)$$

and noting that the (p_L, p_R) values are

$$x: (1,0)/\sqrt{2}$$

$$y: (-1,0)/\sqrt{2}$$

$$x': (0,1)/\sqrt{2}$$

$$y': (0,-1)/\sqrt{2}$$
(3.2)

we see that the allowed operators are precisely the ones that are invariant under

$$K = a_3 \frac{\partial}{\partial a_3} - a_4 \frac{\partial}{\partial a_4}. (3.3)$$

If we set $a_3 = \rho e^{\psi}$, $a_4 = \rho e^{-\psi}$ as in §2.5, then

$$K = \frac{\partial}{\partial \psi}. (3.4)$$

Any K invariant polynomial in the a_i is in fact a polynomial in a_1 and a_2 . (The product a_3a_4 is invariant, but equals a_1a_2 .) Any symmetry of the model must therefore induce a motion of the $a_1 - a_2$ plane – which will turn out to be the eigenvalue phase space of the matrix model.

Now, let us determine the symmetry group of the uncompactified model. This is simply the subgroup of the group of volume-preserving diffeomorphisms of the quadric cone Q which are also K invariant. We recall from (2.38) that the volume form in these coordinates is

$$\Theta = d\psi \ da_1 \ da_2. \tag{3.5}$$

A general K invariant vector field is of the form

$$f(a_1, a_2) \frac{\partial}{\partial a_1} + g(a_1, a_2) \frac{\partial}{\partial a_2} + u(a_1, a_2) \frac{\partial}{\partial \psi}.$$
 (3.6)

Requiring that this preserve Θ , we see that there is no restriction on u, while

$$f\frac{\partial}{\partial a_1} + g\frac{\partial}{\partial a_2} \tag{3.7}$$

must generate a diffeomorphism of the $a_1 - a_2$ plane that preserves the area form da_1da_2 . Thus area-preserving diffeomorphisms of the $a_1 - a_2$ plane appear in the uncompactified theory. The operators

$$u(a_1, a_2) \frac{\partial}{\partial \psi} \tag{3.8}$$

commute with each other, and transform under diffeomorphisms of the $a_1 - a_2$ plane as abelian gauge transformations. The combined structure is thus "gravity" (area-preserving diffeomorphisms) plus abelian gauge theory on the $a_1 - a_2$ plane.

There is an important gap in this discussion, however. While every volume preserving K invariant vector field on the quadric Q determines a vector field (3.6) as just described, the converse is not true. We have to determine what happens to a_3 and a_4 when a given transformation of $\psi - a_1 - a_2$ space is "lifted" to a transformation of the quadric. As $a_3 = \sqrt{a_1 a_2} e^{\psi}$, $a_4 = \sqrt{a_1 a_2} e^{-\psi}$, one

immediately sees that

$$\left(f\frac{\partial}{\partial a_1} + g\frac{\partial}{\partial a_2} + u\frac{\partial}{\partial \psi}\right)a_3 = \frac{a_3}{2} \cdot \left(\frac{f}{a_1} + \frac{g}{a_2}\right) + ua_3$$

$$\left(f\frac{\partial}{\partial a_1} + g\frac{\partial}{\partial a_2} + u\frac{\partial}{\partial \psi}\right)a_4 = \frac{a_4}{2} \cdot \left(\frac{f}{a_1} + \frac{g}{a_2}\right) + ua_4.$$
(3.9)

This is a polynomial vector field in the space of the a_i only if f is divisible by a_1 and g is divisible by a_2 .

The latter requirement means that the symmetries of the theory include not all of the area preserving polynomial vector fields on the $a_1 - a_2$ plane, but only those that preserve the locus $a_1a_2 = 0$. I claim that the $a_1 - a_2$ plane corresponds to the eigenvalue phase space of the matrix model. a_1 and a_2 correspond to p + q and p - q, where q is the matrix eigenvalue and p is its canonical momentum. The locus $a_1a_2 = 0$ corresponds to the fermi surface of the matrix model, that is, it is the surface on which the inverted harmonic oscillator Hamiltonian $H = p^2 - q^2$ vanishes. When we study the matrix model presently, we will see why area preserving diffeormorphisms that preserve the fermi surface arise naturally.

Intuitively, the distinguished role of the locus $a_1a_2 = 0$ comes about because this locus is somewhat analogous to the "branch points" of the projection from the quadric Q to the $a_1 - a_2$ plane. The defining equation $a_3a_4 = a_1a_2$ of the quadric shows that the subspace of Q lying over a generic point in the $a_1 - a_2$ plane is a hyperbola, while over a point with $a_1a_2 = 0$ one has a pair of lines. Because of this difference, a diffeomorphism of the $a_1 - a_2$ plane that does not map the locus $a_1a_2 = 0$ to itself cannot be "lifted" to a K invariant diffeomorphism of Q.

In $\S 2.6$, I have described a hopefully very plausible ansatz according to which general K invariant perturbations of the theory will perturb the defining equation of Q to

$$a_3 a_4 = h(a_1, a_2), (3.10)$$

with h an arbitrary Hamiltonian function. If so, then the "branch locus" would be the curve h=0 in the a_1-a_2 plane. K invariant volume preserving transforma-

tions of the three-manifold (3.10) correspond to area preserving diffeomorphisms of the $a_1 - a_2$ plane that map the locus h = 0 to itself, plus the abelian gauge transformations $u \cdot \partial/\partial \psi$. This would be expected if h corresponds to the one body Hamiltonian of the matrix model. In particular, according to the ansatz, when the cosmological constant is turned on, the hypersurface $a_1a_2 - a_3a_4 = 0$ is deformed to a hypersurface Q_{μ} given by $a_1a_2 - a_3a_4 = \mu$. A K invariant volume preserving diffeomorphism of Q_{μ} is an area preserving diffeomorphism of the $a_1 - a_2$ plane that leaves fixed the curve $a_1a_2 - \mu = 0$. This curve corresponds to the fermi surface of the matrix model at chemical potential μ .

3.1. Symmetries Of The Matrix Model

Let us now analyze the symmetries of the matrix model version of the c=1 theory. Our discussion will be fairly similar to that of [7,13, 14], but we will consider a somewhat larger class of symmetries. The extra symmetries might be considered "trivial" in the matrix model but are important in comparing to the conformal field theory.

Consider the Lagrangian for a free particle moving in the p-q plane:

$$L = \int dt \left(p\dot{q} - H(p,q) \right). \tag{3.11}$$

This can be written

$$L = \int \alpha, \tag{3.12}$$

where α is the one form

$$\alpha = p \ dq - H \ dt. \tag{3.13}$$

A symmetry generator is an arbitrary infinitesimal transformation

$$\delta p = f(p, q, t)$$

$$\delta q = g(p, q, t)$$

$$\delta t = u(p, q, t)$$
(3.14)

that leaves the Lagrangian invariant. To this end, it is not necessary to leave α

invariant. A transformation under which $\alpha \to \alpha + d\beta$ is also a symmetry. Instead of the one form α , we should consider the two form $\omega = d\alpha = dp \ dq - dH \ dt$. A symmetry is a transformation that leaves ω invariant.

Let us consider the matrix model with the standard inverted harmonic oscillator Hamiltonian, $H(p,q) = (p^2 - q^2)/2$. Then $\omega = dp \ dq - (p \ dp - q \ dq)dt$. If we set

$$p' = p \cosh t - q \sinh t$$

$$q' = -p \sinh t + q \cosh t$$
(3.15)

then we find

$$\omega = dp' \ dq'. \tag{3.16}$$

It is therefore easy to identify the vector fields that preserve ω : they are of the form

$$\left(\frac{\partial g(p',q')}{\partial q'}\frac{\partial}{\partial p'} - \frac{\partial g(p',q')}{\partial p'}\frac{\partial}{\partial q'}\right) + u(p',q,t)\frac{\partial}{\partial t}.$$
(3.17)

Note that g is a function of p', q' only but u may also depend on t. This is very similar to the answer that we found in the conformal field theory, with $a_1, a_2 \leftrightarrow p', q'$ and $\psi \leftrightarrow t$. There are two discrepancies: (i) in the conformal field theory u was a function of a_1 and a_2 only, and not ψ ; (ii) in the conformal field theory g was required to obey a certain restriction, such that the surface $a_1a_2 = 0$ is invariant under the symmetry generated by g.

I cannot explain the first discrepancy, but there is a very nice way to parametrize it. Let Θ be the volume form in (p, q, t) space

$$\Theta = dp \ dq \ dt = dp' \ dq' \ dt. \tag{3.18}$$

The transformation in (3.17) preserves Θ as well as ω precisely if u is a function of p', q' only. The invariances of the conformal field theory are thus the transformations that leave invariant the volume form Θ as well as the Lagrangian (3.12) and the surface $a_1a_2 = 0$.

As for that last restriction, which is our second apparent discrepancy, it has a very nice explanation in the matrix model. The (1,0) and (0,1) currents that we found in the conformal field theory are unbroken symmetries of a particular ground state. They should be compared to the symmetries of the matrix model that preserve the ground state. The ground state of the matrix model is the state in which all the single particle levels of H(p,q) < 0 are filled and the others are empty. An area preserving transformation that maps filled states to filled states and empty states to empty states necessarily maps the fermi surface to itself. As $H(p,q) = (p^2 - q^2)/2 = (p'^2 - q'^2)/2$, the area preserving transformations of the p'-q' plane that leave fixed the matrix model ground state are those that leave fixed the fermi surface $p'^2 - q'^2 = 0$. (It must be left fixed as a set, not pointwise.) The fermi surface is obviously isomorphic to the fixed locus $a_1a_2 = 0$ of the conformal field theory, with $a_1 = p' + q'$, $a_2 = p' - q'$.

Thus, we find the dictionary for comparing the conformal field theory to the matrix model: (a_1, a_2, ψ) corresponds to (p' + q', p' - q', t).

Let us give an example of the use of this dictionary. The difference between the left and right moving X momenta, which corresponds to the difference $\partial X - \bar{\partial} X$, annihilates all states in the (uncompactified) conformal field theory. It is mapped under our correspondence to

$$\left. \frac{\partial}{\partial t} \right|_{p',q'},\tag{3.19}$$

which annihilates all matrix model observables since the single particle equation of motion of the matrix model equation of motion is dp'/dt = dq'/dt = 0. The time translation operator of the conformal field theory corresponds to the current $\partial X + \bar{\partial} X$. This is mapped to the canonical transformation of the p' - q' plane generated by the vector field derived from the Hamiltonian $a_1a_2 = p'^2 - q'^2$ – the usual inverted harmonic oscillator, in other words.

We have carried out this discussion at zero cosmological constant, but if the ansatz in §2.6 is correct, then the generalization to nonzero cosmological constant

– and arbitrary perturbation of the matrix model Hamiltonian – is immediate.

Conjectured Interpretation

We will conclude by briefly suggesting a partial interpretation of some of these results. The symmetries of the matrix model preserve both a two form ω and a three form Θ . Since ω depends explicitly on the matrix model Hamiltonian, it seems to be a dynamical variable. On the other hand, Θ may well be more universal. Near the SU(2) point, there seems to be always a volume form analogous to Θ but not necessarily a two form ω . One is tempted to try to think of the theory as a three dimensional theory, with a symmetry group of volume preserving diffeomorphisms, in which one of the fields (analogous to the metric tensor in general relativity) is a two form ω whose equations of motion assert that $d\omega = 0$. The symmetries of a given classical solution would be the subgroup of the group of volume preserving diffeomorphism that leaves fixed both ω and the other degrees of freedom, such as the tachyon field. It would be a challenge to reconcile such a viewpoint with the rich structure that exists upon compactification, for instance near the SU(2) point.

<u>Acknowledgements:</u> I am indebted to R. H. Dijkgraaf, D. J. Gross, A. M. Polyakov, E. Rabinovici, and N. Seiberg for many stimulating discussions.

I understand that I. Klebanov and A. M. Polyakov have recently succeeded in computing directly the current algebra of the chiral spin one discrete fields.

APPENDIX (1): BRST Analysis Of Some Low-Lying States

Our goal in this appendix is to describe the lowest non-trivial examples of some phenomena analyzed theoretically in [9]. First we recall the standard construction of the BRST operator. It is

$$Q = \sum_{n} c_n L_{-n} - \frac{1}{2} \sum_{n,r} c_{-n} c_{-r} b_{n+r} (n-r),$$
 (3.20)

where c_n , b_m , $n, m \in \mathbb{Z}$ are the Fourier modes of ghosts and antighosts, which obey $\{c_n, b_m\} = \delta_{n+m}$, and L_n are the Virasoro generators of matter. The latter are

$$L_n = \frac{1}{2} \sum_{n \in \mathbb{Z}} \left(\alpha_m \alpha_{n-m} + \phi_m \phi_{n-m} \right) + i \sqrt{2} n \phi_n, \tag{3.21}$$

with α_n and ϕ_n (obeying $[\alpha_n, \alpha_m] = [\phi_n, \phi_m] = n\delta_{n+m}$) being the Fourier modes of X and of the Liouville field ϕ . In particular, α_0 and ϕ_0 are the X and ϕ momentum operators.* The Fock vacuum will be labeled $|p,\uparrow\rangle$ or $|p,\downarrow\rangle$, where $p = (\alpha_0, \phi_0)$ labels the momentum eigenvalues, $|p,\downarrow\rangle$ is a state annihilated by b_0 , and $|p,\uparrow\rangle = c_0|p,\downarrow\rangle$.

The above expressions for the L's and Q should all be normal ordered. The normal ordering constants are zero for this two dimensional system (the "tachyon" is a massless particle in D=2).

We will define the ghost number operator, which we will call G, such that $|p,\downarrow\rangle$ has G=0 (from some points of view G=-1/2 is more natural).

Interesting phenomena occur at special values of α_0 and ϕ_0 . We will consider the first two levels. In addition to the points (involving cohomology at G = -1) that were important in this paper, there are other peculiarities of the BRST cohomology that may be important in the future. To exhibit these peculiarities, I will

^{*} In going from states, we will consider in this appendix, to operators, there is a shift $\phi_0 \to \phi_0 - i\sqrt{2}$. The Liouville momentum values given in the text were the values for the operators.

describe the level one situation in full detail. When we get to level two, we will just focus on G = -1. The other peculiarities of level one are repeated at level two (and at all higher levels, according to [9]), but the detailed description would be somewhat long.

3.2. Level One

At level one, discrete states arise at $\alpha_0 = 0$, $\phi_0 = \pm i\sqrt{2}$. We will write the formulas for a more generic situation with arbitrary α_0 and $(\alpha_0^2 + \phi_0^2)/2 = -1$; this condition ensures that L_0 annihilates all the states that we consider. We lose nothing by that restriction, since it is easy to prove that the BRST cohomology arises only at $L_0 = 0$. We will consider the states in order of ascending values of the ghost number.

At level one and G = -1, there is only one state, $b_{-1}|p,\downarrow\rangle$, with

$$Qb_{-1}|p,\downarrow\rangle = \left(\alpha_0\alpha_{-1} + (\phi_0 - i\sqrt{2})\phi_{-1}\right)|p,\downarrow\rangle. \tag{3.22}$$

For G = 0, there are three states, with

$$Qb_{-1}|p,\uparrow\rangle = \left(\alpha_0\alpha_{-1} + (\phi_0 - i\sqrt{2})\phi_{-1}\right)|p,\uparrow\rangle - 2c_{-1}|p,\downarrow\rangle$$

$$Q\alpha_{-1}|p,\downarrow\rangle = \alpha_0c_{-1}|p,\downarrow\rangle$$

$$Q\phi_{-1}|p,\downarrow\rangle = \left(\phi_0 + i\sqrt{2}\right)c_{-1}|p,\downarrow\rangle.$$
(3.23)

For G=1, we get

$$Q\phi_{-1}|p,\uparrow\rangle = \left(\phi_0 + i\sqrt{2}\right)c_{-1}|p,\uparrow\rangle$$

$$Q\alpha_{-1}|p,\uparrow\rangle = \alpha_0c_{-1}|p,\uparrow\rangle$$

$$Qc_{-1}|p,\downarrow\rangle = 0.$$
(3.24)

Finally, at G = 2,

$$Qc_{-1}|p,\uparrow\rangle = 0. (3.25)$$

It is easy to see that the Q cohomology is trivial in this space at generic α_0 .

Something interesting happens at $\alpha_0 = 0$. At that value of α_0 , the L_0 condition permits two values of ϕ_0 , namely $\pm i\sqrt{2}$. The behavior at those two values are completely different. First we consider the value $\phi_0 = i\sqrt{2}$ (which is shifted to $\phi_0 = 0$ when one goes from states to operators).

In this case there is a one dimensional cohomology at G=-1, since $Qb_{-1}|p,\downarrow\rangle=0$ for such p. This state corresponds to the identity operator 1, which is the first of the spin zero, ghost number zero operators that were the main focus of the body of the paper. (G as we have defined it is increased by one in going from states to operators, so states of G=-1 will correspond to operators of G=0, and so on.) At G=0, the cohomology is actually two dimensional. For representatives we can take $\alpha_{-1}|p,\downarrow\rangle$ and $\phi_{-1}|p,\downarrow\rangle+i\sqrt{2}b_{-1}|p,\uparrow\rangle$. The former corresponds to the spin zero, G=1 operator $c\partial X$ and to the spin one, G=0 operator ∂X . The latter is the first of the ghost number zero discrete currents constructed from the bosons only (without the ghosts) and usually considered in discussions of the discrete states. The latter corresponds to the spin zero, G=1 operator $c\partial\phi+\sqrt{2}\partial c$ but does not correspond to any spin one, G=0 current. The reason for this will be explained presently. Finally, to complete the enumeration of the cohomology at this value of the momenta, the G=1 state $\alpha_{-1}|p,\uparrow\rangle$ corresponds to the G=2 spin zero operator $c\partial c\partial X$, but does not correspond to any current.

Relation Between Spin Zero Operators And Currents

Let us now briefly explain why certain of the states just found do not correspond to currents. First of all, given a BRST cohomology class, to find a corresponding BRST invariant spin zero primary field we should find a representative $|\psi\rangle$ of the cohomology class that is of Virasoro highest weight, obeying $L_n|\psi\rangle = 0$, $n \geq 0$. In the cases at hand this is possible; the representatives given above satisfy this condition. In particular, as the highest weight of these states is zero, they correspond to primaries of spin zero. To get a current, we need a highest weight state of spin one. The general strategy for doing this is to set

$$|\alpha\rangle = b_{-1}|\psi\rangle. \tag{3.26}$$

If this vanishes, we cannot proceed further; this is why the G = -1 state $b_{-1}|p,\downarrow\rangle$ does not correspond to a current. Suppose $|\alpha\rangle \neq 0$. Then

$$L_0|\alpha\rangle = L_0 b_{-1} |\psi\rangle = b_{-1} |\psi\rangle = |\alpha\rangle, \tag{3.27}$$

so if $|\alpha\rangle$ is a highest weight state, the highest weight is one, and the operator corresponding to $|\alpha\rangle$ will be a current. Moreover,

$$Q|\alpha\rangle = Qb_{-1}|\psi\rangle = L_{-1}|\psi\rangle. \tag{3.28}$$

Although this is not zero, it will correspond to a total derivative (since acting on operators $L_{-1} \sim \partial/\partial z$); this is good enough. The question that remains is whether $|\alpha\rangle$ is a highest weight vector, that is, whether $L_n|\alpha\rangle$ is zero, or at least a BRST commutator, for n > 0. We have

$$L_n|\alpha\rangle = L_n b_{-1}|\psi\rangle = b_{n-1}|\psi\rangle. \tag{3.29}$$

Now these states may not vanish, but we do at least have

$$Qb_{n-1}|\psi\rangle = L_{n-1}|\psi\rangle = 0, \quad n > 0,$$
 (3.30)

as $|\psi\rangle$ is BRST invariant and highest weight. Therefore, if the BRST invariant states $b_{n-1}|\psi\rangle$ are BRST commutators, then $|\alpha\rangle$ is of highest weight, at least modulo BRST commutators.

For two of the cohomology classes found above, this last condition fails. The obstructions arise for n=1. For $|\psi\rangle = \phi_{-1}|p,\downarrow\rangle + i\sqrt{2}b_{-1}|p,\uparrow\rangle$, we get $b_0|\psi\rangle = -i\sqrt{2}b_{-1}|p,\downarrow\rangle$, which is not a BRST commutator; indeed it represents the now familiar G=-1 cohomology class. For $|\psi\rangle = \alpha_{-1}|p,\uparrow\rangle$, we get $b_0|\psi\rangle = \alpha_{-1}|p,\downarrow\rangle$, the standard discrete state at G=0.

Actually, the only possible obstruction is the one at n = 1 because $L_{n-1} || \psi \rangle$ is an L_0 eigenstate with eigenvalue n - 1, and hence automatically a BRST commutator unless n = 1.

The Other Liouville Dressing

Now we consider what happens for the other Liouville dressing, $\alpha_0 = 0$, $\phi_0 = -i\sqrt{2}$. The important point for the present paper is that nothing interesting happens at G = -1, so we do not get any new operators of spin zero and G = 0. At G = 0, we have $\alpha_{-1}|p,\downarrow\rangle$, which corresponds to the spin zero operator $c\partial X e^{2\sqrt{2}\phi}$ and to the current $\partial X e^{2\sqrt{2}\phi}$; the latter, when combined with right movers, becomes the marginal operator associated with the black hole. At G = 1, the cohomology is two dimensional. One state is $\alpha_{-1}|p,\uparrow\rangle$, which corresponds to the spin zero, G = 2 operator $c\partial c\partial X e^{2\sqrt{2}\phi}$ but not to any spin one current (because of the obstruction described above). The second is $\phi_{-1}|p,\uparrow\rangle$, which corresponds to a spin zero operator $c\partial c\partial \phi e^{2\sqrt{2}\phi}$ of G = 2 and to a current $\partial c\partial \phi e^{2\sqrt{2}\phi}$ of G = 1 that is a spin one conformal field up to a BRST commutator. Finally, at G = 2 we have $c_{-1}|p,\uparrow\rangle$, which corresponds to the spin zero G = 3 field $c\partial c\partial^2 c e^{2\sqrt{2}\phi}$, but not to any current.

3.3. Level Two

At level two, the mass shell condition, which we may as well impose, is $(\alpha_0^2 + \phi_0^2)/2 = -2$. The space of states at G = -1 is three dimensional, generated by $b_{-2}|p,\downarrow\rangle$, $b_{-1}\alpha_{-1}|p,\downarrow\rangle$, and $b_{-1}\phi_{-1}|p,\downarrow\rangle$. We get

$$Qb_{-2}|p,\downarrow\rangle = \left(\alpha_{0}\alpha_{-2} + \frac{1}{2}\alpha_{-1}^{2} + \phi_{0}\phi_{-2} + \frac{1}{2}\phi_{-1}^{2} - 2i\sqrt{2}\phi_{-2} + 3c_{-1}b_{-1}\right)|p,\downarrow\rangle$$

$$Qb_{-1}\alpha_{-1}|p,\downarrow\rangle = \left(\alpha_{0}c_{-1}b_{-1} + \alpha_{0}\alpha_{-1}^{2} + \alpha_{-2} + (\phi_{0} - i\sqrt{2})\phi_{-1}\alpha_{-1}\right)|p,\downarrow\rangle$$

$$Qb_{-1}\phi_{-1}|p,\downarrow\rangle = \left(\alpha_{0}\alpha_{-1}\phi_{-1} + (\phi_{0} - i\sqrt{2})\phi_{-1}^{2} + \phi_{-2} + (\phi_{0} + i\sqrt{2})c_{-1}b_{-1}\right)|p,\downarrow\rangle.$$
(3.31)

The kernel of Q in this space vanishes unless $\alpha_0^2 = 1/2$, that is, $\alpha_0 = \epsilon/\sqrt{2}$ with $\epsilon = \pm 1$. There are then two possible Liouville dressings, with $\phi_0 = \pm 3i/\sqrt{2}$. At $\phi_0 = -3i/\sqrt{2}$, Q remains acyclic at G = -1, as the reader can easily verify

from (3.31). But at $\phi_0 = 3i/\sqrt{2}$, the cohomology is one dimensional, generated by

$$\left(b_{-2} - \frac{\epsilon}{\sqrt{2}}b_{-1}\alpha_{-1} + \frac{i}{\sqrt{2}}b_{-1}\phi_{-1}\right)|p,\downarrow\rangle. \tag{3.32}$$

These states correspond to the spin zero, G=0 operators

$$x = \left(cb + \frac{i}{\sqrt{2}}(\partial X - i\partial\phi)\right) \cdot e^{i(X+i\phi)/\sqrt{2}}$$

$$y = \left(cb - \frac{i}{\sqrt{2}}(\partial X + i\partial\phi)\right) \cdot e^{-i(X-i\phi)/\sqrt{2}}$$
(3.33)

that were discussed in the text and shown to generate multiplicatively the entire ring of spin zero, G=0 operators. They also correspond to the spin one, G=-1 currents $b \cdot \exp(\pm i(X \pm i\phi)/\sqrt{2})$.

The other phenomena that we found at level one all recur at level two. In the text, the important states at $G \ge 0$ were G = 0 states that give rise to currents. At level two, these are

$$\left(\alpha_{-2} - \epsilon\sqrt{2}\alpha_{-1}^2\right)|p,\downarrow\rangle,\tag{3.34}$$

with either Liouville dressing. The corresponding currents are

$$W_{3/2,\epsilon/2}^{\pm} = \left(\partial^2 X - i\epsilon\sqrt{2}(\partial X)^2\right) \exp(i\epsilon X/\sqrt{2} + \phi(2\mp 3)/\sqrt{2}). \tag{3.35}$$

APPENDIX (2): Functions And Vector Fields On Certain Quadrics

In section two we required a knowledge of how polynomial functions and vector fields on the quadric $a_1a_2 - a_3a_4 = 0$ transform under $SO(2,2) \sim SL(2,\mathbb{R}) \times SL(2,\mathbb{R})$. As they transform in finite dimensional representations, and the classification of such representations is discrete, nothing can change if we make a continuous perturbation to the quadric $a_1a_2 - a_3a_4 = \mu$ (which in any case is physically relevant according to the ansatz in §2.6). Moreover, as the discussion is just a question of algebra, we could just as well change the signature and consider the standard sphere

$$\sum_{i=1}^{4} x_i^2 = 1 \tag{3.36}$$

in \mathbb{R}^4 .

The x_i transform under $SU(2) \times SU(2)$ in the (1/2, 1/2) representation. We can make this explicit by writing them as $x^{AA'}$, A, A' = 1, 2; the A and A' indices transform as (1/2, 0) and (0, 1/2), respectively. An n^{th} order polynomial

$$\phi = w_{A_1 A_2 \dots A_n; A'_1 A'_2 \dots A'_n} x^{A_1 A'_1} x^{A_2 A'_2} \dots x^{A_n A'_n}, \tag{3.37}$$

with w symmetric in all A_i and all A'_j , transforms as spin (n/2, n/2). Any polynomial function on the sphere is a linear combination of these, for the following reason. If one were to antisymmetrize in a pair of indices, say A_1 and A_2 , to make $\epsilon_{A_1A_2}x^{A_1A'_1}x^{A_2A'_2}$, then by bose statistics one may as well antisymmetrize in A'_1 and A'_2 . But $\epsilon_{A_1A_2}\epsilon_{A'_1A'_2}x^{A_1A'_1}x^{A_2A'_2} = \sum_i x_i^2$. Such a factor can be discarded, as it equals 1 on the sphere. This justifies the claim made in §2 that the polynomial functions on the sphere transform as

$$\bigoplus_{n=0,1/2,1,\dots} (n,n). \tag{3.38}$$

Regarded as a function on \mathbb{R}^4 (dropping the restriction to $x_i x^i = 1$), ϕ is a solution of the Laplace equation $\Delta \phi = 0$ (where $\Delta = \sum_i \partial^2 / \partial x_i^2$) for the following

reason. $\Delta \phi$ is a polynomial in the x's of degree n-2 that must transform with spin (n/2, n/2); as x has spin (1/2, 1/2), this is impossible. This justifies the claim in §2.6 that every polynomial on the sphere can be extended on \mathbb{R}^4 as a solution of the Laplace equation. Conversely, any polynomial on \mathbb{R}^4 that obeys the Laplace equation is a linear combination of the ϕ 's. To show this, it is enough, by the $SU(2) \times SU(2)$ symmetry and the homogeneity of the Laplace operator, to show that for n > 0, $\left(\sum_i x_i^2\right)^n \cdot (x^{11'})^m$ is not a solution of the Laplace equation. This is easy if one recalls that the Laplacian is $\Delta = \epsilon^{AB} \epsilon^{A'B'} \partial_{AA'} \partial_{BB'}$.

Any polynomial vector field on \mathbb{R}^4 is of the form

$$\sum_{i=1}^{4} f^{i}(x^{j}) \frac{\partial}{\partial x^{i}} \tag{3.39}$$

with each f^i being a function. The $\partial/\partial x_i$ transform as (1/2, 1/2). For each i, the space of possible f^i 's, modulo functions that vanish on the sphere, transforms as $\bigoplus_{n\geq 0}(n,n)$. By combining (1/2,1/2) with $\bigoplus_{n\geq 0}(n,n)$, we get

$$\bigoplus_{n\geq 0} (n,n) \bigoplus_{m\geq 1/2} (m,m) \bigoplus_{r\geq 0} ((r+1,r) \oplus (r,r+1)). \tag{3.40}$$

We want to restrict this to vector fields on the sphere, that is vector fields that leave the surface $\sum_i x_i^2 = 1$ fixed, or in other words vector fields such that $x_i f^i = 0$ on the sphere. In general, $x_i f^i$ would be an arbitrary function on the sphere, transforming as (3.38), so imposing the equation $x_i f^i = 0$ amounts to dropping the first term in (3.40). Thus, the $SU(2) \times SU(2)$ content of the polyomial vector fields on the sphere is actually

$$\bigoplus_{m>1/2} (m,m) \bigoplus_{r\geq 0} ((r+1,r) \oplus (r,r+1)). \tag{3.41}$$

If we want volume preserving vector fields, we must take the orthocomplement

of the vector fields of the type

$$f^i = \partial^i w \tag{3.42}$$

for any function w. The w's would transform like (3.38), but a constant term in w is irrelevant. Removing the vector fields (3.42) thus amounts to removing the first term in (3.41), and thus the volume preserving polynomial vector fields on the sphere transform under $SU(2) \times SU(2)$ as

$$\bigoplus_{r>0} ((r+1,r) \oplus (r,r+1)) \tag{3.43}$$

as was claimed in §2. The various pieces can be described very explicitly. The vector fields

$$w_{A_1,\dots,A_n;A'_1\dots A'_{n+2}} x^{A_1A'_1} \dots x^{A_nA'_n} x^{BA'_{n+1}} \epsilon^{A'_{n+2}B'} \frac{\partial}{\partial x^{BB'}}$$
(3.44)

transform as spin (n/2, n/2 + 1). Volume preserving vector fields of spin (n/2 + 1, n/2) are constructed similarly.

REFERENCES

- J. Goldstone, unpublished; V. G. Kac, "Contravariant Form For Infinite Dimensional Lie Algebras and Superalgebras," in *Group Theoretical Methods* in *Physics*, ed. W. Beiglbock et. al., Lecture Notes in Physics, vol. 94 (Springer-Verlag, 1979); G. Segal, "Unitary Representations Of Some Infinite Dimensional Groups," Comm. Math. Phys. 80 (1981) 301.
- D. J. Gross, I. R. Klebanov, and M. J. Newman, "The Two-Point Correlation Function Of The One Dimensional Matrix Model," Nucl. Phys. B350 (1991) 621; D. J. Gross and I. R. Klebanov, PUPT-1241 (1991).
- 3. A. M. Polyakov, "Self-Tuning Fields And Resonant Correlations in 2D Gravity," Mod. Phys. Lett. **A6** (1991) 635.

- 4. E. Witten, "On String Theory And Black Holes," Phys. Rev. **D44** (1991) 314.
- J. Ellis, N. E. Mavromatos, and D. V. Nanopoulos, "Quantum Coherence And Two-Dimensional Black Holes," CERN-TH.6147/91, CTP-TAMU-41/91.
- 6. D. J. Gross and U. H. Danielson, "On The Correlation Functions Of The Special Operators in c=1 Quantum Gravity," PUPT-1258 (1991).
- 7. G. Moore and N. Seiberg, "From Loops To Fields In Two Dimensional Gravity," Rutgers preprint.
- 8. G. Mandal and S. Wadia, private communication and to appear.
- 9. B. Lian and G. Zuckerman, "New Selection Rules And Physical States in 2D Gravity," Phys. Lett. **B254** (1991) 417, "2D Gravity With c = 1 Matter," Yale University preprint (1991).
- 10. I. Bakas, "The Large N Limit Of Extended Conformal Symmetries," Phys. Lett. B228 (1989) 57, "Area Preserving Diffeomorphisms And Higher Spin Fields In Two Dimensions," in Supermembranes And Physics In 2 + 1 Dimensions, ed. M. Duff et. al (World Scientific, 1990).
- 11. C. N. Pope, L. J. Romans, and X. Shen, "A Brief History of W_{∞} ," in *Strings* 90, ed. R. Arnowitt et. al. (World Scientific, 1991), and references therein.
- L.P. Kadanoff and A. C. Brown, "Correlation Functions On The Critical Lines Of THe Baxter And Ashkin-Teller Models," Ann. Phys. 121 (1979) 318; R. Dijkgraaf, E. Verlinde, and H. Verlinde, "c = 1 Conformal Field Theories on Riemann Surfaces," Comm. Math. Phys. 115 (1988) 649; P. Ginsparg, "Curiosities At c = 1," Nucl. Phys. B295 [FS]21 (1988) 153; S. Chaudhuri and J. A. Schwartz, "A Criterion For Integrably Marginal Operators," Phys. Lett. 219B (1989) 291; A. Giveon, N. Malkin, and E. Rabinovici, "On Discrete Symmetries And Fundamental Domains Of Target Space," Phys. Lett. 238B (1990) 57.

- 13. J. Avan and A. Jevicki, "Classical Integrability And Higher Symmetries of Collective String Field Theory," BROWN-HET-801 (1991).
- 14. D. Minic, J. Polchinski, and Zhu Yang, "Translation-Invariant Backgrounds in 1+1 Dimensional String Theory," UTTG-16-91.