

GUMBEL MDA & TAIL BEHAVIOUR OF INFIMA OF GAUSSIAN PROCESSES

KRZYSZTOF DĘBICKI, ENKELEJD HASHORVA, AND SVYATOSLAV NOVIKOV

ABSTRACT. We study exact high-level asymptotics for probabilities of the form

$$\mathbb{P} \left\{ \inf_{t \in [a, b]} X(t) > u \right\}, \quad u \rightarrow \infty,$$

where X is a centred Gaussian process (GP) with continuous sample paths. As suggested by the findings of [1], the exponential rate is governed by an optimal-measure problem, while the exact prefactor depends on the local geometry of the contact set. The main results treat finite-active regimes in which the optimal measure has finite support. We separate dominated contacts, where residual fluctuations are negligible, from smooth-critical contacts, where a residual derivative gives an explicit quadratic correction, and rough-critical contacts, where a tangent GP contributes the local correction. We also study full-contact regimes through a Laplace-transform reduction and obtain logarithmic refinements for rough fBM. Applications include integrated fractional Brownian motion, a two-interval fractional Brownian phase transition, a fractional Brownian sheet, passive contacts, and smooth and rough endpoint contacts.

1. INTRODUCTION

Let $X(t)$, $t \in \mathcal{T}$ be a centred GP with continuous sample paths on a compact metric space $\mathcal{T}$ and let $R(s, t)$, $s, t \in \mathcal{T}$ be its covariance function. With motivation from [1] in this paper we are interested in the extreme value properties of the infimum $\mathbf{m}_X = \inf_{t \in \mathcal{T}} X(t)$. In order to avoid degenerate cases, we shall assume hereafter that there exists $u_0 > 0$ such that for all $u > u_0$

$$(1) \quad p_{\mathcal{T}}(u) := \mathbb{P} \{ \mathbf{m}_X > u \} > 0.$$

Consider first the simplest case $\mathcal{T} = \{t_1, \dots, t_n\}$, $n \geq 1$ and let $\Sigma = (R(t_i, t_j))_{1 \leq i, j \leq n}$ be nonsingular. It is well-known that the asymptotics of $p_{\mathcal{T}}(u)$ as $u \rightarrow \infty$ are governed by the quadratic programming problem

$$\min_{\mathbf{p} \in \Delta_n} \mathbf{p}^\top \Sigma \mathbf{p}, \quad \Delta_n := \left\{ \mathbf{p} \in [0, \infty)^n : \sum_{i=1}^n p_i = 1 \right\}.$$

Let $\mathbf{p}_*$ be the minimiser and put

$$\sigma_*^2 := (\mathbf{p}_*)^\top \Sigma \mathbf{p}_* > 0, \quad m(t_i) := \frac{(\Sigma \mathbf{p}_*)_i}{\sigma_*^2}, \quad i = 1, \dots, n.$$

The optimality conditions imply that

$$m(t_i) \geq 1, \quad i = 1, \dots, n$$

with equality whenever the i th component of $\mathbf{p}_*$ satisfies $p_{*,i} > 0$. Thus if

$$S := \{t_i \in \mathcal{T} : p_{*,i} > 0\}$$

is the active set, $E = \{t_i \in \mathcal{T} : m(t_i) = 1\}$ is the contact set, and $P = E \setminus S$ is the passive set with ℓ elements, then

$$(2) \quad p_{\mathcal{T}}(u) \sim p_E(u) \sim C \mathbb{P} \{ \min_{t \in S} X(t) > u \}, \quad u \rightarrow \infty,$$

where $C = 1$ if $\ell = 0$, while for $\ell > 0$, C is the orthant probability of the centred Gaussian regression residual at the passive points t_i with $p_{*,i} = 0$; see Lemma 4.2 and [2, Example 6.1 and Theorem 5.1]. Moreover, [3,

Theorem 2.1] implies that the df H of $\mathbf{m}_X$ belongs to the Gumbel max-domain of attraction (GMDA) with scaling function a defined asymptotically by $a(u) = \sigma_*^2/u$, i.e.,

$$(3) \quad \lim_{u \rightarrow \infty} \frac{1 - H(u + xa(u))}{1 - H(u)} = e^{-x}, \quad \forall x \in \mathbb{R}.$$

For $\mathcal{T}$ a compact interval of $\mathbb{R}$, the GMDA of H was established under smoothness assumptions in [1]. For continuous GPs on general compact metric spaces, the GMDA property under the standing non-degeneracy assumption (1) follows from [3, Theorem 2.1].

If H possesses a density $h = H'$, then its canonical scaling is $a(u) = (1 - H(u))/h(u)$ for all u large. The additional condition

$$(4) \quad \lim_{u \rightarrow \infty} a'(u) = 0$$

i.e., the von Mises condition, does not follow from GMDA and absolute continuity alone.

For a general compact parameter space $\mathcal{T}$ our first question concerns the stronger canonical von Mises condition:

Q1. *Under what additional conditions does the canonical scaling of H satisfy the von Mises condition (4)?*

Next, let $\mathcal{M}_1(\mathcal{T})$ denote the set of Borel probability measures on $\mathcal{T}$. For every $\mu \in \mathcal{M}_1(\mathcal{T})$ averaging implies the following upper bound

$$(5) \quad p_{\mathcal{T}}(u) \leq \mathbb{P}\{X_{\mu} > u\}, \quad X_{\mu} := \int_{\mathcal{T}} X(t) \mu(dt).$$

By weak compactness of $\mathcal{M}_1(\mathcal{T})$ and continuity of the covariance function of X , the energy functional $\mu \mapsto \text{Var}(X_{\mu})$ is continuous under weak convergence of probability measures and the minimum is attained. Any minimiser $\mu_* \in \mathcal{M}_1(\mathcal{T})$ is called an *optimal measure* and we set

$$\sigma_*^2 := \min_{\mu \in \mathcal{M}_1(\mathcal{T})} \text{Var}(X_{\mu}).$$

In view of [1], the non-degeneracy assumption (1) implies

$$(6) \quad \sigma_*^2 = \text{Var}(X_{\mu_*}) > 0.$$

Indeed, if $\sigma_*^2 = 0$, then $X_{\mu_*} = 0$ almost surely, which is incompatible with $p_{\mathcal{T}}(u) > 0$ for $u > u_0$.

Next, let μ_* be an optimal measure. In view of the optimality criterion in Lemma A.1, its normalised potential

$$m(t) := \frac{\text{Cov}(X(t), X_{\mu_*})}{\text{Var}(X_{\mu_*})}, \quad t \in \mathcal{T},$$

satisfies

$$m(t) \geq 1, \quad t \in \mathcal{T}, \quad \text{and} \quad m(t) = 1 \quad \text{on} \quad \text{supp}(\mu_*) =: S$$

and hence the support S , which may depend on the choice of optimal measure, is contained in the contact set, i.e.,

$$S \subset E := \{t \in \mathcal{T} : m(t) = 1\}.$$

Moreover, both E and S are compact and further

$$(7) \quad X(t) = m(t)X_{\mu_*} + Z_*(t), \quad t \in \mathcal{T},$$

where the residual process Z_* is a centred GP independent of $X_{\mu_*} \sim N(0, \sigma_*^2)$, see e.g., [1, 3].

Thus S , E , and μ_* are the continuous counterparts of the active set, contact set, and optimal weights in the finite-dimensional problem introduced above.

Optimising the upper bound (5) over μ and using the corresponding large-deviation lower bound [1, 4] gives

$$(8) \quad \lim_{u \rightarrow \infty} \frac{1}{u^2} \log p_{\mathcal{T}}(u) = \lim_{u \rightarrow \infty} \frac{1}{u^2} \log p_S(u) = \lim_{u \rightarrow \infty} \frac{1}{u^2} \log \mathbb{P}\{X_{\mu_*} > u\} = -\frac{1}{2\sigma_*^2} < 0.$$

In this Gaussian setting log-concavity of $p_{\mathcal{T}}$ together with (8) yields the GMDA property, see [3, Section 3.2]. In the case $E = \mathcal{T}$, referred to as the *full-contact regime*, even without knowing the support S explicitly (7) reduces to

$$(9) \quad X(t) = X_{\mu_*} + Z_*(t), \quad t \in \mathcal{T},$$

with Z_* a centred GP independent of $X_{\mu_*} \sim N(0, \sigma_*^2)$.

This class of GP's with full contact is broad. Indeed, if Z_* is a centred GP with continuous sample paths and if $\mu_* \in \mathcal{M}_1(\mathcal{T})$ satisfies

$$\int_{\mathcal{T}} Z_*(t) \mu_*(dt) = 0$$

almost surely, then for an independent Gaussian random variable (rv) $Y \sim N(0, \sigma_*^2)$ the process

$$X(t) := Y + Z_*(t), \quad t \in \mathcal{T}$$

has μ_* as an optimal measure and has full contact set $E = \mathcal{T}$.

In the full-contact regime, integrating (9) against μ_* yields almost surely

$$(10) \quad \int_{\mathcal{T}} Z_*(t) \mu_*(dt) = 0$$

and hence

$$\mathbf{m}_{Z_*} := \inf_{t \in \mathcal{T}} Z_*(t) \leq 0.$$

Moreover with $X_{\mu_*} \sim N(0, \sigma_*^2)$ independent of $\mathbf{m}_{Z_*}$

$$\mathbf{m}_X = X_{\mu_*} + \mathbf{m}_{Z_*}$$

and consequently as shown already in [5] for every $u \in \mathbb{R}$

$$(11) \quad p_{\mathcal{T}}(u) = \frac{\exp(-u^2/(2\sigma_*^2))}{\sqrt{2\pi}\sigma_*} \int_0^\infty \exp\left(-\frac{u\varepsilon}{\sigma_*^2} - \frac{\varepsilon^2}{2\sigma_*^2}\right) \mathbb{P}\{\mathbf{m}_{Z_*} > -\varepsilon\} d\varepsilon.$$

An immediate Abelian consequence of the above identity is

$$\lim_{u \rightarrow \infty} \frac{p_{\mathcal{T}}(u)}{\mathbb{P}\{X_{\mu_*} > u\}} = \mathbb{P}\{\mathbf{m}_{Z_*} = 0\}.$$

If $\mathbb{P}\{\mathbf{m}_{Z_*} = 0\}$ is positive, then both the exact asymptotics of $p_{\mathcal{T}}(u)$ and the von Mises condition (4) holds.

We shall show that (4) is still valid even when $\mathbb{P}\{\mathbf{m}_{Z_*} = 0\} = 0$. For the latter case, the exact formula (11) can be used to transfer the problem to the investigation of $r(\varepsilon) := \mathbb{P}\{\mathbf{m}_{Z_*} > -\varepsilon\}$ as $\varepsilon \downarrow 0$. Indeed, if for some ultimately positive function ℓ , slowly varying at infinity

$$(12) \quad r(\varepsilon) \sim K\varepsilon^\alpha \ell(1/\varepsilon), \quad K > 0, \quad \alpha \geq 0$$

holds as $\varepsilon \downarrow 0$, then by Karamata's Abelian theorem for Laplace transforms and (11) we obtain

$$(13) \quad p_{\mathcal{T}}(u) \sim \frac{K\Gamma(\alpha+1)\sigma_*^{2\alpha+1}}{\sqrt{2\pi}} u^{-(\alpha+1)} \ell(u) \exp\left\{-\frac{u^2}{2\sigma_*^2}\right\}, \quad u \rightarrow \infty.$$

Explicit calculations of the asymptotics of $r(\varepsilon)$, as $\varepsilon \downarrow 0$ are possible in few particular cases, see Example 2.1 below.

Exact asymptotics of $p_{\mathcal{T}}(u)$ are available in particular in the smooth settings of [1] and in special Brownian cases; see [6, 7]. This leads to our second question.

Q2. *Under what conditions can we derive an explicit exact asymptotic expansion of $p_{\mathcal{T}}(u)$ as $u \rightarrow \infty$ in finite-contact regimes?*

We shall address the above question by restricting to the case of the contact set E is finite. This implies that every optimal measure μ_* has finite support and can be written as

$$(14) \quad \mu_* = \sum_{i=1}^k p_{*,i} \delta_{t_i}, \quad p_{*,i} > 0, \quad i = 1, \dots, k, \quad \sum_{i=1}^k p_{*,i} = 1,$$

where δ_{t_i} denotes the Dirac probability measure at t_i .

The support is $S = \{t_1, \dots, t_k\}$ and the weight vector is $\mathbf{p}_* = (p_{*,1}, \dots, p_{*,k})^\top$. If $\Sigma_* = (R(t_i, t_j))_{1 \leq i, j \leq k}$ is nonsingular and $\mathbf{X}_S = (X(t_1), \dots, X(t_k))^\top$, then by Lemma A.1

$$\Sigma_* \mathbf{p}_* = \sigma_*^2 \mathbf{1}, \quad \mathbf{1} = (1, \dots, 1)^\top \in \mathbb{R}^k,$$

which implies

$$(15) \quad m(t) = \mathbf{a}(t)^\top \mathbf{1} \geq 1, \quad X(t) = \mathbf{a}(t)^\top \mathbf{X}_S + Z(t), \quad \mathbf{a}(t) := \Sigma_*^{-1} (R(t_1, t), \dots, R(t_k, t))^\top, \quad t \in \mathcal{T},$$

where Z is a centred GP independent of $\mathbf{X}_S$. The support points of the optimal measure are special since

$$m(t_i) = 1, \quad Z(t_i) = 0, \quad 1 \leq i \leq k.$$

The support points in S are called *active* contacts, while the points in $P = E \setminus S$ are called *passive* contacts. In general the two residual processes Z and Z_* differ since

$$Z_*(t) - Z(t) = \mathbf{a}(t)^\top (\mathbf{X}_S - \mathbf{1} X_{\mu_*}), \quad t \in \mathcal{T},$$

however if S is a singleton, then

$$Z(t) = Z_*(t), \quad t \in \mathcal{T}.$$

The loss-of-dimension phenomenon in (2), i.e., S and not E plays a crucial role in the asymptotics of $p_{\mathcal{T}}(u)$ need not hold for an infinite parameter set, even when E is finite. To see this, let ξ, η be independent standard normal rv's and for $\alpha, \lambda > 0$ set

$$X(t) := \xi(1 + \lambda|t|^\alpha) + \eta t, \quad t \in \mathcal{T} = [-1, 1].$$

For this process, we have

$$\mu_* = \delta_0, \quad S = E = \{0\}, \quad \sigma_*^2 = 1, \quad m(t) - 1 = \lambda|t|^\alpha, \quad Z(t) = \eta t, \quad t \in [-1, 1]$$

and the residual GP Z satisfies the mean-square Hölder condition

$$\text{Var}(Z(t+h) - Z(t+\ell)) \leq c_H |h - \ell|^{2\beta}, \quad t, t+h, t+\ell \in \mathcal{T},$$

with $\beta = c_H = 1$. Direct calculation gives

$$\lim_{u \rightarrow \infty} \frac{p_{\mathcal{T}}(u)}{\mathbb{P}\{\xi > u\}} = \begin{cases} 1, & 0 < \alpha < 2, \\ \left(1 + \frac{1}{2\lambda}\right)^{-1/2}, & \alpha = 2, \\ 0, & \alpha > 2. \end{cases}$$

The three regimes in the above limit reflect the competition between the rate at which $m(t) - 1$ vanishes near 0, controlled by α , and the residual smoothness exponent $\beta = 1$.

Critical contacts balance potential separation of order $|h|^{2\beta}$ with residual fluctuations on the scale $|h|^\beta$. The *smooth-critical* case has quadratic separation and a linear residual tangent, as when $\alpha = 2$ above. Critical contacts with $\beta \in (0, 1)$ are called *rough-critical*. The Brownian endpoint model in Example 3.9 has $\beta = 1/2$ and gives an explicit nontrivial correction factor.

For dominated contacts and the smooth- and rough-critical cases, we obtain exact asymptotics for $p_{\mathcal{T}}(u)$ under local assumptions. Passive orthant factors and smooth quadratic corrections already appear under stronger smoothness assumptions in [1, Theorem 4.1 and equation (4.3)]. Our finite-contact results use local assumptions and also treat rough tangents.

The main contributions are as follows.

- (1) We give an exact finite-contact reduction for dominated support contacts, including the Gaussian orthant factor generated by finitely many passive contacts. Our results cover also general random fields including the fractional Brownian sheet and several other models.
- (2) For a one-point active endpoint, we identify the exact corrections in rough-critical and smooth-critical regimes through a tangent process or a residual derivative.
- (3) In the full-contact regime, we establish the canonical von Mises condition and derive logarithmic refinements for rough fBM using the Laplace-transform representation and small-deviation estimates.

The paper is organised as follows. Section 2 treats the full-contact regime. Section 3 contains the finite-contact results and examples. The proofs are collected in Section 4, and the Appendix records the auxiliary optimal-measure results used in the main text.

2. FULL-CONTACT REGIME

Let $X(t), t \in \mathcal{T}$, $\mu_*, \sigma_*^2, S, E, m(t), t \in \mathcal{T}$ be as in the Introduction and keep our standing assumption $\sigma_*^2 := \text{Var}(X_{\mu_*}) > 0$.

Except for the comparison in the opening example, this section concerns the full-contact regime

$$E = \{t \in \mathcal{T} : m(t) = 1\} = \mathcal{T}.$$

The transfer principle in (9) allows us to reduce the problem of the tail asymptotics of $p_{\mathcal{T}}(u)$ to the study of the small deviation asymptotics of the residual process $Z_*(t), t \in \mathcal{T}$. In particular cases this is a powerful tool for Q1 and for exact full-contact asymptotics. The following fractional Brownian example compares this regime with dominated contact. Throughout the paper we denote by B_H , $H \in (0, 1)$ a standard fractional Brownian motion (fBM).

Example 2.1. Consider B_H , $H \in [1/2, 1)$, $0 < a_1 < b_1 < a_2 < b_2 < \infty$ and set

$$\mathcal{K} := [a_1, b_1] \cup [a_2, b_2], \quad p_H(u) := \mathbb{P} \left\{ \inf_{t \in \mathcal{K}} B_H(t) > u \right\}.$$

Suppose first that $H > 1/2$, then the normalised potential of δ_{a_1} is

$$m_H(t) = \frac{t^{2H} + a_1^{2H} - (t - a_1)^{2H}}{2a_1^{2H}}, \quad t \in \mathcal{K}.$$

Since

$$t^{2H} > a_1^{2H} + (t - a_1)^{2H}, \quad t > a_1,$$

the optimality criterion and its finite-support corollary imply

$$\mu_* = \delta_{a_1}, \quad S = E = \{a_1\}, \quad P = \emptyset.$$

Moreover we have

$$m_H(t) - 1 \sim \frac{H}{a_1}(t - a_1), \quad t \downarrow a_1.$$

For $Z(t) := B_H(t) - m_H(t)B_H(a_1)$ the regression-residual property gives

$$\mathbb{E}\{(Z(t) - Z(s))^2\} \leq |t - s|^{2H}, \quad s, t \in \mathcal{K}.$$

Thus the dominated-contact assumptions hold with $q = 1$ and $\beta = H$. The extension in Remark 3.3 therefore yields

$$p_H(u) \sim \mathbb{P}\{B_H(a_1) > u\} \sim \frac{a_1^H}{\sqrt{2\pi}u} \exp\left\{-\frac{u^2}{2a_1^{2H}}\right\}, \quad u \rightarrow \infty.$$

Now let $H = 1/2$ and write $B := B_{1/2}$. In this case $m_{1/2} \equiv 1$ on $\mathcal{K}$, so $E = \mathcal{K}$. The measure δ_{a_1} is nevertheless the unique optimal measure, because for every $\mu \in \mathcal{M}_1(\mathcal{K})$

$$\text{Var} \left(\int_{\mathcal{K}} B(t) \mu(dt) \right) = a_1 + \int_{a_1}^{b_2} \mu([x, b_2])^2 dx \geq a_1,$$

with equality only for $\mu = \delta_{a_1}$ and hence

$$\sigma_*^2 = a_1, \quad Z_*(t) = B(t) - B(a_1).$$

Set now

$$A := b_1 - a_1, \quad G := a_2 - b_1, \quad C := b_2 - a_2,$$

and let

$$W(t) := B(a_1 + t) - B(a_1), \quad t \geq 0.$$

Then W is a standard Brownian motion independent of $B(a_1)$. Writing

$$\varphi_t(x) := \frac{1}{\sqrt{2\pi t}} \exp\left\{-\frac{x^2}{2t}\right\}, \quad \Phi(x) := \int_{-\infty}^x \varphi_1(y) dy$$

the reflection principle on the first interval and the Markov property across the gap yield

$$\begin{aligned} r_{\mathcal{K}}(\varepsilon) &:= \mathbb{P}\left\{\inf_{t \in [0, A] \cup [A+G, A+G+C]} W(t) > -\varepsilon\right\} \\ &= \int_0^\infty \{\varphi_A(y - \varepsilon) - \varphi_A(y + \varepsilon)\} \int_0^\infty \varphi_G(z - y) \left\{2\Phi\left(\frac{z}{\sqrt{C}}\right) - 1\right\} dz dy. \end{aligned}$$

Consequently, by dominated convergence

$$r_{\mathcal{K}}(\varepsilon) \sim \kappa_{A,G,C} \varepsilon, \quad \varepsilon \downarrow 0,$$

where

$$\kappa_{A,G,C} := \int_0^\infty \frac{2y}{A} \varphi_A(y) \int_0^\infty \varphi_G(z - y) \left\{2\Phi\left(\frac{z}{\sqrt{C}}\right) - 1\right\} dz dy \in (0, \infty).$$

Substitution into (13) yields

$$p_{1/2}(u) \sim \frac{\kappa_{A,G,C} a_1^{3/2}}{\sqrt{2\pi}} u^{-2} \exp\left\{-\frac{u^2}{2a_1}\right\}, \quad u \rightarrow \infty.$$

For a single interval, $\kappa = \sqrt{2/(\pi(b_1 - a_1))}$, recovering the asymptotic in [6]. Thus $H = 1/2$ is a full-contact boundary case, whereas every $H > 1/2$ lies in the dominated regime.

The first result shows that the von Mises condition (4) holds for our full-contact regime without further restrictions.

Theorem 2.2. *If $E = \mathcal{T}$ and $\sigma_*^2 > 0$, then the infimum $\mathfrak{m}_X = \inf_{t \in \mathcal{T}} X(t)$ has a strictly positive C^∞ -density $f_{\mathcal{T}}$ on $\mathbb{R}$ and both*

$$(16) \quad h_{\mathcal{T}}(u) := \frac{f_{\mathcal{T}}(u)}{p_{\mathcal{T}}(u)} = \frac{u}{\sigma_*^2} + o(1), \quad h'_{\mathcal{T}}(u) = \frac{1}{\sigma_*^2} + o(1), \quad u \rightarrow \infty$$

and

$$(17) \quad a_{\mathcal{T}}(u) := \frac{p_{\mathcal{T}}(u)}{f_{\mathcal{T}}(u)} = \frac{\sigma_*^2}{u} + o(u^{-2}), \quad a'_{\mathcal{T}}(u) \sim -\frac{\sigma_*^2}{u^2}, \quad u \rightarrow \infty.$$

The logarithmic asymptotic in (8) was refined for Gaussian Markov processes in [5, Theorem 3.1]. For the rough fBM considered below, [5, Theorem 3.2] already gives the lower correction bound once full support of the optimal measure is known. The matching upper bound follows from the optimal-measure density bound and an L^1 small-deviation estimate.

We focus for the rest of the paper on

$$\mathcal{T} = [a, b], \quad 0 < a < b < \infty$$

unless otherwise stated. For $X = B_H$ with $H \in (0, 1/2)$, [3, Example 2.7 and Lemma A.2] gives $S = E = [a, b]$ and $\mu_*(dt) \geq \eta dt$ on (a, b) for some $\eta > 0$. Our next result identifies the order of the correction to (8).

Proposition 2.3. *If $X = B_H$, $H \in (0, 1/2)$, then there exist constants $D_{\pm} = D_{\pm}(a, b, H)$, with $0 < D_- \leq D_+ < \infty$, such that for all sufficiently large u*

$$-D_+ u^{1/(1+H)} \leq \log \mathbb{P}\left\{\inf_{t \in [a, b]} X(t) > u\right\} + \frac{u^2}{2\sigma_*^2} \leq -D_- u^{1/(1+H)}.$$

3. FINITE CONTACT SETS

We consider $\mathcal{T} = [a, b]$, $-\infty < a < b < \infty$ and assume next that the contact set is finite with

$$S = \{t_1, \dots, t_k\}, \quad E = \{t \in \mathcal{T} : m(t) = 1\} = S \cup P, \quad P = \{s_1, \dots, s_\ell\}.$$

Further we suppose that μ_* is the unique optimal probability measure. The support covariance matrix $\Sigma_* = (R(t_i, t_j))_{1 \leq i, j \leq k}$ is then nonsingular. Indeed, if $\Sigma_* \mathbf{v} = 0$, then

$$\mathbf{1}^\top \mathbf{v} = \sigma_*^{-2} \mathbf{p}_*^\top \Sigma_* \mathbf{v} = 0,$$

and $\mathbf{p}_* + \varepsilon \mathbf{v}$ is a distinct optimal probability vector for all sufficiently small nonzero ε , unless $\mathbf{v} = 0$. Hence μ_* is given from (14) and the finite-support optimality relations imply

$$\Sigma_* \mathbf{p}_* = \sigma_*^2 \mathbf{1}, \quad m(t) = \mathbf{a}(t)^\top \mathbf{1} \geq 1, \quad t \in \mathcal{T},$$

where

$$\mathbf{a}(t) := \Sigma_*^{-1} \mathbf{r}(t), \quad \mathbf{r}(t) := (R(t_1, t), \dots, R(t_k, t))^\top.$$

Writing

$$\mathbf{X}_S = (X(t_1), \dots, X(t_k))^\top$$

the decomposition (15) has a centred Gaussian residual Z , independent of $\mathbf{X}_S$ such that

$$\mathbf{a}(t_i) = \mathbf{e}_i, \quad Z(t_i) = 0, \quad i = 1, \dots, k,$$

where $\mathbf{e}_i$ is the i -th standard basis vector. Setting

$$\mathbf{S}_u := u(\mathbf{X}_S - u\mathbf{1})$$

yields for every $u > 0$

$$(18) \quad X(t) - u = u(m(t) - 1) + Z(t) + \frac{1}{u} \mathbf{a}(t)^\top \mathbf{S}_u, \quad t \in \mathcal{T}.$$

Thus points outside E have a positive deterministic separation $m(t) - 1$. At a passive contact $s \in P$ by contrast $m(s) = 1$ but $Z(s)$ need not vanish, whereas $Z(t_i) = 0$ at every support contact.

3.1. Dominated contacts. A support point t_i is called *dominated* if there exist finite positive constants β_i, C_i, λ_i and $0 < q_i < 2\beta_i$ such that for all sufficiently small admissible h

$$(19) \quad m(t_i + h) - 1 \geq \lambda_i |h|^{q_i},$$

and for all sufficiently small admissible h, ℓ

$$\text{Var}\{Z(t_i + h) - Z(t_i + \ell)\} \leq C_i |h - \ell|^{2\beta_i}.$$

Here and below admissible means that every displayed point belongs to $[a, b]$.

Theorem 3.1. *Under the standing assumptions of this section, if every $t_i \in S$ is dominated, then*

$$\mathbb{P} \left\{ \inf_{t \in [a, b]} X(t) > u \right\} \sim C \mathbb{P}\{X(t_i) > u, i = 1, \dots, k\}, \quad u \rightarrow \infty,$$

with the convention $C = 1$ when $P = \emptyset$ and otherwise $C := \mathbb{P}\{Z(s) > 0, \forall s \in P\} > 0$.

Remark 3.2. The regression decomposition and independence give for $s, t \in \mathcal{T}$

$$\text{Var}(Z(t) - Z(s)) = \text{Var}(X(t) - X(s)) - \text{Var}\left((\mathbf{a}(t) - \mathbf{a}(s))^\top \mathbf{X}_S\right) \leq \text{Var}(X(t) - X(s)).$$

Consequently, in the definition of a dominated contact it is enough to verify the corresponding local mean-square Hölder bound for X .

Remark 3.3. The preceding theorem remains valid when $\mathcal{T} = K$ is a compact subset of $\mathbb{R}^d$. In that case, replace the local dominated-contact assumptions by the following bounds, valid for all sufficiently small admissible $h, h' \in \mathbb{R}^d$

$$m(t_i + h) - 1 \geq \lambda_i \|h\|^{q_i}, \quad \mathbb{E}\{(Z(t_i + h) - Z(t_i + h'))^2\} \leq C_i \|h - h'\|^{2\beta_i}, \quad q_i < 2\beta_i.$$

Here admissible means that all displayed points belong to K . All other assumptions and the conclusion are unchanged, with the infimum taken over K .

Remark 3.4. Let $\mathcal{T} = [a, b]$, where $0 < a < b$, and suppose that $R(a, a) > 0$. If

$$R(t, a) > R(a, a), \quad t \in (a, b],$$

then δ_a is the unique optimal measure and

$$S = E = \{a\}, \quad m(t) = \frac{R(t, a)}{R(a, a)}.$$

If X has stationary increments, $X(0) = 0$ almost surely, and $V(t) := \text{Var}(X(t))$, the preceding condition is equivalent to

$$V(a + h) > V(a) + V(h), \quad 0 < h \leq b - a.$$

If for all sufficiently small $h > 0$

$$V(a + h) - V(a) - V(h) \geq c_0 h^q, \quad V(h) \leq C h^{2\beta}, \quad q < 2\beta,$$

then a is dominated and

$$\mathbb{P} \left\{ \inf_{t \in [a, b]} X(t) > u \right\} \sim \mathbb{P}\{X(a) > u\}, \quad u \rightarrow \infty.$$

This includes fractional Brownian motion with $H > 1/2$, for which $q = 1$ and $\beta = H$.

The same criterion applies to

$$I(t) := \int_0^t Y(s) ds,$$

where Y is a centred stationary Gaussian process with continuous nonnegative covariance r satisfying $r(0) > 0$. Indeed, I has stationary increments, its increment covariance at a is positive and of order h , and its increments are locally mean-square Lipschitz. Thus $q = \beta = 1$. The covariance criterion can also be verified directly for the Riemann–Liouville process

$$X_H^{\text{RL}}(t) := \int_0^t (t - s)^{H - \frac{1}{2}} dB(s), \quad H \in (1/2, 1),$$

for which the covariance separation is of order h and the local increment variance is of order $|h - \ell|^{2H}$. Hence $q = 1$ and $\beta = H$.

Let

$$I_H(t) := \int_0^t B_H(r) dr, \quad H \in (0, 1),$$

where B_H is a standard fractional Brownian motion, and fix $0 < a < b < \infty$. Write

$$R_H(r, s) := \text{Cov}(B_H(r), B_H(s)).$$

For $t > a$ we have

$$\text{Cov}(I_H(a), I_H(t)) - \text{Var}(I_H(a)) = \int_0^a \int_a^t R_H(r, s) ds dr > 0.$$

Thus δ_a is the unique optimal measure and

$$S = E = \{a\}, \quad P = \emptyset.$$

Moreover we have

$$\sigma_*^2 = \frac{a^{2H+2}}{2H+2}, \quad m(a+h) - 1 \sim \frac{H+1}{a} h, \quad h \downarrow 0,$$

while for $s, t \in [a, b]$

$$\text{Var}(I_H(t) - I_H(s)) \leq C|t - s|^2.$$

Hence $q = \beta = 1$, so a is dominated and

$$\mathbb{P} \left\{ \inf_{t \in [a, b]} I_H(t) > u \right\} \sim \mathbb{P}\{I_H(a) > u\} \sim \frac{a^{H+1}}{2\sqrt{\pi(H+1)}u} \exp \left\{ -\frac{H+1}{a^{2H+2}} u^2 \right\}, \quad u \rightarrow \infty.$$

Let X be the centred Gaussian process with covariance

$$R(s, t) = \frac{2st}{s+t}, \quad s, t > 0;$$

see [8]. On $\mathcal{T} = [a, b]$, where $0 < a < b < \infty$

$$m(t) = \frac{R(t, a)}{R(a, a)} = \frac{2t}{a + t} > 1, \quad t > a.$$

Thus δ_a is the unique optimal measure and

$$\mu_* = \delta_a, \quad S = E = \{a\}, \quad \sigma_*^2 = a.$$

For the regression residual $Z(t) = X(t) - m(t)X(a)$

$$m(a + h) - 1 = \frac{h}{2a + h} \geq \frac{h}{2b}, \quad \text{Var}\{Z(t) - Z(s)\} \leq \frac{(t - s)^2}{t + s} \leq \frac{(t - s)^2}{2a}.$$

Hence $q = \beta = 1$ and

$$\mathbb{P} \left\{ \inf_{t \in [a, b]} X(t) > u \right\} \sim \mathbb{P}\{X(a) > u\}, \quad u \rightarrow \infty.$$

Notice that $\lim_{t \rightarrow \infty} R(s, t) = 2s > 0$, so this process does not satisfy the additional condition in [1, Proposition 3.1].

Let X be the fractional Brownian sheet with common Hurst index $H \in (1/2, 1)$ and covariance

$$R(s, t) = \prod_{j=1}^2 \frac{s_j^{2H} + t_j^{2H} - |s_j - t_j|^{2H}}{2}.$$

On $\mathcal{T} = [a, b]^2$, where $0 < a < b < \infty$, put $t_0 = (a, a)$. The normalised potential of δ_{t_0} is

$$m(t) = m_H(t_1)m_H(t_2), \quad m_H(x) := \frac{x^{2H} + a^{2H} - (x - a)^{2H}}{2a^{2H}}.$$

Strict superadditivity of $x \mapsto x^{2H}$ gives $m_H(x) > 1$ for $x > a$. Therefore we obtain

$$\mu_* = \delta_{t_0}, \quad S = E = \{t_0\}, \quad \sigma_*^2 = a^{4H}.$$

Moreover we have

$$m_H(a + x) - 1 = \frac{H}{a}x + o(x), \quad x \downarrow 0$$

and uniformly on $[a, b]^2$

$$\text{Var}\{X(t) - X(s)\} \leq C\|t - s\|^{2H}.$$

The same bound holds for the regression residual. For sufficiently small $h = (h_1, h_2)$ with $h_1, h_2 \geq 0$

$$m(t_0 + h) - 1 \geq \max_{j=1,2} \{m_H(a + h_j) - 1\} \geq c \max(h_1, h_2) \geq \frac{c}{\sqrt{2}}\|h\|$$

for some $c > 0$. Thus $q = 1$, $\beta = H$ and $q < 2\beta$. An application of Remark 3.3 gives

$$\mathbb{P} \left\{ \inf_{t \in [a, b]^2} X(t) > u \right\} \sim \mathbb{P}\{X(a, a) > u\}, \quad u \rightarrow \infty.$$

Example 3.5. Let ξ_1, ξ_2, ζ be independent standard normal rv's, let $\mathcal{T} = [-1, 1]$, and put

$$g(t) := t^2(1 - t^2), \quad a_1(t) := \frac{1 - t + g(t)}{2}, \quad a_2(t) := \frac{1 + t + g(t)}{2}.$$

Define

$$X(t) := a_1(t)\xi_1 + a_2(t)\xi_2 + (1 - t^2)\zeta, \quad t \in [-1, 1].$$

For

$$\mu_* = \frac{1}{2}(\delta_{-1} + \delta_1)$$

we have

$$X_{\mu_*} = \frac{\xi_1 + \xi_2}{2}, \quad \sigma_*^2 = \frac{1}{2}, \quad m(t) = a_1(t) + a_2(t) = 1 + g(t).$$

Hence μ_* is optimal and

$$S = \{-1, 1\}, \quad E = \{-1, 0, 1\}, \quad P = \{0\}.$$

Any other optimal measure must be supported on E . If its masses at $-1, 0, 1$ are x, y, z , respectively, then

$$\text{Var} \left(\int_{\mathcal{T}} X d\mu \right) = (x + y/2)^2 + (z + y/2)^2 + y^2 \geq \frac{1}{2} + y^2.$$

Thus equality is possible only when $y = 0$ and $x = z = 1/2$, proving uniqueness of μ_* .

The support-regression residual is $Z(t) = (1 - t^2)\zeta$. Since

$$g(-1 + h) = 2h + O(h^2), \quad g(1 - h) = 2h + O(h^2),$$

and

$$\text{Var}\{Z(t) - Z(s)\} = (t^2 - s^2)^2 \leq 4|t - s|^2,$$

both support contacts are dominated with $q = \beta = 1$. At the passive contact, $Z(0) = \zeta$, so

$$C = \mathbb{P}\{Z(0) > 0\} = \frac{1}{2}.$$

Consequently, we have

$$\mathbb{P} \left\{ \inf_{t \in [-1, 1]} X(t) > u \right\} \sim \frac{1}{2} \mathbb{P}\{\xi_1 > u, \xi_2 > u\}, \quad u \rightarrow \infty.$$

3.2. Critical rough and smooth contacts. For simplicity, we consider only the case where the optimal measure is a Dirac measure at a single point b , with $R(b, b) > 0$, and

$$E = S = \{b\}.$$

Then $\mu_* = \delta_b$ is the unique optimal measure and

$$m(t) = \frac{R(t, b)}{R(b, b)}, \quad m(b) = 1, \quad m(t) > 1, \quad t \in \mathcal{T} \setminus \{b\}.$$

Since the support is a singleton, there is no distinction between the optimal-measure residual and the support-regression residual. Thus, throughout this subsection

$$Z(t) := X(t) - m(t)X(b), \quad t \in \mathcal{T},$$

is independent of $X(b)$, satisfies $Z(b) = 0$, and has continuous sample paths.

We say that b is a tangent-critical endpoint contact with exponent $\beta \in (0, 1]$, coefficient $\lambda > 0$, and tangent process W , if the following conditions hold. First, for some $C > 0$ and $0 < r_0 \leq b - a$

$$(20) \quad \text{Var}\{Z(b - h) - Z(b - \ell)\} \leq C|h - \ell|^{2\beta}, \quad 0 \leq h, \ell \leq r_0.$$

Second, as $h \downarrow 0$

$$(21) \quad m(b - h) - 1 = \lambda h^{2\beta} + o(h^{2\beta}).$$

Third, for every $L > 0$, as $r \downarrow 0$, the following convergence of finite-dimensional distributions holds:

$$(22) \quad \left\{ r^{-\beta} Z(b - rx), 0 \leq x \leq L \right\} \Rightarrow \{W(x), 0 \leq x \leq L\},$$

where $W = \{W(x), x \geq 0\}$ is a centred GP with continuous sample paths and $W(0) = 0$. When $\beta \in (0, 1)$, we call the contact rough-critical.

A smooth-critical endpoint contact is the differentiable tangent-critical case $\beta = 1$, in which the residual has a linear L^2 -tangent. More precisely, we assume that as $h \downarrow 0$

$$m(b - h) - 1 = \lambda h^2 + o(h^2)$$

for some $\lambda > 0$, and that for some centred Gaussian rv η

$$\frac{Z(b - h)}{h} \longrightarrow \eta \quad \text{in } L^2, \quad h \downarrow 0.$$

We also assume that (20) holds for some $C > 0, 0 < r_0 \leq b - a$ with $\beta = 1$.

Theorem 3.6. *Under the standing assumptions, let $\mathcal{T} = [a, b]$ with $a < b$. Assume that $R(b, b) > 0$, that $\mu_* = \delta_b$ is optimal, and that $E = \{b\}$. If b is a tangent-critical endpoint contact, then $M := \inf_{x \geq 0} \{W(x) + \lambda x^{2\beta}\} \in (-\infty, 0]$ almost surely, and*

$$p_{\mathcal{T}}(u) \sim C_M \mathbb{P}\{X(b) > u\}, \quad C_M := \mathbb{E} \left\{ \exp \left\{ \frac{M}{R(b, b)} \right\} \right\}, \quad u \rightarrow \infty.$$

Corollary 3.7. *Under the standing assumptions assume that $R(b, b) > 0$, that $\mu_* = \delta_b$ is optimal, and that $E = \{b\}$. If b is a smooth-critical endpoint contact and $v^2 := \text{Var}(\eta)$, then*

$$p_{\mathcal{T}}(u) \sim \left\{ \frac{1}{2} + \frac{1}{2} \left(1 + \frac{v^2}{2\lambda R(b, b)} \right)^{-1/2} \right\} \mathbb{P}\{X(b) > u\}, \quad u \rightarrow \infty.$$

Remark 3.8. The left-endpoint versions follow by applying the results to $\tilde{X}(t) := X(a + b - t)$.

For an interior one-point contact t_0 , the local increment bound must hold across both sides. Suppose also that for every $L > 0$

$$\sup_{|x| \leq L} \left| r^{-2\beta} \{m(t_0 + rx) - 1\} - d(x) \right| \longrightarrow 0, \quad r \downarrow 0,$$

and

$$\{r^{-\beta} Z(t_0 + rx), |x| \leq L\} \Rightarrow \{W(x), |x| \leq L\} \quad \text{in } C[-L, L], \quad r \downarrow 0,$$

and that $m(t_0 + h) - 1 \geq c|h|^{2\beta}$ for all sufficiently small $|h|$. The same proof then uses the two-sided variational term

$$M = \inf_{x \in \mathbb{R}} \{W(x) + d(x)\}.$$

In the symmetric smooth case, $W(x) = \eta x$ and $d(x) = \lambda x^2$, so $M = -\eta^2/(4\lambda)$. In the asymmetric case

$$M = \min\{M_-, M_+\}, \quad M_- := \inf_{x \leq 0} \{W(x) + d(x)\}, \quad M_+ := \inf_{x \geq 0} \{W(x) + d(x)\},$$

where the two branches come from the joint two-sided tangent.

Example 3.9. Let B be a standard Brownian motion independent of $\xi \sim N(0, 1)$, and define

$$X(t) := \xi(1 + \lambda t) + \sigma B(t), \quad t \in [0, 1], \quad \lambda, \sigma > 0.$$

Then

$$\mu_* = \delta_0, \quad S = E = \{0\}, \quad m(t) = 1 + \lambda t, \quad Z(t) = \sigma B(t).$$

Brownian scaling implies the following equality in distribution for every $L < \infty$

$$\{r^{-1/2} Z(rx), 0 \leq x \leq L\} \stackrel{d}{=} \{\sigma B(x), 0 \leq x \leq L\}.$$

Thus 0 is a rough-critical left-endpoint contact with $\beta = 1/2$, and

$$M = \inf_{x \geq 0} \{\sigma B(x) + \lambda x\}.$$

Since $-M$ is exponential with rate $2\lambda/\sigma^2$, the left-endpoint version of Theorem 3.6 gives

$$\mathbb{P} \left\{ \inf_{0 \leq t \leq 1} X(t) > u \right\} \sim \frac{2\lambda}{2\lambda + \sigma^2} \mathbb{P}\{\xi > u\}, \quad u \rightarrow \infty.$$

Example 3.10. Fix $0 < c < 1$, $\alpha > 1$ and set

$$X(t) := \frac{B(t)}{t^\alpha}, \quad t \in \mathcal{T} = [c, 1].$$

Since $X(1) = B(1)$ and

$$m(t) = \frac{\text{Cov}(X(t), X(1))}{\text{Var } X(1)} = t^{1-\alpha},$$

the optimality criterion gives

$$\mu_* = \delta_1, \quad E = S = \{1\}, \quad Z(t) = t^{-\alpha} \{B(t) - tB(1)\}.$$

As $h \downarrow 0$

$$m(1-h) - 1 = (\alpha - 1)h + o(h),$$

locally uniformly after replacing h by rx with x in a compact interval. Moreover, for $s \leq t$ near 1

$$\text{Var}(X(t) - X(s)) = \frac{t-s}{t^{2\alpha}} + s(t^{-\alpha} - s^{-\alpha})^2 \leq C|t-s|.$$

The same bound holds for the regression residual Z .

For fixed $L < \infty$ and sufficiently small r , set

$$\tilde{B}_r(x) := r^{-1/2}\{B(1) - B(1-rx)\}, \quad 0 \leq x \leq L.$$

Then $\tilde{B}_r$ is a standard Brownian motion on $[0, L]$, and

$$r^{-1/2}Z(1-rx) = (1-rx)^{-\alpha}\{-\tilde{B}_r(x) + r^{1/2}xB(1)\} \Rightarrow -\tilde{B}(x)$$

in $C[0, L]$, where $\tilde{B}$ is a standard Brownian motion. Hence 1 is rough-critical with

$$\beta = \frac{1}{2}, \quad \lambda = \alpha - 1.$$

By Brownian symmetry, its variational rv may be written as

$$M = \inf_{x \geq 0} \{\tilde{B}(x) + (\alpha - 1)x\}.$$

Since $-M$ is exponential with rate $2(\alpha - 1)$, Theorem 3.6 yields

$$\mathbb{P}\left\{\inf_{t \in [c, 1]} \frac{B(t)}{t^\alpha} > u\right\} \sim \frac{2(\alpha - 1)}{2\alpha - 1} \mathbb{P}\{B(1) > u\}, \quad u \rightarrow \infty.$$

Remark 3.11. Let $Y \sim N(0, \tau^2)$, $\tau > 0$, and let V be a centred Gaussian random element of $C^1[0, 1]$, independent of Y , with $V(1) = 0$ almost surely. Suppose that $m \in C[0, 1]$ satisfies

$$m(1) = 1, \quad m(t) > 1, \quad 0 \leq t < 1, \quad m(1-h) = 1 + \lambda h^2 + o(h^2)$$

for some $\lambda > 0$, and put $X(t) = m(t)Y + V(t)$. Then

$$\mu_* = \delta_1, \quad E = S = \{1\}, \quad \sigma_*^2 = \tau^2, \quad Z = V.$$

Moreover, Fernique's theorem and the fundamental theorem of calculus give

$$\mathbb{E}\{(V(t) - V(s))^2\} \leq |t-s|^2 \mathbb{E}\{\|V'\|_\infty^2\}, \quad \frac{V(1-h)}{h} \longrightarrow -V'(1) \quad \text{in } L^2.$$

Thus Corollary 3.7 applies with $v^2 = \text{Var}(V'(1))$ and yields

$$\mathbb{P}\left\{\inf_{0 \leq t \leq 1} X(t) > u\right\} \sim \left\{\frac{1}{2} + \frac{1}{2} \left(1 + \frac{v^2}{2\lambda\tau^2}\right)^{-1/2}\right\} \mathbb{P}\{Y > u\}, \quad u \rightarrow \infty.$$

4. PROOFS

Standing notation for finite-active proofs: Whenever a finite-support optimal measure is under consideration, we write

$$A_u := \{X(t_i) > u, \ i = 1, \dots, k\}, \quad B_u := \left\{\inf_{t \in [a, b]} X(t) > u\right\},$$

and

$$\mathbb{P}_u(\cdot) := \mathbb{P}(\cdot \mid A_u), \quad \mathbf{S}_u := u(\mathbf{X}_S - u\mathbf{1}).$$

If the passive set is finite, we write

$$P = \{s_1, \dots, s_\ell\}.$$

Since $B_u \subset A_u$, all exact-asymptotic proofs below reduce to proving

$$\lim_{u \rightarrow \infty} \mathbb{P}_u(B_u) = C$$

for the appropriate constant C . The residual process Z is independent of $\mathbf{X}_S$ and remains so after conditioning on A_u . Setting

$$\boldsymbol{\theta} := \Sigma_*^{-1} \mathbf{1} = \frac{\mathbf{p}_*}{\sigma_*^2}$$

we have $\theta_i > 0$ for every i . For $\mathbf{x} \in (0, \infty)^k$ a change of variables in the Gaussian density shows that the conditional density of $\mathbf{S}_u$ under $\mathbb{P}_u$ is

$$f_u(\mathbf{x}) = \frac{g_u(\mathbf{x})}{\int_{(0, \infty)^k} g_u(\mathbf{y}) d\mathbf{y}}, \quad g_u(\mathbf{x}) := \exp \left\{ -\boldsymbol{\theta}^\top \mathbf{x} - \frac{1}{2u^2} \mathbf{x}^\top \Sigma_*^{-1} \mathbf{x} \right\}.$$

Since Σ_*^{-1} is positive definite

$$0 \leq g_u(\mathbf{x}) \leq \exp\{-\boldsymbol{\theta}^\top \mathbf{x}\},$$

and the function on the right is integrable on $(0, \infty)^k$. Dominated convergence therefore gives under $\mathbb{P}_u$

$$(23) \quad \mathbf{S}_u \Rightarrow \boldsymbol{\mathcal{E}} := (\mathcal{E}_1, \dots, \mathcal{E}_k), \quad u \rightarrow \infty,$$

where the coordinates of $\boldsymbol{\mathcal{E}}$ are independent exponential random variables with respective rates $\theta_1, \dots, \theta_k$. In particular, $\mathbf{S}_u$ is tight under $\mathbb{P}_u$.

In the one-point endpoint proofs below we take $t_1 = b$. Then $A_u = \{X(b) > u\}$, $S_u = u(X(b) - u)$, and the preceding convergence holds with exponential rate

$$\theta = \frac{1}{R(b, b)}.$$

The regression decomposition (18) gives

$$\sup_{t \in \mathcal{T}} |u^{-1} X(t) - m(t)| \leq \frac{\sup_{t \in \mathcal{T}} |Z(t)|}{u} + \frac{1}{u^2} \sup_{t \in \mathcal{T}} \|\mathbf{a}(t)\| \|\mathbf{S}_u\|.$$

The first term tends to zero in $\mathbb{P}_u$ -probability because Z has continuous sample paths and is independent of A_u ; the second does so by compactness and the tightness in (23). Hence, if $D \subset [a, b]$ is compact and $\inf_D(m-1) > 0$, then

$$(24) \quad \lim_{u \rightarrow \infty} \mathbb{P}_u \left\{ \inf_{t \in D} X(t) > u \right\} = 1.$$

4.1. Proofs for full contact.

Proof of Theorem 2.2. Set next

$$V := -\inf_{t \in \mathcal{T}} Z_*(t), \quad r(\varepsilon) := \mathbb{P}\{V < \varepsilon\}, \quad \varepsilon > 0,$$

and write $s := \sigma_*^2$. Path continuity and compactness of $\mathcal{T}$ make V finite almost surely, while (10) gives $V \geq 0$. Since $E = \mathcal{T}$

$$\mathbf{m}_X = X_{\mu_*} - V,$$

where $X_{\mu_*} \sim N(0, s)$ is independent of V . Consequently, we have

$$f_{\mathcal{T}}(u) = \frac{1}{\sqrt{2\pi s}} \mathbb{E} \exp \left\{ -\frac{(u+V)^2}{2s} \right\}.$$

This density is strictly positive. Every derivative of the Gaussian density is bounded, so differentiation under the expectation also shows that $f_{\mathcal{T}} \in C^\infty(\mathbb{R})$. With the notation

$$I(u) := \int_0^\infty \exp \left\{ -\frac{u\varepsilon}{s} - \frac{\varepsilon^2}{2s} \right\} r(\varepsilon) d\varepsilon$$

in view of (11)

$$(25) \quad p_{\mathcal{T}}(u) = \frac{e^{-u^2/(2s)}}{\sqrt{2\pi s}} I(u).$$

For every $\delta > 0$, the centred Gaussian random element Z_* has positive probability of belonging to every neighbourhood of zero. Consequently $r(\delta) > 0$. Define the finite measure

$$\nu(d\varepsilon) := e^{-\varepsilon^2/(2s)} r(\varepsilon) d\varepsilon, \quad \varepsilon > 0,$$

and the probability measure

$$\Pi_u(d\varepsilon) := \frac{e^{-u\varepsilon/s} \nu(d\varepsilon)}{I(u)}.$$

For every fixed $\delta > 0$, we have $\nu((0, \delta/2)) > 0$ and

$$I(u) \geq e^{-u\delta/(2s)} \nu((0, \delta/2)), \quad u > 0.$$

Since $0 \leq r \leq 1$, the second moment of ν is finite. Splitting at δ , we therefore obtain

$$\begin{aligned} \int_0^\infty \varepsilon^2 \Pi_u(d\varepsilon) &\leq \delta^2 + \frac{e^{-u\delta/s} \int_0^\infty \varepsilon^2 \nu(d\varepsilon)}{e^{-u\delta/(2s)} \nu((0, \delta/2))} \\ &\leq \delta^2 + C_\delta e^{-u\delta/(2s)}. \end{aligned}$$

Letting first $u \rightarrow \infty$ and then $\delta \downarrow 0$ gives

$$(26) \quad \lim_{u \rightarrow \infty} \int_0^\infty \varepsilon^2 \Pi_u(d\varepsilon) = 0.$$

Set

$$\bar{\varepsilon}_u := \int_0^\infty \varepsilon \Pi_u(d\varepsilon), \quad \tau_u^2 := \text{Var}_{\Pi_u}(\varepsilon).$$

In view of the Cauchy–Schwarz inequality and (26), we have as $u \rightarrow \infty$

$$(27) \quad 0 \leq \bar{\varepsilon}_u \leq \left(\int_0^\infty \varepsilon^2 \Pi_u(d\varepsilon) \right)^{1/2} \rightarrow 0, \quad 0 \leq \tau_u^2 \leq \int_0^\infty \varepsilon^2 \Pi_u(d\varepsilon) \rightarrow 0.$$

Since $0 \leq r \leq 1$, the Gaussian factor provides an integrable bound for the first two derivative integrands. Differentiation under the integral sign therefore gives

$$\frac{d}{du} \log I(u) = -\frac{\bar{\varepsilon}_u}{s}, \quad \frac{d^2}{du^2} \log I(u) = \frac{\tau_u^2}{s^2}.$$

Let

$$\psi(u) := -\log p_{\mathcal{T}}(u).$$

Since $p'_{\mathcal{T}}(u) = -f_{\mathcal{T}}(u)$, we have $\psi'(u) = h_{\mathcal{T}}(u)$. In light of (25), we can write

$$\psi(u) = \frac{u^2}{2s} + \log \sqrt{2\pi s} - \log I(u)$$

implying

$$h_{\mathcal{T}}(u) = \psi'(u) = \frac{u + \bar{\varepsilon}_u}{s}, \quad h'_{\mathcal{T}}(u) = \psi''(u) = \frac{1}{s} - \frac{\tau_u^2}{s^2}.$$

Together with (27), these identities prove (16). Since $a_{\mathcal{T}} = 1/h_{\mathcal{T}}$, the same identities imply

$$a_{\mathcal{T}}(u) = \frac{s}{u + \bar{\varepsilon}_u} = \frac{s}{u} + o(u^{-2}), \quad a'_{\mathcal{T}}(u) = -\frac{h'_{\mathcal{T}}(u)}{h_{\mathcal{T}}(u)^2} = -\frac{s - \tau_u^2}{(u + \bar{\varepsilon}_u)^2} \sim -\frac{s}{u^2},$$

establishing the claim. $\square$

Lemma 4.1. *Assume that $E = \mathcal{T}$ and $\sigma_*^2 > 0$ and set*

$$m_Z := \inf_{t \in \mathcal{T}} Z_*(t), \quad r(\varepsilon) := \mathbb{P}\{m_Z > -\varepsilon\}.$$

If for some $\beta, A_1, A_2, \varepsilon_0 > 0$

$$(28) \quad \exp\{-A_2 \varepsilon^{-1/\beta}\} \leq \mathbb{P}\{m_Z > -\varepsilon\} \leq \exp\{-A_1 \varepsilon^{-1/\beta}\}, \quad 0 < \varepsilon < \varepsilon_0,$$

then there exist constants $0 < C_- \leq C_+ < \infty$ such that for all sufficiently large u

$$(29) \quad -C_+ u^{1/(\beta+1)} \leq \log p_{\mathcal{T}}(u) + \frac{u^2}{2\sigma_*^2} \leq -C_- u^{1/(\beta+1)}.$$

Proof of Lemma 4.1. Write

$$I_u := \int_0^\infty \exp \left\{ -\frac{u\varepsilon}{\sigma_*^2} - \frac{\varepsilon^2}{2\sigma_*^2} \right\} r(\varepsilon) d\varepsilon$$

and put $\kappa := 1/(\beta+1)$ and $\varepsilon_u := u^{-\beta/(\beta+1)}$. For all large u , $2\varepsilon_u < \varepsilon_0$. Splitting at ε_u , using monotonicity of r on the first part and $r \leq 1$ on the second, gives

$$I_u \leq \frac{\sigma_*^2}{u} \left[r(\varepsilon_u) + e^{-u\varepsilon_u/\sigma_*^2} \right] \leq \frac{\sigma_*^2}{u} \left[e^{-A_1 u^\kappa} + e^{-u^\kappa/\sigma_*^2} \right].$$

Hence $\log I_u \leq -c_2 u^\kappa$ for some $c_2 > 0$ and all large u .

For the lower bound, monotonicity and (28) give $r(\varepsilon) \geq r(\varepsilon_u) \geq e^{-A_2 u^\kappa}$ on $[\varepsilon_u, 2\varepsilon_u]$. Consequently, we obtain

$$I_u \geq \varepsilon_u \exp \left\{ -\left(A_2 + \frac{2}{\sigma_*^2} \right) u^\kappa - 1 \right\}$$

for all sufficiently large u ; here the final constant bounds $\varepsilon^2/(2\sigma_*^2)$ uniformly on $[\varepsilon_u, 2\varepsilon_u]$. Since $|\log \varepsilon_u| = O(\log u) = o(u^\kappa)$, we obtain $\log I_u \geq -c_3 u^\kappa$. Finally, (11) gives

$$\log p_{\mathcal{T}}(u) + \frac{u^2}{2\sigma_*^2} = \log I_u - \log(\sqrt{2\pi}\sigma_*),$$

and the claimed two-sided bound follows after adjusting the constants. $\square$

Proof of Proposition 2.3. Let μ_* be an optimal measure. By [3, Example 2.7 and Lemma A.2], there exists $\eta > 0$ such that

$$\int_a^b R(s, t) \mu_*(ds) = \sigma_*^2, \quad t \in [a, b], \quad \mu_*(dt) \geq \eta dt \quad \text{on } (a, b).$$

In particular, $\sigma_*^2 > 0$ and $E = [a, b]$. Define

$$Y := \int_a^b X(t) \mu_*(dt), \quad Z(t) := X(t) - Y, \quad m_Z := \inf_{t \in [a, b]} Z(t).$$

Thus $Z = Z_*$, $Y \sim N(0, \sigma_*^2)$, Y is independent of Z , and

$$\int_a^b Z(t) \mu_*(dt) = 0 \quad \text{a.s.}$$

We first record the small-deviation estimates needed below. For some $c, C, \varepsilon_0 > 0$, and all $0 < \varepsilon < \varepsilon_0$

$$(30) \quad \begin{aligned} \mathbb{P} \left\{ \sup_{0 \leq t \leq b} |X(t)| \leq \varepsilon \right\} &\geq \exp\{-C\varepsilon^{-1/H}\}, \\ \mathbb{P} \left\{ \int_a^b |X(t)| dt \leq \varepsilon \right\} &\leq \exp\{-c\varepsilon^{-1/H}\}. \end{aligned}$$

The first inequality is the sup-norm small-deviation estimate of [9]; the second follows from [10, Theorem 5.7 and equation (5.13), with $N = D = q = 1$].

Put

$$r(\varepsilon) := \mathbb{P}\{m_Z > -\varepsilon\}, \quad \|f\|_1 := \int_a^b |f(t)| dt.$$

If $\sup_{0 \leq t \leq b} |X(t)| \leq \varepsilon/3$, then for $t \in [a, b]$

$$|Z(t)| = \left| \int_a^b \{X(t) - X(s)\} \mu_*(ds) \right| \leq \frac{2\varepsilon}{3}.$$

The first estimate in (30) therefore gives for some $0 < A_2 < \infty$ and all sufficiently small $\varepsilon > 0$

$$r(\varepsilon) \geq \mathbb{P} \left\{ \sup_{0 \leq t \leq b} |X(t)| \leq \frac{\varepsilon}{3} \right\} \geq \exp\{-A_2 \varepsilon^{-1/H}\}.$$

The lower-bound argument in the proof of Lemma 4.1, with $\beta = H$, now gives the required lower correction bound.

For the upper bound, write $s := \sigma_*^2$. We use the change of measure in the proof of [5, Theorem 3.1]. Applied to the independent Gaussian coordinate Y , it gives

$$e^{u^2/(2s)} p_{\mathcal{T}}(u) = \mathbb{E} \left[e^{-uY/s} \mathbf{1}_{\{\inf_{t \in [a,b]} X(t) > 0\}} \right].$$

On the event in the indicator, the lower density bound for μ_* implies

$$Y = \int_a^b X(t) \mu_*(dt) \geq \eta \int_a^b |X(t)| dt = \eta \|X\|_1.$$

Consequently, for every $\varepsilon > 0$

$$e^{u^2/(2s)} p_{\mathcal{T}}(u) \leq e^{-\eta u \varepsilon / s} + \mathbb{P}\{\|X\|_1 \leq \varepsilon\}.$$

Taking $\varepsilon = u^{-H/(1+H)}$, the second estimate in (30) gives

$$e^{u^2/(2s)} p_{\mathcal{T}}(u) \leq e^{-(\eta/s)u^{1/(1+H)}} + e^{-cu^{1/(1+H)}}.$$

The required upper correction bound follows after taking logarithms and adjusting the constant. $\square$

4.2. Proofs for finite contact.

Lemma 4.2. *Assume that*

$$\mu_* = \sum_{i=1}^k p_{*,i} \delta_{t_i}, \quad p_{*,i} > 0, \quad i = 1, \dots, k,$$

is the unique optimal probability measure. If $P = E \setminus S = \{s_1, \dots, s_\ell\}$ is nonempty, then

$$\mathbb{P}\{Z(s_j) \geq 0, j = 1, \dots, \ell\} = \mathbb{P}\{Z(s_j) > 0, j = 1, \dots, \ell\} > 0.$$

Proof. Define next

$$\Delta := (Z(s_1), \dots, Z(s_\ell))^\top$$

and let $L \subseteq \mathbb{R}^\ell$ be the linear support of its centred Gaussian law. Suppose that

$$\mathbb{P}\{\Delta_j > 0, j = 1, \dots, \ell\} = 0.$$

Then $L \cap (0, \infty)^\ell = \emptyset$. Gordan's alternative provides a nonzero vector $\lambda \in [0, \infty)^\ell$ such that

$$\lambda^\top x = 0, \quad x \in L.$$

Consequently, $\lambda^\top \Delta = 0$ almost surely.

Let

$$A := (a_i(s_j))_{\substack{1 \leq i \leq k \\ 1 \leq j \leq \ell}}, \quad c := A\lambda.$$

Since $m(s_j) = \sum_{i=1}^k a_i(s_j) = 1$

$$\sum_{i=1}^k c_i = \sum_{j=1}^\ell \lambda_j m(s_j) = \sum_{j=1}^\ell \lambda_j.$$

As every $p_{*,i} > 0$, for all sufficiently small $\varepsilon > 0$

$$\mu_\varepsilon := \sum_{i=1}^k (p_{*,i} - \varepsilon c_i) \delta_{t_i} + \varepsilon \sum_{j=1}^\ell \lambda_j \delta_{s_j}$$

is a probability measure. Indeed, its coefficients are nonnegative and the preceding identity shows that their sum is one. Writing

$$X_P := (X(s_1), \dots, X(s_\ell))^\top,$$

we have $\mathbf{X}_P = \mathbf{A}^\top \mathbf{X}_S + \mathbf{\Delta}$, and hence

$$\begin{aligned} \int_{\mathcal{T}} X(t) \mu_\varepsilon(dt) - \int_{\mathcal{T}} X(t) \mu_*(dt) &= \varepsilon (\boldsymbol{\lambda}^\top \mathbf{X}_P - \mathbf{c}^\top \mathbf{X}_S) \\ &= \varepsilon \boldsymbol{\lambda}^\top (\mathbf{X}_P - \mathbf{A}^\top \mathbf{X}_S) \\ &= \varepsilon \boldsymbol{\lambda}^\top \mathbf{\Delta} = 0 \quad \text{a.s.} \end{aligned}$$

Thus μ_ε is another optimal measure. It is distinct from μ_* , because $\boldsymbol{\lambda} \neq 0$ and $s_j \notin S$, contradicting uniqueness. Therefore

$$\mathbb{P}\{\Delta_j > 0, j = 1, \dots, \ell\} > 0.$$

In particular, no Δ_j is identically zero. Each coordinate is therefore a nondegenerate Gaussian rv, so

$$\mathbb{P}\{\Delta_j = 0\} = 0, \quad j = 1, \dots, \ell.$$

Consequently, we obtain

$$\mathbb{P}\{\Delta_j \geq 0, j = 1, \dots, \ell\} = \mathbb{P}\{\Delta_j > 0, j = 1, \dots, \ell\} > 0,$$

as claimed. $\square$

Proof of Theorem 3.1. Lemma 4.2 yields

$$C := \mathbb{P}\{Z(s) > 0, s \in P\} > 0$$

and almost surely no passive value is zero. When $P = \emptyset$, these conditions are void and $C = 1$.

For each i , choose

$$\gamma_i \in (q_i/2, \min\{q_i, \beta_i\}), \quad N_i \in 2\mathbb{N}, \quad \gamma_i < \beta_i - 1/N_i.$$

On sufficiently small relative neighbourhoods U_i of t_i , the dominated-contact assumptions and Gaussian moment estimates give

$$m(t_i + h) - 1 \geq \lambda_i |h|^{q_i}, \quad \mathbb{E}\{|Z(t_i + h) - Z(t_i + h')|^{N_i}\} \leq C_{i,N_i} |h - h'|^{N_i \beta_i}.$$

After shrinking U_i , the Kolmogorov–Chentsov theorem and $Z(t_i) = 0$ therefore yield an almost surely finite rv K_i such that

$$|Z(t_i + h)| \leq K_i |h|^{\gamma_i}, \quad t_i + h \in U_i,$$

for the given continuous version.

For $\mathbf{x} \in (0, \infty)^k$ and $z \in C[a, b]$, put

$$H_u(\mathbf{x}, z) := \mathbf{1} \left\{ \inf_{t \in [a, b]} \left[u\{m(t) - 1\} + z(t) + \frac{\mathbf{a}(t)^\top \mathbf{x}}{u} \right] > 0 \right\}.$$

We claim that, for every fixed $\mathbf{x} \in (0, \infty)^k$ as $u \rightarrow \infty$ we have

$$H_u(\mathbf{x}, Z) \longrightarrow \mathbf{1}\{Z(s) > 0, s \in P\} \quad \text{a.s.}$$

Fix a residual path satisfying the preceding bounds and having no zero passive value. If $Z(s) < 0$ at some $s \in P$, evaluating at s , where $m(s) = 1$, shows that $H_u(\mathbf{x}, Z) = 0$ for all large u .

Suppose instead that all passive values are positive. Near each active contact, continuity and $\mathbf{a}(t_i) = \mathbf{e}_i$ give $\mathbf{a}(t)^\top \mathbf{x} \geq x_i/2$. Since $q_i/2 < \gamma_i < q_i$

$$\sup_{r \geq 0} \{K_i r^{\gamma_i} - \lambda_i u r^{q_i}\} = O\left(u^{-\gamma_i/(q_i - \gamma_i)}\right) = o(u^{-1}).$$

Thus the expression defining H_u is bounded below by $x_i/(2u) - o(u^{-1}) > 0$, uniformly on a sufficiently small relative neighbourhood of t_i . Near each passive contact, continuity makes Z bounded below by a positive constant; since $m \geq 1$ and $\mathbf{a}$ is bounded, the same expression is positive there for all large u . The complement of these relative open neighbourhoods is compact and disjoint from E . If nonempty, it has $\inf(m - 1) > 0$, while Z and $\mathbf{a}$ are bounded. Positivity follows there as well, proving the claim.

Using the conditional density f_u and its unnormalised version g_u from the standing notation, independence of Z and $\mathbf{X}_S$ gives

$$\mathbb{P}_u(B_u) = \frac{\int_{(0,\infty)^k} g_u(\mathbf{x}) \mathbb{E}H_u(\mathbf{x}, Z) d\mathbf{x}}{\int_{(0,\infty)^k} g_u(\mathbf{x}) d\mathbf{x}}.$$

The pointwise convergence above and $0 \leq H_u \leq 1$ imply $\mathbb{E}H_u(\mathbf{x}, Z) \rightarrow C$. Since

$$0 \leq g_u(\mathbf{x}) \leq e^{-\boldsymbol{\theta}^\top \mathbf{x}}, \quad g_u(\mathbf{x}) \longrightarrow e^{-\boldsymbol{\theta}^\top \mathbf{x}}, \quad u \rightarrow \infty$$

and every $\theta_i > 0$, dominated convergence yields $\lim_{u \rightarrow \infty} \mathbb{P}_u(B_u) = C$. Finally since $\mathbb{P}(B_u) = \mathbb{P}_u(B_u)\mathbb{P}(A_u)$ the claim follows. $\square$

Proof of Remark 3.3. For each i , choose an even integer $p_i \geq 2$ sufficiently large that

$$p_i \left(\beta_i - \frac{q_i}{2} \right) > d,$$

and then choose

$$\gamma_i \in \left(\frac{q_i}{2}, \min \left\{ q_i, \beta_i - \frac{d}{p_i} \right\} \right).$$

Gaussian moment estimates imply

$$\mathbb{E} \{ |Z(t_i + h) - Z(t_i + h')|^{p_i} \} \leq C_{i,p_i} \|h - h'\|^{p_i \beta_i}.$$

Writing $N(K, \|\cdot\|, \varepsilon)$ for the Euclidean covering number, every compact $K \subset \mathbb{R}^d$ satisfies

$$N(K, \|\cdot\|, \varepsilon) \leq C_K \varepsilon^{-d}, \quad 0 < \varepsilon \leq 1.$$

Therefore, the standard dyadic-net proof of the multidimensional Kolmogorov–Chentsov theorem yields an almost surely finite rv H_i such that

$$|Z(t_i + h)| \leq H_i \|h\|^{\gamma_i}$$

for all sufficiently small admissible h . This is the only dimension-dependent step. Since $q_i/2 < \gamma_i < q_i$, the remainder of the proof is unchanged, with relative Euclidean balls and norms replacing intervals and absolute values. $\square$

Lemma 4.3. Fix $\beta \in (0, 1]$. Let $D \subset [0, R]$ be a nonempty compact set containing 0, where $R \geq 0$. Let $G = \{G(x), x \in D\}$ be a centred GP with continuous sample paths, $G(0) = 0$, and suppose that for some $0 \leq A < \infty$

$$\mathbb{E}\{(G(x) - G(y))^2\} \leq A^2 |x - y|^{2\beta}, \quad x, y \in D.$$

Then there exists a constant $K_\beta < \infty$, depending only on β , such that uniformly in A , R , and D

$$\mathbb{E} \left\{ \sup_{x \in D} |G(x)| \right\} \leq K_\beta A R^\beta, \quad \sup_{x \in D} \text{Var } G(x) \leq A^2 R^{2\beta}.$$

Proof. If $A = 0$ or $R = 0$, the result is immediate from anchoring and continuity. Otherwise, the canonical metric satisfies

$$d_G(x, y) \leq A |x - y|^\beta, \quad N(D, d_G, \varepsilon) \leq 1 + R(A/\varepsilon)^{1/\beta}.$$

Since $G(0) = 0$, the variance bound follows by taking $y = 0$. Dudley's entropy bound and Gaussian symmetry give, with a universal constant K

$$\begin{aligned} \mathbb{E} \left\{ \sup_{x \in D} |G(x)| \right\} &\leq 2K \int_0^{AR^\beta} \sqrt{\log(1 + R(A/\varepsilon)^{1/\beta})} d\varepsilon \\ &= 2K A R^\beta \int_0^1 \sqrt{\log(1 + s^{-1/\beta})} ds \leq K_\beta A R^\beta. \end{aligned}$$

The integral is finite for every $\beta > 0$. $\square$

Lemma 4.4. *Under the assumptions of Theorem 3.6, there exists $r_0 > 0$ such that, with $r_u := u^{-1/\beta}$, we have*

$$\lim_{L \rightarrow \infty} \limsup_{u \rightarrow \infty} \mathbb{P} \left\{ \inf_{Lr_u \leq h \leq r_0} [u^2 \{m(b-h) - 1\} + uZ(b-h)] \leq 0 \right\} = 0.$$

Proof. By (21), after decreasing $r_0 > 0$ if necessary, we may assume that $r_0 \leq b-a$, that the local mean-square Hölder bound holds for $Z(b-h)$ whenever $0 \leq h \leq r_0$, and that

$$m(b-h) - 1 \geq \frac{\lambda}{2} h^{2\beta}, \quad 0 \leq h \leq r_0.$$

Split $[Lr_u, r_0]$ into the nonempty intervals

$$I_{n,u} := [Q_n r_u, 2Q_n r_u] \cap [0, r_0], \quad Q_n := 2^n L, \quad n \geq 0,$$

and put $R_{n,u} := \min\{2Q_n r_u, r_0\}$. By Lemma 4.3 and the increment bound

$$\mathbb{E} \left\{ \sup_{0 \leq h \leq R_{n,u}} |uZ(b-h)| \right\} \leq K_\beta \sqrt{C} u R_{n,u}^\beta \leq C_2 Q_n^\beta,$$

where C_2 is independent of u, n, L . On $I_{n,u}$, the deterministic term is at least $(\lambda/2)Q_n^{2\beta}$. Markov's inequality therefore gives

$$\begin{aligned} \mathbb{P} \left\{ \inf_{h \in I_{n,u}} [u^2 \{m(b-h) - 1\} + uZ(b-h)] \leq 0 \right\} \\ \leq \mathbb{P} \left\{ \sup_{0 \leq h \leq R_{n,u}} |uZ(b-h)| \geq \frac{\lambda}{2} Q_n^{2\beta} \right\} \leq \frac{2C_2}{\lambda} Q_n^{-\beta}. \end{aligned}$$

Summing over the intervals yields

$$\mathbb{P} \left\{ \inf_{Lr_u \leq h \leq r_0} [u^2 \{m(b-h) - 1\} + uZ(b-h)] \leq 0 \right\} \leq \frac{2C_2}{\lambda(1-2^{-\beta})} L^{-\beta},$$

which proves the claim. $\square$

Proof of Theorem 3.6. By the standing reduction, it remains to identify the limit of $\mathbb{P}_u(B_u)$. We use the one-point overshoot convergence stated above and put

$$r_u := u^{-1/\beta}, \quad T_u := \sup_{t \in [a,b]} \frac{-uZ(t) - u^2 \{m(t) - 1\}}{m(t)}.$$

Since $m \geq 1$ and $Z(b) = 0$, we have $T_u \geq 0$. Gaussian regression, compactness and continuity give the exact identity

$$B_u = \{S_u > T_u\}.$$

The rv T_u depends only on Z , so it is independent of S_u and its law is unchanged under $\mathbb{P}_u$.

We first verify that $M > -\infty$ almost surely. Since all processes in (22) are centred Gaussian, taking $r \downarrow 0$ in the endpoint increment bound gives for some $C < \infty$

$$\mathbb{E}\{(W(x) - W(y))^2\} \leq C|x - y|^{2\beta}, \quad x, y \geq 0.$$

Applying Lemma 4.3 on $[0, 2^{n+1}]$ and then Markov's inequality, we obtain a constant $c < \infty$ such that for every $n \geq 0$

$$\mathbb{P} \left\{ \sup_{0 \leq x \leq 2^{n+1}} |W(x)| > \frac{\lambda}{2} 2^{2\beta n} \right\} \leq c 2^{-\beta n}.$$

The right-hand side is summable. Hence, by Borel–Cantelli, almost surely, for all sufficiently large n and every $x \in [2^n, 2^{n+1}]$

$$W(x) + \lambda x^{2\beta} \geq \frac{\lambda}{2} 2^{2\beta n} > 0.$$

Continuity of W on compact intervals therefore gives $M > -\infty$. Since $W(0) = 0$, also $M \leq 0$.

For every fixed $L < \infty$, the expansion (21) gives, uniformly on $0 \leq x \leq L$

$$u^2\{m(b - r_u x) - 1\} \longrightarrow \lambda x^{2\beta}, \quad m(b - r_u x) \longrightarrow 1.$$

Indeed, the error in the first convergence is bounded by

$$L^{2\beta} \sup_{0 < h \leq r_u L} \left| \frac{m(b - h) - 1}{h^{2\beta}} - \lambda \right|.$$

Moreover, the finite-dimensional convergence (22) holds in $C[0, L]$. To see this, choose an even integer p with $p\beta > 1$. The increment bound and Gaussian moments imply, uniformly for all sufficiently large u

$$\mathbb{E}\{|uZ(b - r_u x) - uZ(b - r_u y)|^p\} \leq C_p |x - y|^{p\beta}, \quad 0 \leq x, y \leq L.$$

The Kolmogorov–Chentsov criterion therefore gives tightness. Setting

$$T_{u,L} := \sup_{0 \leq x \leq L} \frac{-uZ(b - r_u x) - u^2\{m(b - r_u x) - 1\}}{m(b - r_u x)},$$

$$D_L := \inf_{0 \leq x \leq L} \{W(x) + \lambda x^{2\beta}\},$$

the continuous mapping theorem gives $T_{u,L} \Rightarrow -D_L$.

Choose r_0 as in Lemma 4.4. Since $E = \{b\}$,

$$d_0 := \inf_{a \leq t \leq b - r_0} \{m(t) - 1\} > 0.$$

As $T_{u,L} \geq 0$, an outside point can change the supremum only if its numerator is positive. Hence, for all sufficiently large u

$$\mathbb{P}\{T_u \neq T_{u,L}\} \leq \mathbb{P}\left\{\inf_{Lr_u \leq h \leq r_0} [u^2\{m(b - h) - 1\} + uZ(b - h)] \leq 0\right\}$$

$$+ \mathbb{P}\left\{\sup_{t \in [a, b]} |Z(t)| \geq ud_0\right\}.$$

The last probability tends to zero by path continuity and compactness. Together with Lemma 4.4, this proves

$$\lim_{L \rightarrow \infty} \limsup_{u \rightarrow \infty} \mathbb{P}\{T_u \neq T_{u,L}\} = 0.$$

Since $D_L \downarrow M > -\infty$, we conclude that $T_u \Rightarrow -M$. Independence and the one-point overshoot limit therefore give, under $\mathbb{P}_u$

$$(S_u, T_u) \Rightarrow (\mathcal{E}, -M),$$

where $\mathcal{E}$ is independent of M and is exponential with rate $1/R(b, b)$. In particular, $\mathbb{P}\{\mathcal{E} = -M\} = 0$, and hence

$$\lim_{u \rightarrow \infty} \mathbb{P}_u(B_u) = \mathbb{P}\{\mathcal{E} > -M\} = \mathbb{E} \exp \left\{ \frac{M}{R(b, b)} \right\}.$$

This proves the assertion. □

Proof of Corollary 3.7. For every fixed $L < \infty$, the quadratic expansion gives uniformly for $0 \leq x \leq L$

$$u^2\{m(b - x/u) - 1\} \longrightarrow \lambda x^2, \quad u \rightarrow \infty.$$

We also have

$$\{uZ(b - x/u), 0 \leq x \leq L\} \Rightarrow \{\eta x, 0 \leq x \leq L\} \quad \text{in finite-dimensional distributions,} \quad u \rightarrow \infty.$$

Indeed, for every fixed $x \geq 0$

$$uZ(b - x/u) = x \frac{Z(b - x/u)}{x/u} \longrightarrow \eta x \quad \text{in } L^2,$$

with the value at $x = 0$ understood as zero. This gives convergence of the finite-dimensional distributions. Together with the assumed local mean-square Lipschitz bound, this verifies the tangent-critical assumptions. Thus b is a tangent-critical endpoint contact with

$$\beta = 1, \quad W(x) = \eta x, \quad x \geq 0.$$

Its variational term is

$$M_{\text{sm}} := \inf_{x \geq 0} \{\eta x + \lambda x^2\} = -\frac{(\max\{-\eta, 0\})^2}{4\lambda}.$$

The conclusion of Theorem 3.6 therefore applies. Writing $s := R(b, b)$, a standard Gaussian calculation gives also when $v = 0$

$$\mathbb{E} \left\{ \exp \left\{ -\frac{(\max\{-\eta, 0\})^2}{4\lambda s} \right\} \right\} = \frac{1}{2} + \frac{1}{2} \mathbb{E} \exp \left\{ -\frac{\eta^2}{4\lambda s} \right\} = \frac{1}{2} + \frac{1}{2} \left(1 + \frac{v^2}{2\lambda s} \right)^{-1/2}.$$

This proves the assertion. $\square$

APPENDIX A. AUXILIARY RESULTS

Let $\mathcal{T}$ be a nonempty compact metric space and let $X(t)$, $t \in \mathcal{T}$ be a centred real-valued GP with almost surely continuous sample paths and continuous covariance function R . For a finite signed measure η on $\mathcal{T}$, define the energy associated with the current covariance kernel R by

$$\mathcal{E}_R(\eta) := \iint_{\mathcal{T}^2} R(s, t) \eta(ds) \eta(dt) \geq 0.$$

For $\mu \in \mathcal{M}_1(\mathcal{T})$ define its potential and energy by

$$\phi_\mu(t) := \int_{\mathcal{T}} R(s, t) \mu(ds), \quad \sigma^2(\mu) := \mathcal{E}_R(\mu).$$

We call μ an optimal measure if

$$\sigma^2(\mu) = \inf_{\nu \in \mathcal{M}_1(\mathcal{T})} \sigma^2(\nu).$$

We use the following standard optimality criterion for minimum-energy probability measures; see [3, Theorem 3.1]. Related formulations appear in [4, Theorem 4.3 and Remark 4.5], [11, Theorem 2.1], and [12, Theorem 1.1].

Lemma A.1. *An optimal measure exists. A probability measure $\mu \in \mathcal{M}_1(\mathcal{T})$ is optimal if and only if*

$$\phi_\mu(t) \geq \sigma^2(\mu), \quad t \in \mathcal{T}.$$

For every optimal μ , equality holds on $\text{supp}(\mu)$. If further $\mathcal{E}_R(\eta) > 0$ for every nonzero finite signed measure η with $\eta(\mathcal{T}) = 0$, then the optimal measure is unique.

Corollary A.2. *Let $S = \{t_1, \dots, t_m\} \subset \mathcal{T}$, assume that $\Sigma = (R(t_i, t_j))_{i,j=1}^m$ is invertible and set*

$$w := \frac{\Sigma^{-1} \mathbf{1}}{\mathbf{1}^\top \Sigma^{-1} \mathbf{1}}, \quad c := \frac{1}{\mathbf{1}^\top \Sigma^{-1} \mathbf{1}}, \quad E := \left\{ t \in \mathcal{T} : \sum_{j=1}^m w_j R(t_j, t) = c \right\}.$$

If all components of w are positive and

$$\sum_{j=1}^m w_j R(t_j, t) \geq c, \quad t \in \mathcal{T},$$

then $\mu_ := \sum_{j=1}^m w_j \delta_{t_j}$ is an optimal measure with support exactly S . If further either $\mathcal{E}_R(\eta) > 0$ for every nonzero finite signed measure η on $\mathcal{T}$ with $\eta(\mathcal{T}) = 0$, or $E = S$, then μ_* is the unique optimal measure. Conversely, if an optimal measure has support exactly S , then it equals $\sum_{j=1}^m w_j \delta_{t_j}$, and w has positive components.*

Proof. Because Σ is an invertible covariance matrix, it is positive definite. Hence $c > 0$, $\mathbf{1}^\top w = 1$, $\Sigma w = c \mathbf{1}$, and $\sigma^2(\mu_*) = c$. The assumed potential inequality and Lemma A.1 give optimality; the strict-energy alternative gives uniqueness.

Suppose instead that $E = S$, and let ν be another optimal measure. Since $\phi_{\mu_*} \geq c$ and both measures have energy c

$$0 \leq \mathcal{E}_R(\nu - \mu_*) = 2c - 2 \int_{\mathcal{T}} \phi_{\mu_*} d\nu \leq 0$$

and hence $\int_{\mathcal{T}}(\phi_{\mu_*} - c) d\nu = 0$. Continuity and nonnegativity imply $\text{supp}(\nu) \subseteq E = S$. Writing $\nu = \sum_j v_j \delta_{t_j}$, we obtain

$$0 = \mathcal{E}_R(\nu - \mu_*) = (v - w)^\top \Sigma(v - w)$$

and positive definiteness gives $v = w$.

For the converse, write the optimal measure as $\mu = \sum_j v_j \delta_{t_j}$ and put $d := \sigma^2(\mu)$. Equality of its potential and energy on its support gives $\Sigma v = d\mathbf{1}$. Together with $\mathbf{1}^\top v = 1$, this yields $d = c$ and $v = w$. Exact support gives $v_j > 0$ for every j . $\square$

REFERENCES

- [1] A. Chakrabarty and G. Samorodnitsky, “Asymptotic behaviour of high Gaussian minima,” *Stochastic Processes and their Applications*, vol. 128, no. 7, pp. 2297–2324, 2018.
- [2] E. Hashorva, “Asymptotic properties of type I elliptical random vectors,” *Extremes*, vol. 10, pp. 175–206, 2007.
- [3] E. Hashorva and S. Novikov, “High minima of Gaussian processes: Overshoots and minimizer locations,” *arXiv preprint arXiv:2607.20714*, 2026.
- [4] R. J. Adler, E. Moldavskaya, and G. Samorodnitsky, “On the existence of paths between points in high level excursion sets of Gaussian random fields,” *The Annals of Probability*, vol. 42, no. 3, pp. 1020–1053, 2014.
- [5] Z. Wu, A. Chakrabarty, and G. Samorodnitsky, “High minima of non-smooth Gaussian processes,” *Electronic Communications in Probability*, vol. 24, pp. Paper No. 53, 1–12, 2019.
- [6] K. Dębicki, E. Hashorva, and L. Ji, “Parisian ruin over a finite-time horizon,” *Science China Mathematics*, vol. 59, no. 3, pp. 557–572, 2016.
- [7] K. Dębicki, E. Hashorva, and Z. Michna, “Distributional and extremal behaviour of Brownian motion with exponential resetting,” *Methodology and Computing in Applied Probability*, vol. 28, no. 3, p. 79, 2026.
- [8] W. V. Li and Q.-M. Shao, “Lower tail probabilities for Gaussian processes,” *The Annals of Probability*, vol. 32, no. 1A, pp. 216–242, 2004.
- [9] W. V. Li and W. Linde, “Existence of small ball constants for fractional brownian motions,” *C. R. Acad. Sci. Paris Sér. I Math.*, vol. 326, no. 11, pp. 1329–1334, 1998.
- [10] M. Lifshits, W. Linde, and Z. Shi, “Small deviations of Gaussian random fields in l_q -spaces,” *Electronic Journal of Probability*, vol. 11, pp. 1204–1233, 2006.
- [11] Z. Selk, “Large deviations for high minima of Gaussian processes with nonnegatively correlated increments,” *Statistics & Probability Letters*, vol. 206, p. 110001, 2024.
- [12] L. Pronzato and A. Zhigljavsky, “Minimum-energy measures for singular kernels,” *Journal of Computational and Applied Mathematics*, vol. 382, p. 113089, 2021.

KRZYSZTOF DĘBICKI, MATHEMATICAL INSTITUTE, UNIVERSITY OF WROCLAW, PL. GRUNWALDZKI 2/4, 50-384 WROCLAW, POLAND

Email address: Krzysztof.Debicki@math.uni.wroc.pl

ENKELEJD HASHORVA, DEPARTMENT OF ACTUARIAL SCIENCE, UNIVERSITY OF LAUSANNE, UNIL-DORIGNY, 1015 LAUSANNE, SWITZERLAND

Email address: Enkelejd.Hashorva@unil.ch

SVYATOSLAV NOVIKOV, DEPARTMENT OF ACTUARIAL SCIENCE, UNIVERSITY OF LAUSANNE, UNIL-DORIGNY, 1015 LAUSANNE, SWITZERLAND

Email address: Svyatoslav.Novikov@unil.ch