COMP3411 Tutorial - Week 9
ILP and Decision Trees

Question 1

a) Find the least general generalisation of the following terms:

```
f(g(a, b), [1, 2, [3, 4], 5], 1 + 2 * 3)
f(g(a, h(x, y)), [1, 2, [3, 4, 5]], 1 + 6)
```

To find the LGG, look for where the terms differ and replace with a variable. Reuse the same variable if the same terms appear in corresponding positions.

```
lgg = f(g(a, X), [1, 2, [3, 4 | Y] | Y], 1 + Z)
subst = [X/{b, h(x, y)}, Y/{[], [5]}, Z/{2*3, 6}]
```

The list is tricky because the inverse substitution Y/{[], [5]} *applies also as* Y/{[5], []} .

b) Find the least general generalisation of the following clauses:

```
q(f(a)) :- p(a, b), r(b, c), r(b, e).
q(f(x)) :- p(x, y), r(y, z), r(w, z).
```

To find the LGG of two clauses, match the head, then find all combinations of possible LGGs between literals that have consistent inverse substitutions.

```
q(f(X)) :- p(X, Y), r(Y, Z), r(W, Z), R(Y, E), r(W, E)
subst = [X/{a, x}, Y/{b, y}, Z/{c, z}, W/{b, w}, E/{e, z}]
```

Question 2

Consider the task of predicting whether children are likely to be hired to play members of the Von Trapp Family in a production of The Sound of Music, based on these data:

height	hair	eyes	hired
short	blond	blue	+
tall	red	blue	+
tall	blond	blue	+
tall	blond	brown	–
short	dark	blue	–
tall	dark	blue	–
tall	dark	brown	–
short	blond	brown	–

a. Compute the information (entropy) gain for each of the three attributes (height, hair, eyes) in terms of classifying objects as belonging to the class, + or – .

There are 3 objects in class '+' and 5 in '-', so the entropy is:

Entropy(parent) = Σ_i P_i $\log_2 P_i$ = -(3/8)log(3/8) - (5/8)log(5/8) = 0.954

Suppose we split on height:

[3,5]

[1,2]   [2,3]

Of the 3 'short' items, 1 is '+' and 2 are '-', so Entropy(short) = -(1/3)log(1/3) - (2/3)log(2/3) = 0.918

Of the 5 'tall' items, 2 are '+' and 3 are '-', so Entropy(tall) = -(2/5)log(2/5) - (3/5)log(3/5) = 0.971

The average entropy after splitting on 'height' is Entropy(height) = (3/8)(0.918) + (5/8)(0.971) = 0.951

The information gained by testing this attribute is: 0.954 - 0.951 = 0.003 (i.e. very little)

If we try splitting on 'hair' we find that the branch for 'dark' has 3 items, all '-' and the branch for 'red' has 1 item, in '+'. Thus, these branches require no further information to make a decision. The branch for 'blond' has 2 '+' and 2 '-' items and so requires 1 bit. That is,

Entropy(hair) = (3/8)(0) + (1/8)(0) + (4/8)(1) = 0.5

and the information gained by testing hair is 0.954 - 0.5 = 0.454 bits.

By a similar calculation, the entropy for testing 'eyes' is (5/8)(0.971) + (3/8)(0) = 0.607, so the information gained is 0.954 - 0.607 = 0.347 bits.

Thus 'hair' gives us the maximum information gain.

b. Construct a decision tree based on the minimum entropy principle.

Since the 'blond' branch for hair still contains a mixed population, we need to apply the procedure recursively to these four items. Note that we now only need to test 'height' and 'eyes' since the 'hair' attribute has already been used. If we split on 'height', the branch for 'tall' and 'short' will each contain one '+' and one '-', so the entropy gain is zero. If we split on 'eyes', the 'blue' brach contains two '+'s and the 'brown' branch two '-'s, so the tree is complete:

Question 3

The Laplace error estimate for pruning a node in a Decision Tree is given by:

$$E = 1 - \frac{n+1}{N+k}$$

where N is the total number of items, n is the number of items in the majority class and k is the number of classes. Given the following sub-tree, should the children be pruned or not? Show your calculations.

[4,7]

[2,1]   [2,6]

[4,7]

[2,1]   [2,6]

Error(Parent) = 1 - (7+1)/(11+2) = 1 - 8/13 = 5/13 = 0.385

Error(Left)   = 1 - (2+1)/(3+2) = 1 - 3/5 = 2/5 = 0.4

Error(Right) = 1 - (6+1)/(8+2) = 1 - 7/10 = 3/10 = 0.3

Backed Up Error = (3/11)*(0.4) + (8/11)*(0.3) = 0.327 < 0.385

Since Error of Parent is larger than Backed Up Error ⟹ Don't Prune

Question 4

Construct a Decision Tree for the following set of examples.

Day	Outlook	Temperature	Humidity	Wind	PlayTennis
D1	Sunny	Hot	High	Weak	No
D2	Sunny	Hot	High	Strong	No
D3	Overcast	Hot	High	Weak	Yes
D4	Rain	Mild	High	Weak	Yes
D5	Rain	Cool	Normal	Weak	Yes
D6	Rain	Cool	Normal	Strong	No
D7	Overcast	Cool	Normal	Strong	Yes
D8	Sunny	Mild	High	Weak	No
D9	Sunny	Cool	Normal	Weak	Yes
D10	Rain	Mild	Normal	Weak	Yes
D11	Sunny	Mild	Normal	Strong	Yes
D12	Overcast	Mild	High	Strong	Yes
D13	Overcast	Hot	Normal	Weak	Yes
D14	Rain	Mild	High	Strong	No

What class is assigned to the instance {D15, Sunny, Hot, High, Weak}?

1.

(ii) $Values(Temperature) = \{hot, mild, cool\}$

$S = [9+, 5-]$

$S_{hot} \leftarrow [2+, 2-]$

$S_{mild} \leftarrow [4+, 2-]$

$S_{cool} \leftarrow [3+, 1-]$

$Gain(S, Temperature) = Entropy(S) - \sum_{v=\{hot, mild, cool\}} \frac{|S_v|}{|S|} Entropy(S_v)$

$= Entropy(S) - \frac{4}{14} Entropy(S_{hot}) - \frac{6}{14} Entropy(S_{mild}) - \frac{4}{14} Entropy(S_{cool})$

$= 0.940 - \frac{4}{14} \times 1.00 - \frac{6}{14} \times 0.918 - \frac{4}{14} \times 0.232$

$= 0.029$

$Entropy(S_{hot}) = Entropy([2+, 2-]) = -\frac{2}{4}log_2\frac{2}{4} - \frac{2}{4}log_2\frac{2}{4} = 1.00$

$Entropy(S_{mild}) = Entropy([4+, 2-]) = -\frac{4}{6}log_2\frac{4}{6} - \frac{2}{6}log_2\frac{2}{6} = 0.918$

$Entropy(S_{cool}) = Entropy([3+, 1-]) = -\frac{3}{4}log_2\frac{3}{4} - \frac{1}{4}log_2\frac{1}{4} = 0.232$

(iii) $Values(Humidity) = \{high, normal\}$

$S = [9+, 5-]$

$S_{high} \leftarrow [3+, 4-]$

$S_{normal} \leftarrow [6+, 1-]$

$Gain(S, Humidity) = Entropy(S) - \sum_{v=\{high, normal\}} \frac{|S_v|}{|S|} Entropy(S_v)$

$= Entropy(S) - \frac{7}{14} Entropy(S_{high}) - \frac{7}{14} Entropy(S_{normal})$

$= 0.940 - \frac{7}{14} \times 0.985 - \frac{7}{14} \times 0.592$

$= 0.152$

$Entropy(S_{high}) = Entropy([3+, 4-]) = -\frac{3}{7}log_2\frac{3}{7} - \frac{4}{7}log_2\frac{4}{7} = 0.985$

$Entropy(S_{mild}) = Entropy([6+, 1-]) = -\frac{6}{7}log_2\frac{6}{7} - \frac{1}{7}log_2\frac{1}{7} = 0.592$

(iv) $Values(Wind) = \{weak, strong\}$

$S = [9+, 5-]$

$S_{weak} \leftarrow [6+, 2-]$

$S_{strong} \leftarrow [3+, 3-]$

$Gain(S, Wind) = Entropy(S) - \sum_{v=\{weak, strong\}} \frac{|S_v|}{|S|} Entropy(S_v)$

$= Entropy(S) - \frac{8}{14} Entropy(S_{weak}) - \frac{6}{14} Entropy(S_{strong})$

$= 0.940 - \frac{8}{14} \times 0.811 - \frac{6}{14} \times 1.00$

$= 0.048$

$Entropy(S_{weak}) = Entropy([6+, 2-]) = -\frac{6}{8}log_2\frac{6}{8} - \frac{6}{8}log_2\frac{6}{8} = 0.811$

$Entropy(S_{strong}) = Entropy([3+, 3-]) = -\frac{3}{6}log_2\frac{3}{6} - \frac{3}{6}log_2\frac{3}{6} = 1.00$

2.

(i) $Values(Temperature) = \{hot, mild, cool\}$

$S_{sunny} = [2+, 3-]$

$S_{sunny,hot} \leftarrow [0+, 2-]$

$S_{sunny,mild} \leftarrow [1+, 1-]$

$S_{sunny,cool} \leftarrow [1+, 0-]$

$Gain(S_{sunny}, Temperature) = Entropy(S_{sunny}) - \sum_{v=\{hot,mild,cool\}} \frac{|S_{sunny,v}|}{|S_{sunny}|} Entropy(S_{sunny,v})$

$= Entropy(S) - \frac{2}{5} Entropy(S_{sunny,hot}) - \frac{2}{5} Entropy(S_{sunny,mild}) - \frac{1}{5} Entropy(S_{sunny,cool})$

$= 0.971 - \frac{2}{5} \times 0.00 - \frac{2}{5} \times 1.00 - \frac{1}{5} \times 0.00$

$= 0.571$

$Entropy(S_{sunny}) = Entropy([2+, 3-]) = -\frac{2}{5} log_2 \frac{2}{5} - \frac{3}{5} log_2 \frac{3}{5} = 0.971$

$Entropy(S_{sunny,hot}) = Entropy([0+, 2-]) = -\frac{0}{2} log_2 \frac{0}{2} - \frac{2}{2} log_2 \frac{2}{2} = 0.00$

$Entropy(S_{sunny,mild}) = Entropy([1+, 1-]) = -\frac{1}{2} log_2 \frac{1}{2} - \frac{1}{2} log_2 \frac{1}{2} = 1.00$

$Entropy(S_{sunny,cool}) = Entropy([1+, 0-]) = -\frac{1}{1} log_2 \frac{1}{1} - \frac{0}{1} log_2 \frac{0}{1} = 0.00$

(ii) $Values(Humidity) = \{high, normal\}$

$S_{sunny} = [2+, 3-]$

$S_{sunny,high} \leftarrow [0+, 3-]$

$S_{sunny,normal} \leftarrow [2+, 0-]$

$Gain(S, Humidity) = Entropy(S_{sunny}) - \sum_{v=\{high,normal\}} \frac{|S_{sunny,v}|}{|S_{sunny}|} Entropy(S_{sunny,v})$

$= Entropy(S_{sunny}) - \frac{3}{5} Entropy(S_{sunny,high}) - \frac{2}{5} Entropy(S_{sunny,normal})$

$= 0.971 - \frac{3}{5} \times 0.00 - \frac{2}{5} \times 0.00$

$= 0.971$

$Entropy(S_{sunny,high}) = Entropy([0+, 3-]) = -\frac{0}{3} log_2 \frac{0}{3} - \frac{3}{3} log_2 \frac{3}{3} = 0.00$

$Entropy(S_{sunny,mild}) = Entropy([2+, 0-]) = -\frac{2}{2} log_2 \frac{2}{2} - \frac{0}{2} log_2 \frac{0}{2} = 0.00$

(iii) $Values(Wind) = \{weak, strong\}$

$S_{sunny} = [2+, 3-]$

$S_{weak} \leftarrow [1+, 2-]$

$S_{strong} \leftarrow [1+, 1-]$

$Gain(S, Wind) = Entropy(S_{sunny}) - \sum_{v=\{weak,strong\}} \frac{|S_{sunny,v}|}{|S_{sunny}|} Entropy(S_{sunny,v})$

$= Entropy(S) - \frac{3}{5} Entropy(S_{sunny,weak}) - \frac{2}{5} Entropy(S_{strong})$

$= 0.971 - \frac{3}{5} \times 0.918 - \frac{2}{5} \times 1.00$

$= 0.020$

$Entropy(S_{weak}) = Entropy([1+, 2-]) = -\frac{1}{3} log_2 \frac{1}{3} - \frac{1}{3} log_2 \frac{1}{3} = 0.918$

$Entropy(S_{strong}) = Entropy([1+, 1-]) = -\frac{1}{2} log_2 \frac{1}{2} - \frac{1}{2} log_2 \frac{1}{2} = 1.00$

3.

(i) $Values(Temperature) = \{hot, mild, cool\}$

$S_{rain} = [2+, 3-]$

$S_{rain,hot} \leftarrow [0+, 0-]$

$S_{rain,mild} \leftarrow [2+, 1-]$

$S_{rain,cool} \leftarrow [1+, 1-]$

$Gain(S_{rain}, Temperature) = Entropy(S_{rain}) - \sum_{v=\{hot,mild,cool\}} \frac{|S_{rain,v}|}{|S_{rain}|} Entropy(S_{rain,v})$

$= Entropy(S) - \frac{0}{5} Entropy(S_{rain,hot}) - \frac{3}{5} Entropy(S_{rain,mild}) - \frac{2}{5} Entropy(S_{rain,cool})$

$= 0.971 - \frac{0}{5} \times 0.00 - \frac{3}{5} \times 0.918 - \frac{2}{5} \times 1.00$

$= 0.020$

$Entropy(S_{rain}) = Entropy([3+, 2-]) = -\frac{3}{5} log_2 \frac{3}{5} - \frac{2}{5} log_2 \frac{2}{5} = 0.971$

$Entropy(S_{rain,hot}) = Entropy([0+, 0-]) = -\frac{0}{0} log_2 \frac{0}{0} - \frac{0}{0} log_2 \frac{0}{0} = 0.00$

$Entropy(S_{rain,mild}) = Entropy([2+, 1-]) = -\frac{2}{3} log_2 \frac{2}{3} - \frac{1}{3} log_2 \frac{1}{3} = 0.918$

$Entropy(S_{rain,cool}) = Entropy([1+, 1-]) = -\frac{1}{2} log_2 \frac{1}{2} - \frac{1}{2} log_2 \frac{1}{2} = 1.00$