


Bound Entanglement

Arne Hansen

ETH Zurich

02/2014

Entanglement



... peculiar,
non-classical correlation
arising from the tensor
product structure in
combined quantum
systems

Bound Entanglement

Bound Entanglement

non-distillable entanglement

Bound Entanglement

Bound Entanglement

non-distillable entanglement

Existence Explicite example of a bound entangled state

Bound Entanglement

Bound Entanglement

non-distillable entanglement

Existence Explicite example of a bound entangled state

Use Why should we be concerned about bound entanglement?

Bound Entanglement

Bound Entanglement

non-distillable entanglement

Existence Explicite example of a bound entangled state

Use Why should we be concerned about bound entanglement?

Detection How do we decide for a given state whether it is bound entangled?

1 Existence Bound Entanglement

- The PPT Criterion
- The range criterion
- First BE state
- Distillable states
- Summary

2 Applications of BE

3 Testing bound entanglement

4 Notions of Quantum Repeaters

5 Question

The PPT Criterion

(Peres Criterion)

Definition

The partial transpose of a density matrix $\rho \in \mathcal{D}(\mathcal{H}_A \otimes \mathcal{H}_B)$ is defined as

$$\rho^\Gamma := (\mathcal{I}_A \otimes T_B)\rho \quad (1)$$

The PPT Criterion

(Peres Criterion)

Definition

The partial transpose of a density matrix $\rho \in \mathcal{D}(\mathcal{H}_A \otimes \mathcal{H}_B)$ is defined as

$$\rho^\Gamma := (\mathcal{I}_A \otimes T_B)\rho \quad (1)$$

Matrix elements in the tensor basis


In the product basis

$$(\rho^\Gamma)_{ikjl} = \langle i \otimes k | \rho^\Gamma | j \otimes l \rangle \quad (2)$$

$$= \langle i \otimes l | \rho | j \otimes k \rangle = \rho_{iljk} \quad (3)$$


The PPT Criterion

(Peres Criterion)



The PPT Criterion

(Peres Criterion)




Entanglement criterion

$$\rho_{AB} \in \mathcal{S} \subset \mathcal{H}_A \otimes \mathcal{H}_B \Rightarrow \rho_{AB} \in \Gamma \quad (1)$$

The PPT Criterion

(Peres Criterion)




Entanglement criterion

$$\rho_{AB} \in \mathcal{S} \subset \mathcal{H}_A \otimes \mathcal{H}_B \Rightarrow \rho_{AB} \in \Gamma \quad (1)$$

The PPT Criterion

(Peres Criterion)




Entanglement criterion

$$\rho_{AB} \notin \Gamma \Rightarrow \rho_{AB} \text{ is entangled} \quad (1)$$

The PPT Criterion

(Peres Criterion)



Entanglement criterion


$$\rho_{AB} \notin \Gamma \Rightarrow \rho_{AB} \text{ is entangled} \quad (1)$$

Accuracy

The criterion is accurate in $2 \otimes 2$ and $2 \otimes 3$.
But not in higher dimensions. [HHH96]

The PPT Criterion

(Peres Criterion)



Group of Entanglement Tests

Any Positive but not Completely Positive Map ('channel') serves as a similar entanglement test.

The Range Criterion [Hor97]

Decomposition of separable states

$\forall \rho \in \mathcal{S} :$

$$\rho = \sum_{i=1}^{n_A} \sum_{j=1}^{n_B} p_{ij} |\phi_i\rangle\langle\phi_i| \otimes |\psi_j\rangle\langle\psi_j| \quad (1)$$

$$|\phi_i\rangle \in \mathcal{H}_A, |\psi_j\rangle \in \mathcal{H}_B$$

$$n_A < \dim \mathcal{H}_A, n_B < \dim \mathcal{H}_B$$

The Range Criterion [Hor97]

Decomposition of separable states

$\forall \rho \in \mathcal{S} :$

$$\rho = \sum_{i=1}^{n_A} \sum_{j=1}^{n_B} p_{ij} |\phi_i\rangle\langle\phi_i| \otimes |\psi_j\rangle\langle\psi_j| \quad (1)$$

$$|\phi_i\rangle \in \mathcal{H}_A, |\psi_j\rangle \in \mathcal{H}_B$$

$$n_A < \dim \mathcal{H}_A, n_B < \dim \mathcal{H}_B$$

→ features of convex sets, Caratheodories Thm [Car11]

The Range Criterion [Hor97]

Decomposition of separable states

$\forall \rho \in \mathcal{S} :$

$$\rho = \sum_{i=1}^{n_A} \sum_{j=1}^{n_B} p_{ij} |\phi_i\rangle\langle\phi_i| \otimes |\psi_j\rangle\langle\psi_j| \quad (1)$$

$$|\phi_i\rangle \in \mathcal{H}_A, |\psi_j\rangle \in \mathcal{H}_B$$

$$n_a < \dim \mathcal{H}_A, n_B < \dim \mathcal{H}_B$$

The range of an operator

$$\text{Ran } \rho := \{|\psi\rangle \in \mathcal{H}, \exists |\phi\rangle \in \mathcal{H} : |\psi\rangle = \rho|\phi\rangle\} \quad (2)$$

The Range Criterion [Hor97]

The range of separable density matrices

Any separable density matrix $\rho \in \mathcal{S} \subset \mathcal{H}_A \otimes \mathcal{H}_B$ can be written as a convex combination of pure products states

$$\begin{aligned}\rho &= \sum_{i,j} p_{ij} |\psi_i\rangle\langle\psi_i| \otimes |\phi_j\rangle\langle\phi_j| = \sum_{i,j} p_{ij} |\psi_i \otimes \phi_j\rangle\langle\psi_i \otimes \phi_j| \\ \rho^\Gamma &= (\mathcal{I} \otimes T)\rho = \sum_{i,j} |\psi_i\rangle\langle\psi_i| \otimes \underbrace{|\phi_j\rangle\langle\phi_j|^T}_{|\phi_j^*\rangle\langle\phi_j^*|} \\ &= \sum_{ij} p_{ij} |\psi_i \otimes \phi_j^*\rangle\langle\psi_i \otimes \phi_j^*|\end{aligned}$$

and the set of product states $\{|\psi_i \otimes \phi_j\rangle\}_{ij}$ ($\{|\psi_i \otimes \phi_j^*\rangle\}_{ij}$) span the range of ρ (ρ^Γ)

The Range Criterion [Hor97]

Entanglement test

- find product vectors that span the range of ρ
- show that these vectors do **not** span the range of ρ^Γ

The first bound entangled state

$$\rho_a = \frac{1}{8a+1} \begin{bmatrix} a & 0 & 0 & 0 & a & 0 & 0 & 0 & a \\ 0 & a & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & a & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & a & 0 & 0 & 0 & 0 & 0 \\ a & 0 & 0 & 0 & a & 0 & 0 & 0 & a \\ 0 & 0 & 0 & 0 & 0 & a & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{1+a}{2} & 0 & \frac{\sqrt{1-a^2}}{2} \\ 0 & 0 & 0 & 0 & 0 & 0 & 0 & a & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{\sqrt{1-a^2}}{2} & 0 & \frac{1+a}{2} \end{bmatrix} \quad a \in (0, 1)$$

The first bound entangled state

$$\rho_a = \frac{1}{8a+1} \begin{bmatrix} a & 0 & 0 & 0 & a & 0 & 0 & 0 & a \\ 0 & a & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & a & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & a & 0 & 0 & 0 & 0 & 0 \\ a & 0 & 0 & 0 & a & 0 & 0 & 0 & a \\ 0 & 0 & 0 & 0 & 0 & a & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{1+a}{2} & 0 & \frac{\sqrt{1-a^2}}{2} \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{2}{0} & a & \frac{2}{0} \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{\sqrt{1-a^2}}{2} & 0 & \frac{1+a}{2} \end{bmatrix} \quad a \in (0, 1)$$

The first bound entangled state

$$\rho_a = \frac{1}{8a+1} \begin{bmatrix} a & 0 & 0 & 0 & a & 0 & 0 & 0 & a \\ 0 & a & 0 & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & a & 0 & 0 & 0 & 0 & 0 & 0 \\ 0 & 0 & 0 & a & 0 & 0 & 0 & 0 & 0 \\ a & 0 & 0 & 0 & a & 0 & 0 & 0 & a \\ 0 & 0 & 0 & 0 & 0 & a & 0 & 0 & 0 \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{1+a}{2} & 0 & \frac{\sqrt{1-a^2}}{2} \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{2}{0} & a & \frac{2}{0} \\ 0 & 0 & 0 & 0 & 0 & 0 & \frac{\sqrt{1-a^2}}{2} & 0 & \frac{1+a}{2} \end{bmatrix} \quad a \in (0, 1)$$

Properties of ρ_a

- PPT
- ρ_a and ρ_a^Γ violate the range criterion

PPT and Distillability

Definition

A density matrix ρ is called distillable if

$$\exists \text{ LOCC } \Lambda : \mathcal{D}((\mathcal{H}_A \otimes \mathcal{H}_B)^{\otimes n}) \rightarrow \mathcal{D}(\mathbb{C}^2) \quad \text{s.t. :} \\ \Lambda(\rho^{\otimes n}) = |\Phi^+\rangle\langle\Phi^+| \quad \text{for some large } n$$

Distillation of PPT states

Cannot distill qubits from PPT density matrices ρ and $\rho^{\otimes N}$

Non-distillable entanglement

Any PPT entangled state is not distillable.

Distillation of PPT states

Distilling $\rho^{\otimes n}$

Assume ρ is distillable:

$$\Lambda(\rho^{\otimes n}) = \frac{1}{M} \sum_i (A_i \otimes B_i) \rho^{\otimes n} (A_i^\dagger \otimes B_i^\dagger) \quad \text{is entangled}$$

$$\Rightarrow \quad \exists \, i_0 \text{ s.t.: } \underbrace{(A_{i_0} \otimes B_{i_0}) \rho^{\otimes n} (A_{i_0}^\dagger \otimes B_{i_0}^\dagger)}_{=:\rho_{i_0}} \quad \text{is entangled}$$

Distillation of PPT states

Distilling $\rho^{\otimes n}$

Assume ρ is distillable:

$$\Lambda(\rho^{\otimes n}) = \frac{1}{M} \sum_i (A_i \otimes B_i) \rho^{\otimes n} (A_i^\dagger \otimes B_i^\dagger) \quad \text{is entangled}$$

$$\Rightarrow \exists \textcolor{red}{i}_0 \text{ s.t.: } \underbrace{(A_{\textcolor{red}{i}_0} \otimes B_{\textcolor{red}{i}_0}) \rho^{\otimes n} (A_{\textcolor{red}{i}_0}^\dagger \otimes B_{\textcolor{red}{i}_0}^\dagger)}_{=:\rho_{\textcolor{red}{i}_0}} \quad \text{is entangled}$$

$$A_{i_0} = |0\rangle\langle\psi_A| + |1\rangle\langle\phi_A| \quad B_{i_0} = |0\rangle\langle\psi_B| + |1\rangle\langle\phi_B|$$

$$P_A(P_B) := \text{projector onto } \langle\psi_A, \phi_A\rangle(\langle\psi_B, \phi_B\rangle)$$

Distillation of PPT states

Distilling $\rho^{\otimes n}$

Assume ρ is distillable:

$$\Lambda(\rho^{\otimes n}) = \frac{1}{M} \sum_i (A_i \otimes B_i) \rho^{\otimes n} (A_i^\dagger \otimes B_i^\dagger) \quad \text{is entangled}$$

$$\Rightarrow \exists i_0 \text{ s.t.: } \underbrace{(A_{i_0} \otimes B_{i_0}) \rho^{\otimes n} (A_{i_0}^\dagger \otimes B_{i_0}^\dagger)}_{=:\rho_{i_0}} \quad \text{is entangled}$$

$$\Rightarrow \rho' := P_A \otimes P_B \rho^{\otimes n} P_A \otimes P_B \quad \text{is entangled}$$

Distillation of PPT states

Distilling $\rho^{\otimes n}$

Assume ρ is distillable:

$$\Lambda(\rho^{\otimes n}) = \frac{1}{M} \sum_i (A_i \otimes B_i) \rho^{\otimes n} (A_i^\dagger \otimes B_i^\dagger) \quad \text{is entangled}$$

$$\Rightarrow \exists i_0 \text{ s.t.: } \underbrace{(A_{i_0} \otimes B_{i_0}) \rho^{\otimes n} (A_{i_0}^\dagger \otimes B_{i_0}^\dagger)}_{=:\rho_{i_0}} \quad \text{is entangled}$$

$$\Rightarrow \rho' := P_A \otimes P_B \rho^{\otimes n} P_A \otimes P_B \quad \text{is NPT}$$

Distillation of PPT states

Distilling $\rho^{\otimes n}$

Assume ρ is distillable:

$$\Lambda(\rho^{\otimes n}) = \frac{1}{M} \sum_i (A_i \otimes B_i) \rho^{\otimes n} (A_i^\dagger \otimes B_i^\dagger) \quad \text{is entangled}$$

$$\Rightarrow \exists i_0 \text{ s.t.: } \underbrace{(A_{i_0} \otimes B_{i_0}) \rho^{\otimes n} (A_{i_0}^\dagger \otimes B_{i_0}^\dagger)}_{=:\rho_{i_0}} \quad \text{is entangled}$$

$$\Rightarrow \rho' := P_A \otimes P_B \rho^{\otimes n} P_A \otimes P_B \quad \text{is NPT}$$

$$\Rightarrow \exists \psi \text{ s.t.: } \langle \psi | (\rho')^{\Gamma_B} | \psi \rangle < 0$$

Distillation of PPT states

Distilling $\rho^{\otimes n}$

Assume ρ is distillable:

$$\Lambda(\rho^{\otimes n}) = \frac{1}{M} \sum_i (A_i \otimes B_i) \rho^{\otimes n} (A_i^\dagger \otimes B_i^\dagger) \quad \text{is entangled}$$

$$\Rightarrow \exists i_0 \text{ s.t.: } \underbrace{(A_{i_0} \otimes B_{i_0}) \rho^{\otimes n} (A_{i_0}^\dagger \otimes B_{i_0}^\dagger)}_{=: \rho_{i_0}} \quad \text{is entangled}$$

$$\Rightarrow \rho' := P_A \otimes P_B \rho^{\otimes n} P_A \otimes P_B \quad \text{is NPT}$$

$$\Rightarrow \exists \psi \text{ s.t.: } \langle \psi | (\rho')^{\Gamma_B} | \psi \rangle < 0$$

$$\Rightarrow \langle \psi | \rho^{\Gamma_B} | \psi \rangle < 0$$

$$\Rightarrow \rho \text{ is NPT}$$

Bound entanglement

a first summary

def entanglement that cannot be distilled \rightarrow not a sufficient resource for many quantum informational protocols

Bound entanglement

a first summary

def entanglement that cannot be distilled \rightarrow not a sufficient resource for many quantum informational protocols

construction PPT entangled states (any PPT is not distillable)

Bound entanglement

a first summary

def entanglement that cannot be distilled \rightarrow not a sufficient resource for many quantum informational protocols

construction PPT entangled states (any PPT is not distillable)

existence only in higher dimensions (smallest BE systems in $3 \otimes 3$ and $4 \otimes 2$)

Bound entanglement

a first summary

def entanglement that cannot be distilled \rightarrow not a sufficient resource for many quantum informational protocols

construction PPT entangled states (any PPT is not distillable)

existence only in higher dimensions (smallest BE systems in $3 \otimes 3$ and $4 \otimes 2$)

detection need a stronger criterion than PPT, e.g. range criterion

Quantum Key Distribution

with bound entangled resources

Bound entangled resources can be employed to distill secret key [HHHO05].

Deciding entanglement for PPT


PPT all NPT states are entangled

range criterion compare the range of ρ and ρ^Γ

extendibility check extendibility of ρ

- 1 Existence Bound Entanglement
- 2 Applications of BE
- 3 Testing bound entanglement
- 4 Notions of Quantum Repeaters
 - Swap Operation on States
 - Composition of Channels
 - Hybrid Approach
- 5 Question


Endeavour: Extending entanglement



Endeavour: Extending entanglement




Endeavour: Extending entanglement




Swap Operator on States




Swap Operator on States



Swap Operator on States



Swap Operator on States




Swap operation

$$\text{swap}(\rho_A, \rho_B) = \frac{1}{N} \text{Tr}_{BB'} \left[(|\Phi^+\rangle\langle\Phi^+|_{BB'} \otimes \mathbb{1}_{AC}) (\rho_1 \otimes \rho_2) \right]$$

Composing Channels

corresponds to swapping CJ associated states




Composing Channels

corresponds to swapping CJ associated states




Composing Channels

corresponds to swapping CJ associated states




Composing Channels

corresponds to swapping CJ associated states




Composing Channels

corresponds to swapping CJ associated states



Composing Channels

corresponds to swapping CJ associated states




Swap in terms of channels

$$\tau^{-1}(\text{swap}(\rho_1, \rho_2)) = \tau^{-1}(\rho_2) \circ \tau^{-1}(\rho_1)$$

(flipping systems or equivalently transposing channels if necessary, up to normalization)

Hybrid approach



Hybrid approach



Hybrid approach



Hybrid approach



Hybrid formulation

$$\text{swap}(\rho_1, \rho_2) = [\mathcal{I}_A \otimes \tau^{-1}(\rho_2)] (\rho_1) = [\mathcal{I}_C \otimes \tau^{-1}(\rho_1)^T] (\rho_2)$$

(flipping systems or equivalently transposing channels if necessary, up to normalization)

Swapped Bound Entanglement

Entanglement of swapped states from BE resources?

States: Image of the Swap operator

$$\text{swap} : \text{BE states} \times \text{BE states} \rightarrow \mathcal{S}$$

OR

$$\text{swap}(\text{BE states}, \text{BE states}) \cap \mathcal{D} \setminus \mathcal{S} \neq \emptyset$$

¹Defined by their image: $\Lambda_{\Gamma}(\mathcal{D}) \subset \Gamma$

Swapped Bound Entanglement

Entanglement of swapped states from BE resources?

States: Image of the Swap operator

$$\text{swap} : \text{BE states} \times \text{BE states} \rightarrow \mathcal{S}$$

OR

$$\text{swap}(\text{BE states}, \text{BE states}) \cap \mathcal{D} \setminus \mathcal{S} \neq \emptyset$$

Composed Channels

If $\Lambda_1 : \mathcal{D}(A) \rightarrow \mathcal{D}(B)$ and $\Lambda_2 : \mathcal{D}(B) \rightarrow \mathcal{D}(C)$ are PPT channels¹,
is $\Lambda_2 \circ \Lambda_1 : \mathcal{D}(A) \rightarrow \mathcal{D}(C)$ separable?

¹Defined by their image: $\Lambda_\Gamma(\mathcal{D}) \subset \Gamma$

Bibliography I

C. Carathéodory.

Über den variabilitätsbereich der fourier'schen konstanten von positiven harmonischen funktionen.

Rendiconti del Circolo Matematico di Palermo, 32:193–217, 1911.

Michał Horodecki, Paweł Horodecki, and Ryszard Horodecki.

Separability of mixed states: necessary and sufficient conditions.

Physics Letters A, 223(1–2):1 – 8, 1996.

Karol Horodecki, Michał Horodecki, Paweł Horodecki, and Jonathan Oppenheim.

Secure key from bound entanglement.

Phys. Rev. Lett., 94:160502, Apr 2005.

Bibliography II

Pawel Horodecki.

Separability criterion and inseparable mixed states with positive partial transposition.

Physics Letters A, 232(5):333 – 339, 1997.