

Logistic Regression

Data Science: Jordan Boyd-Graber University of Maryland

Logistic Regression: Objective Function

$$\ell \equiv \ln p(Y|X,\beta) = \sum_{j} \ln p(y^{(j)}|x^{(j)},\beta)$$

$$= \sum_{j} y^{(j)} \left(\beta_0 + \sum_{i} \beta_i x_i^{(j)}\right) - \ln \left[1 + \exp\left(\beta_0 + \sum_{i} \beta_i x_i^{(j)}\right)\right]$$
(2)


Logistic Regression: Objective Function

$$\ell \equiv \ln p(Y|X,\beta) = \sum_{j} \ln p(y^{(j)}|x^{(j)},\beta)$$

$$= \sum_{j} y^{(j)} \left(\beta_{0} + \sum_{i} \beta_{i} x_{i}^{(j)}\right) - \ln \left[1 + \exp\left(\beta_{0} + \sum_{i} \beta_{i} x_{i}^{(j)}\right)\right]$$
(2)


Training data (y,x) are fixed. Objective function is a function of β ... what values of β give a good value.

Convexity



- Convex function
- Doesn't matter where you start, if you "walk up" objective


Convexity



- Convex function
- Doesn't matter where you start, if you "walk up" objective
- Gradient!


Goal

Optimize log likelihood with respect to variables β




Goal

Optimize log likelihood with respect to variables eta




Goal

Optimize log likelihood with respect to variables eta




Goal

Optimize log likelihood with respect to variables eta




Goal

Optimize log likelihood with respect to variables eta




Goal

Optimize log likelihood with respect to variables eta




Goal

Optimize log likelihood with respect to variables eta




Goal

Optimize log likelihood with respect to variables eta




Goal

Optimize log likelihood with respect to variables eta



Goal

Optimize log likelihood with respect to variables β



Luckily, (vanilla) logistic regression is convex