

A-level

ENVIRONMENTAL SCIENCE

Paper 1 - Generated practice paper Set 02

Mark scheme

120 marks: Questions 1-10 (95 marks) plus one essay from Question 11 (25 marks).

Marking guidance

- The number in brackets after each marking point is the mark it earns; the note above the points says how many are needed.
- Where a question asks for a fixed number of points (eg *two reasons*), credit only the first points given.
- Alternative correct wording and other relevant, accurate points are credited.
- For levels-of-response questions, first determine the level using the descriptors, then position the mark within the level using the indicative content.
- Calculations: credit correct working (method marks) even if the final answer is wrong; incorrect answers with no working score 0.

Textbook references: every question below cites the printed page numbers in R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018) from which the content is drawn, together with the AQA 7447 specification reference.

Question	Marking guidance	Marks
01.1	<i>1 mark for the correct answer.</i> • $16.0 - 5.0 = \text{£}11 \text{ million}$ (1)	1
01.2	<i>1 mark for each point.</i> • It is rarely practical to reduce emissions to zero and the impact of low-level emissions may be acceptable, so an acceptable level must be decided (1) • ALARA: emissions should be As Low As Reasonably Achievable (1) • BATNEEC: the Best Available Technology Not Entailing Excessive Cost is chosen to achieve this (1)	3
01.3	<i>1 mark for each point.</i> • As industries expand or cities grow the number of pollutant sources increases (1) • So the total emissions / combined concentration rises even if each source is unchanged, and higher standards are needed (1)	2

Question	Marking guidance	Marks
01.4	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) • Critical Pathway Analysis: predicting movement of mobile persistent pollutants (state of effluent, wind, currents, geology, pH / oxygen and solubility, bioaccumulation, food sources, half-lives) so monitoring focuses on sites at risk; mainly radioactive discharges; checked by sampling • Critical Group Monitoring: exposure of the most-exposed group monitored before health effects; if their risk is acceptable everyone else is safer; emissions reduced if not • Emission control strategies: location (marine discharges, downwind of cities, not over aquifers) and timing (avoiding temperature inversions, low river flow) • Principles: polluter pays (incentive depends on penalties and enforcement), precautionary principle (assume harm until shown otherwise; ignorance no excuse), ALARA / BATNEEC and the non-linear cost-efficiency relationship • Selection of methods: production prevention (desulfurisation), prevention of release (precipitators, catalytic converters), post-release remediation (oil spill clean-up, phytoremediation), alternative processes (electric vehicles, renewables, pyrethroids) • Evaluation: prevention usually cheaper and more effective than remediation; costs rise steeply near 100%; CPA relies on understanding of pathways; enforcement and international agreement needed for transboundary pollutants; judgement about acceptable risk is political as well as scientific • Justified conclusion	9
	Model answer - Level 3 (8 marks)	
	Deciding how much to control pollution starts with predicting where a pollutant will go. Critical Pathway Analysis uses the state of the effluent, wind and currents, geology, the effect of pH and oxygen on solubility, bioaccumulation and local food sources to predict whether a discharge will disperse harmlessly or concentrate somewhere sensitive; it is mainly used for radioactive discharges and means monitoring can focus on the sites at risk rather than sampling everywhere. Critical Group Monitoring then checks the exposure of the people most at risk from their lifestyle before any health effects occur: if their risk is acceptable everyone else's is lower, and if not, emissions are reduced.	

Question	Marking guidance	Marks
	The choice of control is guided by principles. The polluter pays principle gives a financial incentive to prevent damage, but only if penalties are high and polluters cannot escape responsibility. The precautionary principle assumes a waste is harmful until research shows otherwise, so ignorance is no defence. Because the cost of control rises steeply as emissions approach zero, emissions are set As Low As Reasonably Achievable using the Best Available Technology Not Entailing Excessive Cost. Where emission location and timing can be chosen (marine outfalls, downwind of towns, not over aquifers; avoiding temperature inversions) severity is reduced cheaply.	
	Overall, prevention of production (fuel desulfurisation) or of release (electrostatic precipitators, catalytic converters) is more effective than post-release remediation such as oil-spill clean-up, which never recovers all the pollutant. CPA depends on understanding pathways, and transboundary pollutants need international enforcement, but together these principles give a rational basis for deciding acceptable risk rather than aiming for an unaffordable zero.	
	Weaker answer - Level 1 (3 marks)	
	Pollution can be controlled in lots of ways. Critical Pathway Analysis looks at where pollution goes and Critical Group Monitoring looks at people. The polluter pays principle means the person who pollutes has to pay. The precautionary principle means being careful. Pollution can be prevented, or cleaned up afterwards with things like booms for oil spills. Electric cars are an alternative process. Pollution control can be expensive so ALARA is used. All of these are good ways of controlling pollution.	
	What makes the difference	
	• The Level 3 answer explains <i>how</i> each principle works (why penalties matter, why costs rise near zero emissions); the weak answer only names them.	
	• The Level 3 answer groups the material logically (predict, decide, select) rather than listing everything at the same weight.	
	• 'Evaluate' needs comparison and a judgement: prevention vs remediation, CPA's dependence on understanding pathways, and a conclusion about acceptable risk. The weak answer ends with 'all of these are good'.	
	• Precise examples (desulfurisation, precipitators, catalytic converters, temperature inversions) show applied knowledge; 'things like booms' does not.	
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 243-245, 303-304; AQA spec 3.4.3.1, 3.4.3.3.	

Question	Marking guidance	Marks
02.1	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) • Stages of scientific method illustrated by ozone: identification (Antarctic UV data), hypothesis (Rowland-Molina 1974), data collection / analysis (Dobson units, satellites, balloons confirming CIO), proposed solutions, enactment (Vienna 1985, Montreal 1987) • Evidence of success: ozone hole area and depth stabilised and beginning to shrink; CFC concentrations in the atmosphere falling; UV levels not rising further; global ozone loss limited to about 4% • Reasons for success: international recognition of serious consequences (skin cancer); agreement of nearly every country; legally binding targets with a fund for developing countries; relatively few producers / products; alternatives (HCFCs, HFCs, pump sprays, hydrocarbons) already available and cheap; no lifestyle change needed • Limits: persistence of CFCs means recovery takes decades; some ODSs still used; HFC replacements are greenhouse gases • Climate change contrast: greenhouse gases come from almost every human activity (energy, transport, agriculture, industry) in every country; fossil fuels underpin economies so control requires major lifestyle and economic change • Agreements (Kyoto, Paris) have weaker / non-binding targets, disputes between developed and developing nations; emissions still rising; natural fluctuations, time delays and tipping points make cause-effect less clear to the public • Judgement: ozone control was a technical substitution problem; climate change is a systemic one, so the ozone success is only partly transferable	9
	Model answer - Level 3 (9 marks)	
	Ozone depletion followed the scientific method almost textbook-fashion: Antarctic UV and ozone measurements identified a problem, Rowland and Molina's 1974 hypothesis explained it, Dobson-unit, satellite and balloon data confirmed chlorine monoxide in the stratosphere, and the Montreal Protocol enacted a solution. It has largely worked: CFC concentrations are falling, the ozone hole has stabilised and is starting to shrink, and global ozone loss was limited to about 4%.	

Question	Marking guidance	Marks
	Success came from particular conditions. The consequence - skin cancer - was clear and frightening, nearly every country agreed, targets were legally binding with a fund to help developing countries, there were relatively few producers and products, and alternatives (HCFCs, HFCs, hydrocarbons, pump sprays) already existed, so substitution cost little. Even so recovery will take decades because CFCs are so persistent, some ozone-depleting substances are still in use, and the HFC replacements are themselves greenhouse gases.	
	Climate change has none of these advantages. Greenhouse gases come from almost every activity - energy, transport, agriculture, industry - in every country, and fossil fuels underpin entire economies, so control means changing lifestyles and economies rather than swapping one chemical for another. Kyoto and Paris set weaker, largely non-binding targets, developed and developing nations dispute responsibility, and natural fluctuations, time lags and tipping points make cause and effect less obvious to the public. Emissions are still rising. The ozone story shows international agreement can work, but it was a technical substitution problem; climate change is systemic, so its lessons transfer only partly.	
	Weaker answer - Level 2 (4 marks)	
	Ozone depletion was caused by CFCs. Scientists found the hole over Antarctica and the Montreal Protocol banned CFCs, so the ozone layer is now recovering. This was successful because everyone agreed and there were alternatives to CFCs. Climate change has not been tackled as well. There have been agreements like Kyoto and Paris but carbon dioxide is still going up. This is because fossil fuels are used everywhere and countries do not agree. So ozone depletion was solved better than climate change.	
	What makes the difference	
	• Level 2 gives the right outline but Level 3 supports it with evidence (falling CFC levels, 4% loss, Dobson units, ClO detection) and explains each reason for the difference.	
	• Level 3 acknowledges the limits of the ozone success (persistence, HFCs) - balance is expected at the top level.	
	• The final judgement in Level 3 characterises the difference (substitution vs systemic problem); Level 2 just restates the question.	
02.2	<i>1 mark.</i> • Vienna Convention for the Protection of the Ozone Layer - no legally binding goals (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 108-113, 105-106; AQA spec 3.2.1.3, 3.2.1.2.	
03.1	<i>Correct answer only.</i> • B - Steam washing (hot water / steam jets kill organisms in the sediment) (1)	1
03.2	<i>1 mark for each point.</i> • Chemicals cause the oil molecules to join together / solidify (1) • Producing more solid material that can be collected / lifted more easily than liquid oil (1)	2

Question	Marking guidance	Marks
03.3	<i>1 mark for each point.</i> • Air contains about 210 cm³ of oxygen per litre whereas water at 20 °C holds a maximum of about 6 cm³ per litre (less than 3% of the amount in air) (1) • So a small amount of oxygen-consuming activity (decomposition, respiration) or warming can remove most of the oxygen; many aquatic organisms are near the lower end of their range of tolerance (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 254-262; AQA spec 3.4.3.2.6-3.4.3.2.7.	
04.1	<i>Any 3 from the points below - 1 mark each, maximum 3.</i> • Raise river banks / build sea walls, flood barriers or wave screens (1) • Permeable urban surfaces (gravel, soil) instead of concrete / tarmac to slow runoff (1) • River flow management: upstream dams / low soil dams / afforestation / maintained flood plains to delay runoff (1) • Raised buildings on stilts or floating houses (1) • Pumping rainwater from land below river level (1) • Managed retreat - abandoning land where protection costs exceed value (eg Happisburgh) (1)	3
04.2	<i>1 mark for each point.</i> • Where the cost of flood prevention is greater than the value of what is protected, or defences would be ineffective (1) • Land is abandoned as water levels rise / coast erodes; defences used only where land is very important (1)	2
04.3	<i>Any 2 from the points below - 1 mark each, maximum 2.</i> • Road surfaces deform as tar melts - need tarmac with a higher melting point (1) • Rail track expands and buckles above the temperature it was pre-stretched for (27 °C in UK) - relay with higher pre-stress (1) • Bridge damage from high river flows / debris; landslides on deforested slopes (1)	2
04.4	<i>1 mark for each point.</i> • CO₂ is captured from the flue gases (or carbon removed from the fuel before combustion) (1) • Transported by pipeline / ship / tanker and stored in depleted oil or gas fields, aquifers, or used for secondary oil recovery (1) • Limitation: developmental / expensive, uses energy (reducing efficiency), risk of leakage, limited storage sites (1)	3
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 99-100, 105-108; AQA spec 3.2.1.2.	
05.1	<i>1 mark for each point.</i> • t-test (or Mann-Whitney U if not normally distributed) (1) • Comparing two means of counted / measured data from two independent samples (1)	2
05.2	<i>1 mark for each point.</i> • The difference is statistically significant (less than 1% probability that it arose by chance) (1) • The null hypothesis (no difference between fields) is rejected; the organic field has significantly more earthworms (1)	2

Question	Marking guidance	Marks
05.3	1 mark for each point. • A significant difference does not show a causal relationship (1) • Other variables (soil type, moisture, pH, cultivation history) may differ between the two fields and cause the difference (1)	2
05.4	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) • Scientific methodology: identifying the topic, reviewing existing knowledge, formulating a testable hypothesis and null hypothesis, designing the study, analysing and drawing conclusions, planning further research • Sample location: random (grid coordinates / numbered sites) or systematic (transects, fixed intervals) to avoid bias; interval chosen so variation is detected without excess data • Sample timing: diurnal, seasonal, weather-related and long-term changes; sampling on different occasions; interval matched to rate of change • Sample size and number: larger and more samples for non-homogeneous variables; running mean and standard deviation show when the mean is reliable; preliminary study to decide • Standardisation and control of variables; accuracy, precision, reliability, representativeness; anomalies and repeats; calibration of equipment; appropriate techniques and risk management • Statistical analysis: choosing the test (Spearman, chi-squared, t-test, Mann-Whitney), significance and p-values, error bars, correlation vs causation • Evaluation: trade-offs between cost, time, disturbance and reliability; limitations of sub-sampling; justified conclusion	9
	Model answer - Level 3 (8 marks)	
	A valid investigation begins with a clear, testable hypothesis and a null hypothesis, informed by existing knowledge, so that the data collected can actually accept or reject it. The first design decision is sample location: random sampling (grid coordinates or numbered sites) or systematic sampling along transects avoids the bias that would come from choosing convenient sites, and a preliminary study shows the interval needed to detect variation without collecting unnecessary data.	

Question	Marking guidance	Marks
	Timing matters because many variables change diurnally, seasonally or with weather, so samples on different occasions are needed and the interval must match the rate of change. Sample size and number depend on how variable the population is: a running mean shows when enough samples have been taken and the standard deviation shows the scatter around the mean. Techniques must be standardised so results from different sites, times or researchers are comparable, other variables controlled or monitored, and equipment calibrated so that precise readings are also accurate; anomalies should be repeated rather than ignored.	
	Finally the analysis must be chosen to suit the data - Spearman's rank for correlation, chi-squared for frequencies, a t-test for two means or Mann-Whitney U for medians - so that significance ($p < 0.05$) can be assessed, remembering that a significant correlation does not prove causation. Every decision trades reliability against cost, time and disturbance; sub-samples can only estimate the whole, so the most valid study is the one whose sampling design makes its limitations explicit.	
	Weaker answer - Level 2 (5 marks)	
	When planning fieldwork you need a hypothesis. You should choose sites randomly so it is not biased, for example using random numbers. You need to take enough samples and repeat them so the results are reliable. The method should be the same each time. You should measure accurately with calibrated equipment. Then you do a statistical test such as a t-test to see if the results are significant. If p is less than 0.05 the result is significant. This makes the conclusions valid.	
	What makes the difference	
	• Level 2 is sound but generic: it says 'enough samples' where Level 3 explains how a running mean and standard deviation tell you how many.	
	• Level 3 covers timing and the choice between statistical tests; Level 2 mentions one test only.	
	• The evaluation is missing from Level 2 - there is no discussion of trade-offs or of correlation vs causation, so it cannot reach the top level however accurate it is.	
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 390-393, 420-423; AQA spec 3.7.1, 3.7.2.3.	
06.1	<i>A correct final answer (with unit where asked) scores full marks. If the answer is wrong, award 1 mark for each correct step of working shown.</i> • No reservoir range = $3.9 - 0.7 = 3.2$ m; with reservoir = $2.7 - 1.7 = 1.0$ m (1) • Reduction = 2.2 m (1)	2
06.2	<i>1 mark for each point.</i> • After heavy rainfall surplus water is stored, reducing peak flows / flooding downstream (1) • During dry periods the dam is opened to release water and maintain river levels (1)	2
06.3	<i>1 mark for each point.</i> • Deforestation / urbanisation reduce infiltration and interception (1) • So runoff reaches the river faster after rain and less water is stored to maintain flow in dry periods (1)	2

Question	Marking guidance	Marks
06.4	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) • Site factors: topography - narrow exit from a large deep basin so a small dam holds a large volume; impermeable, strong bedrock without faults or seismic activity; large catchment with regular high rainfall and low evaporation; catchment land use with low pollution risk (pesticides, industry) and low soil erosion to avoid sedimentation; forest / vegetation in the flooded area decaying anaerobically releasing methane; access for workers and materials and distance to demand; value of the land lost (urban, conservation, agricultural) - loss-benefit analysis differs between countries • Alternatives for supply: rainwater collection, rivers, estuary barrages, desalination (energy-intensive so only where freshwater is inadequate), unexploited aquifers (N Africa, Kenya), inter-basin transfers (China, Australia, Wales-England), artificial aquifer recharge using surplus wet-season water, river-regulation reservoirs, afforestation to even out flow • Conservation strategies: low-water appliances, dual-flush toilets, xeriscaping, drip irrigation, grey-water re-use, leak repair (about 20% lost in the UK), metering • Treatment / pollution control allowing re-use of contaminated sources: sewage treatment, buffer strips, mine drainage control • Evaluation: reservoirs give large reliable supplies and flood control but flood land, cause habitat loss and methane; conservation is cheap and quick but limited by behaviour; transfers create conflicts between regions; desalination has high energy / carbon cost; combination needed as demand grows • Justified conclusion	9
	Model answer - Level 3 (8 marks)	
	A reservoir site is judged first on topography: a narrow exit from a large, deep basin lets a small, cheap dam hold a large volume. The bedrock must be impermeable so water is not lost and strong enough, without faults or seismic activity, to carry the dam. The catchment should be large with regular, high rainfall and low evaporation, and its land uses should not pose a pollution risk from industry or pesticides or cause the soil erosion that would silt the reservoir. Flooding forest or vegetation causes anaerobic decay and methane. Access for workers and materials, distance to the area of demand, and the value of the land lost - urban areas and nature reserves protected, farmland less so - complete a loss-benefit analysis that different countries weigh differently.	

Question	Marking guidance	Marks
	Supply can also be increased without new reservoirs: rainwater collection in cities, estuary barrages, desalination where freshwater is genuinely inadequate, newly found aquifers in North Africa and Kenya, inter-basin transfers as in China and from Wales to England, artificial recharge of aquifers with wet-season surplus, river-regulation reservoirs and afforestation to even out flow. Demand can be cut by low-water appliances, xeriscaping, drip irrigation, grey-water re-use, metering and repairing the leaks that lose about a fifth of UK supply, and treatment lets contaminated sources be re-used.	
	Reservoirs give large, reliable supplies and flood control but flood land and habitats and release methane; conservation is cheap and quick but limited by behaviour; transfers create conflicts between regions; desalination carries a high energy and carbon cost. As demand grows, the sustainable answer is conservation first, then a combination of sources matched to each region.	
	Weaker answer - Level 2 (4 marks)	
	A reservoir needs a large catchment area with lots of rain, impermeable rock so the water does not drain away, and a narrow valley so the dam is small. It should be near the people who need the water and not flood important land. Other ways to get water include desalination, which is expensive, and collecting rainwater. Water can be conserved by fixing leaks, using water meters and having low-flush toilets. All of these help make water supplies more sustainable.	
	What makes the difference	
	• Level 2 lists correct site factors but without the reasons (why impermeable, why the valley shape matters, why land use in the catchment matters, methane from flooded vegetation).	
	• Level 3 covers far more alternatives (aquifer recharge, river regulation, transfers, afforestation) and evaluates each; Level 2 evaluates only desalination.	
	• Level 3 ends with a prioritised judgement; Level 2 ends with 'all of these help'.	
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 122-127; AQA spec 3.2.2.4-3.2.2.5.	
07.1	<i>Correct answer only.</i> • Aerobic decomposition produces little methane and returns nutrients to the soil (1)	1
07.2	<i>1 mark for each point.</i> • It may contain concentrated heavy metals / toxic residues (1) • It is stored in solid form at high pH / in hazardous-waste landfill so the metals stay insoluble (1)	2
07.3	<i>Any 2 from the points below - 1 mark each, maximum 2.</i> • Impermeable polymer / clay liner to stop leachate escaping (1) • Collection and treatment of leachate (1) • Collection of methane and use as fuel (1) • Daily covering of waste / final capping and restoration (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 279-285; AQA spec 3.4.3.2.12.	

Question	Marking guidance	Marks
08.1	<i>Correct answer only.</i> • B - Tropospheric ozone (formed by reactions of NO _x in sunlight) (1)	1
08.2	<i>1 mark for each point.</i> • Primary pollutant: released directly by human activities, eg sulfur dioxide from burning coal / smelting sulfide ores; NO _x from high-temperature combustion (1) • Secondary pollutant: produced by chemical reactions between primary pollutants, often with non-pollutants (water, oxygen, sunlight) (1) • Sulfur dioxide is oxidised by ozone to sulfur trioxide which dissolves to form sulfuric acid; NO _x dissolve to form nitric acid (1) • The acids (sulfurous, sulfuric, nitric) are secondary pollutants; ozone itself is a secondary pollutant of NO _x photochemistry (1)	4
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 237-238; AQA spec 3.4.1.	
09.1	<i>Correct answer only.</i> • Oxygen (1)	1
09.2	<i>1 mark for each point.</i> • A change to one process causes other processes to alter as a direct result (chains of linked processes / feedback) (1) • There is still a lot that is not understood about how atmospheric processes work (1)	2
09.3	<i>1 mark for each point.</i> • The surface warmed in the tropics heats the air above; warm winds (south-westerlies from the Caribbean) carry heat energy to higher latitudes (1) • Winds carry water vapour (evaporated from warm oceans) to areas that would otherwise receive little precipitation (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 86-88; AQA spec 3.2.1.1.	

Question	Marking guidance	Marks
10.1	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) • Context: Lasky's principle - vast quantities exist at low grade; reserves vs resource; COOG affected by price, technology, energy costs • Improved exploration: higher resolution multispectral satellite sensors, IR spectroscopy, gravimetry, magnetometry, seismic, resistivity, portable field analysis, targeted trial drilling • Mechanisation allowing deeper underground mines and larger open-cast mines • Low-energy extraction of low-grade ores: bioleaching with acidophilic bacteria and fungi, phytomining, iron displacement, bacterial adsorption of rare earths, polymer adsorption from seawater • Previously inaccessible deposits: polymetallic nodules, deep sea sediments, Antarctica (currently protected) - environmental and legal issues • Extending use: recycling (advantages, difficulties), Cradle to Cradle design, product lifetime extension, substitution of scarce materials, circular economy • Evaluation: environmental impacts vs benefits; energy costs; economics; which strategies are most sustainable; conclusion	9
	Model answer - Level 3 (8 marks)	
	Lasky's principle means that as grade falls, the quantity of metal available rises enormously, so depletion of high-grade deposits does not mean running out - it means the cut-off ore grade must fall, which depends on price, technology and energy costs. Better exploration finds what remains: high-resolution multispectral and infrared satellite sensors, gravimetry, magnetometry, seismic and resistivity surveys and portable field analysers narrow the search so expensive trial drilling is targeted. Mechanisation allows deeper underground mines and larger open-cast pits, and previously inaccessible deposits - polymetallic nodules and deep-sea sediments, and Antarctica if its protection were ever lifted - become possible, with serious environmental and legal problems.	

Question	Marking guidance	Marks
	The most sustainable route to low-grade ores is low-energy extraction: bioleaching with acidophilic bacteria and fungi, phytomining, iron displacement of copper, bacterial adsorption of rare earths and polymer adsorption from seawater all avoid the energy of crushing and smelting vast tonnages. Equally important is extending the use of metal already mined: recycling, Cradle-to-Cradle design so products can be disassembled, longer product lifetimes and substitution of scarce metals - the circular economy approach.	
	Each strategy has costs: deep-sea mining and Antarctica risk unique habitats, deeper mines need more energy, recycling is limited by dispersal and mixed alloys. The strategies that mimic natural low-temperature processes and keep metals in use are the most sustainable, and exploration and new mining should be the last resort rather than the first.	
	Weaker answer - Level 1 (3 marks)	
	When high grade ores run out we can mine lower grade ores instead, but this uses more energy. Better technology like satellites can find new deposits. We could mine the deep sea or Antarctica. Bacteria can be used to extract metals from low grade ore. Recycling metals means we do not need to mine as much. These strategies will help secure supplies of metals in the future.	
	What makes the difference	
	• Level 1 names most strategy types but Level 3 explains each (what the surveys measure, why low-temperature extraction matters, what Lasky's principle implies).	
	• Level 3 weighs environmental and energy costs and ranks the strategies; Level 1 offers no evaluation.	
	• Framing the answer with the concept of cut-off ore grade shows understanding rather than recall.	
10.2	<i>1 mark.</i> • All of the material that exists in the lithosphere, including that which will never be technologically or economically exploitable (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 130-147; AQA spec 3.2.3.	

Question	Marking guidance	Marks
11.1	Discuss the difficulties in monitoring and predicting global climate change and evaluate the methods used to overcome them. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) • Nature of the problem: interconnected atmosphere-hydrosphere-biosphere systems; changes over different time and spatial scales; natural fluctuations (solar output, orbit) masking or exaggerating anthropogenic change; time delay between cause and effect (ocean heat capacity) • Limitations of historic data: sparse global coverage, unsophisticated equipment, heat island bias, no atmospheric composition records • Proxy data and their reliability: dendrochronology, coral growth rings, pollen in dated sediments, ice cores (gas bubbles, isotope ratios, volcanic ash layers, compaction problems) • Modern monitoring: long-term data sets (Mauna Loa), electronic equipment, satellites (LEO vs geostationary), Argo floats, remote sensing of ice thickness / area / sea level • Feedback mechanisms and tipping points adding uncertainty; ecological uncertainty (species survival, distribution, fragmentation) • Computer models: how they are tested and refined; importance of accurate representative data; scenario-based predictions and ranges • Uncertainty over ENSO / Conveyor / jet stream responses • Evaluation and conclusion: confidence in the overall trend and human cause despite detailed uncertainty; role of peer review and international bodies (IPCC) Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 89-90, 100-104, 10; AQA spec 3.2.1.2, 3.1.1.3.	25

Question	Marking guidance	Marks
11.2	Explain how feedback mechanisms and tipping points influence the rate of global climate change and discuss the implications for the strategies needed to control it. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) • Negative feedback: increased low-level cloud (albedo), increased photosynthesis and sequestration; hydrological cycle evaporation-precipitation • Positive feedback: melting permafrost (methane), reduced albedo of snow / ice / sea ice, methane hydrate release, ocean acidification and coral loss, forest and peat fires, soil decomposition, water vapour, cirrus clouds • Tipping points: self-sustaining processes; why control must precede them; examples • Implications: urgency and precautionary principle; need for deep cuts in greenhouse gases (energy, transport, buildings, agriculture, forestry, CCS) rather than only adaptation; geoengineering risks • Difficulty of prediction because feedbacks interact at different scales; models; international agreements and per capita footprints • Judgement on whether current strategies are adequate Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 103-107, 371-372; AQA spec 3.2.1.2, 3.6.1.	25