

A-level

ENVIRONMENTAL SCIENCE

Paper 2 - Generated practice paper Set 01

Mark scheme

120 marks: Questions 1-10 (95 marks) plus one essay from Question 11 (25 marks).

Marking guidance

- The number in brackets after each marking point is the mark it earns; the note above the points says how many are needed.
- Where a question asks for a fixed number of points (eg *two reasons*), credit only the first points given.
- Alternative correct wording and other relevant, accurate points are credited.
- For levels-of-response questions, first determine the level using the descriptors, then position the mark within the level using the indicative content.
- Calculations: credit correct working (method marks) even if the final answer is wrong; incorrect answers with no working score 0.

Textbook references: every question below cites the printed page numbers in R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018) from which the content is drawn, together with the AQA 7447 specification reference.

Question	Marking guidance	Marks
01.1	<i>A correct final answer (with unit where asked) scores full marks. If the answer is wrong, award 1 mark for each correct step of working shown.</i> • $(15 - 6) / 6 \times 100$ (1) • = 150% (1)	2
01.2	<i>1 mark for each point.</i> • Cells shed by the newts into the water contain their DNA (1) • Water samples are analysed for DNA sequences unique to the species (1)	2
01.3	<i>1 mark for each point.</i> • Catches depend on trapping effort, weather and newt activity on the night, so they vary from year to year (1) • Trapping only samples one pond and one part of the population, whereas eDNA surveys all 20 ponds with a standard method (1)	2

Question	Marking guidance	Marks
01.4	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) • Image databases of unique markings (tiger stripes, whale shark spots, dolphin fins) giving territory size, movements, lifespan and social groups; limited to species with unique features • Motion-sensitive and infrared cameras, CCTV of nests with less disturbance; marking (rings, wing tags) • DNA: eDNA to detect presence (great crested newts, invasive carp); DNA databases to identify regional populations and trace timber, ivory and fish • Auditory monitoring: bat detectors and sonograms for presence, abundance and activity • Position monitoring: radio, GPS and acoustic transmitters; geolocator tags for small birds; data recorders; carrier systems - ROVs / drones, AUVs, balloons, aircraft, satellites, animals • Satellites: visible light for land use and deforestation, infrared for vegetation and temperature, radar for wind, waves, sea level and oil; GPS for tracking • Evaluation: greater coverage, less disturbance and continuous data vs cost, battery / weight limits, calibration, need for ground-truthing, ethical issues of tagging; indirect evidence and traditional methods still needed • Justified conclusion	9
	Model answer - Level 3 (9 marks)	
	New technologies have transformed what can be monitored and how often. Image databases of unique markings - tiger stripes, whale-shark spots, dolphin fin scars - let individuals be recognised, giving territory size, movements, lifespan and social structure without repeated capture, though only for species with unique features. Motion-sensitive and infrared cameras and CCTV record activity day and night with far less disturbance than observers, especially at nests.	

Question	Marking guidance	Marks
	Genetics adds detection where nothing can be seen: eDNA in water reveals great crested newts or invasive carp from shed cells, and DNA databases identify which regional population timber, ivory or fish came from, supporting enforcement. Bat detectors and sonograms give presence, abundance and activity of species heard but not seen. Position monitoring uses radio, GPS and acoustic transmitters, or tiny geolocator tags for birds too small to carry a transmitter, while data recorders on drones, AUVs, balloons, aircraft, satellites or the animals themselves collect abiotic data. Satellites map land use and deforestation in visible light, vegetation density and temperature in infrared, and wind, waves, sea level and oil slicks with radar that works through cloud.	
	The gains are coverage, continuity and reduced disturbance. The limits are cost, battery life and weight, the need to calibrate sensors and ground-truth remote data, ethical questions about tagging, and the fact that many species still require traditional methods and indirect evidence. New technology therefore complements rather than replaces fieldwork, but for rare, remote or nocturnal species it has made effective conservation possible where it was not before.	
	Weaker answer - Level 1 (3 marks)	
	New technologies help monitor wildlife. Cameras can take pictures of animals at night. GPS collars can track where animals go. Satellites can see deforestation from space. DNA can tell which animals are present. Bat detectors can hear bats. These help conservation because we know where animals are and how many there are. They can be expensive though.	
	What makes the difference	
	• Level 1 lists technologies with no detail of how they work or what data they give; Level 3 links each to the information it provides (territory size, presence, activity).	
	• Level 3's evaluation considers several limitations (calibration, ground-truthing, ethics, species coverage) and reaches a balanced judgement; Level 1 has one word - 'expensive'.	
	• Structure: Level 3 groups by imagery, genetics, sound, position and remote sensing; the list in Level 1 has no organisation.	
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 402-407; AQA spec 3.1.2.3.5-3.1.2.3.6.	
02.1	<i>1 mark for each point.</i> • Surveys monitor photosynthesis rates, biomass, soil water content and the spread of pests / diseases so inputs are applied only where needed (1) • Yield mapping of different parts of a field controls fertiliser application in each area for future crops (1) • This maximises yield while avoiding wastage / reducing leaching and nutrient pollution (1)	3
02.2	<i>1 mark for each point.</i> • Machinery, pesticides, fertilisers, improved breeds and infrastructure (transport, refrigeration) are expensive (1) • They are mainly available in affluent societies; poorer farmers cannot afford the inputs (1)	2

Question	Marking guidance	Marks
02.3	<i>1 mark.</i> • Provides an income that allows producers to afford basic rights such as water, education, health care and food (1)	1
02.4	<i>1 mark for each point.</i> • Nitrates are leached into water bodies used as sources of potable water (1) • High nitrate levels cause blue baby syndrome (methaemoglobinaemia) and nitrates may be carcinogenic (1)	2
02.5	<i>Any 2 from the points below - 1 mark each, maximum 2.</i> • Anaerobic digestion by microbes in rice padi fields (1) • Livestock intestines (ruminant digestion) (1) • Anaerobic decay of manure / slurry (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 329-332; AQA spec 3.5.1.3-3.5.1.4.	
03.1	<i>1 mark for each point.</i> • Plants produce chemicals (often alkaloids) that are toxic to the herbivores that would eat them (1) • In carefully controlled amounts these chemicals have beneficial physiological effects in humans (1)	2
03.2	<i>1 mark for each point.</i> • Only a very small proportion of species have been studied for medicinal substances, so it is not known which species will be useful (1) • Conserving the habitat protects all the species in it, including those not yet studied and the species they depend on (1)	2
03.3	<i>Any 2 from the points below - 1 mark each, maximum 2.</i> • Marsupials (kangaroo / wallaby): young develop in the pouch so are easier to study than a human foetus - developmental problems (1) • Hippopotamus: hipposudoric acid skin secretion (sunscreen / antimicrobial) - burns treatment (1) • Marine sponges: proteins preventing graft rejection - organ transplants (1)	2

Question	Marking guidance	Marks
03.4	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) • Resources: wood, fibres, oils, fuels, food; new food species adapted to local conditions (morama bean, yeheb, perennial maize, Kernza, bison, eland) • Biomimetics: whale flipper tubercles, shark skin, termite-mound ventilation, bird bones, gecko adhesion, burrs / Velcro, lotus self-cleaning, spider silk • Medicines and physiological research; genetic resources - crop wild relatives, Vavilov centres, gene pools, inbreeding • Ecosystem services: atmospheric composition, hydrological cycle, biogeochemical cycles, soil maintenance; interspecies relationships - food, pollination, seed dispersal, habitat provision • Evaluation: services go unnoticed because of dynamic equilibrium; unknown future value; economic vs moral / intrinsic arguments; conservation also needed for aesthetic and cultural reasons • Justified conclusion	9
	Model answer - Level 3 (9 marks)	
	Other species supply resources we use directly: wood, fibres such as cotton and wool, oils, fuels and food. Because most farmed species were domesticated long ago and are often kept where they are poorly adapted, wild species such as the drought-resistant morama bean, the yeheb tree that grows on poor soils, perennial maize that needs no ploughing, and bison that graze grassland too poor for cattle offer new, better-adapted food sources. Wild species also carry genetic resources: crop wild relatives in Vavilov centres hold genes for disease resistance and salt tolerance that inbred commercial varieties lack.	
	Biomimetics copies adaptations into technology - humpback flipper tubercles in turbine blades, shark skin in friction-reducing coatings, termite-mound convection in naturally ventilated buildings, hollow bird bones in lightweight roofs. Medicines such as taxol, quinine, AZT and diosgenin came from wild species, and physiological research on marsupials, hippo skin and sponge proteins is helping with foetal development, burns and transplant rejection. Since only a tiny fraction of species has been studied, the unknown future value argues for conserving whole habitats.	

Question	Marking guidance	Marks
	Beyond resources, other species provide services we could not replace: regulation of atmospheric gases, evapotranspiration driving rainfall, the carbon, nitrogen and phosphorus cycles run by microbes, soil formation by decomposers, and pollination, seed dispersal and habitat provision. These are unnoticed precisely because they are in dynamic equilibrium, so their economic value is underestimated. On balance the resource and service arguments alone justify conservation, but they are strongest when combined with the intrinsic and aesthetic case, because many species with no known use still sustain the systems we depend on.	
	Weaker answer - Level 1 (3 marks)	
	Biodiversity is important because we get lots of things from other species. We get wood, cotton and food from plants and animals. Some medicines come from plants, like aspirin from willow trees. Animals can help design things, like Velcro was copied from seeds. Other species also do things for us like bees pollinating crops. If we lose species we might lose things that could be useful in the future. So we should conserve biodiversity.	
	What makes the difference	
	Both answers cover resources, medicines, biomimetics and services, but Level 3 develops each with specific, accurate detail (which species, what property, why it matters).	
	The question asks 'to what extent'. Level 3 weighs resource arguments against intrinsic value and reaches a reasoned conclusion; Level 1 asserts 'so we should conserve'.	
	Level 3 explains <i>why</i> services are undervalued (dynamic equilibrium) - an explanation, not just a list.	
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 10-19; AQA spec 3.1.2.1.	
04.1	<i>2 marks for each point (partial credit for a partly correct point).</i> - EDGE: Evolutionarily Distinct and Globally Endangered - few close relatives, often the last species of their genus / family (Bactrian camel, pygmy hippo, northern bald ibis); loss would remove a unique branch of evolution / genes (2) - Endemic: found in only one area (often islands - Galapagos, Madagascar, Seychelles: Aldabra giant tortoise, red ruffed lemur); if lost locally there is no population elsewhere to recolonise (2) - Keystone: disproportionately important for the structure of the community relative to its abundance - predation of dominant species, seed dispersal, habitat creation (jaguar, forest elephants keeping paths and water holes open, wolves, beavers); losing them causes loss of many other species (2) - Flagship: high public profile (tiger, panda, elephant, orang-utan) raising funds and support that also protects the whole habitat and less charismatic species (fungi, bats, moths) (2)	8
04.2	<i>1 mark for each point.</i> - A fragmented population may consist of isolated groups each too small to be viable (resources, inbreeding) even if total numbers are large (1) - A species confined to a single area (eg bonobos in one country) is vulnerable to any local change / conflict, unlike chimpanzees spread across many countries (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 28-29; AQA spec 3.1.2.3.1.	

Question	Marking guidance	Marks
05.1	<i>Up to 4 marks in total; award marks in proportion to the completeness of each point.</i> • Success: Cactoblastis moth introduced from South America controlled prickly pear cactus in Australia; Encarsia parasitic wasps control greenhouse whitefly (up to 2) • Problems: cane toads introduced to Australia to control sugar cane beetles did not control them but ate many other species and are toxic to predators; rosy wolfsnail introduced to Polynesia to control African land snails ate endemic Partulid snails (48 species extinct); harlequin ladybirds introduced to Europe eat native ladybirds (up to 2) • Lesson: control agents should be specialist feeders on the pest (up to 2)	4
05.2	<i>1 mark for each point.</i> • Release an artificial sex scent attracting males (or females) of the pest species (1) • Used to monitor when the pest arrives so pesticide is timed correctly, or (in enclosed greenhouses) to trap enough of one sex to prevent breeding (1)	2
05.3	<i>1 mark for each point.</i> • Regular low doses kill only the most sensitive bacteria so resistant strains survive and multiply (1) • Agricultural use exceeds medical use, so resistance spreads (1) • Resistant zoonotic bacteria (E. coli, Salmonella, Campylobacter) can transfer to humans and are harder to treat; hence reduced use of antibiotics is a sustainability goal (1)	3
05.4	<i>1 mark.</i> • Grow nectar-rich flowering plants alongside crops; restrict pesticides harmful to bees / spray when not in flower; introduce bee hives (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 316-320; AQA spec 3.5.1.1.	
06.1	<i>Correct answer only.</i> • B (1)	1
06.2	<i>Any 4 from the points below - 1 mark each, maximum 4.</i> • Harvesting no faster than the MSY - resource never depleted (1) • Mixed species plantations - more food choices, relationships and niches (1) • Indigenous species - support indigenous insects, birds (1) • Mixed age structure - varied light, temperature, wind so more species; continuous cover and yield (1) • Selective logging - small clearings, less disturbance, long-term timber benefit (1) • Retention of dead wood, buffer zones, corridors (1)	4
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 360-369; AQA spec 3.5.3.	
07.1	<i>Any 2 from the points below - 1 mark each, maximum 2.</i> • Hydrological cycle (1) • Carbon cycle (1) • Nitrogen cycle (1) • Atmospheric circulation (1) • Thermohaline circulation (1) • Photosynthesis (1)	2

Question	Marking guidance	Marks
07.2	<i>1 mark for each point.</i> - Enzymes reduce the activation energy of reactions (1) - So reactions such as photosynthesis, decomposition and nitrogen fixation do not need high temperatures (1)	2
07.3	<i>Any 1 from the points below - 1 mark each, maximum 1.</i> - Haber process (nitrogen fixation) (1) - Incineration of wastes (decomposition) (1) - Many manufacturing processes (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 373; AQA spec 3.6.2.	
08.1	<i>Correct answer only.</i> Its mass (1)	1
08.2	<i>1 mark for each point.</i> - The 24-hour rotation period means the surface is not exposed to sunlight for long, so temperature extremes / excessive heating and cooling are reduced (1) - The axis is tilted relative to the orbit so the amount of insolation at each latitude changes through the year, producing seasonal variations in conditions (1)	2
08.3	<i>1 mark for each point.</i> - The distance from the Sun gives surface temperatures between 0 and 35 °C in most areas: too close and water would boil / evaporate, too far and it would freeze (1) - Atmospheric pressure (produced by the mass of gases held by gravity) is high enough to stop the water boiling / evaporating away; at much lower pressure there would be no liquid water (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 5-7; AQA spec 3.1.1.1.	
09.1	<i>1 mark.</i> Cross-bred offspring are healthier / more productive because the problems caused by homozygous recessive genes in inbred varieties are reduced (heterosis) (1)	1
09.2	<i>1 mark for each point.</i> - Zebu tolerate heat but have a low milk yield; Ayrshire have a high milk yield but are not heat tolerant (1) - Cross-breeding combined the desirable characteristics: heat tolerance with a high milk yield (1)	2
09.3	<i>Any 2 from the points below - 1 mark each, maximum 2.</i> - Replacement of valuable animals that die with genetically identical individuals (1) - Replacement of herds culled during a disease outbreak (1) - Production of large numbers of individuals with desirable characteristics (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 321-322; AQA spec 3.5.1.2.	

Question	Marking guidance	Marks
10.1	<i>Any 4 from the points below - 1 mark each, maximum 4.</i> • Zooxanthellae are photosynthetic algae living inside the polyp cells (1) • The algae absorb CO₂ and excretory wastes from the polyp for photosynthesis and growth (efficient nutrient recycling) (1) • They supply the polyp with oxygen and glucose - over 95% of its food energy (1) • In return the algae gain a protected habitat and nutrients they could not absorb from the nutrient-deficient water (1) • Polyps also feed on plankton captured by cilia and nematocysts (1)	4
10.2	<i>1 mark for each point.</i> • Corals deposit a skeleton of calcium carbonate (calcite) (1) • Each carbonate molecule requires the absorption of a molecule of carbon dioxide, so reefs are a carbon sink (1)	2
10.3	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) • Importance: cover 1% of ocean area but support over a quarter of marine species; fisheries (fish, lobsters, conch); medicines (AZT, Ara-C, dolastatin, secosteroids, bryostatin, trabectedin, bone grafts); climate control as carbon sink; erosion protection of low-lying islands; tourism income • Threats: physical damage (anchors, divers, litter, coral extraction), collection of ornaments (CITES Appendix II), sedimentation from rivers and coastal development, pollution (oil, nutrients causing algal shading, sewage, pesticides, sunscreen), ocean acidification, bleaching from warming, overfishing (giant triton / crown of thorns starfish), introduced species (red lionfish), loss of mangroves and seagrass nurseries • Conservation: marine protected areas / national parks (Great Barrier Reef, Belize, Chagos), mooring buoys, diver education, No Take Zones which can increase catches elsewhere, sustainable ecotourism and fishing, protection of mangroves and seagrass, erosion and pollution control, planning control of ports and resorts • Fishing controls: closed seasons, minimum and maximum catchable sizes, bans on dynamite, trawling and small-mesh nets • Evaluation: local measures cannot control global threats (warming, acidification) which need international action; enforcement in remote areas is difficult; conflicts with local livelihoods; large vs small NTZs • Justified conclusion	9
	Model answer - Level 3 (9 marks)	

Question	Marking guidance	Marks
	Tropical coral reefs cover about one per cent of the oceans yet support over a quarter of marine species, a diversity built on the symbiosis between polyps and zooxanthellae that recycles nutrients in nutrient-poor water. They matter to people for fisheries of fish, lobsters and conch; for medicines such as AZT, Ara-C, dolastatin, secosteroids, bryostatin and trabectedin discovered in reef organisms; as a carbon sink, since every carbonate unit in coral skeleton absorbs a molecule of carbon dioxide; for dissipating wave energy that would erode low-lying islands; and for tourism income.	
	They are also unusually vulnerable because they need narrow abiotic conditions. Anchors, divers, litter and coral extraction kill polyps; ornaments are collected; sediment from rivers and coastal development smothers them; oil, sewage and nutrients cause algal shading; ocean acidification hinders skeleton formation; warming causes bleaching; overfishing of giant tritons lets crown-of-thorns starfish destroy reefs; introduced lionfish eat native fish; and loss of mangroves and seagrass removes the nurseries and sediment traps reefs depend on.	
	Marine protected areas such as the Great Barrier Reef, Belize and Chagos combine mooring buoys, diver education, no-take zones that raise catches outside them, sustainable tourism and protection of linked habitats, while fishing controls set closed seasons, minimum and maximum sizes and ban dynamite, trawling and small mesh. These work well against local threats where they are enforced, and no-take zones have measurable benefits, but they cannot touch warming and acidification, which need international action, enforcement in remote areas is weak, and restrictions conflict with local livelihoods. Local conservation buys time; only global climate action can secure the reefs.	
	Weaker answer - Level 1 (3 marks)	
	Coral reefs are important because lots of fish live there and people catch them for food. Tourists visit reefs to dive and this brings money. Reefs also protect the coast from waves. Reefs are threatened by climate change which causes bleaching, and by pollution and damage from boats. They are conserved by making marine protected areas where fishing is banned. This helps but climate change is still a problem.	
	What makes the difference	
	• Level 1 has the right headings but Level 3 explains each (why reefs are diverse, why they are a carbon sink, how each threat acts).	
	• Level 3 uses named reserves and specific management measures; Level 1 gives one generic measure.	
	• Level 3's evaluation separates local from global threats and reaches a nuanced judgement; Level 1 states that climate change is 'still a problem'.	
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 57-62; AQA spec 3.1.2.3.4.	

Question	Marking guidance	Marks
11.1	Agriculture involves the control of abiotic and biotic factors to maximise food production. Discuss how this control can be achieved and evaluate its environmental impacts. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) • Species selection: market, range of tolerance, controllability • Abiotic control: temperature, light, water, soil fertility / nutrients, aeration, salinity, pH, CO₂, topography, relief, wind - methods and examples • Biotic control: pest groups, cultural control, biological control (successes and failures), sterile males, pheromones, genetic resistance, pesticides (properties), antibiotics, pollinators, soil biota • Manipulation of food species: population density, monocultures, selective breeding, asexual reproduction, cross-breeding, AI / embryo transfer, GM; energetics - intensive vs extensive, energy subsidies, ratios, FCR, food chain losses • Impacts: habitat loss / change, reduced biodiversity, introduced species, pesticide and nutrient pollution, GHGs, hydrological changes, soil erosion; positive: plagioclimax habitats • Social, economic and political influences shaping methods • Sustainable strategies and evaluation; conclusion Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 306-337; AQA spec 3.5.1.	25

Question	Marking guidance	Marks
11.2	Fishing is the last large-scale human hunting activity. Discuss the environmental impacts of fishing and evaluate the methods used to make the exploitation of fish populations sustainable. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) • Marine productivity: light, nutrients, upwelling; where fisheries occur • Methods: demersal / pelagic trawling, purse seining, drift nets, long lines, shellfish traps - effectiveness, selectivity, energy, impacts • Impacts: overfishing (r vs k selection, catchable vs breeding size), changed age structure, by-catch (albatross, porpoise, turtles, shrimp), ghost fishing, seabed / reef / seagrass damage, dynamite, food web effects (sandeels, tritons) • Population dynamics and MSY: Russell formula, difficulties of data, indicators from catches (CPUE, mean size, age), research on eggs and larvae • Management: quotas, mesh design, escape panels, TEDs, pingers, hook design, effort limits, method bans, NTZs, closed seasons, size limits, protected individuals, population seeding, biodegradable gear; EU CFP, IWC, CCAMLR • Evaluation with examples (cod, orange roughy, herring recovery, NTZ spillover); role of aquaculture; judgement Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 338-351; AQA spec 3.5.2.	25