

A-level

ENVIRONMENTAL SCIENCE

Paper 2 - Generated practice paper Set 02

Mark scheme

120 marks: Questions 1-10 (95 marks) plus one essay from Question 11 (25 marks).

Marking guidance

- The number in brackets after each marking point is the mark it earns; the note above the points says how many are needed.
- Where a question asks for a fixed number of points (eg *two reasons*), credit only the first points given.
- Alternative correct wording and other relevant, accurate points are credited.
- For levels-of-response questions, first determine the level using the descriptors, then position the mark within the level using the indicative content.
- Calculations: credit correct working (method marks) even if the final answer is wrong; incorrect answers with no working score 0.

Textbook references: every question below cites the printed page numbers in R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018) from which the content is drawn, together with the AQA 7447 specification reference.

Question	Marking guidance	Marks
01.1	<i>A correct final answer (with unit where asked) scores full marks. If the answer is wrong, award 1 mark for each correct step of working shown.</i> • By-catch = $5.9 + 1.9 = 7.8$ t; total = 8.4 t (1) • $7.8 / 8.4 \times 100 = 92.9\%$ (accept 93%) (1)	2
01.2	<i>1 mark.</i> • The nets have a very small mesh size so few non-target animals can escape (1)	1
01.3	<i>1 mark for each point.</i> • They breed at an old age (skate about 11 years; Greenland shark 100-150 years) and produce few young (1) • They may reach a catchable size before they are sexually mature, so fishing can remove the whole breeding population before it has bred (1) • Populations recover slowly, unlike r-selected species (herring, mackerel) which breed young and produce many young; they are caught as by-catch in demersal trawls so fishing continues even though they have no commercial value (1)	3

Question	Marking guidance	Marks
01.4	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) • Impacts: population decline of k-selected species, by-catch (immature target fish, over-quota fish, non-commercial species; drift nets - whales, dolphins, turtles; longlines - albatrosses; pair trawls - porpoises), ghost fishing, seabed damage by demersal trawls, coral and seagrass damage, dynamite fishing, food-web effects • Catch quotas: limit weight landed; work well for single-species pelagic shoals but in mixed fisheries over-quota fish are discarded dead so the quota fails to protect them • Gear design: mesh size and square-mesh escape panels let small fish escape (diagonal mesh closes under drag); turtle exclusion devices; acoustic pingers for dolphins; curved hooks, decoys, sinkers and night setting to protect albatrosses; biodegradable / radio-tracked traps to reduce ghost fishing • Restricted effort: limits on boat size, engine power and days at sea; decommissioning; restricted methods - drift-net bans, demersal trawling bans on sensitive seabeds (Rockall Bank, Hatton Bank) • Evaluation: enforcement at sea is difficult; some methods reduce catch or raise costs; quotas need reliable stock data; effectiveness depends on species (single vs mixed shoals) and on international cooperation as fish migrate; as local stocks decline fishing moves to more distant waters • Justified conclusion weighing effectiveness of different methods	9
	Model answer - Level 3 (8 marks)	
	Fishing's impacts go beyond depleting target species: k-selected species that breed late and produce few young collapse and recover slowly; by-catch of immature target fish, over-quota fish and non-commercial species is thrown back dead, with drift nets killing whales, dolphins and turtles, longlines drowning albatrosses and pair trawls porpoises; lost gear keeps ghost fishing; demersal trawls destroy sea fans, corals and seagrass; and removing any species alters its competitors, prey and predators.	

Question	Marking guidance	Marks
	Catch quotas limit the weight landed and work well for single-species pelagic shoals where fishing stops when the quota is reached, but in mixed demersal fisheries over-quota fish are discarded dead, so the quota does not protect them. Gear design tackles by-catch directly: mesh sizes and square-mesh escape panels let small fish out because, unlike diagonal mesh, they do not close under drag; turtle exclusion devices, acoustic pingers for dolphins, curved hooks, decoys, sinkers and night setting protect turtles, cetaceans and albatrosses; biodegradable and radio-tracked traps limit ghost fishing. Restricting effort - boat size, engine power, days at sea, decommissioning - reduces total pressure, and banning drift nets or demersal trawling on sensitive seabeds such as Rockall Bank protects habitats.	
	Effectiveness depends on enforcement at sea, which is difficult, on reliable stock data for quotas, on the species and fishery type, and on international cooperation because fish migrate; some measures cut catches or raise costs, and as local stocks fall fleets simply move further away. Gear modifications are the most effective per unit cost because they reduce damage without reducing legal catch; quotas and effort limits only work where they are enforced.	
	Weaker answer - Level 2 (5 marks)	
	Fishing damages the environment by overfishing, catching by-catch such as dolphins and damaging the seabed with trawling. Quotas limit how many fish can be caught so stocks can recover. Bigger mesh sizes let small fish escape so they can grow and breed. Turtle exclusion devices let turtles escape from nets. Some fishing methods can be banned in sensitive areas. No-take zones let fish breed. These methods help but they are hard to enforce at sea and some fishermen ignore them.	
	What makes the difference	
	• Level 2 covers impacts and methods soundly but without the mechanisms (why square mesh works, why quotas fail in mixed fisheries).	
	• Level 3 evaluates each method on its own terms and then compares them; Level 2's evaluation is a single general sentence.	
	• Named examples (Rockall Bank, albatross measures, ghost-fishing controls) give Level 3 the range the top band expects.	
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 341-345; AQA spec 3.5.2.2.	
02.1	<i>1 mark.</i> • Development that meets the needs of the present generation without compromising the ability of future generations to meet their own needs (1)	1
02.2	<i>1 mark for each point.</i> • Negative feedback resists change / returns the system towards its previous state (1) • eg increased temperature increases cloud cover and albedo, or higher CO ₂ increases photosynthesis (1) • Positive feedback increases the change / makes it self-sustaining (1) • eg melting permafrost releasing methane, declining albedo as ice melts, forest and peat fires (1)	4

Question	Marking guidance	Marks
02.3	1 mark. Many species with a complex food web so the loss of one has a small effect; agroecosystems have few species and lack detritivores, pollinators and pest predators that maintain stability (1)	1
02.4	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) - Principles of natural systems: dynamic equilibria and negative feedback; diverse systems resistant to change; low energy-density renewable energy (solar-driven hydrological, carbon and nitrogen cycles); low-temperature processes using enzymes; material cycles with wastes as raw materials; small number of abundant elements; biodegradable non-toxic wastes - Human systems: non-renewable energy, high temperatures (Haber process, incineration, manufacturing), linear resource use, dispersal of minerals and mixtures that are hard to separate, persistent toxins (PCBs, organochlorines, toxic metals in electronics) - Applications: renewable energy and heat pumps, bioleaching, recycling and design for re-use, industrial symbiosis, organic and integrated agriculture, retaining hedgerows and pollinators, circular economy - Human responses that override feedback (fertilisers, irrigation, pesticides) and positive feedback triggered by human activity (permafrost, methane hydrates, fires); tipping points - Barriers: population size and affluence, attitudes of urban populations to distant resources, economics, need for international cooperation, existing infrastructure - Justified conclusion on how far natural principles can be applied	9
	Model answer - Level 3 (8 marks)	
	Natural systems have remained sustainable because they run on low energy-density renewable energy, mainly sunlight driving the hydrological, carbon and nitrogen cycles; their reactions occur at low temperatures because enzymes lower activation energies; materials cycle so the waste of one process is the raw material of the next; only a few abundant elements are used; and wastes are biodegradable and non-toxic. Negative feedback keeps them in dynamic equilibrium and high diversity makes them resistant to change.	

Question	Marking guidance	Marks
	Human systems contrast on every point: non-renewable fossil fuels, high-temperature processes such as the Haber process and incineration, linear use that disperses minerals and mixes materials that cannot be separated, and persistent toxins like PCBs and the metals in electronics. We also override negative feedback - when degraded soil reduces yields we add fertiliser, irrigation and pesticides - and trigger positive feedback such as permafrost melting.	
	Copying natural principles is already possible in many areas: renewable energy and heat pumps, bioleaching at low temperature, recycling and design for disassembly, industrial symbiosis, legumes and organic matter instead of manufactured nitrate, and retaining hedgerows for pest predators - the basis of the circular economy. The limits are population size and affluence, the attitude of urban consumers who never see depleted resources, cost, and the need for international cooperation. So the principles can make individual activities far more sustainable, but only a large-scale, cooperative shift in lifestyle would make the whole human system sustainable.	
	Weaker answer - Level 2 (5 marks)	
	Natural systems use renewable energy from the Sun and recycle materials, so nothing runs out and there is no waste. Humans use fossil fuels which will run out and produce waste that is toxic. Natural processes happen at low temperatures but the Haber process needs high temperatures. Humans could copy natural systems by using renewable energy like solar and wind, recycling more and using less fertiliser. This would make human activities more sustainable. However it would be expensive and people would have to change their lifestyle.	
	What makes the difference	
	• Level 2 states the contrast correctly but does not explain the mechanisms (enzymes, negative feedback, diversity, dispersal of minerals) that Level 3 uses.	
	• Level 3 applies the principles to specific technologies (bioleaching, heat pumps, industrial symbiosis, legumes); Level 2 stays at 'recycle more'.	
	• 'Evaluate the extent' - Level 3 distinguishes individual activities from the whole system in its conclusion; Level 2's evaluation is a single sentence about cost.	
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 370-374; AQA spec 3.6.1-3.6.3.	
03.1	<i>Correct answer only.</i> • Waste products that are the raw materials for other processes (1)	1
03.2	<i>1 mark for each point.</i> • Bacteria fix nitrogen at low temperature using enzymes and energy ultimately from sunlight (renewable) (1) • The Haber process needs high temperature and pressure powered by non-renewable fossil fuels, releasing CO ₂ (1)	2
03.3	<i>1 mark for each point.</i> • Incineration uses high temperatures / fuel and releases pollutants; nutrients are not returned as usable raw materials (1) • Decomposition at low temperature recycles nutrients as the raw materials for other natural processes (material cycle) (1)	2

Question	Marking guidance	Marks
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 373; AQA spec 3.6.2-3.6.3.	
04.1	<i>Correct answer only.</i> • B (1)	1
04.2	<i>Any 4 from the points below - 1 mark each, maximum 4.</i> • Deforestation (1) • Ploughing of grassland / expansion of farmland (1) • Reservoir creation / flooding (1) • Mineral extraction, especially open-cast mining (1) • Urban expansion (1)	4
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 20-26; AQA spec 3.1.2.2.	
05.1	<i>1 mark for each point.</i> • Stable tropical climate over long periods gave time for species to evolve and adapt to local abiotic and biotic factors; many rare species with small ranges (1) • High light, warm temperatures and high rainfall all year allow continuous photosynthesis and growth, supporting rich food webs with no seasonal food shortage (no hibernation / migration) (1) • Stratification (emergent, canopy, understorey, shrub, ground layers) and epiphytes create many niches; intense competition for light drives specialisation (1) • Biotic pressures (predation, inter-species relationships) drive evolution of specialised pollination / seed dispersal relationships; plants flower and fruit at different times (1)	4
05.2	<i>1 mark for each point.</i> • Adapted to stable conditions so they have narrow ranges of tolerance; a small change can move conditions outside the range (1) • Many species are rare with small ranges / small populations (1)	2
05.3	<i>1 mark for each point.</i> • No cold winter kills herbivores so plants defend themselves with toxic chemicals (alkaloids) that are medicinally active in controlled doses (quinine, diosgenin, curare) (1) • Animals without teeth / claws (frogs, reptiles) rely on chemical defences or venoms (fer-de-lance venom - captopril) (1)	2
05.4	<i>1 mark for each point.</i> • Over half the Amazon's rainfall comes from transpiration within the forest; clearance reduces downwind rainfall (1) • Rapid decay recycles nutrients so soils are shallow / poor; canopy and roots prevent erosion of the thin soil - clearance leads to rapid erosion and infertility (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 52-55; AQA spec 3.1.2.3.4.	
06.1	<i>1 mark for each point.</i> • Yield increases as inputs increase (1) • But each extra unit of input produces a smaller increase in yield than the previous one (the curve flattens / law of diminishing returns) (1)	2

Question	Marking guidance	Marks
06.2	<i>A correct final answer (with unit where asked) scores full marks. If the answer is wrong, award 1 mark for each correct step of working shown.</i> • Second unit: $70 - 50 = 20$ units (1) • Fifth unit: $94 - 90 = 4$ units (1)	2
06.3	<i>1 mark for each point.</i> • The first units of input on each area give the largest yield increases (1) • Concentrating inputs means later units add very little, whereas the same inputs spread thinly all work at the steep part of the curve (1)	2
06.4	<i>1 mark for each point.</i> • Some areas have much greater potential productivity so investment there is sensible (1) • Affluent societies can afford to buy more inputs while poorer communities that would benefit most cannot afford them (economic reasons) (1)	2
06.5	<i>1 mark for each point.</i> • Yield per unit area is very high because of large inputs (1) • But the extra yield per unit of input is small, so efficiency (yield per unit input / energy ratio) is low (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 324-325; AQA spec 3.5.1.2.	
07.1	<i>Any 2 from the points below - 1 mark each, maximum 2.</i> • Potato bean (N America): high protein content (1) • Morama bean (southern Africa): protein content similar to soya, drought resistant (1) • Yeheb tree (Somalia): edible nuts, drought resistant, grows on poor soils (1) • Perennial maize (Mexico) / Kernza: no need to buy seed each year, less ploughing so less soil erosion (1) • American bison: graze grassland too poor for cattle (1) • Common eland: browse and graze so can be kept with cattle without severe competition (1)	2
07.2	<i>1 mark for each point.</i> • Indigenous species are better adapted to local conditions such as climate, pests / diseases and soil (1) • So they have higher survival rates / productivity is not reduced, as it is when introduced species are kept in areas to which they are not adapted (1)	2
07.3	<i>1 mark.</i> • They grow in soils with a high salt content, so their genes could be used in breeding programmes to increase the salt tolerance of crops grown on soils salinised by irrigation (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 11; AQA spec 3.1.2.1.1.	
08.1	<i>1 mark for each point.</i> • Interconnected biological processes recycle nutrient elements so small amounts support life indefinitely without depletion; waste products of one process are raw materials of another so wastes do not build up (1) • Transpiration returned water vapour to the atmosphere increasing rainfall elsewhere (1) • Allowing more plant growth, further stabilising climate and soils (1)	3

Question	Marking guidance	Marks
08.2	<i>Any 3 from the points below - 1 mark each, maximum 3.</i> • Lack of ancient historical / instrumental data (1) • Reliance on proxy data (tree rings, pollen) of limited reliability for ancient conditions (1) • Lack of sophisticated equipment / inability to measure many factors (1) • Limited coordination between researchers; lack of data collection in many areas (oceans, poles) (1)	3
08.3	<i>1 mark for each point.</i> • The ratio of oxygen-18 to oxygen-16 in the ice indicates the temperature when the snow fell (1) • Air bubbles trapped in ice give the composition of the past atmosphere (CO₂, methane) (1) • Ratio of carbon-14 to carbon-12 dates organic material (eg pollen in sediments) (1) • More magnesium is incorporated into calcite at higher temperatures, so the ratio estimates past sea temperature (1)	4
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 8-9; AQA spec 3.1.1.2-3.1.1.3.	
09.1	<i>Correct answer only.</i> • B - radio waves do not travel through seawater; acoustic or satellite tags transmit position (1)	1
09.2	<i>2 marks for each point (partial credit for a partly correct point).</i> • Shows survival, dispersal and breeding of released individuals so success can be assessed and future releases improved (2) • Radio / GPS tags or collars; leg rings / wing tags on birds; camera traps; DNA from droppings to confirm identity and diet (2)	4
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 393-394, 402-407; AQA spec 3.1.2.3.5-3.1.2.3.6.	
10.1	<i>1 mark for each point.</i> • About 10% of land is already cultivated and 25% grazed (1) • Most remaining land is too dry, wet, cold, hot or steep, is covered in ice, or has no soil (1)	2
10.2	<i>1 mark for each point.</i> • Land becomes too expensive for local people to rent / is used to grow export crops for more affluent societies (1) • So people with less money lose access to the land and food they used to produce (1)	2
10.3	<i>1 mark for each point.</i> • It uses natural processes to supply nutrients (legumes, recycled organic matter, soil biota), control pests and provide ecological services (1) • So that pesticides and artificial fertilisers become unnecessary rather than just being banned (1)	2

Question	Marking guidance	Marks
10.4	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) • Unsustainable features: reliance on chemical pesticides (resistance, non-target impacts, bans), non-renewable rock phosphate, fossil-fuel-intensive nitrate manufacture, over-exploitation of rivers / groundwater, salinisation, loss of soil organic matter, greenhouse gases from fuel, rice and livestock, loss of species providing ecosystem services • Pest control: cultural methods (weeding, mulching, crop rotation, barrier crops, culling), biological control, predator habitats, polyculture / companion crops, integrated control, reduced antibiotic use • Nutrients: recycling organic matter, crop rotation, cultivation of legumes, conservation of soil biota, reduced artificial fertiliser especially nitrate, low tillage • Water: low water-use crops, reservoirs and aquifer recharge, drip irrigation rather than sprays • Soil and carbon: low-tillage farming to maintain organic matter, reduced machinery, renewable energy, drought-tolerant rice varieties so fields are drained earlier, high-carbohydrate ground feed for cattle reduces methane • Biodiversity: conserve habitats of crop wild relatives, seed banks, retain hedgerows, ditches, ponds, woodland; agri-environment schemes • Evaluation: growing population (perhaps 12 billion) needs more food; lower trophic level diets; obstacles - cost, need for support, gaining acceptance of new techniques; learning from past land degradation (fertile crescent, US mid-west) • Judgement: no single strategy sufficient; combination of ecological principles and new technology; justified conclusion	9
	Model answer - Level 3 (8 marks)	

Question	Marking guidance	Marks
	Present agriculture relies on inputs that cannot last: chemical pesticides to which pests evolve resistance and which harm non-target species, rock phosphate that is non-renewable, nitrate fertiliser made with fossil-fuel energy, and irrigation that over-draws rivers and groundwater and salinises soil, while ploughing loses soil organic matter and livestock, rice and machinery release greenhouse gases. Each can be replaced by strategies based on ecological principles. Pests are controlled culturally by weeding, mulching, rotation, barrier crops and culling, by biological control and predator habitats, by polyculture and companion crops, and by integrated control that uses chemicals only as a last resort. Nutrients come from recycled organic matter, rotation with legumes and conserved soil biota, so artificial fertiliser, especially nitrate, can be reduced. Water is saved with low water-use crops, reservoirs and aquifer recharge and drip irrigation; soil carbon is kept by low tillage and less machinery, with renewable energy for what remains; methane falls with rice varieties that allow earlier drainage and with ground, carbohydrate-rich cattle feed. Habitats of crop wild relatives, seed banks, hedgerows, ditches, ponds and woodland and agri-environment schemes protect the biodiversity that supplies these services.	
	Food demand will rise with a population that may reach 12 billion, there is little new land, and imports can reduce food availability where it is grown, so yield cannot be ignored. Some strategies cost more or need farmers to accept new techniques, and past land degradation from the fertile crescent to the American mid-west shows what happens when they are not adopted. No single strategy suffices; a combination of ecological principles with new technology is the only route to feeding more people without exhausting the resources food depends on.	
	Weaker answer - Level 1 (2 marks)	
	To make agriculture sustainable farmers should use fewer chemicals. Organic farming does not use pesticides or artificial fertilisers. Crop rotation and manure keep the soil fertile. Farmers should not waste water. Hedges should be kept for wildlife. This would make farming more sustainable but it might produce less food.	
	What makes the difference	
	• Level 1 gives general advice; Level 3 identifies each unsustainable feature and matches it with specific strategies, showing understanding of why they work.	
	• Level 3's evaluation considers population growth, land limits, cost, adoption and historical evidence; Level 1's is a single clause.	
	• Note that Level 3 avoids defining organic farming as 'not using chemicals' - the textbook stresses it is the positive use of natural processes.	
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 335-337; AQA spec 3.5.1.5.	

Question	Marking guidance	Marks
11.1	Fishing is the last large-scale human hunting activity. Discuss the environmental impacts of fishing and evaluate the methods used to make the exploitation of fish populations sustainable. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) • Marine productivity: light, nutrients, upwelling; where fisheries occur • Methods: demersal / pelagic trawling, purse seining, drift nets, long lines, shellfish traps - effectiveness, selectivity, energy, impacts • Impacts: overfishing (r vs k selection, catchable vs breeding size), changed age structure, by-catch (albatross, porpoise, turtles, shrimp), ghost fishing, seabed / reef / seagrass damage, dynamite, food web effects (sandeels, tritons) • Population dynamics and MSY: Russell formula, difficulties of data, indicators from catches (CPUE, mean size, age), research on eggs and larvae • Management: quotas, mesh design, escape panels, TEDs, pingers, hook design, effort limits, method bans, NTZs, closed seasons, size limits, protected individuals, population seeding, biodegradable gear; EU CFP, IWC, CCAMLR • Evaluation with examples (cod, orange roughy, herring recovery, NTZ spillover); role of aquaculture; judgement Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 338-351; AQA spec 3.5.2.	25

Question	Marking guidance	Marks
11.2	Discuss the methods that can be used to conserve biodiversity and evaluate their effectiveness. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) • Setting priorities: IUCN Red List criteria, EDGE, endemic, keystone, flagship species, population dispersal • Legislation and protocols: Wildlife and Countryside Act, designated areas (SSSI to MCZ), CITES Appendices, IWC, EU CFP, ITTO - successes and limits of enforcement • Captive breeding and release: selection, difficulties, techniques, release site criteria, hard / soft release, monitoring; seed banks • Habitat conservation: creation, design (area, shape, corridors, age structure, colonisation), abiotic and biotic management, plagioclimax maintenance, population control, reintroductions • Habitat case studies (woodland, rainforest, coral reefs, islands, mangroves, Antarctica) illustrating threats and conservation efforts • Monitoring and new technologies informing decisions • Evaluation: examples of success (Iberian lynx, Hawaiian goose, Scilly rats, Yellowstone) and failure (ITTO, tigers); cost, politics, climate change; judgement on the most effective combination Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 26-71; AQA spec 3.1.2.3.	25
	Model answer - Level 5 (23 marks)	
	Biodiversity is lost through direct exploitation, habitat destruction, introduced species, pollution and changes to abiotic factors, so its conservation needs methods that address each cause, at scales from a single species to international law. Their effectiveness varies with the threat, the species and the resources available.	

Question	Marking guidance	Marks
	Deciding what to conserve comes first. The IUCN Red List uses population size, area, decline rate and probability of extinction to prioritise species, and endemic, keystone, flagship and evolutionarily distinct species receive weight because their loss has wider consequences. This is effective at directing scarce resources, but it favours well-studied vertebrates over invertebrates and plants.	
	Legislation and agreements protect species and habitats. In the UK the Wildlife and Countryside Act, SSSIs, NNRs and Marine Conservation Zones, and in Europe SACs, SPAs and the Habitats Directive, restrict damaging activities; internationally CITES controls trade in threatened species, the IWC regulates whaling, the Ramsar Convention protects wetlands and the Rio Convention on Biological Diversity commits nations to conservation. Their effectiveness depends on enforcement: CITES has reduced trade in ivory and rhino horn but poaching continues where enforcement is weak; the IWC moratorium allowed some whale populations to recover but Japan and Norway continued whaling; designations protect against development but not against pollution or climate change crossing boundaries.	
	In-situ habitat conservation and management are the most fundamental methods because they keep species with the community and abiotic conditions they depend on. Management uses ecological understanding: maintaining plagioclimaxes such as chalk grassland and heathland by grazing or burning, coppicing and pollarding woodland, controlling water levels, light and grazing, designing reserves with the right area, shape, biological corridors and buffer zones, and creating new habitats such as wetlands in worked-out gravel pits. Case studies show it works - coppiced woodland restored dormice and butterflies, wetland creation increased bitterns - but reserves are often small and fragmented and need permanent funding.	
	Ex-situ methods rescue species for which in-situ protection has failed. Captive breeding faces problems of space, diet, breeding triggers, social groups and small gene pools, addressed with studbooks, artificial insemination, embryo transfer and cryopreservation; seed banks such as the Millennium Seed Bank store plant genetic diversity cheaply. The Arabian oryx, California condor and pool frog show captive breeding can bring species back, but only where protected habitat exists for hard or soft release with post-release monitoring, and it is expensive and works for few species.	
	Ecological monitoring - image databases, eDNA, tracking, satellites - measures whether any of this is working and has improved the targeting of effort. Overall, legislation is effective when enforced, habitat management is the most effective and widely applicable method, and captive breeding is a last resort that succeeds only in combination with habitat protection. The Living Planet Index continues to fall, however, which shows that conservation methods are winning individual battles while the drivers of loss - population, consumption, land-use change and climate change - remain unaddressed.	
	What makes the difference	
	The essay is structured by type of method and evaluates the effectiveness of each in turn, with named examples for both successes and failures.	

Question	Marking guidance	Marks
	• It draws on several specification areas (biodiversity, conservation legislation, habitat management, captive breeding, ecological processes, monitoring) - the synoptic breadth Level 5 requires.	
	• The conclusion ranks the methods and then steps back to the bigger picture (Living Planet Index, underlying drivers) - a supported judgement rather than a summary.	
	• A Level 3 essay would describe the same methods accurately but with fewer examples, little evaluation of enforcement or cost, and a conclusion that all methods are important.	