

A-level

ENVIRONMENTAL SCIENCE

Paper 2 - Generated practice paper Set 09

Mark scheme

120 marks: Questions 1-10 (95 marks) plus one essay from Question 11 (25 marks).

Marking guidance

- The number in brackets after each marking point is the mark it earns; the note above the points says how many are needed.
- Where a question asks for a fixed number of points (eg *two reasons*), credit only the first points given.
- Alternative correct wording and other relevant, accurate points are credited.
- For levels-of-response questions, first determine the level using the descriptors, then position the mark within the level using the indicative content.
- Calculations: credit correct working (method marks) even if the final answer is wrong; incorrect answers with no working score 0.

Textbook references: every question below cites the printed page numbers in R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018) from which the content is drawn, together with the AQA 7447 specification reference.

Question	Marking guidance	Marks
01.1	<i>Correct answer only.</i> • B - radio waves do not travel through seawater; acoustic or satellite tags transmit position (1)	1
01.2	<i>2 marks for each point (partial credit for a partly correct point).</i> • Shows survival, dispersal and breeding of released individuals so success can be assessed and future releases improved (2) • Radio / GPS tags or collars; leg rings / wing tags on birds; camera traps; DNA from droppings to confirm identity and diet (2)	4
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 393-394, 402-407; AQA spec 3.1.2.3.5-3.1.2.3.6.	
02.1	<i>1 mark for each point.</i> • k-selected: very slow growth, does not breed until 30, few eggs, so the population cannot replace losses (1) • Fish reach catchable size (20) before they breed (30) so fishing can remove the entire future breeding population (1) • Deep-water aggregations easily located and caught by trawling (1) • Recovery takes decades so fisheries have had to be closed (1)	4
02.2	<i>1 mark for each point.</i> • Herring breed at 2 years - the same age they become catchable - so there is always a breeding population and many young (1) • Cod are catchable at 2 but do not mature until 4, so heavy fishing removes most fish before they breed (1)	2
02.3	<i>1 mark for each point.</i> • Larger numbers of smaller / younger fish are caught, or bigger boats, nets and sonar increase catch effort (1) • Masks the decline of the breeding population until a sudden collapse (1)	2

Question	Marking guidance	Marks
02.4	<i>1 mark for each point.</i> - Neither is a target species but both are caught in demersal trawls; skate now extinct in the Baltic and critically endangered elsewhere (1) - Very late maturity (skate 11 years, Greenland shark 100-150 years, living over 400) and few young mean even low by-catch mortality exceeds replacement (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 341-342, 348-349; AQA spec 3.5.2.2.	
03.1	<i>1 mark for each point.</i> - Small clearings were farmed for a few years then abandoned as nutrients declined; the forest recolonised and soil fertility recovered before the area was cleared again (1) - Population growth shortens the gap between clearances so the forest and soil do not recover (1) - Clearings enlarge and degrade; less regrowth (1)	3
03.2	<i>1 mark for each point.</i> - Rainforest soils are thin and nutrient-poor (nutrients held in the biomass); once cleared, fertility falls in a few years and yields drop, so more land is cleared (1) - Abandoned land may not be recolonised if no forest survives nearby; erosion of exposed soil (1)	2
03.3	<i>1 mark for each point.</i> - Mining (aluminium, iron, gold, copper) destroys habitat directly and pollutes rivers with turbid drainage, cyanide and mercury from processing (1) - Reservoirs (eg Tucurui, 2850 km²) flood large areas, block migratory fish and river dolphins; decomposing vegetation produces acidic, deoxygenated water; fluctuating downstream flows (1) - Droughts increase frequency, area and duration of fires (long fires boil sap and kill trees); cloud forests lose humidity as clouds form higher so frogs and epiphytes decline; some models predict replacement by savannah (1)	3
03.4	<i>1 mark for each point.</i> - A creditor country or conservation charity cancels / buys part of a debtor country's foreign debt (1) - In return the country guarantees to protect an area of rainforest (eg USA - Guatemala, Brazil; WWF - Costa Rica, Philippines) (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 55-57; AQA spec 3.1.2.3.4.	
04.1	<i>1 mark for each point.</i> - Ecological footprint: the area of the Earth's surface (global hectares) needed to provide the resources used by a person / population / activity and to absorb the wastes produced (1) - Biocapacity: the biological productivity of an area (forests, cropland, grazing, fisheries, built land) - its ability to supply resources and absorb wastes, expressed in global hectares (1)	2
04.2	<i>1 mark for each point.</i> - A hectare of land / sea with the world-average biological productivity (1) - Biocapacities of real areas vary widely so a standard unit allows footprints and biocapacities of different places to be compared (1)	2

Question	Marking guidance	Marks
04.3	<i>1 mark for the correct answer.</i> • $8.1 / 1.7 = 4.8$ (accept 4.7-4.8) (1)	1
04.4	<i>1 mark for each point.</i> • Higher HDI is associated with a larger footprint (USA 8.1 vs Bangladesh 0.8), though the UK achieves the same HDI as the USA with about half the footprint (1) • Global footprint (2.8) exceeds global biocapacity (1.7) so humanity uses resources faster than they regenerate and CO₂ accumulates (overshoot) (1) • Earth Overshoot Day is the date each year when a year's worth of resources / waste absorption has been used up; it falls earlier each year (8 August in 2016) (1)	3
04.5	<i>Any 2 from the points below - 1 mark each, maximum 2.</i> • Renewable energy and energy conservation (carbon footprint is the largest component) (1) • Reduced meat consumption / food waste; local seasonal food (1) • Circular economy: repair, re-use, recycling; longer product lives (1) • Public transport, efficient buildings (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 379-380; AQA spec 3.6.4.2.	
05.1	<i>Up to 4 marks in total; award marks in proportion to the completeness of each point.</i> • Rotation: each crop has its own pests; growing a different crop each year (4-5 year cycle) means pests left at the end of a season die before that crop returns, so populations cannot build up early (up to 2) • Beetle banks / hedgerows provide habitat and overwintering sites for natural predators (ground beetles, ladybirds) that eat aphids (up to 2) • Rotation also evens out nutrient demands and includes legumes (up to 2)	4
05.2	<i>1 mark for each point.</i> • Integrated control gives almost the same yield as chemical control (18.1 vs 18.5) with far fewer applications (1-2 vs 6) (1) • It retains most of the biodiversity of cultural control (26 vs 31 species) whereas chemical control alone leaves only 8 (1) • Cultural control alone loses about 12% yield; integrated control combines cultural methods, resistant varieties and carefully timed specific non-persistent pesticides only when essential - best balance of yield, cost and impact (1)	3
05.3	<i>1 mark for each point.</i> • Large numbers of males are sterilised with gamma radiation and released (1) • Females of many insects mate only once and store sperm, so mating with a sterile male produces no offspring; enough sterile males reduce fertile matings below replacement level (screw worm fly, Mediterranean fruit fly) (1) • Only the target species' females mate with them so non-target species are unaffected; fails if sterilised males behave abnormally or re-colonisation occurs (1)	3
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 314-318; AQA spec 3.5.1.1.	

Question	Marking guidance	Marks
06.1	<i>1 mark for each point.</i> • Timber: building construction, furniture, tools, telegraph poles, concrete shuttering (oak, beech, mahogany, teak, conifer softwood) (1) • Fuel: fuelwood and charcoal - still the main energy source for most people in LEDCs (1) • Food: fruit, nuts, bushmeat, livestock fodder; crops of forest origin (coffee, cacao, bananas, Brazil nuts); pigs and chickens were forest animals (1) • Fibres: paper (wood pulp cellulose), cotton, viscose / rayon (1) • Medicines: quinine (cinchona), taxol (yew), aspirin (willow); most species not yet researched (1)	5
06.2	<i>1 mark for each point.</i> • Wood is mainly cellulose which is hard to digest, so carbon stored in wood has a long residence time; total forest carbon (vegetation plus soil) is about double that in the atmosphere (1) • Growing trees sequester CO₂ by photosynthesis and release oxygen; forest soils hold more carbon than the trees (especially boreal) (1) • Planting / protecting forests removes CO₂ and reduces the enhanced greenhouse effect (1)	3
06.3	<i>1 mark for each point.</i> • Highest transpiration of any ecosystem increases downwind rainfall; interception increases evaporation; deep organic soils retain water and delay runoff so river flow is more even and flooding reduced (1) • Low albedo absorbs sunlight; heat stored and released at night reduces day-night temperature extremes; canopy microclimate: lower light, wind, higher humidity (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 360-364; AQA spec 3.5.3.1.	

Question	Marking guidance	Marks
07.1	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) • Why needed: growing population (to 12 billion), land degradation since the fertile crescent, reliance on artificial fertilisers, pesticides and unsustainable irrigation; limits to new land and imports • Pest control: cultural methods (weeding, mulching, rotation, barrier / companion crops, predator habitats, biological control, sterile males, pheromones), resistant varieties, integrated control, reduced antibiotics • Nutrients: natural processes - nitrogen-fixing legumes, decomposition, recycling organic matter (manure, compost), crop rotation, permaculture, conservation of soil biota and mycorrhizae, reduced artificial fertiliser; phosphate recycling • Energy: natural processes instead of Haber-process fertiliser, low tillage, less machinery, renewable energy, low food miles, reduced methane (cattle diets, rice varieties) • Water: low water-use crops, drip irrigation, reservoirs and aquifer recharge, avoiding saline water; soil: organic matter maintenance, erosion control • Biodiversity: retaining hedgerows, ditches, ponds, woodlands; CWR conservation and seed banks; agri-environment schemes • Social: control of impacts on rural communities, Fairtrade, local food; consumer choices • Evaluation: yields vs sustainability, cost, need for policy support; judgement	9
	Model answer - Level 3 (8 marks)	
	Sustainability matters because the population may reach 12 billion, land has been degraded since farming began in the fertile crescent, and current yields depend on artificial fertilisers, pesticides and irrigation that draws down groundwater, with little new land to cultivate. Pest control can move from chemicals to cultural methods - weeding, mulching, rotation, barrier and companion crops, habitats for predators - biological control, sterile males and pheromones, resistant varieties and integrated control, with fewer antibiotics in livestock.	

Question	Marking guidance	Marks
	Nutrients can come from natural processes: legumes fixing nitrogen, recycling manure and compost, rotation, permaculture and conserving soil biota and mycorrhizae, reducing manufactured nitrate and the non-renewable rock phosphate it accompanies. Energy use falls with low tillage, less machinery, renewable power, low food miles and lower methane from cattle diets and drought-tolerant rice. Water is conserved with low water-use crops, drip irrigation, reservoirs and aquifer recharge and by avoiding saline water; soil by maintaining organic matter and controlling erosion. Retaining hedgerows, ditches, ponds and woodland, conserving crop wild relatives in seed banks and agri-environment schemes protect the biodiversity that supplies pollination and pest control, while Fairtrade, local food and consumer choices address the social side.	
	The tension is between yield and sustainability: some strategies cost more or yield less in the short term and need policy support to be adopted. But since the unsustainable inputs themselves depend on finite fossil fuels and phosphate, the strategies that use natural processes are the only ones that can feed a larger population indefinitely.	
	Weaker answer - Level 2 (4 marks)	
	Agriculture can be made more sustainable by using less pesticides and using biological control instead, using manure instead of artificial fertilisers, growing legumes which add nitrogen to the soil, and using crop rotation. Drip irrigation saves water. Farmers can keep hedgerows to provide habitats for wildlife. Organic farming uses these methods. These strategies reduce pollution and keep the soil healthy so food can be produced in the future.	
	What makes the difference	
	• Level 2 lists valid strategies but does not explain why they are needed (population growth, land degradation, dependence on finite inputs) or group them by the problem they solve.	
	• Level 3 covers energy, soil, biodiversity and social strategies as well as pests, nutrients and water.	
	• Level 3 discusses the yield-sustainability tension and the need for policy support; Level 2 has no evaluation.	
07.2	<i>1 mark.</i> • Agricultural systems designed to mimic natural ecosystems - perennial crops, mixed species, recycling of nutrients, minimal external inputs (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 335-337; AQA spec 3.5.1.5.	

Question	Marking guidance	Marks
08.1	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) • Context: wild catches have not increased since about 1990 as stocks are fully or over-exploited; aquaculture output growing rapidly and approaching the wild catch; fish are the last large-scale hunted food • Advantages of aquaculture: control of limiting factors, selective breeding, gender control, high yields per area; fish have low FCRs (no thermoregulation / support) compared with livestock; herbivores (tilapia, carp) and molluscs very efficient; polyculture, IMTA, aquaponics, rice-fish • Limits: carnivorous species (salmon, shrimp) need fishmeal from wild fish so still depend on fishing and deplete forage fish (sandeels - puffins); some species (bluefin tuna) will not breed in captivity; consumer preference for carnivores • Environmental impacts: habitat loss (mangroves), organic pollution, pesticides, antibiotics and resistance, lice, escapes and gene pool contamination, predator control; and how they can be reduced • Fishing can be sustainable with quotas, gear design, NTZs, size limits, effort limits; MSY management needs good data • Judgement: aquaculture can supply much of the growth in demand, especially herbivores and shellfish, but cannot fully replace fishing; both need sustainable management	9
	Model answer - Level 3 (9 marks)	
	Wild catches have not increased since about 1990 because most stocks are fully or over-exploited, while aquaculture has grown rapidly and now approaches the wild catch, so the question is how far it can take over. Its advantages are real: limiting factors, breeding, gender and disease can be controlled, yields per area are high, and fish have low food conversion ratios because they need not support their weight or keep warm; herbivores such as tilapia and carp and filter-feeding molluscs need no fishmeal at all.	

Question	Marking guidance	Marks
	Its limits are equally real. The carnivorous species consumers prefer - salmon, shrimp - are fed fishmeal from wild forage fish, so their farming still depends on fishing and depletes species such as sandeels on which puffins rely; some species, such as bluefin tuna, will not breed in captivity. Farms destroy mangroves, release organic waste, pesticides and antibiotics that breed resistance, spread lice, allow escapes that contaminate wild gene pools and provoke predator control, though siting, closed systems, polyculture and vaccines can reduce these. Meanwhile fishing itself can be sustainable if quotas based on good stock data, gear design, size and effort limits and no-take zones are enforced.	
	Aquaculture can therefore supply most of the growth in demand, especially through herbivorous fish and shellfish, and can relieve pressure on wild stocks, but it cannot fully replace fishing while carnivorous species depend on wild fishmeal and some species cannot be farmed. Both must be managed sustainably; neither is a substitute for the other.	
	Weaker answer - Level 1 (2 marks)	
	Aquaculture is farming fish instead of catching them. Fish farms can produce a lot of fish and they do not run out like wild fish do. But fish farms cause pollution and diseases can spread to wild fish. Salmon farms feed the salmon on fish which have to be caught. So aquaculture can replace some fishing but not all of it.	
	What makes the difference	
	• Level 1 reaches a reasonable conclusion but supports it with two sentences; Level 3 gives the context (flat wild catch since 1990), the advantages in detail and the impacts with ways to reduce them.	
	• Level 3 distinguishes between species types (herbivores and molluscs vs carnivores) - the key to 'to what extent'.	
	• Level 3 also considers whether fishing can be made sustainable, which the comparison requires.	
08.2	<i>1 mark.</i> • A species that is given artificial food (pellets) rather than relying on natural food in the water, eg salmon, shrimp (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 338-359; AQA spec 3.5.2.	
09.1	<i>1 mark for each point.</i> • The new opportunity is overwhelmingly attractive (economic benefit) (1) • Lack of knowledge / understanding of the impacts (1) • The impacts occur elsewhere or much later, so are not evident to those who benefit (1)	3
09.2	<i>1 mark for each point.</i> • Cultural belief that resources should be exploited to satisfy needs only (1) • Problems of over-exploitation become obvious quickly as local supplies dwindle, so behaviour changes (1)	2
09.3	<i>1 mark for each point.</i> • Resources are brought in from other areas (1) • If one area is depleted, exploitation moves to another area (eg timber, fish, metals, energy) without the consumers noticing (1)	2

Question	Marking guidance	Marks
09.4	1 mark for each point. - Easier access to energy has made many new activities possible (1) - The scale of activities has greatly increased (1) - Toxic wastes are produced that cannot be decontaminated by natural processes (1)	3
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 370-371; AQA spec 3.6.1.	
10.1	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) - Resources: wood, fibres (cotton, wool, silk), oils, fuels (wood, biofuels), food; new food species (morama bean, perennial cereals, eland, bison) better adapted to local conditions - Biomimetics: vehicle design, architecture, structures, adhesion, materials, ultrasound - Medicines from plants and animals (examples); most species unstudied - argument for habitat conservation - Physiological research using wildlife species; wildlife as pest control agents (ladybirds, <i>Encarsia</i>, <i>Cactoblastis</i>, Bt) - Genetic resources: crop wild relatives, Vavilov centres, gene pools - Ecosystem services: atmospheric composition, hydrological and biogeochemical cycles, soil maintenance, pollination, seed dispersal, habitat provision - Ethical / aesthetic / recreational arguments; the unknown value of species not yet investigated (precautionary principle) - Evaluation: which arguments are most persuasive to decision makers; economic valuation vs intrinsic value; judgement	9
	Model answer - Level 3 (8 marks)	
	Human society depends on other species first as resources: wood, fibres such as cotton, wool and silk, oils, fuels and, above all, food, with wild species such as the morama bean, perennial cereals, eland and bison offering new crops and livestock better adapted to local conditions. Biomimetics copies adaptations into vehicle design, architecture, structures, adhesives, materials and ultrasound; medicines such as taxol, quinine and AZT came from plants and animals, and physiological research on marsupials, hippos and sponges continues to yield medical insights. Wild species are pest-control agents - ladybirds, <i>Encarsia</i>, <i>Cactoblastis</i>, <i>Bacillus thuringiensis</i> - and crop wild relatives in Vavilov centres hold the genes future crops will need.	

Question	Marking guidance	Marks
	Less visibly, other species provide the services that keep the planet habitable: regulating atmospheric composition, driving the hydrological and biogeochemical cycles, maintaining soil, pollinating crops, dispersing seeds and providing habitat. Most species have never been investigated, so their value is unknown, and there are ethical, aesthetic and recreational reasons for conservation that owe nothing to usefulness.	
	Which arguments persuade decision-makers? Economic valuation of resources and services speaks the language of policy and has been most effective, but it undervalues the services that go unnoticed because they operate in dynamic equilibrium and it can never capture the value of species not yet studied. The strongest case combines the precautionary logic of unknown value with the intrinsic argument: conserving biodiversity protects both what we know we need and what we cannot yet know.	
	Weaker answer - Level 1 (3 marks)	
	Biodiversity is important for humans in many ways. We get food, wood and cotton from plants and animals. Many medicines come from plants. Bees pollinate our crops so we need to conserve them. Animals and plants are also nice to look at and people like to visit nature. We should conserve biodiversity because we might need species in the future.	
	What makes the difference	
	• Level 1 names categories without examples; Level 3 gives specific, accurate examples in every category.	
	• Level 3 includes ecosystem services, genetic resources and pest control that Level 1 omits.	
	• 'Discuss' at the top level means weighing the arguments against each other - Level 3 evaluates which arguments persuade and why; Level 1 does not.	
10.2	<i>1 mark.</i> • The variety and abundance of life (species, genes, habitats) in an area / on Earth (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 10-19; AQA spec 3.1.2.1.	

Question	Marking guidance	Marks
11.1	Discuss the extent to which agriculture can be made sustainable while feeding a growing human population. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) • Need: population possibly 12 billion; humans raise carrying capacity; trophic level of diet • Impacts to be reduced: habitat loss, drainage, nutrient enrichment, introduced species, pesticide and nutrient pollution, nitrate toxicity, greenhouse gases, hydrological changes, soil erosion • Social, economic and political influences: consumer choices, technology availability, grants, guaranteed markets, surpluses, quotas, set-aside, agri-environment schemes • Strategies: cultural and biological pest control, integrated control, organic nutrient supply, legumes, low tillage, water conservation, CWR conservation, renewable energy, methane reduction, habitat retention • Limits: little new land, imports reducing local availability, cost and acceptance of new techniques, past land degradation • Justified conclusion Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 328-337; AQA spec 3.5.1.3-3.5.1.5.	25

Question	Marking guidance	Marks
11.2	Discuss the causes and consequences of deforestation and evaluate the extent to which forests can be exploited sustainably. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) • Importance: resources (timber, fuel, food, fibres, medicines) and ecosystem services (atmospheric regulation, carbon store, hydrological regulation, climate, soil conservation, habitat, recreation) • Causes: agriculture (subsistence and cash crops), mineral extraction, reservoirs, urbanisation, transport, unsustainable timber, fuel and fodder exploitation above MSY; trends by region • Consequences: loss of resources, biodiversity and species, fragmentation; hydrological changes; soil impacts; climate impacts (albedo, carbon, rainfall) • Productivity vs biodiversity: non-indigenous species, monocultures, close planting, simple age structure, clear-felling • Sustainable management: MSY harvest rates, mixed indigenous species, mixed age structure, selective logging, traditional coppicing / pollarding, FSC certification, community involvement, protected areas, debt-for-nature, afforestation programmes (China); Rio Bravo case study • Evaluation: economics of slow-growing hardwoods, enforcement, demand from MEDCs, ITTO failures; judgement Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 50-57, 360-369, 386-387; AQA spec 3.5.3, 3.1.2.3.4.	25