

A-level

ENVIRONMENTAL SCIENCE

Paper 2 - Generated practice paper Set 19

Mark scheme

120 marks: Questions 1-10 (95 marks) plus one essay from Question 11 (25 marks).

Marking guidance

- The number in brackets after each marking point is the mark it earns; the note above the points says how many are needed.
- Where a question asks for a fixed number of points (eg *two reasons*), credit only the first points given.
- Alternative correct wording and other relevant, accurate points are credited.
- For levels-of-response questions, first determine the level using the descriptors, then position the mark within the level using the indicative content.
- Calculations: credit correct working (method marks) even if the final answer is wrong; incorrect answers with no working score 0.

Textbook references: every question below cites the printed page numbers in R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018) from which the content is drawn, together with the AQA 7447 specification reference.

Question	Marking guidance	Marks
01.1	<i>A correct final answer (with unit where asked) scores full marks. If the answer is wrong, award 1 mark for each correct step of working shown.</i> • Extensive cereal 5.0; intensive cereal 3.0 (1) • Intensive beef 0.1 (1)	2
01.2	<i>1 mark for each point.</i> • Intensive cereal is most productive per hectare (60 GJ) but less efficient (3.0) than extensive (5.0) (1) • Each extra unit of input gives a smaller extra yield (law of diminishing returns) (1) • Beef has very low productivity and efficiency because energy is lost at an extra trophic level (1)	3
01.3	<i>Up to 3 marks in total; award marks in proportion to the completeness of each point.</i> • Any input that increases productivity but requires the use of energy (usually fossil fuel) (up to 2) • Manufacture of nitrate fertiliser (Haber process) and pesticides; fuel for machinery; pumping irrigation water; grain drying; food processing and transport; embodied energy of machinery (up to 2)	3
01.4	<i>1 mark for each point.</i> • Because of diminishing returns, the first units of input on unfertilised land give the largest yield increases (1) • Moving inputs from intensively farmed land (where extra input adds little) to extensively farmed land gives a greater total yield - though land, cost and distribution limit this (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 324-327; AQA spec 3.5.1.2.	

Question	Marking guidance	Marks
02.1	<i>2 marks for each point (partial credit for a partly correct point).</i> - Tracking: continuous data on movements, territory size, migration routes and habitat use without disturbance, over large areas and long periods; acoustic tags in seawater; geolocators for small birds; more representative than occasional sightings (2) - DNA: identifies individuals, gene pool size / inbreeding and regional populations (tracing ivory, timber, fish); eDNA detects rare or hidden aquatic species (great crested newt, invasive carp) from shed cells without capture (2) - Image recognition: software identifies individuals from unique markings (tiger stripes, whale shark spots, fin damage) in camera-trap / drone / satellite images, giving population size, lifespan, social structure from large image sets quickly and objectively (2) - Acoustic: bat detectors, hydrophones and sonograms record species that are hard to see (bats, cetaceans, insects), giving presence, abundance and activity continuously, day and night (2)	8
02.2	<i>1 mark for each point.</i> - Data collected continuously / over large areas / at night rather than at a few times and places a researcher can visit (1) - Less disturbance so animals behave naturally; larger sample sizes; objective rather than subjective identification (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 402-406; AQA spec 3.1.2.3.6.	
03.1	<i>1 mark for each point.</i> - Coordinating global data on biodiversity conservation (Red List) (1) - Increasing understanding of the importance of biodiversity (1) - Deploying nature-based solutions to global challenges in climate, food and sustainable development (1)	3
03.2	<i>1 mark for each point.</i> - Decline = $(12\,000 - 2100) / 12\,000 \times 100 = 82.5\%$ (1) - Decline over 80% meets the Critically Endangered criterion even though habitat area (400 km²) and adults (2100) only meet Endangered (1) - A species is placed in the highest category for which it meets any one criterion, so Critically Endangered (1)	3
03.3	<i>1 mark for each point.</i> - Categorising a species requires a great deal of research (population, range, trends) and resources; about 85 000 of millions of species assessed (1) - Key species in habitats under particular threat (eg lemurs indicating the state of Madagascar's forests) (1) - Evolutionarily unique species / EDGE species; endemic species; keystone species; flagship species; species with fragmented populations (1)	3
03.4	<i>1 mark.</i> Iberian lynx: Critically Endangered to Endangered after captive breeding and protection of two habitat areas (population trebled in 15 years); Fregate Island beetle after rat eradication (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 26-27; AQA spec 3.1.2.3.1.	

Question	Marking guidance	Marks
04.1	<i>1 mark for each point.</i> • Giant tritons are predators of the crown-of-thorns starfish which eats coral polyps (1) • Removing tritons allowed starfish populations to explode, destroying large areas of reef (food web / trophic cascade) (1)	2
04.2	<i>1 mark for each point.</i> • Fixed mooring buoys so boats do not anchor on coral (1) • Education of divers / snorkellers not to touch coral or kick up sediment; bans on spear fishing and souvenir collection (1) • Fishing controls: no-take zones (Chagos, Great Barrier Reef, Lundy, Lamlash Bay), seasons, minimum and maximum catch sizes, bans on dynamite / small-mesh nets / trawling (1) • Protection of turtle nesting beaches; control of coastal development; protection of mangroves and seagrass; artificial reefs (reef balls, sunk ships / oil rigs) (1)	4
04.3	<i>1 mark for each point.</i> • No sunlight so no symbiotic algae; polyps rely only on captured food so have far less energy and grow very slowly (1) • Damage from trawls takes hundreds of years to repair; the slow-growing, low-fecundity fish (orange roughy) they support are easily over-exploited (1)	2
04.4	<i>1 mark for each point.</i> • Deep-water trawling (also by-catch); oil and gas exploration in deeper water; ocean acidification (1) • Protected areas banning trawling, eg Darwin Mounds SAC off NW Scotland, Oculina Bank off Florida (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 61-64; AQA spec 3.1.2.3.4.	
05.1	<i>2 marks for each point (partial credit for a partly correct point).</i> • Normal diagonal mesh closes as the net fills / stretches so small fish cannot escape; square-mesh panels do not close so juveniles escape to grow and breed; TEDs let turtles out; larger mesh sizes generally (2) • Minimum size lets fish reach breeding age before capture (eg UK lobster minimum carapace above breeding size, cod 35 cm); maximum size / V-notching egg-carrying females protects large, highly fecund breeding adults (Hebridean lobsters over 145 mm, Nassau grouper) (2) • NTZs (Lundy, Lamlash Bay, Chagos, Great Barrier Reef) protect breeding populations and habitats; surplus young spill over and increase surrounding catches; damaged communities recover (2) • Closed seasons, ideally in the breeding season, let fish grow and breed; fishermen may oppose because shoaling breeders are easy to catch (2)	8
05.2	<i>1 mark for each point.</i> • Pelagic species form single-species shoals so fishing stops when the quota is reached (1) • In mixed fisheries many species are caught together; once one species' quota is reached it must be discarded (dead) while fishing for others continues (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 344-348; AQA spec 3.5.2.2.	

Question	Marking guidance	Marks
06.1	<i>1 mark for each point.</i> • Trees (hazel, oak) are cut to ground level every 3-30 years and allowed to regrow from the stump (1) • About 0.5 ha cut each year on a rotation (eg 8 years) so the wood is a patchwork of areas of different ages (1) • Each age has its own community: newly cut areas get high light for wildflowers and butterflies; older areas give dense cover for nesting birds / dormice (1) • Pollarding cuts above browsing height so deer / livestock cannot eat the regrowth (1)	4
06.2	<i>1 mark for each point.</i> • Wolves (top predator, exterminated by 1926) reduced deer populations and changed their behaviour (1) • Fewer young trees eaten so trees and riverside vegetation recovered (1) • More trees allowed beaver populations to increase; beaver dams increased wetland area (1) • Wetland species (birds, fish, invertebrates) became more common; overall biodiversity rose, though wolves cause problems for livestock farmers (1)	4
06.3	<i>1 mark for each point.</i> • Undesirable species (introduced predators / competitors, or natural succession species such as scrub, conifers in the New Forest heathland) reduce survival of the target species (1) • Removal restores suitable conditions (eg heathland for sand lizards and smooth snakes) or reduces predation / competition (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 47-49; AQA spec 3.1.3.3-3.1.3.4.	
07.1	<i>Up to 6 marks in total; award marks in proportion to the completeness of each point.</i> • Organic advantages: often waste products, locally available and cheap; add humus improving structure and water retention; increase soil biota; slow release reduces leaching (up to 2) • Organic disadvantages: nutrient composition cannot be controlled; released slowly so needs long-term planning; bulky with high water content so expensive to transport; cannot be applied to a growing crop; odour / pathogens (up to 2) • Inorganic advantages: composition controlled to match crop needs; nutrients released rapidly; concentrated and easy to apply (up to 2) • Inorganic disadvantages: high energy (fossil fuel) manufacture (Haber process); no organic matter so humus and biota decline; some toxic to worms; high solubility so leached causing eutrophication; raw materials (phosphate rock) finite (up to 2)	6

Question	Marking guidance	Marks
07.2	<i>Up to 4 marks in total; award marks in proportion to the completeness of each point.</i> • Crops grown in a circulating nutrient solution (inert medium) in greenhouses; nutrient concentrations monitored and adjusted so nutrients are never limiting; abiotic factors controlled (up to 2) • Roots are small (all in contact with nutrients) so more growth goes into the harvested crop; no soil-borne pathogens or weeds; crop stays fresh with roots (up to 2) • Disadvantages: high inputs of nutrients and energy (heating, lighting, pumping); high level of technical knowledge; only economic for high-value crops (up to 2)	4
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 310-312; AQA spec 3.5.1.1.	
08.1	<i>Any 2 from the points below - 1 mark each, maximum 2.</i> • Hedgerows (1) • Hay meadows (1) • Grazed moorland / heathland (1) • Chalk grassland (1)	2
08.2	<i>1 mark for each point.</i> • Drainage produces more aerobic soils so wetland plants and the animals they support cannot survive (1) • Fertilisers increase growth of tall species such as grasses which out-compete smaller plants such as wildflowers (1)	2
08.3	<i>1 mark.</i> • The introduced (non-indigenous) predator may eat other species as well as the pest (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 328-329; AQA spec 3.5.1.3.	
09.1	<i>1 mark for each point.</i> • Negative (1) • Positive (1) • Negative (1) • Positive (1)	4
09.2	<i>1 mark for each point.</i> • The response to a change acts to re-establish / return the system to the previous equilibrium (1) • So the change does not become self-sustaining unless a large unusual event overwhelms the negative feedback (1)	2
09.3	<i>1 mark for each point.</i> • eg the possible release of methane hydrate at the end of the Permian era (1) • Overwhelmed the negative feedback mechanisms, causing climate change and a mass extinction / a new equilibrium (1)	2
09.4	<i>1 mark for each point.</i> • Soil degradation, reduced water availability and loss of natural pest predators would reduce yields (negative feedback curtailing the activity) (1) • Instead of reducing the activity, farmers use fertilisers, irrigation and pesticides, which further damage the natural systems that support food production (1)	2

Question	Marking guidance	Marks
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 371-372; AQA spec 3.6.1.	
10.1	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) • Succession: pioneers, seres (hydrosere, lithosere, psammosere), modification of abiotic conditions, climax communities controlled by climate; deflected succession and plagioclimaxes • Application: deciding whether to manage (plagioclimax habitats need grazing, mowing, burning, coppicing, pollarding continued) or leave alone (climax habitats); habitat creation planning the stages; controlling scrub / secondary succession; understanding that new habitats change over time • Population dynamics: birth and death rates, density-dependent and independent factors, carrying capacity, sigmoid growth, predator-prey cycles, r- and k-selection, MSY • Application: setting sustainable harvest levels (fisheries, whaling, forestry); recognising k-selected species need strict protection and recover slowly; deciding when culling is needed (deer without wolves, introduced species) or when releases / captive breeding are needed (low breeding rate); predicting effects of reintroducing predators • Monitoring population size, distribution, survival rate and age structure to assess success; adaptation and range of tolerance guiding abiotic management • Evaluation: uncertainties (natural fluctuations, unknown carrying capacity); examples (Yellowstone wolves, Scottish deer, chalk grassland); judgement	9
	Model answer - Level 3 (8 marks)	
	Succession is the sequence by which pioneers colonise bare ground, modify the abiotic conditions and are replaced by later seres until a climax community controlled by climate is reached; where grazing, mowing, burning or coppicing halts it, a plagioclimax forms. This tells a conservation manager whether to intervene: chalk grassland, heathland and coppiced woodland are plagioclimaxes that disappear without continued management, whereas a climax habitat is best left alone, and habitat creation must plan the stages, control scrub and manage secondary succession.	

Question	Marking guidance	Marks
	Population dynamics - birth and death rates, density-dependent and independent factors, carrying capacity, sigmoid growth, predator-prey cycles and r- and k-selection - inform decisions of a different kind. Maximum sustainable yield sets harvest levels in fisheries, whaling and forestry; k-selected species that breed late and slowly need strict protection and recover slowly; culling may be justified where predators are absent, as with Scottish red deer without wolves, or where introduced species threaten natives, while the reintroduction of wolves to Yellowstone shows how restoring a predator changes vegetation through the food web. Monitoring population size, distribution, survival and age structure measures success, and knowledge of adaptation and ranges of tolerance guides how abiotic conditions are managed.	
	The limits are uncertainty: natural fluctuations, unknown carrying capacities and the difficulty of predicting how communities respond. Ecological understanding cannot remove that uncertainty, but without it conservation decisions - to manage or not, to cull or protect, how much to harvest - are guesses.	
	Weaker answer - Level 2 (5 marks)	
	Succession is when a habitat changes over time from pioneer species to a climax community. Conservationists need to understand this because some habitats like heathland will turn into woodland if they are not managed by grazing or burning. Population dynamics is about how populations grow and are limited by factors like food and predators. This helps conservation because you can work out how many animals can be taken without the population declining, and when a population needs protecting. Monitoring populations shows if conservation is working.	
	What makes the difference	
	• Level 2 explains both concepts correctly and links them to decisions, but generally; Level 3 uses the technical vocabulary (seres, plagioclimax, k-selection, MSY) with named cases (Yellowstone wolves, Scottish deer, chalk grassland).	
	• Level 3 covers more applications (habitat creation, culling introduced species, adaptation and tolerance) and evaluates the uncertainties.	
	• Structure: Level 3 gives each concept its own paragraph with the decisions it informs, then a conclusion.	
10.2	<i>1 mark.</i> • Any taxonomic group of organisms classified together by similarity (eg species, genus, family) (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 72-85; AQA spec 3.1.3.	

Question	Marking guidance	Marks
11.1	Explain how the physical features of the Earth and the activities of living organisms have combined to create and maintain conditions suitable for life. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) • Physical features: mass / gravity retaining an atmosphere and giving pressure for liquid water; distance from the Sun and the Goldilocks zone; speed of rotation reducing temperature extremes; tilt producing seasons; magnetic field deflecting the solar wind; the Moon stabilising the orbit and producing tides • Importance of liquid water: solvent, transport, temperature control, anomalous expansion, high specific heat capacity, habitats, absorption of UV before the ozone layer • Development of life: Archaea at vents; photosynthesis; oxygen reacting with iron then accumulating; ozone layer allowing colonisation of land; aerobic organisms • Life changing conditions: carbon sequestration offsetting solar brightening, biogeochemical cycles, transpiration and rainfall, regulation of atmospheric gases • Methods of studying past conditions: proxy data, isotope ratios, ice cores, Mg:Ca ratios; understanding of continental drift and ocean currents • Links to current threats (climate change, ozone depletion) and a justified conclusion about the interdependence of physical and biological factors Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 5-9; AQA spec 3.1.1.	25

Question	Marking guidance	Marks
11.2	Discuss how an understanding of energy flow and the manipulation of food species can be used to increase food production sustainably. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) • Energy flow: autotrophs and heterotrophs; losses by reflection, transmission, respiration; 100 000 to 1 000 units; few food chains beyond four levels; plant food most productive; grazing animals where crops impossible; pigs on wastes • Energetics: intensive vs extensive, law of diminishing returns, energy subsidies (fertiliser, pesticides, irrigation, machinery, drying, processing, transport), energy ratios, food conversion ratios, dependence on fossil fuels • Control of genetics: asexual reproduction / cuttings / cloning, selective breeding and breeds, cross-breeding and hybrid vigour, AI and embryo transfer • GM: Roundup Ready soya, Bt crops, golden rice, omega-3 oilseed rape; advantages; risks (gene transfer, antibiotic markers, allergies, costs to LEDC farmers, loss of local diversity) • Sustainability links: energy dependence, trophic level of diet, feeding 12 billion people • Justified conclusion on the balance between productivity, efficiency and sustainability Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 321-328; AQA spec 3.5.1.2.	25