

A-level

ENVIRONMENTAL SCIENCE

Paper 2 - Generated practice paper Set 37

Mark scheme

120 marks: Questions 1-10 (95 marks) plus one essay from Question 11 (25 marks).

Marking guidance

- The number in brackets after each marking point is the mark it earns; the note above the points says how many are needed.
- Where a question asks for a fixed number of points (eg *two reasons*), credit only the first points given.
- Alternative correct wording and other relevant, accurate points are credited.
- For levels-of-response questions, first determine the level using the descriptors, then position the mark within the level using the indicative content.
- Calculations: credit correct working (method marks) even if the final answer is wrong; incorrect answers with no working score 0.

Textbook references: every question below cites the printed page numbers in R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018) from which the content is drawn, together with the AQA 7447 specification reference.

Question	Marking guidance	Marks
01.1	<i>1 mark for each point.</i> • Trees (hazel, oak) are cut to ground level every 3-30 years and allowed to regrow from the stump (1) • About 0.5 ha cut each year on a rotation (eg 8 years) so the wood is a patchwork of areas of different ages (1) • Each age has its own community: newly cut areas get high light for wildflowers and butterflies; older areas give dense cover for nesting birds / dormice (1) • Pollarding cuts above browsing height so deer / livestock cannot eat the regrowth (1)	4
01.2	<i>1 mark for each point.</i> • Wolves (top predator, exterminated by 1926) reduced deer populations and changed their behaviour (1) • Fewer young trees eaten so trees and riverside vegetation recovered (1) • More trees allowed beaver populations to increase; beaver dams increased wetland area (1) • Wetland species (birds, fish, invertebrates) became more common; overall biodiversity rose, though wolves cause problems for livestock farmers (1)	4
01.3	<i>1 mark for each point.</i> • Undesirable species (introduced predators / competitors, or natural succession species such as scrub, conifers in the New Forest heathland) reduce survival of the target species (1) • Removal restores suitable conditions (eg heathland for sand lizards and smooth snakes) or reduces predation / competition (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 47-49; AQA spec 3.1.3.3-3.1.3.4.	

Question	Marking guidance	Marks
02.1	<i>1 mark for each point.</i> • Less dead organic matter / leaf litter so less humus and slower soil formation; fewer nutrients returned (1) • Loss of canopy and litter exposes soil to rain splash and wind (1) • Decomposing roots no longer bind soil so erosion, gullyng and landslides increase (especially on slopes) (1) • Reduced organic matter and biota lower infiltration and water retention (1)	4
02.2	<i>1 mark for each point.</i> • Reduced interception and transpiration so less water returned to the atmosphere and less rainfall downwind (1) • Increased runoff: more flooding after rain, lower flows in dry periods; sedimentation of rivers (1) • Increased albedo of cleared land (reflects more sunlight) (1) • Reduced carbon sequestration and loss of the carbon reservoir (CO₂ released by burning / decay) contributing to climate change (1)	4
02.3	<i>Any 2 from the points below - 1 mark each, maximum 2.</i> • Fragmentation of remaining forest into isolated patches with non-viable populations / edge effects (1) • Loss of inter-species relationships (pollinators, seed dispersers) causing further declines (1) • Loss of genetic resources / potential medicines before they are discovered (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 368-369; AQA spec 3.5.3.3.	
03.1	<i>1 mark for each point.</i> • Island communities evolved from few colonising species in isolation, often without mammal predators or large herbivores (1) • So indigenous species lack adaptations (eg ground-nesting birds, flightlessness) to survive introduced cats, rats, pigs, goats (1)	2
03.2	<i>2 marks for each point (partial credit for a partly correct point).</i> • Competitor: grey squirrel (North America) out-competes red squirrel - digests acorns, larger, wins nest sites; rhododendron shades ground and releases toxins preventing native regrowth (2) • Predator: American mink (escaped fur farms) reduced water vole range; cane toads in Australia; cats / rats / pigs on islands eating ground-nesting birds; Nile perch in Lake Victoria ate cichlids to extinction (2) • Pathogen: squirrel pox virus carried by grey squirrels kills red squirrels; crayfish plague carried by signal crayfish kills white-clawed crayfish; Dutch elm disease, ash dieback, sudden oak death imported on plants / soil (2)	6
03.3	<i>1 mark for each point.</i> • Closely related introduced species (sika deer; domestic cats) cross-breed producing fertile hybrids (1) • The natural gene pool is changed / diluted with genes it would not naturally contain, so the pure indigenous species may disappear (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 24-25; AQA spec 3.1.2.2.	

Question	Marking guidance	Marks
04.1	<i>1 mark for each point.</i> • B: frost-free period (150 days) exceeds the 120 needed whereas A (110) is too short - late frosts would kill seedlings (1) • B is warmer / longer thermal growing season because south-facing slopes receive more insolation; valley floors are frost pockets where cold dense air collects (1) • But B has lower rainfall and a steeper slope (erosion / machinery difficulties) so irrigation / contour cultivation may be needed (1)	3
04.2	<i>Any 3 from the points below - 1 mark each, maximum 3.</i> • Greenhouses / polytunnels heated by solar gain or burning gas / paraffin (ventilated to prevent overheating) (1) • Woven plastic cloches / fleece over rows to retain warm air near the ground (1) • Burning oil / gas or using large fans in orchards to prevent frost damage to blossom (1) • Site selection (aspect, avoiding frost pockets); heated buildings and shelter for livestock (1)	3
04.3	<i>1 mark for each point.</i> • Mammals and birds use food energy to maintain a constant body temperature (thermoregulation) (1) • Reducing heat loss (shelter, heated buildings) means less food energy is used for heat and more is available for growth (1)	2
04.4	<i>1 mark for each point.</i> • Long days increase milk production and egg laying; poultry grow best with short days; sheep mate as days shorten in autumn (1) • Artificial lighting extends the growing season for greenhouse crops or simulates autumn in spring to give sheep a second mating / lambing season (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 306-309; AQA spec 3.5.1.1.	
05.1	<i>Up to 4 marks in total; award marks in proportion to the completeness of each point.</i> • Extensive: low inputs of energy, food and chemicals so low cost and low environmental impact; sustainable; but low yield per area and dependent on natural productivity (up to 2) • Intensive: control of all limiting factors gives very high yield and profit per area; but high inputs (fishmeal from wild fish, energy), disease and lice at high density, pollution from wastes, antibiotic / pesticide use, escapes (up to 2) • Judgement depends on species, market and site (up to 2)	4
05.2	<i>1 mark for each point.</i> • Adults with desirable characteristics (fast growth, shape, colour) stripped of eggs and milt which are mixed for fertilisation (1) • Eggs raised in aerated freshwater tanks, dead / diseased eggs removed daily; fry fed fishmeal pellets and moved through tanks as they grow (1) • After 12-18 months smolts transferred to seawater cages, fed pellets until harvest at 3-5 kg (1)	3

Question	Marking guidance	Marks
05.3	<i>1 mark for each point.</i> • About 10% of the world's mangroves have been cleared for coastal shrimp lagoons (up to 100 ha), removing nurseries and coastal protection (1) • Lagoons need water exchange to remove wastes and aeration to prevent deoxygenation as wastes decompose (1) • Eyestalk ablation stimulates breeding; high densities increase disease so antibiotics / pesticides used (1)	3
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 352-356; AQA spec 3.5.2.3.	
06.1	<i>Any 2 from the points below - 1 mark each, maximum 2.</i> • Hydrological cycle (1) • Carbon cycle (1) • Nitrogen cycle (1) • Atmospheric circulation (1) • Thermohaline circulation (1) • Photosynthesis (1)	2
06.2	<i>1 mark for each point.</i> • Enzymes reduce the activation energy of reactions (1) • So reactions such as photosynthesis, decomposition and nitrogen fixation do not need high temperatures (1)	2
06.3	<i>Any 1 from the points below - 1 mark each, maximum 1.</i> • Haber process (nitrogen fixation) (1) • Incineration of wastes (decomposition) (1) • Many manufacturing processes (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 373; AQA spec 3.6.2.	
07.1	<i>A correct final answer (with unit where asked) scores full marks. If the answer is wrong, award 1 mark for each correct step of working shown.</i> • $(60 \times 45) / 9$ (1) • $= 300$ (1)	2
07.2	<i>Any 4 from the points below - 1 mark each, maximum 4.</i> • Marked individuals mix freely / randomly with the rest of the population (1) • No births, deaths, immigration or emigration between samples (1) • Marking does not affect survival or behaviour (eg make them more visible to predators) (1) • Marks do not wear off / fall off (1) • All individuals have the same chance of being caught (traps do not attract or repel marked animals) (1)	4
07.3	<i>1 mark for each point.</i> • Fewer marked beetles would survive to be recaptured (1) • So the proportion marked in the second sample would be too low and the population would be overestimated (1)	2
07.4	<i>1 mark for each point.</i> • Enough time for marked beetles to mix with the population (1) • Not so long that births / deaths / migration or loss of marks change the population (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 401-402; AQA spec 3.7.2.1.	

Question	Marking guidance	Marks
08.1	<i>2 marks for each point (partial credit for a partly correct point).</i> - Tracking: continuous data on movements, territory size, migration routes and habitat use without disturbance, over large areas and long periods; acoustic tags in seawater; geolocators for small birds; more representative than occasional sightings (2) - DNA: identifies individuals, gene pool size / inbreeding and regional populations (tracing ivory, timber, fish); eDNA detects rare or hidden aquatic species (great crested newt, invasive carp) from shed cells without capture (2) - Image recognition: software identifies individuals from unique markings (tiger stripes, whale shark spots, fin damage) in camera-trap / drone / satellite images, giving population size, lifespan, social structure from large image sets quickly and objectively (2) - Acoustic: bat detectors, hydrophones and sonograms record species that are hard to see (bats, cetaceans, insects), giving presence, abundance and activity continuously, day and night (2)	8
08.2	<i>1 mark for each point.</i> - Data collected continuously / over large areas / at night rather than at a few times and places a researcher can visit (1) - Less disturbance so animals behave naturally; larger sample sizes; objective rather than subjective identification (1)	2
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 402-406; AQA spec 3.1.2.3.6.	
09.1	<i>1 mark for each point.</i> - Isolation makes colonisation difficult: only marine species, seabirds, plants with floating seeds or chance arrivals (tortoises on logs, wind-blown birds) reach them (1) - Whole taxa (terrestrial mammals, amphibians) may be absent so few original colonists (1) - Colonists evolve in isolation under different selection pressures to fill the empty niches, producing unique species (Galapagos, Madagascar lemurs, Hawaii) (1) - Endemics have small ranges and populations so are vulnerable to extinction (1)	4
09.2	<i>1 mark for each point.</i> - The birds evolved without mammal predators so nest on the ground / in burrows with no defence (1) - Rats eat eggs and chicks; numbers fell to very low levels (45 and 12 pairs) (1) - After eradication breeding success rose about 7-12 fold, confirming predation was the main limiting factor (1)	3
09.3	<i>1 mark for each point.</i> - Introduced mosquitoes carry avian malaria to which endemic birds have no resistance; birds survive only above about 1500 m where it is too cold for malaria (1) - Warming allows mosquitoes / malaria to spread to higher altitudes, shrinking the safe range (1)	2
09.4	<i>1 mark.</i> Control of developments and visitor numbers (biosecurity, quarantine of imports); protected areas; captive breeding of endemics (1)	1

Question	Marking guidance	Marks
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 64-65; AQA spec 3.1.2.3.4.	
10.1	Levels of response Level 3 (7-9 marks): Answer is comprehensive, relevant and logically structured. Shows detailed knowledge and understanding of the relevant scientific ideas and applies them accurately to the context. Evaluation/analysis is well supported by evidence and a clear judgement or conclusion is reached. Level 2 (4-6 marks): Answer is mostly relevant and shows some structure. Knowledge and understanding of relevant ideas is sound but application to the context or evaluation may be incomplete or lack detail. A judgement may be made but is not fully supported. Level 1 (1-3 marks): Answer is basic, with limited structure. Some relevant knowledge is shown but there are omissions or errors. Little application to the context and little or no evaluation. 0 marks: No relevant content. Indicative content (credit other relevant, accurate points) • Succession: pioneers, seres (hydrosere, lithosere, psammosere), modification of abiotic conditions, climax communities controlled by climate; deflected succession and plagioclimaxes • Application: deciding whether to manage (plagioclimax habitats need grazing, mowing, burning, coppicing, pollarding continued) or leave alone (climax habitats); habitat creation planning the stages; controlling scrub / secondary succession; understanding that new habitats change over time • Population dynamics: birth and death rates, density-dependent and independent factors, carrying capacity, sigmoid growth, predator-prey cycles, r- and k-selection, MSY • Application: setting sustainable harvest levels (fisheries, whaling, forestry); recognising k-selected species need strict protection and recover slowly; deciding when culling is needed (deer without wolves, introduced species) or when releases / captive breeding are needed (low breeding rate); predicting effects of reintroducing predators • Monitoring population size, distribution, survival rate and age structure to assess success; adaptation and range of tolerance guiding abiotic management • Evaluation: uncertainties (natural fluctuations, unknown carrying capacity); examples (Yellowstone wolves, Scottish deer, chalk grassland); judgement	9
	Model answer - Level 3 (8 marks)	
	Succession is the sequence by which pioneers colonise bare ground, modify the abiotic conditions and are replaced by later seres until a climax community controlled by climate is reached; where grazing, mowing, burning or coppicing halts it, a plagioclimax forms. This tells a conservation manager whether to intervene: chalk grassland, heathland and coppiced woodland are plagioclimaxes that disappear without continued management, whereas a climax habitat is best left alone, and habitat creation must plan the stages, control scrub and manage secondary succession.	

Question	Marking guidance	Marks
	Population dynamics - birth and death rates, density-dependent and independent factors, carrying capacity, sigmoid growth, predator-prey cycles and r- and k-selection - inform decisions of a different kind. Maximum sustainable yield sets harvest levels in fisheries, whaling and forestry; k-selected species that breed late and slowly need strict protection and recover slowly; culling may be justified where predators are absent, as with Scottish red deer without wolves, or where introduced species threaten natives, while the reintroduction of wolves to Yellowstone shows how restoring a predator changes vegetation through the food web. Monitoring population size, distribution, survival and age structure measures success, and knowledge of adaptation and ranges of tolerance guides how abiotic conditions are managed.	
	The limits are uncertainty: natural fluctuations, unknown carrying capacities and the difficulty of predicting how communities respond. Ecological understanding cannot remove that uncertainty, but without it conservation decisions - to manage or not, to cull or protect, how much to harvest - are guesses.	
	Weaker answer - Level 2 (5 marks)	
	Succession is when a habitat changes over time from pioneer species to a climax community. Conservationists need to understand this because some habitats like heathland will turn into woodland if they are not managed by grazing or burning. Population dynamics is about how populations grow and are limited by factors like food and predators. This helps conservation because you can work out how many animals can be taken without the population declining, and when a population needs protecting. Monitoring populations shows if conservation is working.	
	What makes the difference	
	• Level 2 explains both concepts correctly and links them to decisions, but generally; Level 3 uses the technical vocabulary (seres, plagioclimax, k-selection, MSY) with named cases (Yellowstone wolves, Scottish deer, chalk grassland).	
	• Level 3 covers more applications (habitat creation, culling introduced species, adaptation and tolerance) and evaluates the uncertainties.	
	• Structure: Level 3 gives each concept its own paragraph with the decisions it informs, then a conclusion.	
10.2	<i>1 mark.</i> • Any taxonomic group of organisms classified together by similarity (eg species, genus, family) (1)	1
	Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 72-85; AQA spec 3.1.3.	

Question	Marking guidance	Marks
11.1	Using tropical coral reefs as an example, discuss how the ecological features of a habitat determine both its importance and its vulnerability to human activities. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) • Ecological features: cnidarian polyps, cilia, nematocysts, zooxanthellae symbiosis, limestone skeletons building reefs, complex structure and food web; specific abiotic requirements (temperature, light, salinity, low turbidity) • Importance: quarter of marine species, fisheries, medicines, carbon sink, erosion protection, tourism • Vulnerability: narrow abiotic tolerance means sensitivity to warming, acidification, sedimentation, nutrient pollution, oil, bleaching; slow growth so slow recovery; physical damage; overfishing and food-web effects (crown of thorns); introduced species; dependence on mangroves and seagrass • Conservation: MPAs, No Take Zones, fishing controls, tourism management, protection of linked habitats, CITES • Comparison with other habitats (rainforests, mangroves, Antarctica) where relevant • Evaluation of local vs global threats and justified conclusion Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 57-62; AQA spec 3.1.2.3.4.	25

Question	Marking guidance	Marks
11.2	Discuss the causes and consequences of deforestation and evaluate the extent to which forests can be exploited sustainably. Levels of response Level 5 (21-25 marks): Comprehensive and detailed knowledge and understanding drawn from across the specification, applied accurately to the question. Argument is coherent, well-structured and supported by a wide range of relevant, well-chosen examples and quantitative detail where appropriate. Interconnections between topics are made explicitly and a clear, justified conclusion is reached. Level 4 (16-20 marks): Detailed knowledge and understanding with good application to the question. Argument is logically organised with relevant examples; some links between different areas of the specification are made. Conclusions are drawn but may be less fully justified. Level 3 (11-15 marks): Sound knowledge and understanding, mostly applied to the question. Some structure and relevant examples, but coverage may be uneven or descriptive rather than analytical. Limited linking of topics. Level 2 (6-10 marks): Basic knowledge and understanding with partial relevance. Answer is largely descriptive with few examples and limited structure; some errors or omissions. Level 1 (1-5 marks): Limited, fragmented knowledge with little relevance to the question. Little structure; examples absent or inappropriate. 0 marks: No relevant content. Indicative content (students are not expected to cover all of these; credit other relevant material) • Importance: resources (timber, fuel, food, fibres, medicines) and ecosystem services (atmospheric regulation, carbon store, hydrological regulation, climate, soil conservation, habitat, recreation) • Causes: agriculture (subsistence and cash crops), mineral extraction, reservoirs, urbanisation, transport, unsustainable timber, fuel and fodder exploitation above MSY; trends by region • Consequences: loss of resources, biodiversity and species, fragmentation; hydrological changes; soil impacts; climate impacts (albedo, carbon, rainfall) • Productivity vs biodiversity: non-indigenous species, monocultures, close planting, simple age structure, clear-felling • Sustainable management: MSY harvest rates, mixed indigenous species, mixed age structure, selective logging, traditional coppicing / pollarding, FSC certification, community involvement, protected areas, debt-for-nature, afforestation programmes (China); Rio Bravo case study • Evaluation: economics of slow-growing hardwoods, enforcement, demand from MEDCs, ITTO failures; judgement Source: R. Genn, Environmental Science A level AQA (Insight & Perspective, 2018), pp. 50-57, 360-369, 386-387; AQA spec 3.5.3, 3.1.2.3.4.	25